

University of Groningen

Iron chelators inhibit amyloid-beta-induced production of lipocalin 2 in cultured astrocytes

Dekens, Doortje W; De Deyn, Peter P; Sap, Friederike; Eisel, Ulrich L M; Naudé, Petrus J W

Published in:
Neurochemistry International

DOI:
[10.1016/j.neuint.2019.104607](https://doi.org/10.1016/j.neuint.2019.104607)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Dekens, D. W., De Deyn, P. P., Sap, F., Eisel, U. L. M., & Naudé, P. J. W. (2020). Iron chelators inhibit amyloid-beta-induced production of lipocalin 2 in cultured astrocytes. *Neurochemistry International*, 132, [104607]. <https://doi.org/10.1016/j.neuint.2019.104607>

Copyright


Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Iron chelators inhibit amyloid- β -induced production of lipocalin 2 in cultured astrocytes

Doortje W. Dekens^{a,b}, Peter P. De Deyn^{a,c}, Friederike Sap^b, Ulrich L.M. Eisel^{b,d,1},
Petrus J.W. Naudé^{a,b,*,1}

^a Department of Neurology and Alzheimer Center Groningen, University Medical Center Groningen, University of Groningen, Hanzeplein 1, 9713 GZ, Groningen, the Netherlands

^b Department of Molecular Neurobiology, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, Nijenborgh 7, 9747 AG, Groningen, the Netherlands

^c Laboratory of Neurochemistry and Behavior, Biobank, Institute Born-Bunge, University of Antwerp, Universiteitsplein 1, BE-2610, Antwerp, Belgium

^d University Center of Psychiatry & Interdisciplinary Center of Psychopathology of Emotion Regulation, University Medical Center Groningen, University of Groningen, Hanzeplein 1, 9713 GZ, Groningen, the Netherlands

ARTICLE INFO

Keywords:

Neutrophil gelatinase-associated lipocalin (NGAL)
Iron metabolism
Neuroinflammation
Ferritin
Deferoxamine
Deferiprone

ABSTRACT

Lipocalin 2 (Lcn2) has been implicated to play a role in various neurodegenerative diseases, and normalizing its overexpression may be of therapeutic potential. Iron chelators were found to reduce Lcn2 levels in certain animal models of CNS injury. Focusing on Alzheimer's disease (AD), we found that the iron chelators deferoxamine and deferiprone inhibited amyloid- β (A β)-induced Lcn2 production in cultured primary astrocytes. Accordingly, A β -exposure increased astrocytic ferritin production, indicating the possibility that A β induces iron accumulation in astrocytes. This effect was not significantly modulated by Lcn2. Known neuroprotective effects of iron chelators may rely in part on normalization of Lcn2 levels.

1. Introduction

Lipocalin 2 (Lcn2, also known as neutrophil gelatinase-associated lipocalin (NGAL)) is involved in several physiological processes including inflammation, iron metabolism, cell death and cell survival. Increased Lcn2 levels were found in the central nervous system (CNS) of patients with neurodegenerative diseases, including Alzheimer's disease (AD) and Parkinson's disease. Moreover, mechanistic studies showed that Lcn2 may contribute to their pathophysiology (Kim et al., 2016; Mesquita et al., 2014; Naudé et al., 2012). Regarding AD, it was shown that amyloid- β (A β) induces Lcn2 production in cultured primary astrocytes, and that Lcn2 sensitizes primary neurons and astrocytes to A β -induced cell death (Mesquita et al., 2014; Naudé et al., 2012). Astrocytes appear to be the major producers of Lcn2 in the brain (Kim et al., 2016; Mesquita et al., 2014). The reported neurotoxic effects of Lcn2 indicate that inhibition of Lcn2 overexpression may be a promising therapeutic strategy for different CNS conditions.

Iron chelators such as deferoxamine and deferiprone have been shown to exert neuroprotective effects (Belaidi and Bush, 2016), maybe

partly via reducing the brain iron accumulation that characterizes many CNS conditions. Interestingly, deferoxamine was found to decrease Lcn2 levels in certain animal models of CNS injury (Dong et al., 2013; Zhao et al., 2016). However, it is still unknown if iron chelators may reduce Lcn2 production in the context of AD.

The aim of this study was to explore (1) whether the iron chelators deferoxamine and deferiprone are able to inhibit A β ₁₋₄₂-induced Lcn2 production in cultured astrocytes, and (2) whether A β may affect astrocytic iron metabolism, and the potential effect of Lcn2 hereon by comparing A β -treated wild-type (WT) and Lcn2 knock-out (Lcn2 KO) astrocytes.

2. Methods

Primary astrocytes were obtained from newborn (P0-P3) WT and Lcn2 KO (Berger et al., 2006) mouse pups, according to a protocol approved by the local and national animal ethics committees (DEC6659A and CCD-AVD105002016630). Astrocytes were cultured as

* Corresponding author. Department of Molecular Neurobiology, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, Nijenborgh 7, 9747 AG, Groningen, the Netherlands.

E-mail addresses: d.w.dekens@rug.nl (D.W. Dekens), p.p.de.deyn@umcg.nl (P.P. De Deyn), friederike.sap@web.de (F. Sap), u.l.m.eisel@rug.nl (U.L.M. Eisel), p.j.w.naude@umcg.nl (P.J.W. Naudé).

¹ Shared last author.

<https://doi.org/10.1016/j.neuint.2019.104607>

Received 9 July 2019; Received in revised form 3 November 2019; Accepted 21 November 2019

Available online 21 November 2019

0197-0186/© 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

described previously (Naudé et al., 2012). Six hours before treatment, medium was exchanged for medium containing 5% fetal bovine serum. Human recombinant A β ₁₋₄₂ (A-1002-1, rPeptide) was prepared as described previously (Granic et al., 2010). Before use, the A β stock solution (100 μ M in DMEM) was allowed to oligomerize for 6 h at 4 °C (Ahmed et al., 2010). The oligomeric state of A β was confirmed with non-reducing SDS-PAGE Western blotting. Astrocytes were treated with 1 μ M A β , 10 ng/ml interleukin 1 beta (IL-1 β) or 100 ng/ml lipopolysaccharide (LPS), or were co-treated with 1 μ M A β and either 0–150 μ M deferoxamine (D9533, Sigma-Aldrich), 0–500 μ M deferiprone (S4067, SelleckChem), 0–200 μ M bathocuproine disulfonic acid (B1125, Sigma-Aldrich) or 0–25 μ M tetrathiomolybdate (323446, Sigma-Aldrich) for the indicated periods of time. Collection of proteins and Western blotting were performed as described previously (Naudé et al., 2012). Primary antibodies used include anti-Lcn2 (ab63929, Abcam, 1:1000), anti-ferritin (ab75973, Abcam, 1:1000) and anti-actin (691002, MP Biomedicals, 1:500,000). All treatments were performed three times in duplicate or triplicate.

3. Results

Firstly, it was confirmed that A β ₁₋₄₂ induced Lcn2 production and secretion by astrocytes (Fig. 1a and b). Intracellular Lcn2 levels peaked 36 h after A β ₁₋₄₂ treatment ($p < 0.0001$). This corresponds to kinetics of Lcn2-induction upon TNF- α , IL-1 β and LPS-stimulation ((Naudé et al., 2012) and Suppl. Fig. 1a and b). Secondly, deferoxamine significantly reduced A β -induced Lcn2 production, after 36 h co-incubation ($p < 0.0001$, Fig. 1c and d). The inhibitory effect of deferoxamine on A β -induced Lcn2 production was confirmed with another iron chelator; deferiprone (Suppl. Fig. 1c and d).

The finding that the Lcn2-inducing effects of A β can be suppressed by iron chelators, points to the possibilities that (1) A β may provoke iron accumulation in astrocytes, and (2) this disturbance in iron metabolism correlates with the induction of Lcn2 expression. As shown in Fig. 1e, A β indeed increased ferritin protein levels in WT and Lcn2 KO astrocytes ($p < 0.05$ at 36 h, compared to control), indicating an increase in astrocytic iron accumulation upon A β exposure, independent of endogenous Lcn2 production. Although increased astrocytic iron levels might be an important co-factor in the induction of Lcn2, it appeared that iron alone is not sufficient to induce Lcn2 upregulation (Suppl. Fig. 1e).

4. Discussion

Results from this study suggest that iron chelators are potent inhibitors of A β -induced Lcn2 production in astrocytes, which may contribute to their reported neuroprotective effects. Interestingly, it was proposed that iron-loaded deferiprone (unlike deferoxamine) may bind to Lcn2, after which the iron-deferiprone-Lcn2 complex is excreted from the body (Zughaier et al., 2014). Certain iron chelators, i.e. deferiprone, might thus not only affect Lcn2 production but also its removal from the body.

The modulation of A β -induced Lcn2 production by iron chelators further suggests that A β may act in part via increasing iron levels in astrocytes (also illustrated in Fig. 1f). This is supported by our result showing that A β causes an increase in astrocytic ferritin levels. This is the first study to our best knowledge that indicates iron accumulation in astrocytes upon direct A β -stimulation. This is in accordance with the previously reported A β -induced iron accumulation in microglia ((McCarthy et al., 2018) and Suppl. Fig. 1f) and a neuronal cell line (Wan et al., 2011). Future experiments are required to confirm the finding in astrocytes, including direct read-outs of iron accumulation. Moreover, further investigations are needed to elucidate the role of disturbed iron metabolism in A β -induced astrocyte activation and Lcn2 production. Namely, while the current results may suggest a potential involvement of disturbed iron metabolism in A β -induced Lcn2


production, it is possible that iron is not essential, and that other factors and pathways are also involved. In addition, more work is required to determine whether deferoxamine and deferiprone inhibit A β -induced Lcn2 production by chelating iron, or also via alternative pathways. For example, it is known that deferoxamine and deferiprone are not entirely specific for iron but are also able to chelate copper, suggesting that their effects might partly rely on chelation of copper. Interestingly, we found that A β -induced Lcn2 production can be modulated by certain copper chelators: while bathocuproine disulfonic acid (a membrane impermeable copper chelator) did not affect Lcn2 protein levels, tetrathiomolybdate (a membrane permeable copper chelator) was shown to significantly reduce intracellular Lcn2 levels (Suppl. Fig. 1g–j). The observed inhibitory effect of tetrathiomolybdate on Lcn2 production may be explained by a previous finding from Spisni et al. (2009), showing that copper treatment results in increased Lcn2 secretion from cultured neurons. It thus appears that deferoxamine and deferiprone are not the only chelators that can affect Lcn2 production, and that possibly different biometals might influence Lcn2 production.

Finally, although Lcn2 is known to play a role in iron regulation and is able to mediate both cellular iron import and export, no effect of Lcn2 was found on A β -induced ferritin protein production in astrocytes when comparing A β -treated WT and Lcn2 KO astrocytes (despite a previously reported effect of Lcn2 on ferritin mRNA expression (Mesquita et al., 2014)). This finding indicates that Lcn2 may not significantly affect A β -mediated changes in iron metabolism in astrocyte cultures. Interestingly however, Lcn2 appeared to significantly aggravate brain iron accumulation in mouse models of hemorrhagic stroke and AD (Dekens et al., 2018; Ni et al., 2015). In a mouse model of AD, Lcn2 promoted iron accumulation in A β plaques and neuronal layers of the hippocampus (Dekens et al., 2018). However, the exact cellular localization of accumulated iron remains to be determined in more detail. For instance, previous studies suggested that also microglia tend to accumulate high levels of iron under inflammatory conditions (Holland et al., 2018; McCarthy et al., 2018; Thomsen et al., 2015; Urrutia et al., 2013). As such, iron accumulation in AD (which is in part mediated by Lcn2) might occur mostly in specific cell types and structures, including plaques, neurons and microglia. Astrocytes might be less prone to (Lcn2-mediated) iron accumulation (Rathore et al., 2012; Urrutia et al., 2013), which would be in line with the similar ferritin levels in A β -treated WT vs. Lcn2 KO astrocyte cultures that were found here. It should be emphasized that the current study is a short report, warranting further investigation of Lcn2-mediated brain iron regulation in various other experimental conditions. For example, effects of Lcn2 on astrocytic iron metabolism might surface when more ferric and/or ferrous iron would be supplemented to the cell culture medium. Moreover, it is important to recognize that brain iron metabolism depends on intricate communication between different brain cell types (You et al., 2017). Therefore, it would be of great relevance to study iron metabolism in co-/slice-cultures and animals, rather than in single cell type cultures.

Iron chelators are promising therapeutic possibilities for various neurodegenerative diseases and CNS conditions. Their beneficial effects might depend in part on normalization of Lcn2 protein levels.

Ethical approval

All applicable international, national, and institutional guidelines for the care and use of animals were followed. All procedures performed in studies involving animals were in accordance with EU regulations (EU Directive, 2010/63/EU for animal experiments), and were approved by the local (University of Groningen, DEC6659A) and Dutch national (CCD-AVD105002016630) animal ethics committees. This work does not contain any studies with human participants performed by any of the authors.



(caption on next page)

Fig. 1. The iron chelator deferoxamine blocks A β -induced astrocytic Lcn2 production, and indicates that A β induces a disturbance in astrocytic iron metabolism. **a-b** Intracellular (a, controlled for actin) and secreted (b) Lcn2 protein levels in primary WT astrocytes treated with 1 μ M A β for 0–48 h. **c-d** Intracellular (c, controlled for actin) and secreted (d) Lcn2 protein levels in primary WT astrocytes treated with 1 μ M A β and 0–150 μ M deferoxamine (DFO) for 36 h. **e** Intracellular ferritin protein levels (controlled for actin) in primary WT and Lcn2 KO astrocytes treated with 1 μ M A β for 0–72 h. **f** Proposed connection between A β , iron and Lcn2, with uncertain points indicated in grey. Bars depict the mean and standard error of the mean (SEM). Representative blots are shown below graphs. Tested with one-way ANOVA with Dunnett's multiple comparison post-hoc test to compare conditions to their respective control condition. * p < 0.05, ** p < 0.01 and *** p < 0.0001 compared to the respective control conditions.

Funding

This study was funded by grants from the Internationale Stichting Alzheimer Onderzoek (ISAO#06511 to ULME and PPDD), ZonMW Deltaplan Dementie Memorabel (733050815 and 733050501 to PJWN, PPDD and ULME), Alzheimer Research Center Groningen, IAP Network P7/16 funding of the Belgian Federal Science Policy Office, Methusalem excellence grant of the Flemish Government and University Research Fund of the University of Antwerp (to PPDD), NeuroSearch Antwerp (to PPDD and PJWN), Alzheimer Nederland (WE. 13-2015-19 to PJWN), The Research School of Behavioural and Cognitive Neurosciences (to DWD), and Stichting Hadders-De Cock (2017-30 to DWD).

Declarations of competing interest

None.

Acknowledgments

We thank Wanda Douwenga, Jan Keijser, Kunja Slopsema, Wendy Kaspers, Roelie Veenstra-Wiegman, Benjamin Otten, Robin Kremer, Harm Ruesink and Margo Jansen for their excellent technical assistance.

Abbreviations

A β	amyloid- β
AD	Alzheimer's disease
Def	deferiprone
DFO	deferoxamine
DMEM	Dulbecco's Modified Eagle Medium
IL-1 β	interleukin 1 beta
Lcn2	lipocalin 2
Lcn2 KO	lipocalin 2 knock-out
LPS	lipopolysaccharide
NGAL	neutrophil gelatinase-associated lipocalin
WT	wild type

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.neuint.2019.104607>.

References

Ahmed, M., Davis, J., Aucoin, D., Sato, T., Ahuja, S., Aimoto, S., Elliott, J.I., Van Nostrand, W.E., Smith, S.O., 2010. Structural conversion of neurotoxic amyloid- β (1–42) oligomers to fibrils. *Nat. Struct. Mol. Biol.* 17, 561–567. <https://doi.org/10.1038/nsmb.1799>.

Belaïdi, A.A., Bush, A.I., 2016. Iron neurochemistry in Alzheimer's disease and Parkinson's disease: targets for therapeutics. *J. Neurochem.* 139 (Suppl. 1), 179–197. <https://doi.org/10.1111/jnc.13425>.

Berger, T., Togawa, A., Duncan, G.S., Elia, A.J., You-Ten, A., Wakeham, A., Fong, H.E.H.,

Cheung, C.C., Mak, T.W., 2006. Lipocalin 2-deficient mice exhibit increased sensitivity to *Escherichia coli* infection but not to ischemia-reperfusion injury. *Proc. Natl. Acad. Sci. U.S.A.* 103, 1834–1839. <https://doi.org/10.1073/pnas.0510847103>.

Dekens, D.W., Naudé, P.J.W., Keijser, J.N., Boerema, A.S., De Deyn, P.P., Eisel, U.L.M., 2018. Lipocalin 2 contributes to brain iron dysregulation but does not affect cognition, plaque load, and glial activation in the J20 Alzheimer mouse model. *J. Neuroinflammation* 15, 330. <https://doi.org/10.1186/s12974-018-1372-5>.

Dong, M., Xi, G., Keep, R.F., Hua, Y., 2013. Role of iron in brain lipocalin 2 upregulation after intracerebral hemorrhage in rats. *Brain Res.* 1505, 86–92. <https://doi.org/10.1016/j.brainres.2013.02.008>.

Granic, I., Masman, M.F., Kees Mulder, C., Nijholt, I.M., Naude, P.J.W., de Haan, A., Borbély, E., Penke, B., Luiten, P.G.M., Eisel, U.L.M., 2010. LPYFDa neutralizes amyloid-beta-induced memory impairment and toxicity. *J. Alzheimers Dis.* JAD 19, 991–1005. <https://doi.org/10.3233/JAD-2010-1297>.

Holland, R., McIntosh, A.L., Finucane, O.M., Mela, V., Rubio-Araiz, A., Timmons, G., McCarthy, S.A., Gun'ko, Y.K., Lynch, M.A., 2018. Inflammatory microglia are glycolytic and iron retentive and typify the microglia in APP/PS1 mice. *Brain Behav. Immun.* 68, 183–196. <https://doi.org/10.1016/j.bbi.2017.10.017>.

Kim, B.-W., Jeong, K.H., Kim, Jae-Hong, Jin, M., Kim, Jong-Heon, Lee, M.-G., Choi, D.-K., Won, S.-Y., McLean, C., Jeon, M.-T., Lee, H.-W., Kim, S.R., Suk, K., 2016. Pathogenic upregulation of glial lipocalin-2 in the Parkinsonian dopaminergic system. *J. Neurosci. Off. J. Soc. Neurosci.* 36, 5608–5622. <https://doi.org/10.1523/JNEUROSCI.4261-15.2016>.

McCarthy, R.C., Sosa, J.C., Gardeck, A.M., Baez, A.S., Lee, C.-H., Wessling-Resnick, M., 2018. Inflammation-induced iron transport and metabolism by brain microglia. *J. Biol. Chem.* 293, 7853–7863. <https://doi.org/10.1074/jbc.RA118.001949>.

Mesquita, S.D., Ferreira, A.C., Falcao, A.M., Sousa, J.C., Oliveira, T.G., Correia-Neves, M., Sousa, N., Marques, F., Palha, J.A., 2014. Lipocalin 2 modulates the cellular response to amyloid beta. *Cell Death Differ.* 21, 1588–1599. <https://doi.org/10.1038/cdd.2014.68>.

Naudé, P.J.W., Nyakas, C., Eiden, L.E., Ait-Ali, D., van der Heide, R., Engelborghs, S., Luiten, P.G.M., De Deyn, P.P., den Boer, J.A., Eisel, U.L.M., 2012. Lipocalin 2: novel component of proinflammatory signaling in Alzheimer's disease. *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 26, 2811–2823. <https://doi.org/10.1096/fj.11-202457>.

Ni, W., Zheng, M., Xi, G., Keep, R.F., Hua, Y., 2015. Role of lipocalin-2 in brain injury after intracerebral hemorrhage. *J. Cereb. Blood Flow Metab.* 35, 1454–1461. <https://doi.org/10.1038/jcbfm.2015.52>.

Rathore, K.L., Redensek, A., David, S., 2012. Iron homeostasis in astrocytes and microglia is differentially regulated by TNF- α and TGF- β 1. *Glia* 60, 738–750. <https://doi.org/10.1002/glia.22303>.

Spisni, E., Valerij, M.C., Manerba, M., Strillacci, A., Polazzi, E., Mattia, T., Griffoni, C., Tomasi, V., 2009. Effect of copper on extracellular levels of key pro-inflammatory molecules in hypothalamic GN11 and primary neurons. *Neurotoxicology (Little Rock)* 30 (4), 605–612. <https://doi.org/10.1016/j.neuro.2009.03.005>.

Thomsen, M.S., Andersen, M.V., Christoffersen, P.R., Jensen, M.D., Lichota, J., Moos, T., 2015. Neurodegeneration with inflammation is accompanied by accumulation of iron and ferritin in microglia and neurons. *Neurobiol. Dis.* 81, 108–118. <https://doi.org/10.1016/j.nbd.2015.03.013>.

Urrutia, P., Aguirre, P., Esparza, A., Tapia, V., Mena, N.P., Arredondo, M., González-Billault, C., Núñez, M.T., 2013. Inflammation alters the expression of DMT1, FPN1 and hepcidin, and it causes iron accumulation in central nervous system cells. *J. Neurochem.* 126, 541–549. <https://doi.org/10.1111/jnc.12244>.

Wan, L., Nie, G., Zhang, J., Luo, Y., Zhang, P., Zhang, Z., Zhao, B., 2011. β -Amyloid peptide increases levels of iron content and oxidative stress in human cell and *Caenorhabditis elegans* models of Alzheimer disease. *Free Radic. Biol. Med.* 50, 122–129. <https://doi.org/10.1016/j.freeradbiomed.2010.10.707>.

You, L.-H., Yan, C.-Z., Zheng, B.-J., Ci, Y.-Z., Chang, S.-Y., Yu, P., Gao, G.-F., Li, H.-Y., Dong, T.-Y., Chang, Y.-Z., 2017. Astrocyte hepcidin is a key factor in LPS-induced neuronal apoptosis. *Cell Death Dis.* 8, e2676. <https://doi.org/10.1038/cddis.2017.93>.

Zhao, J., Xi, G., Wu, G., Keep, R.F., Hua, Y., 2016. Deferoxamine attenuated the upregulation of lipocalin-2 induced by traumatic brain injury in rats. *Acta Neurochir. Suppl.* 121, 291–294. https://doi.org/10.1007/978-3-319-18497-5_50.

Zughaier, S.M., Stauffer, B.B., McCarty, N.A., 2014. Inflammation and ER stress down-regulate BDH2 expression and dysregulate intracellular iron in macrophages. *J. Immunol. Res.* 2014. <https://doi.org/10.1155/2014/140728>.