

University of Groningen

The SLC25 Carrier Family

Kunji, Edmund R S; King, Martin S; Ruprecht, Jonathan J; Thangaratnarajah, Chancievan

Published in:
 Physiology

DOI:
[10.1152/physiol.00009.2020](https://doi.org/10.1152/physiol.00009.2020)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kunji, E. R. S., King, M. S., Ruprecht, J. J., & Thangaratnarajah, C. (2020). The SLC25 Carrier Family: Important Transport Proteins in Mitochondrial Physiology and Pathology. *Physiology*, 35(5), 302-327. <https://doi.org/10.1152/physiol.00009.2020>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The SLC25 Carrier Family: Important Transport Proteins in Mitochondrial Physiology and Pathology

Members of the mitochondrial carrier family (SLC25) transport a variety of compounds across the inner membrane of mitochondria. These transport steps provide building blocks for the cell and link the pathways of the mitochondrial matrix and cytosol. An increasing number of diseases and pathologies has been associated with their dysfunction. In this review, the molecular basis of these diseases is explained based on our current understanding of their transport mechanism.

mitochondrial physiology; mitochondrial disease; impaired transport mechanism; pathological mutations; bioenergetics

Introduction

The inner membrane of mitochondria is highly impermeable to molecules and ions, which is a key property required for energy conversion in oxidative phosphorylation. Therefore, a large number of transport proteins and channels are required to transport molecules and ions across this membrane to link cytosolic and mitochondrial metabolism, and to provide compounds for building and maintenance of the mitochondrion and the cell. In fact, all major food groups pass through the mitochondrion as part of central metabolism, including the degradation products of fats, sugars, and proteins, as well as nucleotides, vitamins, and inorganic ions (FIGURE 1). Most of the transport steps are carried out by members of the mitochondrial carrier family (SLC25), the topic of this review, but there are also other transporter families, such as the mitochondrial pyruvate carrier (SLC54) (19, 61), sideroflexins (SLC56) (87, 127, 170), and mitochondrial ABC transporters (98, 148). Mitochondrial carriers provide key transport steps in a variety of metabolic pathways, such as the oxidation of degradation products of fats and sugars, the degradation, synthesis, and interconversion of amino acids, and the synthesis of iron sulfur clusters and heme, but also in ion homeostasis, mitochondrial macromolecular synthesis, heat production, mitochondrial dynamics, signaling, cellular differentiation, development, and cell death (89, 123). The genes of 53 mitochondrial carriers of the SLC25 family have been identified in the human genome (89, 123), based on their shared sequence features (137, 147) and structural properties (128, 141, 143, 144). Most carriers are found in the inner membrane of the mitochondrion, but an adenine nucleotide

Edmund R. S. Kunji,¹
Martin S. King,¹
Jonathan J. Ruprecht,¹ and
Chancievan Thangaratnarajah²
¹Medical Research Council Mitochondrial Biology Unit,
University of Cambridge, Cambridge, United Kingdom; and
²Groningen Biomolecular Sciences and Biotechnology Institute,
Membrane Enzymology, University of Groningen, Groningen,
The Netherlands
ek@mrc-mbu.cam.ac.uk

transporter (SLC25A17) has been located in the peroxisome, where it supplies ATP for energy-requiring processes (161). Moreover, two highly divergent carriers, SLC25A46 and SLC25A50 (MTCH2), have been localized to the outer membrane of the mitochondrion (2, 172), where they might have some involvement in mitochondrial dynamics and apoptosis, respectively, but no transport function has been assigned to them.

Mitochondrial carriers are functional as monomers (10–12, 32, 91). The only confirmed exception is the mitochondrial aspartate/glutamate carrier, which is a structural dimer through the interactions of the NH₂-terminal calcium regulatory domains, but the carrier domains, which are involved in transport, function separately (see further below) (157). Recently, a claim was made for the bovine mitochondrial ADP/ATP carrier being monomeric and dimeric using native mass spectrometry of intact membranes (28), but the reported mass deviates significantly from those observed by others (7, 62, 153). To account for the difference, the carrier was proposed to have an unusually large number of modifications, which have not been observed in other studies (7, 153) or in the structure (128). Furthermore, the species observed by native mass spectrometry lacks three tightly bound cardiolipin molecules (28), which are observed consistently in other studies (10, 14, 96, 128, 141, 142), and thus does not represent the native ADP/ATP carrier.

Given the central role of mitochondrial carriers in cellular metabolism and physiology, it is not surprising that mutations in mitochondrial carriers have been associated with a large number of pathologies. Due to recent advances in sequencing technologies, this number is likely to rise rapidly,

as new links to disease are discovered frequently, including for mitochondrial carriers that have no assigned function yet. Although originally thought to be relatively rare, it is now clear that pathologies involving mitochondrial carriers may be among the most prevalent of all mitochondrial diseases. For example, citrin deficiency, which is caused by disease variants of the mitochondrial aspartate/glutamate carrier (SLC25A13), has a very high frequency in Far Eastern populations (146). Current estimates, based on pathogenic allele frequencies, are 1:17,000 in Japan and China, 1:9,000 in Taiwan, and 1:50,000 in Korea, but the disease has now also been discovered in other populations, making it panethnic (38, 122). With few exceptions most pathologies are inherited in an autosomal recessive manner. The phenotypic manifestation of these pathologies is highly variable, starting at different

ages, affecting various organs in different ways, and is dependent on the type of mutation, i.e., deletion, missense, nonsense, inversion, or splice-site mutation. The pathologies can be broadly divided into developmental, metabolic, and neuromuscular diseases. Understanding the molecular basis for these diseases is vital for diagnosis and prognosis of the disease, and for development of effective treatments and therapies.

In this review, we provide a comprehensive overview of the role of mitochondrial carriers in physiology and pathology. We describe all pathogenic missense mutations that have been identified to date in the context of new insights into the structural mechanism of these transport proteins. This analysis can be used to predict with better accuracy which mutations are likely to be pathological, resulting in a dysfunctional mitochondrial carrier.




FIGURE 1. The role of the human mitochondrial carrier family (SLC25) in metabolism and mitochondrial function
 Schematic representation of the mammalian mitochondrion. Shown in green are the electron transfer chain complexes. Bottom to top: glycerophosphate dehydrogenase, fatty acid-dehydrogenase-electron transfer flavoprotein, dihydroorotate dehydrogenase, and complex I to IV with cytochrome c. Shown in blue is the dimer of ATP synthase, and in red/orange the mitochondrial pyruvate carrier heterodimer (MPC). Shown in purple and yellow are unidentified and identified mitochondrial carriers, respectively. AAC1–4, ADP/ATP carriers; AGC1–2, aspartate/glutamate carriers; APC1–4, ATP-Mg/Pi carriers; BAC, basic amino acid carrier; CAC, carnitine-acylcarnitine carrier; CIC, citrate carrier; DIC, dicarboxylate carrier; GC1–2, glutamate carriers; GLYC, glycine carrier; MFRN1–2, mitoferrins; ODC, oxoadipate carrier; OGC, oxoglutarate carrier; ORC1–2, ornithine carriers; PIC, phosphate carrier; SAMC, S-adenosylmethionine carrier; TPC, thiamine pyrophosphate carrier; UCP1, uncoupling protein; UCP2, uncoupling-like protein 2; CS, cytosol; IS, intermembrane space; OM, outer membrane; IM, inner membrane; MM, mitochondrial matrix; Q, ubiquinone; Pi, phosphate; ALA, aminolevulinic acid.

The data presented in this review may also help to identify new pathological variants and associated mitochondrial diseases.

The Role of Mitochondrial Carriers in Physiology and Pathology

With its 53 members, the mitochondrial carrier family SLC25 is the largest solute transporter family in humans. Here, we first briefly review the function of the known human mitochondrial carriers in cellular physiology and pathology, but for more detailed information the reader is referred to Ref. 124.

Nucleotide Transport

Mitochondria require nucleotides for a host of important functions, such as the synthesis of ATP, mitochondrial RNA and DNA synthesis, enzymatic reactions, and regulation. The best-known member of the SLC25 family is the mitochondrial ADP/ATP carrier, also called adenine nucleotide translocase or translocator (ANT). The carrier imports ADP into the mitochondrial matrix, where it is converted to ATP by ATP synthase, and exports

the newly synthesized ATP to the cytosol, where it fuels the metabolic energy-requiring processes that are vital for cell survival (7, 83, 90, 144). There are four different paralogs in humans, AAC1, AAC2, AAC3, and AAC4 (SLC25A4, SLC25A5, SLC25A6, and SLC25A31, respectively), which are expressed in a tissue-dependent manner (FIGURE 1A) (31, 40, 88, 115). The mitochondrial ADP/ATP carriers carry out equimolar exchange of ADP and ATP, and thus do not alter the total adenine nucleotide pool in the mitochondrial matrix. The mitochondrial ADP/ATP carrier has been extensively characterized structurally (FIGURE 2A) (128, 141–144). In the absence of adenine nucleotides and in the presence of fatty acids, the mitochondrial ADP/ATP carrier may function as an uncoupling protein, transporting protons (13, 20), but a molecular mechanism is lacking.

The human mitochondrial ADP/ATP carrier AAC1 (SLC25A4) has three associated pathologies, which are quite different in nature: 1) late-onset dominant progressive external ophthalmoplegia with mitochondrial DNA deletions (OMIM 609283) (77), 2) recessive mitochondrial DNA depletion syndrome (OMIM 615418) (53, 125), and 3) early




FIGURE 2. Structures of the mitochondrial ADP/ATP carrier and the calcium-regulated ATP-Mg/phosphate carrier and aspartate/glutamate carrier
Structures of three different mitochondrial carriers, based on PDB entries 1OKC (128) and 4C9Q (141) for the carrier domains and 4P5W (157) and 4ZCU (59) for the calcium-regulatory domains. A: the mitochondrial ADP/ATP carrier. B: the mitochondrial ATP-Mg/Pi carrier consists of three domains: 1) NH₂-terminal calcium-regulatory domain with four EF-hands (EF1–4), each binding calcium, 2) amphipathic helix, and 3) COOH-terminal carrier domain. C: the aspartate/glutamate carrier also has a three-domain structure, but with a different order: 1) NH₂-terminal calcium-regulatory domain, 2) carrier domain, and 3) COOH-terminal amphipathic helix. The NH₂-terminal domain has eight EF-hand folds, but only EF-hand 2 is capable of binding calcium, which together with EF-hands 1 and 3 forms a calcium-responsive mobile unit. EF-hands 4–8 have evolved to form a static dimerization interface. The structures are shown in a cartoon representation, colored from the NH₂-terminus in blue to the COOH-terminus in red. Also shown are the canonical substrates as well as calcium ions (green), magnesium ion (chartreuse), and protons (white) in sphere representations. IS, intermembrane space; OM, outer membrane; IM, inner membrane; MM, mitochondrial matrix.

onset dominant de novo mutations leading to mitochondrial disease, varying from lethal to mild severity (OMIM 103220) (81, 158). The complexity of the genetics underlying these three diseases has not been fully explained. As will be shown below, the carrier has all the features required for its transporter function, so the dominant inheritance cannot be explained by dimerization, as was thought previously (91). This notion is supported further by the fact that the early onset dominant disease mutations involve key functional residues on the inside of the carrier, which cannot be involved in dimerization (158). Every day, the ADP/ATP carriers transport approximately our own body weight in ADP and ATP across the inner membrane of mitochondria, meaning that the dominant inheritance could well be explained by haplo-insufficiency. In agreement, the early onset de novo dominant mutations affect highly conserved residues that are central to the transport mechanism (p.Lys33Gln, p.Arg80His, and p.Arg235Gly), whereas late-onset domain mutations affect non-conserved residues on the periphery of the protein without an obvious role in the mechanism (e.g., p.Ala90Asp, p.Asp104Gly, Leu98Pro, and Ala114Pro). Not fitting this genetic explanation are the recessive mutations, which lead to mitochondrial DNA deletions and mitochondrial disease only in homozygotes, whereas the heterozygotes are unaffected (53), but they may give rise to biogenesis rather than functional issues (e.g., p.Ala123Asp, p.Arg263Pro). However, it would be impossible for humans to live without functional ADP/ATP carriers, since cells would be reliant on glycolysis and fermentation alone, providing low yields of ATP. The oxidative phosphorylation of sugars, fats, and amino acids in mitochondria is critically dependent on ADP/ATP carriers. This notion is supported by the fact that the powerful inhibitors carboxyatractyloside and bongkreic acid kill humans, showing that human life cannot sustain itself without mitochondrial ADP/ATP transport. There are three other paralogs of the mitochondrial ADP/ATP carrier, AAC2, AAC3, and AAC4, which are expressed at different levels and/or may be upregulated to compensate for the missing transport activity in homozygous recessive patients. In addition, there are the four paralogs of the ATP-Mg/Pi carrier and some vitamin transporters, such as the CoA transporter (SLC25A42) (51) and thiamine pyrophosphate carrier (SLC25A19) (39), which can also transport adenine nucleotides in addition to their key substrates. The complexity of adenine nucleotide transport might explain viability in the homozygous-recessive cases, and the variation in severity and onset of the pathogenicity in the heterozygous-dominant cases (81, 158). The mitochondrial ADP/ATP carrier can also transport the

deoxy forms of ADP and ATP, which are required for mitochondrial DNA replication, which could explain the role of disease variants in mitochondrial DNA deletions and maintenance, aside from the production of ATP. Finally, very little is known about the effect of these mutations on the biogenesis of the disease variants. Therefore, other genetic mechanisms, such as gain of function through aggregation, are also pursued as an explanation (100, 101). However, it is clear that this important issue is unresolved, requiring a more complete picture of adenine nucleotide transport in mitochondria.

The net import and export of adenine nucleotides is carried out by mitochondrial ATP-Mg/Pi carriers, which exchange phosphate for adenine nucleotides, coupled to magnesium or protons, in an electroneutral way (8, 36, 45). This transport activity allows the mitochondrion to respond to changes in energetic demand and to replenish adenine nucleotide pools after mitochondrial division and macromolecular synthesis (8, 26, 45). There are three human paralogs, APC1 (SLC25A24), APC2 (SLC25A23), and APC3 (SLC25A25), which are calcium-regulated (5, 45), and a fourth paralog, APC4 (SLC25A41), which is not (160) (FIGURE 1B). APC1–3 have three domains: 1) an NH₂-terminal calcium-regulatory domain with four calcium-binding EF-hands, 2) an amphipathic helix, and 3) a COOH-terminal carrier domain (59) (FIGURE 2B). In the current model of calcium regulation, the amphipathic helix is bound to the regulatory domain in the presence of calcium, allowing transport by the carrier domain to occur (59, 169). In the absence of calcium, the amphipathic helix is released from the regulatory domain and binds to the carrier domain, leading to inhibition of transport (58, 60). APC4 lacks both the NH₂-terminal calcium-regulatory domain and the amphipathic helix (160), which accounts for the absence of calcium regulation.

Dysfunction of the mitochondrial Mg-ATP/phosphate carrier APC1 (SLC25A24) leads to Fontaine Progeroid Syndrome, also described as Gorlin-Chaudhry-Moss or Fontaine-Farriaux syndrome (43, 138, 145, 168) (OMIM 612289). The mutations are spontaneous rather than inherited and lead to a developmental disease, which is characterized by prenatal and postnatal growth retardation, failure to thrive, a lack of subcutaneous adipose tissue, premature closure of certain skull bones (coronal craniosynostosis), and short distal phalanges of the fingers and toes. Other clinical features include abnormal hair patterns, skin agenesis, umbilical hernia, and progeroid facial appearance. APC1 variants are expressed in mitochondria, and they affect both the mitochondrial morphology and cell viability. There is also a decrease in mitochondrial ATP

synthesis in fibroblasts, but only under stress conditions (168). The disease variants have not been characterized with respect to folding and transport function. Interestingly, the paralog APC3 (SLC25A25) has been implicated in left-right determination during development and is regulated by TRPP2 ion channels (64), showing that the Mg-ATP/phosphate carriers are also playing an important role in development.

There are two pyrimidine nucleotide carriers in humans (PNC1, SLC25A33 and PNC2, SLC25A36), which are required for mitochondrial DNA and RNA synthesis and breakdown (37, 52) (FIGURE 1C). SLC25A33 transports uracil, thymine, and cytosine (deoxy)nucleoside di- and tri-phosphates by an antiport mechanism, whereas SLC25A36 translocates both cytosine and uracil (deoxy)nucleoside mono-, di-, and tri-phosphates using a uniport or antiport mechanism (37, 52). Both carriers also transport guanine nucleotides, but not adenine (deoxy)-nucleotides. It has also been claimed that the pyrimidine carriers are involved in the uptake of zinc ions (85), even though several dedicated zinc transporters are likely to exist in mitochondria. In yeast, a separate mitochondrial GTP/GDP carrier has been identified (162), but an equivalent transporter has not been identified in humans. The most closely related carriers by sequence comparison are SLC25A51 and SLC25A52 (137), but no experimental data are available to support this notion. So far, no disease variants of PNC1 and 2 have been identified.

Amino Acid Transport

Amino acids need to be transported into mitochondria for mitochondrial protein synthesis and for amino acid interconversion and degradation, which can generate metabolic energy. Several mitochondrial amino acid transporters have been identified, but some are still missing, importantly those for tryptophan, tyrosine, phenylalanine, methionine, glutamine, asparagine, and cysteine (FIGURE 1D).

The mitochondrial aspartate/glutamate carriers import glutamate and a proton, and they export aspartate from the mitochondrial matrix (9, 16, 18). AGC1 (SLC25A12, aralar) and AGC2 (SLC25A13, citrin) are expressed in excitable and non-excitable tissues, respectively, and play important roles in the malate-aspartate shuttle (FIGURE 1E), gluconeogenesis, purine and pyrimidine synthesis, the urea cycle (AGC2) (FIGURE 1F), and myelin synthesis (AGC1) (126). They have an unusual three-domain structure, consisting of 1) a NH₂-terminal calcium-regulatory domain, 2) a carrier domain, and 3) a COOH-terminal amphipathic helix (157) (FIGURE 2C). The NH₂-terminal domain has eight EF-hand domains, but only EF-hand 2 is capable of

binding calcium, whereas EF-hands 4–8 are involved in dimerization, generating a structural homo-dimer (FIGURE 2C). In the calcium-bound state, the amphipathic helix is bound to the regulatory domain, whereas in the absence of calcium the amphipathic helix is released (157), but the mechanism of regulation is not fully resolved.

Mutations of the gene coding for the liver paralog AGC2 (SLC25A13) causes neonatal and/or adult-onset type-II citrullinemia (OMIM 603859), an autosomal recessive disease characterized by hyperammonemia and citrullinemia, because of a dysfunctional urea cycle, as well as neuropsychiatric symptoms and fatty liver disease in later life (86, 122). Mutations of the gene coding for the brain-specific paralog AGC1 (SLC25A12) lead to early infantile epileptic encephalopathy due to hypomyelination (OMIM 603667). This is caused by the lack of mitochondrial aspartate export, which is required for the synthesis of *N*-acetyl-aspartate, a precursor for myelin synthesis (167).

Two related proteins, the mitochondrial glutamate carriers GC1 (SLC25A22) and GC2 (SLC25A18), are involved in the import of glutamate together with a proton but do not export aspartate (FIGURE 1D) (50). The glutamate carriers do not have the calcium-regulatory elements found in aspartate/glutamate carriers. Mutations in SLC25A22 (OMIM 609302) cause neonatal epileptic encephalopathy with suppression bursts (97, 110) or migrating partial seizures in infancy with poor developmental prognosis (129). The most likely reason for the disease phenotype is the role of mitochondrial glutamate transport in sustaining glutamate homeostasis in astrocytes (55).

The glycine carrier GLYC (SLC25A38) is involved in the import of glycine into mitochondria, where it reacts enzymatically with succinyl-CoA to form aminolevulinic acid (FIGURE 1G) (102). The gene was flagged because a disease variant leads to autosomal recessive sideroblastic anemia (OMIM 610819), caused by the inability of blood cells to synthesize heme due to defective glycine transport (48). Aminolevulinic acid is transported by an unknown transporter into the cytosol, where it is used as a precursor for porphyrin synthesis. The resulting coproporphyrinogen III is then transported into mitochondria by an unknown transporter for incorporation of iron and insertion into heme-containing proteins.

The ornithine carriers ORC1 and ORC2 (SLC25A15 and SLC25A2) catalyze the exchange of ornithine and citrulline, which links the fixation of ammonia in mitochondria to the urea cycle (FIGURE 1H) (47). The substrate specificity and binding of these two paralogs have been studied in detail (111). Dysfunction of ORC1 (SLC25A15) leads to HHH syndrome (OMIM 603861), characterized

by hyperornithinemia, hyperammonemia, and homocitrullinuria, due to an impaired urea cycle, which is required for the deamination of amino acids (22, 65). ORC2 (SLC25A2) (OMIM 608157) may be responsible for the milder phenotype in HHH patients, secondary to a gene redundancy effect. The basic amino acid carrier BAC (SLC25A29) is closely related to ORC1 and 2, but transports arginine, lysine, homoarginine, methylarginine, and, to a much lesser extent, ornithine and histidine (131), and has thus far no pathology associated with it.

Recently, the mitochondrial carrier for branched-chain amino acids (SLC25A44) was identified (FIGURE 1D) (171). The branched-chain amino acids valine, leucine, and isoleucine can be degraded to provide metabolic energy and are required for the synthesis of proteins in mitochondria. The carrier was discovered in brown adipose tissue upon cold exposure, where branched-chain amino acids can be used as a fuel for thermogenesis.

Vitamin Transport

Many vitamins need to be transported into mitochondria, where they serve as co-factors or as donors of key functional groups in the enzymatic reactions of the mitochondrial matrix (FIGURE 1I).

The mitochondrial S-adenosylmethionine carrier SAMC (SLC25A26) imports S-adenosylmethionine into mitochondria, which is required for methylation reactions of DNA, RNA, and protein in the mitochondrial matrix, and exports the product S-adenosylhomocysteine (3). A missense mutation in the gene coding for this protein causes intra-mitochondrial methylation deficiency (OMIM 611037) in agreement with this function, leading to oxidative phosphorylation deficiency (82).

The human thiamine pyrophosphate transporter TPC (SLC25A19) was initially identified as a deoxynucleotide transporter (39), but its main function is the transport of thiamine pyrophosphate, which is an important co-factor in dehydrogenase reactions (99). Defective transport of thiamine pyrophosphate is the cause of Amish microcephaly, which is characterized by profound congenital microcephaly, delayed psychomotor development, and lactic and alpha-ketoglutaric aciduria (OMIM 606521) (78). A second mutation was reported in siblings of non-Amish background (OMIM 613710), expanding the phenotypes associated with the *SLC25A19* gene. These patients showed recurrent episodes of flaccid paralysis and encephalopathy associated with bilateral striatal necrosis and chronic progressive polyneuropathy caused by a missense mutation in *SLC25A19* (155).

Mitochondria require folate for one-carbon metabolism and flavins for electron transfer steps in the respiratory chain. The transport of folate and

flavin have both been assigned to a single mitochondrial carrier (SLC25A32) (154, 159). The substrate binding site has the typical features of an adenine binding pocket, agreeing with the flavin assignment (94, 136, 137). When mutated, the disease variant of this carrier causes exercise intolerance (OMIM 610815), and riboflavin supplementation proved to be beneficial to the patients, indicating that the most likely substrate of this carrier is flavin rather than folate (149).

Many reactions in the mitochondrial matrix, such as dehydrogenase activities, require coenzyme A (CoA) as co-factor. Coenzyme A is synthesized outside of mitochondria and must be transported into the mitochondrial matrix. The CoA transporter (SLC25A42) was initially identified based on database searches and showed expression in all tissues, with the highest levels detected in adipose tissue, and high levels detected in hypothalamus and all brain coronal sections (57). The disease variant of the CoA transporter (OMIM 610823) causes mitochondrial myopathy with muscle weakness and lactic acidosis, whereas other tissues and cognitive functions are not impaired (150). Another report described variable clinical manifestations, including lactic acidosis, developmental regression, and epilepsy (4).

Before *SLC25A42* was identified as the CoA transporter, this transport activity was putatively assigned to a protein encoded by the *SLC25A16* gene (133). Both *SLC25A16* and *SLC25A42* are phylogenetically related to the yeast protein Leu5p and are capable of complementing the yeast knockout strain (51). The function and kinetic parameters of *SLC25A42* were determined in transport assays with substrate specificities restricted to CoA, dephospho-CoA, ADP, and adenosine 3',5'-diphosphate (51). The human protein encoded by the *SLC25A16* gene has been identified through a possible association with a thyroid disease called Grave's disease, but it has not been demonstrated that it can transport CoA (173). To date only a single homozygous mutation has been reported in the *SLC25A16* gene (OMIM 139080), causing a nail disorder of the hand with different severity levels of onychodystrophy (79), but a direct correlation to function has not been confirmed either.

Inorganic Ion Transport

Inorganic ions also need to be transported into the mitochondrial matrix, where they function as co-factors, as substrates for enzymatic reactions, and as regulators. There are likely to be transporters of other families as well as channels involved in inorganic ion transport, but they are beyond the scope of this review.

The mitoferrins *SLC25A37* (MFRN1) and *SLC25A28* (MFRN2) have been proposed to

transport iron ions into mitochondria for incorporation into heme and iron-sulfur cluster synthesis, as well as other functions (FIGURE 1J) (113, 151). MFRN1 is highly expressed in differentiating erythroid cells and in other tissues at low levels, whereas MFRN2 is expressed ubiquitously in non-erythroid tissues (29, 151). Abnormal MFRN1 expression might contribute to erythropoietic protoporphyria phenotype, in agreement with this notion (165).

The mitochondrial phosphate carrier PIC (SLC25A3) imports inorganic phosphate for the synthesis of ATP (140) together with a proton (FIGURE 1K) (95). There are two alternative splicing variants showing different kinetic parameters and different expression profiles: whereas isoform A is highly expressed in heart, skeletal muscle, diaphragm (49), and pancreas (67), isoform B is expressed in all tested tissues, i.e., lung, kidney, brain, thymus, liver, heart, skeletal muscle, and diaphragm (49), albeit at lower levels compared with isoform A (67). The phosphate carrier is fundamental in maintaining the inorganic phosphate pool in the mitochondrial matrix, but there are also other carriers capable of transporting phosphate, such as the dicarboxylate carrier DIC (SLC25A10) and the mitochondrial ATP-Mg/Pi carriers APC1 (SLC25A24), APC2 (SLC25A23), APC3 (SLC25A25), and APC4 (SLC25A41) (5, 45, 160). Phosphate carrier deficiency leads to lactic acidosis, hypertrophic cardiomyopathy, muscular hypotonia, and early mortality (OMIM 600370), in agreement with the notion that its transport activity is the main pathway for phosphate import for ATP synthesis (15, 105, 106).

Fatty Acid Transport

The carnitine/acylcarnitine carrier CAC (SLC25A20) is a key component of the carnitine cycle and imports acyl-carnitine into mitochondria for fatty acid β -oxidation and exports carnitine (FIGURE 1L) (66, 70). Mutations cause carnitine/acylcarnitine carrier deficiency, an autosomal recessive disorder characterized by severe, neonatal onset with cardiomyopathy or a milder phenotype with hypoglycemia, but no cardiomyopathy (OMIM 613698) (66). The inability to transport fatty acid chains into mitochondria makes the patients dependent on carbohydrates and amino acids for energy metabolism.

Uncoupling Protein

The uncoupling protein UCP1 (SLC25A7) is predominantly found in brown adipose tissue of neonatal mammals (6, 118, 120) but is also found in the supraclavicular and the neck regions in adults in later life (FIGURE 1M) (116). UCP1 dissipates the proton motive force, short-circuiting the mitochondrion, which leads to the production of heat.

The generation of heat from the oxidation of brown adipose fat protects the newly born against cold stress of vital organs. UCP1 is activated by fatty acids and inhibited by purine nucleotides (134), but the mechanism is still debated (23, 33, 44, 72, 84, 119). UCP1 is monomeric and binds three cardiolipins and a single purine nucleotide (96). Based on sequence analysis, UCP1 has retained all of the key features of mitochondrial carriers, indicating that it operates by a conventional carrier-like mechanism (33, 96). The transport of protons induced by fatty acids is relatively slow (44), which would fit a transporter rather than a channel mechanism, but a molecular mechanism has not been resolved. Although there are other closely related proteins (see below), UCP1 is most likely the only one involved in thermogenesis. There are no known disease states associated with mutations in the gene coding for UCP1.

Dicarboxylate Transport

Dicarboxylates (FIGURE 1N) need to be continuously exchanged across the mitochondrial inner membrane for many different pathways, such as the tricarboxylic acid cycle, malate/aspartate shuttle, gluconeogenesis, and amino acid metabolism.

The mitochondrial dicarboxylate carrier DIC (SLC25A10) is involved in the transport of malonate, malate, succinate, sulfate, thiosulphate, and phosphate by electroneutral exchange (46, 73). The carrier is involved in gluconeogenesis and ureogenesis, the metabolism of sulfur compounds, as well as de novo fatty acid synthesis (109). The mitochondrial oxoglutarate carrier OGC (SLC25A11) exchanges cytosolic malate for 2-oxoglutarate from the mitochondrial matrix and plays an important role in the malate-aspartate shuttle, the oxoglutarate-isocitrate shuttle, and gluconeogenesis (69). Mutations in the gene might be correlated to metastatic paragangliomas (21).

The oxodicarboxylate carrier ODC (SLC25A21) imports 2-oxoadipate and exports 2-oxoglutarate, playing a central role in the catabolism of lysine, hydroxylysine, and tryptophan (FIGURE 1O) (48). Oxodicarboxylate carrier deficiency (OMIM 607571) is associated with mitochondrial DNA depletion and spinal muscular atrophy-like disease, most likely caused by the accumulation of toxic amino acid breakdown products (17).

There are several closely related sequences to UCP1, such as UCP2 (SLC25A8), UCP3 (SLC25A9), UCP4 (SLC25A27), UCP5 (SLC25A14), and UCP6 (SLC25A30), but they are likely to be transporters of carboxylic acids (54, 163), which is in agreement with their close phylogenetic relationship to dicarboxylate transporters (123) and with the properties of their substrate binding sites (137). Many unresolved questions remain with respect to their




FIGURE 3. Mitochondrial carriers have a threefold pseudo-symmetrical structure

A: aligned amino acid sequences of the three repeats of the human ADP/ATP carrier AAC1 (ANT1). Symmetrically conserved residues are shown in the consensus sequence and as bars, when present in at least two out of three repeats. B and C: comparative model of the human ADP/ATP carrier, based on PDB:1OKC (128), viewed from the membrane and the intermembrane space, respectively. Shown also is the threefold pseudo-symmetrical axis, symbolized by an equilateral triangle. Odd-numbered (H1, H3, H5), matrix (h12, h34, h56), linker (I12, I34, I56), and even-numbered helices (H2, H4, H6) are shown in primary colors for the core elements and in gray for the gate elements (142). The black spheres with roman numerals show the positions of the three contact points of the substrate binding site (94, 136). The example, triplet 8–113–210, is indicated by a rectangle across the three repeats in A and by cyan spheres in B. IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

molecular properties and their role in thermogenesis and metabolism. There are also other potential dicarboxylate carriers, such as SLC25A34 and SLC25A35, but their role in metabolism has not been clarified.

Tricarboxylate Transport

The tricarboxylate or citrate carrier (SLC25A1) catalyzes the electroneutral exchange of tricarboxylates (citrate, isocitrate) for another tricarboxylate, a dicarboxylate or phosphoenolpyruvate (68, 76, 103). An important physiological function is the export of citrate from the mitochondria to the cytosol for the production of acetyl CoA, which is a starting point for lipid, dolichol, ubiquinone, and sterol synthesis (see references in Ref. 103), and acetylation reactions (112) (FIGURE 1P). Citrate carrier deficiency (OMIM 190315), which is hallmarked by combined D-2- and L-2-hydroxyglutaric aciduria, leads to neonatal-onset epileptic encephalopathy with severe muscular weakness, respiratory distress, and lack of psychomotor development resulting in early death (30, 42, 114, 121, 132, 152), which is most likely due to the severe biosynthetic deficiencies (103). Many of the missense mutations have been characterized with respect to the function of SLC25A1 (27, 42, 103, 130).

Apoptosis

SLC25A50 (MTCH2) is a partially characterized mitochondrial carrier, which acts as a receptor-like

protein for the truncated BH3-interacting domain death agonist protein in the outer membrane of mitochondria, as part of the apoptosis pathway (FIGURE 1Q) (172). MTCH2 is likely to have a similar topology as other mitochondrial carriers (135), but unusually it is found in the mitochondrial outer membrane. A transported substrate has not yet been identified, if this protein has a transporter function at all, but MTCH2 was subsequently found to be required and sufficient for lipid homeostasis shifts (139).

Mitochondrial Dynamics

The partially characterized mitochondrial carrier SLC25A46 is most likely involved in mitochondrial dynamics (FIGURE 1R). Overexpression leads to mitochondrial fragmentation, whereas knockdown results in hyperfilamentous mitochondria and mitochondrial hyperfusion, likely resulting from decreased fission. Loss of SLC25A46 was not associated with changes in total ATP concentration, mitochondrial DNA content, or membrane potential (2). SLC25A46 could interact with proteins associated with mitochondrial dynamics, such as OPA1 and MFN2, and with components of the mitochondrial contact site and mitochondrial cristae organizing system complex, which plays a role in cristae maintenance (71). SLC25A46 might also interact with components of an endoplasmic reticulum membrane protein complex involved in lipid transfer to




FIGURE 4. Alternating access transport mechanism for the mitochondrial ADP/ATP carrier

Lateral view from the membrane of the mitochondrial ADP/ATP carrier in the cytoplasmic state (left) and matrix state (right). Shown are the cytoplasmic state of the bovine ADP/ATP carrier (PDB:1OKC) (128) and the matrix state of the ADP/ATP carrier of *Thermothelomyces thermophila* (PDB:6GCI) (142). The water-accessible surfaces are shown in light blue. Also indicated are the three main functional features: cytoplasmic gate, substrate binding site, and matrix gate. The black spheres with roman numerals are the contact points of the substrate binding site (94, 136). Shown in green are residues that most likely bind the adenine nucleotide substrates. IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

mitochondria, which are also required for cristae growth and maintenance. SLC25A46 is found in the outer membrane of mitochondria (71), which is atypical for mitochondrial carriers except for MTCH2 (see above). It is possible that the carrier has evolved away from the canonical transporter function, since, so far, no substrates have been identified. Disease variants of SLC25A46 (OMIM 610826) lead to hereditary motor and sensory neuropathy type VIB (2) and in severe cases to death in infancy (1, 164).

Uncharacterized Mitochondrial Carriers

The functions of a large number of mitochondrial carriers have not yet been assigned, limiting our understanding of their role in human physiology and pathology. Among them are SLC25A34, SLC25A35, SLC25A39, SLC25A40, SLC25A43, SLC25A45, SLC25A47, SLC25A48, SLC25A49 (MTCH1), SLC25A51 (MCART1), SLC25A52 (MCART2), and SLC25A53 (MCART6), which is roughly a quarter of the total. No substrates have been identified for SLC25A46 and SLC25A50 (MTCH2) (see above), although they have a role in




FIGURE 5. Structural changes in the transport cycle of the human mitochondrial ADP/ATP carrier

View from the intermembrane space (A) and lateral view from the membrane of the human ADP/ATP carrier (B) in the cytoplasmic state (left) and matrix state (right). For the import of ADP, conformational changes involve the simultaneous outward rotation of the core elements, shown in primary colors, and inward rotation of the gate elements, shown in gray, in a coordinated way (142). For the export of ATP, the converse happens. Substrate binding increases the probability of the state interconversion (156). The structural models are based on the structures of the cytoplasmic state of the bovine ADP/ATP carrier (PDB:1OKC) (128) and the matrix state of the ADP/ATP carrier of *Thermothelomyces thermophila* (PDB:6GCI) (142). The black spheres with roman numerals are the contact points of the substrate binding site (94, 136). IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

mitochondrial dynamics and apoptosis, respectively. There is still a lot of debate on the role of UCP2 (SLC25A8), UCP3 (SLC25A9), UCP4 (SLC25A27), UCP5 (SLC25A14), and UCP6 (SLC25A30) in human physiology. In some other cases, the function of particular carriers has been disputed, for example SLC25A32, which has been described as a folate or flavin transporter. Counting these, the number of carriers that are not yet fully characterized is closer to a third of the total.

The Molecular Basis of Pathogenic Missense Mutations

Missense mutations in 16 different carriers lead to human disease, but it is likely that this number will increase substantially as more links are discovered through genome and exome sequencing. These mutations can impair the structure and mechanism of the carrier, but can also cause issues with biogenesis, i.e., the expression, targeting, insertion, and folding of the disease variant. The impact of the vast majority of these missense mutations on the structure, function, and biogenesis of different carrier has not been studied experimentally. To discriminate between these different scenarios, it is important to understand the transport mechanism in detail. Recently, good progress has been made facilitating this assessment, although many details still need to be worked out. To explain the molecular impact of the pathogenic mutations on the function of mitochondrial carriers, it is important to explain their basic structure and transport mechanism first.

Structures of Mitochondrial Carriers

Mitochondrial carriers consist of three homologous repeats of ~100 amino acid residues each (FIGURE 3A) (147). The three repeats fold up into a threefold pseudo-symmetrical fold, noted first in the projection structure of the yeast ADP/ATP carrier Aac3p (92). This study also demonstrated that the carrier had a monomeric structure and a translocation path for substrates through the center of the protein (92). The first atomic structure of the bovine ADP/ATP carrier provided evidence for the basic structural topology of all mitochondrial carriers (128). Each repeat or domain has an

odd-numbered helix (H1, H3, H5), a matrix loop of variable length, a matrix helix (h12, h34, h56), a linker helix (l12, l34, l56), and an even-numbered helix (H2, H4, H6) (128) (FIGURE 3, A AND B). The domains are linked by cytoplasmic loops, which are located in the intermembrane space, together with the NH₂ and COOH termini. The structure was locked by the specific inhibitor carboxyatractyloside in the cytoplasmic state in which the central cavity is open to the intermembrane space for binding of ADP (FIGURES 4 AND 5) (128). Although there are some structural differences, the same basic structural fold was observed for the yeast ADP/ATP carrier Aac2p and Aac3p, even though they share only ~50% identity with the bovine carrier (141). More recently, the atomic structure of the mitochondrial ADP/ATP carrier inhibited by bongkreikic acid has been solved, locked in the matrix state in which the central cavity is open to the mitochondrial matrix, for binding of ATP (FIGURES 4 AND 5) (142, 144). The central cavity in both states is positively charged, primed for binding of the negatively charged adenine nucleotides (128, 142, 144). The residues of the substrate binding site have been identified by computational methods, using chemical and distance constraints in comparative models (94, 136), deviation of symmetry (95, 137), or molecular dynamics simulations (35, 107, 166). The three main contact points involved in substrate binding are located on the even-numbered helices (FIGURES 4 AND 5) (94, 136). The substrate binding site is located at the bottom of a water-filled cavity in both states, which corresponds to the middle of the membrane (94, 136). There are two gates on either side of the carrier that regulate access to the central binding site. In the cytoplasmic state, the matrix gate is closed and the cytoplasmic gate is open, whereas, in the matrix state, the cytoplasmic gate is closed and the matrix gate is open (FIGURES 4 AND 5) (142–144). Each closed gate is ~15-Å thick, an important insulation layer to prevent the leak of protons and other ions in the presence of a 180-mV membrane potential. These gates contain salt-bridge networks of positively and negatively charged amino acid residues and other features, which will be discussed later.

FIGURE 6. Pathogenic mutations observed in disease variants of mitochondrial carriers

Aligned triplets of the 16 mitochondrial carriers associated with developmental, metabolic, and neuromuscular diseases. The mutations that have a severe effect on function are shown in red boxes, whereas those that have milder effects are in yellow. Mutations in blue boxes have been identified by genetic analysis, but their effect has not been studied experimentally. At the bottom are the three residue numbers that form a triplet in the human ADP/ATP carrier (SLC25A4). At the top are the conserved structural and functional features of mitochondrial carriers. The triplet is labeled by the one-letter code of the most conserved residue or by the most common property: π , small amino acids; Φ , hydrophobic amino acids; ξ , hydrophilic amino acids; Ω , aromatic amino acids; or by X for any amino acid. The black spheres with roman numerals are the contact points of the substrate binding site (94, 136). H6 in the ADP/ATP carrier is one residue shorter than other carriers and lacks a residue in triplet 90. The matrix loops (indicated by the black bar), as well as the cytoplasmic loops and NH₂ and COOH terminus have been omitted.

Transport Mechanism of Mitochondrial Carriers

Comparison of the two inhibited states has revealed the basic structural mechanism of transport and has completed our understanding of the importance of key conserved sequence features of the SLC25 family for the transport mechanism (FIGURE 5) (142–144). The two inhibitors occupy the proposed substrate binding site of the carrier, preventing substrate binding, but also induce slight conformational changes, which lock the carrier permanently in an abortive state (142). These distortions can be corrected for structurally to achieve a closer approximation of the unliganded

states, which are used in this review. A morph between the unliganded states passes through an occluded state, where access to the substrate binding site is blocked from both sides of the membrane, which is a requirement for an alternating access transport mechanism (142).

Further analysis shows that most of the domain structure is conserved between the two states, i.e., the odd-numbered helix, the matrix helix, the linker helix and a third of the even-numbered helix (142–144). These parts of the domain are called the core elements and are shown in primary colors blue, yellow, and red for the first, second, and third domain, respectively (FIGURE 5). In contrast, a significant change




FIGURE 7. Small amino acid residues on the odd-numbered helices

View from the intermembrane space (A) and lateral view from the membrane (B) of the human ADP/ATP carrier in the cytoplasmic state (left) and the matrix state (right). The structural models are based on the structures of the cytoplasmic state of the bovine ADP/ATP carrier (128) and the matrix state of the ADP/ATP carrier of *Thermothelomyces thermophila* (142). The carrier is shown in surface representation and the helices in cartoon representation with the core elements in primary colors and the gate elements in gray. Glycine or small residues in the interface with the preceding helix are shown in pink, whereas those in the interface with the following helix are shown in magenta. The sequence motif is $\pi G\pi\pi\pi Gxx\pi xxx\pi$, where G stands for glycine and π for small amino acids (142–144). When the carrier changes from the cytoplasmic state to the matrix state, the inter-helical distances become smaller on the cytoplasmic side of the carrier to facilitate the formation of the cytoplasmic network, requiring small residues in the helical interfaces. IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

occurs in the position of the COOH-terminal regions of the even-numbered helices in the state interconversion (143, 144). These parts of the structure are called the gate elements, and they are shown in gray (FIGURE 5). The hinge points of these movements turned out to be the contact points of the proposed substrate binding site (142–144), identified previously (94, 136). Opening or closing of the matrix side of the carrier involves the rotation of the three core elements as rigid bodies, whereas opening or closing of the cytoplasmic side requires the rotation of the three gate elements. In this way, access to the central substrate binding site from one or the other side of the membrane is alternated (FIGURE 5). The substrate binding site is the fulcrum of these movements (142–144). Thus

state interconversion requires the coordinated movement of six elements simultaneously, making the mitochondrial carriers one of the most dynamic of all transport proteins. These movements are facilitated by the transmembrane helices being held together only by relatively weak van der Waals interactions. Next, the sequence features of mitochondrial carriers are examined to determine why they are crucial to the structure and transport mechanism, since they are often altered in pathogenic variants.

Key Sequence Features Affected by Pathogenic Mutations

The easiest way to present the sequence features of mitochondrial carriers is to use their threefold pseudo-symmetry (FIGURE 3). The three repeats




FIGURE 8. Key amino acid residues of the matrix gate

View from the intermembrane space (A) and lateral view from the membrane (B) of the human ADP/ATP carrier in the cytoplasmic state (left) and matrix state (right). The structures are described in the legend to FIGURE 7. The key residues shown belong to the sequence motif P_x[DE]_{xx}[RK]_{xxx}Q on the odd-numbered helices. The proline residues (orange) are found at the pronounced kink in the odd-numbered helices, bringing the negatively charged (red) and positively charged (blue) residues together to form the matrix network in the cytoplasmic state. Underneath one of the salt bridges is a glutamine residue (green) that forms hydrogen bonds with both residues (glutamine brace), but in other carriers one, two, or three glutamine braces can be found (FIGURE 6) (141). IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

are homologous to each other, meaning that residues that are in the same position in each repeat are symmetry-related and thus mostly identical or similar in physiochemical properties (137). These residues can be grouped together in a triplet of symmetry-related residues, which facilitates their comparison (FIGURE 3A). Here, we have used the residue numbering of the human ADP/ATP carrier 1 (AAC1, SLC25A4), also known as ANT1, to relate the triplet to the original sequence. For example, the highlighted triplet (cyan sphere) in FIGURE 3A contains *residue 8 in repeat 1, residue 113 in repeat 2, and residue 210 in repeat 3* (FIGURE 3). For convenience, the entire triplet is named by the residue number in the first repeat in SLC25A4, meaning that the example above would be called *triplet 8*. Given the high sequence identity within the SLC25 family, the same triplets in carriers with different functions can be compared to look for ones that are conserved throughout the family and are therefore universally important for the structure and mechanism of mitochondrial carriers (137).




FIGURE 9. Detailed view of one of the three binding sites for cardiolipin

One cardiolipin molecule is shown in ball-and-stick representation (cdl802, PDB: 2C3E), which is bound on the surface of the carrier and spans the inter-domain interface. The carrier is shown in cartoon representation with transmembrane H4 (yellow and gray) and matrix helix h56 (red) enhanced. The two phosphate groups of cardiolipin, which are linked by a glycerol moiety, form hydrogen bonds with the amide groups (128) and interact with the positively charged ends of the helix dipoles of the NH₂-terminal ends of the matrix helices (cardiolipin binding site I) and the even-numbered helices (cardiolipin binding site II) (24, 141, 142). The four fatty acid chains of cardiolipin interact with the surface of the carrier in a non-specific way via van der Waals interactions (41). Residues in the conserved cardiolipin binding site I and II are shown as green and purple sticks, respectively. The interface between domain 2 and 3 is shown as a dashed line.

FIGURE 6 shows the triplets of all conserved helical features for the 16 mitochondrial carriers that have been linked to human disease, as well as the reported pathogenic mutations. The mutations that have been experimentally verified to have a severe effect on function are shown in red boxes, whereas those that have milder effects are in yellow ones. Mutations shown in blue boxes have been flagged in genetic analysis, but their effect on function has not been studied experimentally. It is clear that mutations affect a large number of triplets. Next, starting from the NH₂-terminus, these features will be presented by highlighting the properties of the triplets in relation to the known structural mechanism of transport (FIGURE 6). The analysis focuses on the conserved helical parts of the carriers, since the loops are highly variable in length and sequence, and do not play an important role in the transport mechanism. The most conserved amino acid residues of the triplets are indicated by the one-letter amino acid code, whereas the most common chemical and physical properties are also indicated by a one-letter code: π for small residues (Gly, Ala, Ser, Pro, Cys, Val, Thr), Φ for hydrophobic residues (Val, Ile, Leu, Phe, Trp, Tyr, Cys, Ala, and Met), Ω for aromatic residues (Trp, Tyr, Phe), ξ for hydrophilic residues (Asn, Gln, Glu, Asp, Lys, Arg, His, Ser, Thr), and X for any residues.

Small Amino Acid Residues on the Odd-Numbered Helices

The odd-numbered helices H1, H3, and H5 have a strong kink, giving them an L shape (FIGURES 3 AND 4) (128). The NH₂-terminal parts are transmembrane and contain a large number of glycine and other small residues, which are often mutated in disease variants of the carriers (FIGURE 6). The extended sequence motif is $\pi G \pi x \pi G x x \pi x x x \pi$, where G stands for glycine, π for small amino acids, and x for any amino acid residue (142–144) (FIGURE 7). These residues can be divided further into two categories: 1) small residues in the interface with the preceding helix, i.e., triplets 14, 18, 22, and 26 (pink residues, FIGURE 7), and 2) glycine or small residues in the interface with the following helix, i.e., triplets 15 and 19, which form the GxxxG motif, and triplet 16 (magenta residues, FIGURE 7) (141–144).

When the carrier transitions from the cytoplasmic state to the matrix state (FIGURES 4 AND 5), the cytoplasmic side of the carrier closes and the transmembrane helices come together. The reason is that the cytoplasmic side closes because the gate elements rotate inward, allowing the cytoplasmic network to form (see below). At the same time, the core elements rotate outward, and simultaneously

the odd helices move inward on the cytoplasmic side. The glycine and other small residues of the $\pi G\pi x\pi Gxx\pi xxx\pi$ motif are in these crucial interhelical interfaces, allowing movement of the gate elements across the surface of the odd-numbered helices and the close proximity of the helices in the matrix state. A large number of pathogenic mutations are observed in these triplets, and functional and structural analysis shows that these residues are likely to be important for the mechanism of mitochondrial carriers in general (25, 33, 90). Small residues can also be found on the even-numbered helices, as will be discussed below.

The other triplets found on the transmembrane parts of the odd-numbered helices preserve their strong amphipathic properties. *Triplets 8, 9, 10, 12, 13, 17, 20, 21, 24, and 25* contain mostly generic hydrophobic residues (Φ symbols, FIGURE 6),

since they point toward the hydrophobic core of the membrane. *Triplets 11, 23, and 27* contain generic hydrophilic residues (ξ symbols, FIGURE 6) and point toward the water-filled cavity. Pathogenic mutations are sporadically observed in these triplets, most likely when they are replaced with a residue with the opposite properties. For instance, a charged residue for a hydrophobic one, a large residue for a small one, or a substitution with a proline residue in the middle of a helix, which could break the helix. These features could be important for both the function and biogenesis of the carriers.

Key Amino Acid Residues of the Matrix Gate

The next important motif on the odd-numbered helices is a highly conserved symmetrical feature $Px[DE]xx[RK]xxxQ$ (FIGURE 6). As mentioned




FIGURE 10. Amino acid residues involved in cardiolipin binding

View from the intermembrane space (A) and lateral view from the membrane (B) of the human ADP/ATP carrier in the cytoplasmic state (left) and matrix state (right). The structures are described in the legend to FIGURE 7. There are three highly conserved binding sites for cardiolipin in the mitochondrial ADP/ATP carrier. The residues of cardiolipin binding site I belong to the $[YF]xG$ motif (green), whereas those of cardiolipin binding site II belong to the $[YWF][RK]G$ motif (128, 141, 142). IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

above, the odd-numbered helices have a strong kink of ~50 degrees (FIGURE 5). At this kink is a highly conserved proline residue (P-kink, *triplet 28*), the first residue of the motif (128), but it can be replaced by a serine residue (141). These residues break the hydrogen bond arrangement, allowing the kink to occur, and a network of interactions between residues in the domain help to stabilize it (141) (FIGURE 8). The kinks bring the COOH-terminal ends of the odd-numbered helices together in the center of carrier in the cytoplasmic state, where the negatively charged residues (red residues, *triplet 30*) and positively charged residues (blue residues, *triplet 33*) form an ionic interaction network (117). This interaction network can be seen in the structure of the cytoplasmic state (128, 141), now called the matrix salt-bridge network (137). A glutamine residue (*triplet 37*) functions as

a brace of one of the salt-bridge interactions of the matrix network (glutamine-brace) (green residue, FIGURE 8) (141). Only one of the three domains of the ADP/ATP carrier has a glutamine brace, but other carriers have up to three glutamine braces (FIGURE 6). The salt-bridge interactions and glutamine braces together determine the overall interaction energy of the matrix network. Although not conserved between different carriers, residues in *triplets 34* and *38* are in the translocation path, sealing the carrier to the mitochondrial matrix (FIGURE 6). Together, all these residues form the matrix gate, which is closed in the cytoplasmic state and open in the matrix state (FIGURES 5 AND 6). In addition, there are triplets with hydrophobic residues (*triplet 29, 32, 36*), hydrophilic residues (*triplet 34, 38*), and residues with varied properties (*triplet 31*). This part of the carrier is one of the




FIGURE 11. Amino acid residues important for the domain structures on the matrix side

View from the intermembrane space (A) and lateral view (B) from the membrane of the human ADP/ATP carrier in the cytoplasmic state (left) and matrix state (right). The structures are described in the legend to FIGURE 7. The positively charged residue of the E-R link I (blue), which is located in the matrix gate area, interacts with the negatively charged residue of the E-R link II, which is located on the matrix helices. The NH₂-terminal and the fourth residues of the linker helices are most commonly glycine residues (magenta), although the latter can be replaced by other small residues. IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

most important for their function, and a large number of pathogenic variants have been identified in this region (FIGURE 6).

Amino Acid Residues Involved in Cardiolipin Binding

Three cardiolipin molecules are tightly bound to mitochondrial carriers and are important for their stability and function, as observed by phosphorous NMR experiments (14), crystallographic analyses (128, 141, 142), lipid analysis (10, 93, 96), disease models (56, 74, 104), thermostability analysis (34), and transport assays (63, 75, 107). The phosphate groups of cardiolipin are bonded to the NH₂-terminal ends of the matrix and even-numbered helices, bridging the inter-domain interface (FIGURE 9). Preceding them are highly conserved symmetrical sequence motifs [YF]XG (*triplets 51–53*, cardiolipin binding site I) and [YWF][RK]G

(*triplets 71–73*, cardiolipin binding site II), respectively (FIGURES 6 AND 10). The glycine residues of these motifs (*triplets 53 and 73*) are in the loop to helix transition, where they function as helix breakers, but serine, asparagine, or threonine can also play this role (24). This loop-to-helix transition is crucial for binding of the phosphate groups of cardiolipin via hydrogen bonds (128) and electrostatic interactions with the helix dipoles (141). Many pathogenic mutations affect these glycine residues (FIGURE 6). The aromatic residue of cardiolipin binding site I (*triplet 51*) is involved in stabilization of the domain on the matrix side (108), whereas the aromatic residue of cardiolipin binding site II (*triplet 71*) is involved in binding the fork of the lipid moiety (128, 141, 142). Although not supported by the structures, molecular dynamics simulations show that the positive charged residue (*triplet 72*) might be involved in binding the




FIGURE 12. Amino acid residues of the substrate binding site
View from the intermembrane space (A) and lateral view (B) from the membrane of the human ADP/ATP carrier in the cytoplasmic state (left) and matrix state (right). The structures are described in the legend to FIGURE 7. The black spheres with roman numerals are the contact points of the substrate binding site, which are involved in binding of the substrates (94, 136). Other residues in this site (green) may either bind substrate directly or may allow the binding. IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

phosphate moiety via electrostatic interactions (41). *Triplet 71–72* have pathogenic mutations, supporting their importance in stability and function of the carriers.

Amino Acid Residues Important for the Stability of the Domain Structure

Analysis of all of the polar interactions of the ADP/ATP carrier show that there are no conserved polar interactions between the transmembrane helices (141), which agrees with a mechanism where the transmembrane helices move relative to each other in the state interconversion (FIGURE 5) (142). The only highly conserved interaction is in the domain structure between a positively charged residue on the odd-numbered helices (E-R link I, *triplet 35*) and a negatively charged residue on the matrix helices (E-R link II, *triplet 65*) (FIGURES 6

AND 11). In the known structures of the ADP/ATP carrier, one or two interactions are evident, supporting the unusual shape of the domain structure (128, 141, 142). One of the residues of E-R link I in repeat I is a leucine, which cannot be involved in a direct interaction with the glutamine residue in E-R link II (FIGURE 6). However, a preceding residue, Arg31, which is located one turn of a helix away, is in bonding distance, thus fulfilling the same role (FIGURE 6). On the basis of sequence analysis, it is likely that all three domains have this interaction in other carriers, which it has been shown to be important for function (108).

Another residue important for the stability is a tyrosine residue in cardiolipin binding site I (*triplet 51*), which forms extra interactions and seals the domain toward the mitochondrial




FIGURE 13. Small amino acid residues on the even-numbered helices View from the intermembrane space (A) and lateral view (B) from the membrane of the human ADP/ATP carrier in the cytoplasmic state (left) and matrix state (right). The structures are described in the legend to FIGURE 7. The small residues (chartreuse) in the interhelical interfaces with the odd-number helices facilitate conformational changes. Some residues are larger, such as the phenylalanine on H6, because their side chains face the membrane in both conformations. IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

matrix (FIGURE 10). The E-R link often contains pathogenic mutations, further supporting its importance.

Residues of the matrix helices are strongly amphipathic, with polar and charged residues facing the mitochondrial matrix (*triplets* 56, 59, 60, 63, 64) and hydrophobic residues facing the membrane (*triplets* 54, 55, 58, 61, 62) (FIGURE 6). Small residues are also required for the stability of the domain structure as helix breakers, such as glycine residues (*triplet* 53, 66, 73) or small residues pointing toward other residues (*triplet* 57, 69) (FIGURE 11). These areas are affected by pathogenic mutations, likely because the mutations cause changes to amino acids with opposite biophysical properties.

Amino Acid Residues of the Substrate Binding Site

A single substrate binding site can be found in the central cavity, approximately in the middle of the membrane. The binding site consists of residues that are directly involved in binding of the substrate, such as the contact points (FIGURES 4, 5, AND 12), but also of residues that allow the binding of the substrate in the water-filled cavity. This area is a hyper-variable region and contains a large number of asymmetric residues (137). Most of these residues can be found on the even-numbered helices at *triplets* 77, 80 (contact points), 81, 84, and possibly 85. There are many disease variants that have mutations in this area and that have been shown to affect function in functional




FIGURE 14. Amino acid residues of the cytoplasmic gate

View from the intermembrane space (A) and lateral view (B) from the membrane of the human ADP/ATP carrier in the cytoplasmic state (left) and matrix state (right). The structures are described in the legend to FIGURE 7. The key residues belong to the sequence motif ξ [FY]xx[YF][DE]xx[RK]. The negatively charged (red) and positively charged (blue) residues together form the cytoplasmic network in the matrix state. Underneath are aromatic residues (orange), which are part of the hydrophobic plug that closes the cytoplasmic gate. The aromatic residue preceding the negatively charged residue is the tyrosine brace (Y-brace) (142–144). Preceding the aromatic residues are hydrophilic residues (ξ), which form the ceiling of the substrate binding site. IS, intermembrane space; IM, inner membrane; MM, mitochondrial matrix.

studies. There are also triplets that often contain proline residues, such as *triplets* 76 and 83, which can be found on either side of the contact points (*triplet* 80) and may facilitate the curvature or the relative movement of the helices.

Small Amino Acid Residues on the Even-Numbered Helices

As explained earlier, the state interconversion requires the presence of small residues in the interhelical interfaces. Several can be found on the even-numbered helices that facilitate these movements, such as the aforementioned *triplet* 76, and a π xxx π motif, formed by *triplets* 86 and 90 (FIGURE 6 AND 13) (142). The even-numbered helices are also highly amphipathic, containing hydrophobic residues in *triplets* 74, 75, 78, 79, 82, 87, and 91, facing the membrane, and hydrophilic residues facing the cavity, such as the substrate binding site residues, and *triplets* 88, 94, and 97 (FIGURE 6).

Amino Acid Residues of the Cytoplasmic Gate

The COOH-terminal ends of the even-numbered helices contain highly conserved symmetrical sequence motifs ξ [FY]xx[YF][DE]xx[RK] (*triplets* 88–96). Together, they form the cytoplasmic gate (FIGURE 6). The negatively charged residues (*triplet* 93) and positively charged residues (*triplet* 96) form the cytoplasmic salt-bridge network when the carrier is in the matrix state (80, 142, 144), whereas this network is disrupted in the cytoplasmic state (red and blue residues, FIGURE 14) (141). The preceding aromatic residue (*triplet* 92), most often a tyrosine residue, can form hydrogen-bond interactions with the negatively charged residue of the neighboring domain (orange residue, FIGURE 14). This interaction is called the tyrosine brace, and mitochondrial carriers have one to three of these interactions, modifying the overall interaction energy of this network. Together with *triplet* 89, it also doubles up as a hydrophobic layer when the carrier is in the matrix state. *Triplet* 88, which is quite variable, is part of the cytoplasmic gate and also forms part of the ceiling of the substrate binding site (green residues, FIGURE 14). These residues are often mutated in disease variants.

Concluding Remarks

Mitochondrial carriers are highly dynamic transporters, which interconvert between a cytoplasmic state and a matrix state by using six dynamic elements, comprising three core elements and three gate elements. They are among the smallest transporters in nature, yet they transport some of the

largest molecules, such as adenine nucleotides, S-adenosyl methionine, flavins, and acyl-carnitines. They do so without significant proton leak, because of a matrix and cytoplasmic gate, both with salt-bridge networks and braces, and other residues that provide an insulation layer. All carriers have a single central substrate binding site with three contact points, which is alternately accessible from one side of the membrane or the other, key properties of an alternating access transport mechanism. A large number of pathogenic mutations have been identified, which cause a range of metabolic, neuromuscular, and developmental diseases. The vast majority of them can be explained because they affect key structural and functional features of mitochondrial carriers. Some of them are found in loop regions, which have not been included in this analysis, and others in extra domains, such as the regulatory domain of the aspartate/glutamate carrier, which have been explained before (FIGURE 2C) (157). The few that are remaining often involve mutations that introduce different properties from those of the original residue (e.g., p.Asp69Tyr in SLC25A1, p.Cys23Arg in SLC25A20, or p.Thr56Pro in SLC25A22), which could impair the structure and function but also the biogenesis of the carrier. To understand these diseases, it is really important to discriminate between these different options, but the majority of these pathogenic mutations have not been studied experimentally. Given the fact that so many amino acid residues are important for the function of mitochondrial carriers and that the SLC25 family is the largest solute carrier family in humans, it is likely that many more disease variants will be discovered. ■

E. R. Kunji, M. S. King, and J. J. Ruprecht thank the Medical Research Council UK for funding (program grant MC_UU_00015/1). C. Thangaratnarajah received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 847675.

No conflicts of interest, financial or otherwise, are declared by the author(s).

E.R.K., M.S.K., J.J.R., and C.T. conceived and designed research; E.R.K., M.S.K., J.J.R., and C.T. analyzed data; E.R.K. and M.S.K. interpreted results of experiments; E.R.K. and M.S.K. prepared figures; E.R.K. and C.T. drafted manuscript; E.R.K., M.S.K., J.J.R., and C.T. edited and revised manuscript; E.R.K., M.S.K., J.J.R., and C.T. approved final version of manuscript.

References

1. Abrams AJ, Fontanesi F, Tan NBL, Buglo E, Campeanu IJ, Rebelo AP, Kornberg AJ, Phelan DG, Stark Z, Zuchner S. Insights into the genotype-phenotype correlation and molecular function of SLC25A46. *Hum Mutat* 39: 1995–2007, 2018. doi:10.1002/humu.23639.

2. Abrams AJ, Hufnagel RB, Rebelo A, Zanna C, Patel N, Gonzalez MA, Campeanu IJ, Griffin LB, Groenewald S, Strickland AV, Tao F, Spezziani F, Abreu L, Schüle R, Caporali L, La Morgia C, Marasca A, Liguori R, Lodi R, Ahmed ZM, Sund KL, Wang X, Krueger LA, Peng Y, Prada CE, Prows CA, Schorry EK, Antonellis A, Zimmerman HH, Abdul-Rahman OA, Yang Y, Downes SM, Prince J, Fontanesi F, Barrientos A, Németh AH, Carelli V, Huang T, Zuchner S, Dallman JE. Mutations in SLC25A46, encoding a UGO1-like protein, cause an optic atrophy spectrum disorder. *Nat Genet* 47: 926–932, 2015. doi:10.1038/ng.3354.
3. Agrimi G, Di Noia MA, Marobbio CM, Fiermonte G, Lasorsa FM, Palmieri F. Identification of the human mitochondrial S-adenosylmethionine transporter: bacterial expression, reconstitution, functional characterization and tissue distribution. *Biochem J* 379: 183–190, 2004. doi:10.1042/bj20031664.
4. Almannai M, Alasmari A, Alqasbi A, Faqeih E, Al Mutairi F, Alotaibi M, Samman MM, Eyaid W, Aljadhay YI, Shamseldin HE, Craigen W, Alkuraya FS. Expanding the phenotype of SLC25A42-associated mitochondrial encephalomyopathy. *Clin Genet* 93: 1097–1102, 2018. doi:10.1111/cge.13210.
5. Aprille JR. Mechanism and regulation of the mitochondrial ATP-Mg/P(i) carrier. *J Bioenerg Biomembr* 25: 473–481, 1993. doi:10.1007/BF01108404.
6. Aquila H, Link TA, Klingenberg M. The uncoupling protein from brown fat mitochondria is related to the mitochondrial ADP/ATP carrier. Analysis of sequence homologies and of folding of the protein in the membrane. *EMBO J* 4: 2369–2376, 1985. doi:10.1002/j.1460-2075.1985.tb03941.x.
7. Aquila H, Misra D, Eulitz M, Klingenberg M. Complete amino acid sequence of the ADP/ATP carrier from beef heart mitochondria. *Hoppe Seylers Z Physiol Chem* 363: 345–349, 1982.
8. Austin J, Aprille JR. Carboxyatractyloside-insensitive influx and efflux of adenine nucleotides in rat liver mitochondria. *J Biol Chem* 259: 154–160, 1984.
9. Azzi A, Chappell JB, and Robinson BH. Penetration of the mitochondrial membrane by glutamate and aspartate. *Biochem Biophys Res Commun* 29: 148–152, 1967. doi:10.1016/0006-291x(67)90556-6.
10. Bamber L, Harding M, Butler PJG, Kunji ERS. Yeast mitochondrial ADP/ATP carriers are monomeric in detergents. *Proc Natl Acad Sci USA* 103: 16224–16229, 2006. doi:10.1073/pnas.0607640103.
11. Bamber L, Harding M, Monné M, Slotboom DJ, Kunji ERS. The yeast mitochondrial ADP/ATP carrier functions as a monomer in mitochondrial membranes. *Proc Natl Acad Sci USA* 104: 10830–10834, 2007. doi:10.1073/pnas.0703969104.
12. Bamber L, Slotboom DJ, Kunji ERS. Yeast mitochondrial ADP/ATP carriers are monomeric in detergents as demonstrated by differential affinity purification. *J Mol Biol* 371: 388–395, 2007. doi:10.1016/j.jmb.2007.05.072.
13. Bertholet AM, Chouchani ET, Kazak L, Angelin A, Fedorenko A, Long JZ, Vidoni S, Garrity R, Cho J, Terada N, Wallace DC, Spiegelman BM, Kirichok Y. H⁺ transport is an integral function of the mitochondrial ADP/ATP carrier. *Nature* 571: 515–520, 2019. doi:10.1038/s41586-019-1400-3.
14. Beyer K, Klingenberg M. ADP/ATP carrier protein from beef heart mitochondria has high amounts of tightly bound cardiolipin, as revealed by 31P nuclear magnetic resonance. *Biochemistry* 24: 3821–3826, 1985. doi:10.1021/bi00336a001.
15. Bhoj EJ, Li M, Ahrens-Nicklas R, Pyle LC, Wang J, Zhang VW, Clarke C, Wong LJ, Sondheimer N, Ficcioglu C, Yudkoff M. Pathologic variants of the mitochondrial phosphate carrier SLC25A3: two new patients and expansion of the cardiomyopathy/skeletal myopathy phenotype with and without lactic acidosis. *JIMD Rep* 19: 59–66, 2015. doi:10.1007/8904_2014_364.
16. Bisaccia F, De Palma A, Palmieri F. Identification and purification of the aspartate/glutamate carrier from bovine heart mitochondria. *Biochim Biophys Acta* 1106: 291–296, 1992. doi:10.1016/0005-2736(92)90008-A.
17. Boczonadi V, King MS, Smith AC, Olahova M, Bansagi B, Roos A, Eyassu F, Borchers R, Ramesh V, Lochmüller H, Polvikoski T, Whittaker RG, Pyle A, Griffin H, Taylor RW, Chinnery PF, Robinson AJ, Kunji ERS, Horvath R. Mitochondrial oxodipicolinate carrier deficiency is associated with mitochondrial DNA depletion and spinal muscular atrophy-like disease. *Genet Med* 20: 1224–1235, 2018. doi:10.1038/gim.2017.251. A correction for this article is available at https://doi.org/10.1038/s41436-019-0506-1.
18. Brand MD, Chappell JB. Glutamate and aspartate transport in rat brain mitochondria. *Biochem J* 140: 205–210, 1974. doi:10.1042/bj1400205.
19. Bricker DK, Taylor EB, Schell JC, Orsak T, Boutron A, Chen YC, Cox JE, Cardon CM, Van Vranken JG, Dephoure N, Redin C, Boudina S, Gygi SP, Brivet M, Thummel CS, Rutter J. A mitochondrial pyruvate carrier required for pyruvate uptake in yeast, *Drosophila*, and humans. *Science* 337: 96–100, 2012. doi:10.1126/science.1218099.
20. Brustovetsky N, Klingenberg M. The reconstituted ADP/ATP carrier can mediate H⁺ transport by free fatty acids, which is further stimulated by mersalyl. *J Biol Chem* 269: 27329–27336, 1994.
21. Buffet A, Morin A, Castro-Vega LJ, Habarou F, Lussey-Lepoutre C, Letouzé E, Lefebvre H, Guilhem I, Haissaguerre M, Raingard I, Padilla-Girola M, Tran T, Tchara L, Bertherat J, Amar L, Ottolenghi C, Burnichon N, Gimenez-Roqueplo AP, Favier J. Germline mutations in the mitochondrial 2-oxoglutarate/malate carrier SLC25A11 gene confer a predisposition to metastatic paragangliomas. *Cancer Res* 78: 1914–1922, 2018. doi:10.1158/0008-5472.CAN-17-2463.
22. Camacho JA, Obie C, Biery B, Goodman BK, Hu CA, Almashanu S, Steel G, Casey R, Lambert M, Mitchell GA, Valle D. Hyperornithinaemia-hyperammonaemia-homocitrullinuria syndrome is caused by mutations in a gene encoding a mitochondrial ornithine transporter. *Nat Genet* 22: 151–158, 1999. doi:10.1038/9658.
23. Cannon B, Nedergaard J. What Ignites UCP1? *Cell Metab* 26: 697–698, 2017. doi:10.1016/j.cmet.2017.10.012.
24. Cappello AR, Curcio R, Valeria Miniero D, Stipani I, Robinson AJ, Kunji ERS, Palmieri F. Functional and structural role of amino acid residues in the even-numbered transmembrane alpha-helices of the bovine mitochondrial oxoglutarate carrier. *J Mol Biol* 363: 51–62, 2006. doi:10.1016/j.jmb.2006.08.041.
25. Cappello AR, Miniero DV, Curcio R, Ludovico A, Daddabbo L, Stipani I, Robinson AJ, Kunji ERS, Palmieri F. Functional and structural role of amino acid residues in the odd-numbered transmembrane alpha-helices of the bovine mitochondrial oxoglutarate carrier. *J Mol Biol* 369: 400–412, 2007. doi:10.1016/j.jmb.2007.03.048.
26. Cavero S, Traba J, Del Arco A, Satrustegui J. The calcium-dependent ATP-Mg/Pi mitochondrial carrier is a target of glucose-induced calcium signalling in *Saccharomyces cerevisiae*. *Biochem J* 392: 537–544, 2005. doi:10.1042/BJ20050806.
27. Chaouch A, Porcelli V, Cox D, Edvardson S, Scarcia P, De Grassi A, Pierri CL, Cossins J, Laval SH, Griffin H, Müller JS, Evangelista T, Töpf A, Abicht A, Huebner A, von der Hagen M, Bushby K, Straub V, Horvath R, Elpeleg O, Palace J, Senderek J, Beeson D, Palmieri L, Lochmüller H. Mutations in the mitochondrial citrate carrier SLC25A1 are associated with impaired neuromuscular transmission. *J Neuromuscul Dis* 1: 75–90, 2014. doi:10.3233/JND-140021.
28. Chorev DS, Baker LA, Wu D, Beilsten-Edmands V, Rouse SL, Zeev-Ben-Mordehai T, Jiko C, Samudrin F, Gerle C, Khalid S, Stewart AG, Matthews SJ, Grünewald K, Robinson CV. Protein assemblies ejected directly from native membranes yield complexes for mass spectrometry. *Science* 362: 829–834, 2018. doi:10.1126/science.aau0976.
29. Chung J, Anderson SA, Gwynn B, Deck KM, Chen MJ, Langer NB, Shaw GC, Huston NC, Boyer LF, Datta S, Paradar PN, Li L, Wei Z, Lambert AJ, Sahr K, Wittig JG, Chen W, Lu W, Galy B, Schläeger TM, Hentze MW, Ward DM, Kaplan J, Eisenstein RS, Peters LL, Paw BH. Iron regulatory protein-1 protects against mitoferrin-1-deficient porphyria. *J Biol Chem* 289: 7835–7843, 2014. doi:10.1074/jbc.M114.547778.
30. Cohen I, Staretz-Chacham O, Wormser O, Perez Y, Saada A, Kadir R, Birk OS. A novel homozygous SLC25A1 mutation with impaired mitochondrial complex V: Possible phenotypic expansion. *Am J Med Genet A* 176: 330–336, 2018. doi:10.1002/ajmg.a.38574.
31. Cozens AL, Runswick MJ, Walker JE. DNA sequences of two expressed genes for human mitochondrial ADP/ATP translocase. *J Mol Biol* 206: 261–280, 1989. doi:10.1016/0022-2836(89)90477-4.
32. Crichton PG, Harding M, Ruprecht JJ, Lee Y, Kunji ERS. Lipid, detergent, and Coomassie Blue G-250 affect the migration of small membrane proteins in blue native gels: mitochondrial carriers migrate as monomers not dimers. *J Biol Chem* 288: 22163–22173, 2013. doi:10.1074/jbc.M113.484329.
33. Crichton PG, Lee Y, Kunji ERS. The molecular features of uncoupling protein 1 support a conventional mitochondrial carrier-like mechanism. *Biochimie* 134: 35–50, 2017. doi:10.1016/j.biochi.2016.12.016.
34. Crichton PG, Lee Y, Ruprecht JJ, Cerson E, Thangaratnarah C, King MS, Kunji ERS. Trends in thermostability provide information on the nature of substrate, inhibitor, and lipid interactions with mitochondrial carriers. *J Biol Chem* 290: 8206–8217, 2015. doi:10.1074/jbc.M114.616607.
35. Dehez F, Pebay-Peyroula E, Chipot C. Binding of ADP in the mitochondrial ADP/ATP carrier is driven by an electrostatic funnel. *J Am Chem Soc* 130: 12725–12733, 2008. doi:10.1021/ja8033087.
36. del Arco A, Satrustegui J. Identification of a novel human subfamily of mitochondrial carriers with calcium-binding domains. *J Biol Chem* 279: 24701–24713, 2004. doi:10.1074/jbc.M401417200.
37. Di Noia MA, Todisco S, Cirigliano A, Rinaldi T, Agrimi G, Iacobazzi V, Palmieri F. The human SLC25A33 and SLC25A36 genes of solute carrier family 25 encode two mitochondrial pyrimidine nucleotide transporters. *J Biol Chem* 289: 33137–33148, 2014. doi:10.1074/jbc.M114.618088.
38. Dimmock D, Maranda B, Dionisi-Vici C, Wang J, Kleppe S, Fiermonte G, Bai R, Hainline B, Hamosh A, O'Brien WE, Scaglia F, Wong LJ. Citrin deficiency, a perplexing global disorder. *Mol Genet Metab* 96: 44–49, 2009. doi:10.1016/j.ymgme.2008.10.007.
39. Dolce V, Fiermonte G, Runswick MJ, Palmieri F, Walker JE. The human mitochondrial deoxynucleotide carrier and its role in the toxicity of nucleoside antivirals. *Proc Natl Acad Sci USA* 98: 2284–2288, 2001. doi:10.1073/pnas.031430998.

40. Dolce V, Scarcia P, Iacopetta D, Palmieri F. A fourth ADP/ATP carrier isoform in man: identification, bacterial expression, functional characterization and tissue distribution. *FEBS Lett* 579: 633–637, 2005. doi:10.1016/j.febslet.2004.12.034.
41. Duncan AL, Ruprecht JJ, Kunji ERS, Robinson AJ. Cardiolipin dynamics and binding to conserved residues in the mitochondrial ADP/ATP carrier. *Biochim Biophys Acta Biomembr* 1860: 1035–1045, 2018. doi:10.1016/j.bbamem.2018.01.017.
42. Edvardson S, Porcelli V, J alas C, Soiferman D, Kellner Y, Shaag A, Korman SH, Pierr i CL, Scarcia P, Fraenkel ND, Segel R, Schechter A, Frumkin A, Pines O, Saada A, Palmieri L, Elpeleg O. Agenesis of corpus callosum and optic nerve hypoplasia due to mutations in SLC25A1 encoding the mitochondrial citrate transporter. *J Med Genet* 50: 240–245, 2013. doi:10.1136/jmedgenet-2012-101485.
43. Ehmke N, Graul-Neumann L, Smorag L, Koenig R, Segebrecht L, Magoulas P, Scaglia F, Kilic E, Hennig AF, Adolphs N, Saha N, Fauler B, Kalscheuer VM, Hennig F, Altmüller J, Netzer C, Thiele H, Nürnberg P, Yigit G, Jäger M, Hecht J, Krüger U, Mielke T, Krawitz PM, Horn D, Schuelke M, Mundlos S, Bacino CA, Bonnen PE, Wollnik B, Fischer-Zirnsak B, Kornak U. De novo mutations in SLC25A24 cause a craniosynostosis syndrome with hypertrichosis, progeroid appearance, and mitochondrial dysfunction. *Am J Hum Genet* 101: 833–843, 2017. doi:10.1016/j.ajhg.2017.09.016.
44. Fedorenko A, Lishko PV, Kirichok Y. Mechanism of fatty-acid-dependent UCP1 uncoupling in brown fat mitochondria. *Cell* 151: 400–413, 2012. doi:10.1016/j.cell.2012.09.010.
45. Fiermonte G, De Leonardi F, Todisco S, Palmieri L, Lasorsa FM, Palmieri F. Identification of the mitochondrial ATP-Mg/Pi transporter. Bacterial expression, reconstitution, functional characterization, and tissue distribution. *J Biol Chem* 279: 30722–30730, 2004. doi:10.1074/jbc.M400445200.
46. Fiermonte G, Dolce V, Arrigoni R, Runswick MJ, Walker JE, Palmieri F. Organization and sequence of the gene for the human mitochondrial dicarboxylate carrier: evolution of the carrier family. *Biochem J* 344: 953–960, 1999. doi:10.1042/bj3440953.
47. Fiermonte G, Dolce V, David L, Santorelli FM, Dionisi-Vici C, Palmieri F, Walker JE. The mitochondrial ornithine transporter. Bacterial expression, reconstitution, functional characterization, and tissue distribution of two human isoforms. *J Biol Chem* 278: 32778–32783, 2003. doi:10.1074/jbc.M302317200.
48. Fiermonte G, Dolce V, Palmieri L, Ventura M, Runswick MJ, Palmieri F, Walker JE. Identification of the human mitochondrial oxodicarboxylate carrier. Bacterial expression, reconstitution, functional characterization, tissue distribution, and chromosomal location. *J Biol Chem* 276: 8225–8230, 2001. doi:10.1074/jbc.M009607200.
49. Fiermonte G, Palmieri L, Dolce V, Lasorsa FM, Palmieri F, Runswick MJ, Walker JE. The sequence, bacterial expression, and functional reconstitution of the rat mitochondrial dicarboxylate transporter cloned via distant homologs in yeast and *Caenorhabditis elegans*. *J Biol Chem* 273: 24754–24759, 1998. doi:10.1074/jbc.273.38.24754.
50. Fiermonte G, Palmieri L, Todisco S, Agrimi G, Palmieri F, Walker JE. Identification of the mitochondrial glutamate transporter. Bacterial expression, reconstitution, functional characterization, and tissue distribution of two human isoforms. *J Biol Chem* 277: 19289–19294, 2002. doi:10.1074/jbc.M201572200.
51. Fiermonte G, Paradies E, Todisco S, Marobbio CM, Palmieri F. A novel member of solute carrier family 25 (SLC25A42) is a transporter of coenzyme A and adenosine 3',5'-diphosphate in human mitochondria. *J Biol Chem* 284: 18152–18159, 2009. doi:10.1074/jbc.M109.014118.
52. Floyd S, Favre C, Lasorsa FM, Leahy M, Trigiant e G, Stroebel P, Marx A, Loughran G, O'Callaghan K, Marobbio CM, Slotboom DJ, Kunji ERS, Palmieri F, O'Connor R. The insulin-like growth factor-I-mTOR signaling pathway induces the mitochondrial pyrimidine nucleotide carrier to promote cell growth. *Mol Biol Cell* 18: 3545–3555, 2007. doi:10.1091/mbc.e06-12-1109.
53. Fontanesi F, Palmieri L, Scarcia P, Lodi T, Donnini C, Limongelli A, Tiranti V, Zeviani M, Ferrero I, Viola AM. Mutations in AAC2, equivalent to human adPEO-associated ANT1 mutations, lead to defective oxidative phosphorylation in *Saccharomyces cerevisiae* and affect mitochondrial DNA stability. *Hum Mol Genet* 13: 923–934, 2004. doi:10.1093/hmg/ddh108.
54. Gorgoglione R, Porcelli V, Santoro A, Daddabbo L, Voza A, Monné M, Di Noia MA, Palmieri L, Fiermonte G, Palmieri F. The human uncoupling proteins 5 and 6 (UCP5/SLC25A14 and UCP6/SLC25A30) transport sulfur oxyanions, phosphate and dicarboxylates. *Biochim Biophys Acta Bioenerg* 1860: 724–733, 2019. doi:10.1016/j.bbabi.2019.07.010.
55. Goubert E, Mircheva Y, Lasorsa FM, Melon C, Profilo E, Sutura J, Becq H, Palmieri F, Palmieri L, Anikstejn L, Molinari F. Inhibition of the mitochondrial glutamate carrier SLC25A22 in astrocytes leads to intracellular glutamate accumulation. *Front Cell Neurosci* 11: 149, 2017. doi:10.3389/fncel.2017.00149.
56. Haghghi A, Haack TB, Atiq M, Mottaghi H, Haghghi-Kakhki H, Bashir RA, Ahting U, Feichtinger RG, Mayr JA, Rötig A, Lebre AS, Klopstock T, Dworschak A, Pulido N, Saeed MA, Saleh-Gohari N, Holzerova E, Chinnery PF, Taylor RW, Prokisch H. Sengers syndrome: six novel AGK mutations in seven new families and review of the phenotypic and mutational spectrum of 29 patients. *Orphanet J Rare Dis* 9: 119, 2014. doi:10.1186/s13023-014-0119-3.
57. Haitina T, Lindblom J, Renström T, Fredriksson R. Fourteen novel human members of mitochondrial solute carrier family 25 (SLC25) widely expressed in the central nervous system. *Genomics* 88: 779–790, 2006. doi:10.1016/j.ygeno.2006.06.016.
58. Harborne SP, King MS, Crichton PG, Kunji ERS. Calcium regulation of the human mitochondrial ATP-Mg/Pi carrier SLC25A24 uses a locking pin mechanism. *Sci Rep* 7: 45383, 2017. doi:10.1038/srep45383.
59. Harborne SP, Ruprecht JJ, Kunji ERS. Calcium-induced conformational changes in the regulatory domain of the human mitochondrial ATP-Mg/Pi carrier. *Biochim Biophys Acta* 1847: 1245–1253, 2015. doi:10.1016/j.bbabi.2015.07.002.
60. Harborne SPD, Kunji ERS. Calcium-regulated mitochondrial ATP-Mg/Pi carriers evolved from a fusion of an EF-hand regulatory domain with a mitochondrial ADP/ATP carrier-like domain. *IUBMB Life* 70: 1222–1232, 2018. doi:10.1002/iub.1931.
61. Herzig S, Raemy E, Montessuit S, Veuthey JL, Zamboni N, Westermann B, Kunji ERS, Martinou JC. Identification and functional expression of the mitochondrial pyruvate carrier. *Science* 337: 93–96, 2012. doi:10.1126/science.1218530.
62. Hirst J, Kunji ERS, Walker JE. Comment on "Protein assemblies ejected directly from native membranes yield complexes for mass spectrometry". *Science* 366: eaaw9830, 2019. doi:10.1126/science.aaw9830.
63. Hoffmann B, Stöckl A, Schlame M, Beyer K, Klingenberg M. The reconstituted ADP/ATP carrier activity has an absolute requirement for cardiolipin as shown in cysteine mutants. *J Biol Chem* 269: 1940–1944, 1994.
64. Hofherr A, Seger C, Fitzpatrick F, Busch T, Michel E, Luan J, Osterried L, Linden F, Kramer-Zucker A, Wakimoto B, Schütze C, Wiedemann N, Artati A, Adamski J, Walz G, Kunji ERS, Montell C, Watnick T, Köttgen M. The mitochondrial transporter SLC25A25 links ciliary TRPP2 signaling and cellular metabolism. *PLoS Biol* 16: e2005651, 2018. doi:10.1371/journal.pbio.2005651.
65. Hommes FA, Roesel RA, Metoki K, Hartlage PL, Dyken PR. Studies on a case of HHH-syndrome (hyperammonemia, hyperornithinemia, homocitrullinuria). *Neuropediatrics* 17: 48–52, 1986. doi:10.1055/s-2008-1052499.
66. Huizing M, Iacobazzi V, Ijst L, Savelkoul P, Ruitenbeek W, van den Heuvel L, Indiveri C, Smeitink J, Trijbels F, Wanders R, Palmieri F. Cloning of the human carnitine-acylcarnitine carrier cDNA and identification of the molecular defect in a patient. *Am J Hum Genet* 61: 1239–1245, 1997. doi:10.1086/301628.
67. Huizing M, Ruitenbeek W, van den Heuvel LP, Dolce V, Iacobazzi V, Smeitink JA, Palmieri F, Trijbels JM. Human mitochondrial transmembrane metabolite carriers: tissue distribution and its implication for mitochondrial disorders. *J Bioenerg Biomembr* 30: 277–284, 1998. doi:10.1023/A:1020501021222.
68. Iacobazzi V, Lauria G, Palmieri F. Organization and sequence of the human gene for the mitochondrial citrate transport protein. *DNA Seq* 7: 127–139, 1997. doi:10.3109/10425179709034029.
69. Iacobazzi V, Palmieri F, Runswick MJ, Walker JE. Sequences of the human and bovine genes for the mitochondrial 2-oxoglutarate carrier. *DNA Seq* 3: 79–88, 1992. doi:10.3109/10425179209034000.
70. Indiveri C, Tonazzi A, Palmieri F. Identification and purification of the carnitine carrier from rat liver mitochondria. *Biochim Biophys Acta* 1020: 81–86, 1990. doi:10.1016/0005-2728(90)90096-M.
71. Janer A, Prudent J, Paupe V, Fahiminiya S, Majewski J, Sgarioto N, Des Rosiers C, Forest A, Lin ZY, Gingras AC, Mitchell G, McBride HM, Shoubridge EA. SLC25A46 is required for mitochondrial lipid homeostasis and cristae maintenance and is responsible for Leigh syndrome. *EMBO Mol Med* 8: 1019–1038, 2016. doi:10.15252/emmm.201506159.
72. Jezek P, Jaburek M, and Garlid KD. Channel character of uncoupling protein-mediated transport. *FEBS Lett* 584: 2135–2141, 2010. doi:10.1016/j.febslet.2010.02.068.
73. Johnson RN, Chappell JB. The transport of inorganic phosphate by the mitochondrial dicarboxylate carrier. *Biochem J* 134: 769–774, 1973. doi:10.1042/bj1340769.
74. Jordens EZ, Palmieri L, Huizing M, van den Heuvel LP, Sengers RC, Dörner A, Ruitenbeek W, Trijbels FJ, Vålsson J, Sigfusson G, Palmieri F, Smeitink JA. Adenine nucleotide translocator 1 deficiency associated with Sengers syndrome. *Ann Neurol* 52: 95–99, 2002. doi:10.1002/ana.10214.
75. Kadenbach B, Mende P, Kolbe HV, Stipani I, Palmieri F. The mitochondrial phosphate carrier has an essential requirement for cardiolipin. *FEBS Lett* 139: 109–112, 1982. doi:10.1016/0014-5793(82)80498-5.
76. Kaplan RS, Mayor JA, Johnston N, Oliveira DL. Purification and characterization of the reconstitutively active tricarboxylate transporter from rat liver mitochondria. *J Biol Chem* 265: 13379–13385, 1990.

77. Kaukonen J, Juselius JK, Tiranti V, Kytälä A, Zeviani M, Comi GP, Keränen S, Peltonen L, Suomalainen A. Role of adenine nucleotide translocator 1 in mtDNA maintenance. *Science* 289: 782–785, 2000. doi:10.1126/science.289.5480.782.
78. Kelley RI, Robinson D, Puffenberger EG, Strauss KA, Morton DH. Amish lethal microcephaly: a new metabolic disorder with severe congenital microcephaly and 2-ketoglutaric aciduria. *Am J Med Genet* 112: 318–326, 2002. doi:10.1002/ajmg.10529.
79. Khan S, Ansar M, Khan AK, Shah K, Muhammad N, Shahzad S, Nickerson DA, Bamshad MJ, Santos-Cortez RLP, Leal SM, Ahmad W. A homozygous missense mutation in SLC25A16 associated with autosomal recessive isolated fingernail dysplasia in a Pakistani family. *Br J Dermatol* 178: 556–558, 2018. doi:10.1111/bjd.15661.
80. King MS, Kerr M, Crichton PG, Springett R, Kunji ERS. Formation of a cytoplasmic salt bridge network in the matrix state is a fundamental step in the transport mechanism of the mitochondrial ADP/ATP carrier. *Biochim Biophys Acta* 1857: 14–22, 2016. doi:10.1016/j.bbabi.2015.09.013. A corrigendum for this article is available at <https://doi.org/10.1016/j.bbabi.2016.09.013>.
81. King MS, Thompson K, Hopton S, He L, Kunji ERS, Taylor RW, Ortiz-Gonzalez XR. Expanding the phenotype of de novo SLC25A4-linked mitochondrial disease to include mild myopathy. *Neurol Genet* 4: e256, 2018. doi:10.1212/NXG.0000000000000256.
82. Kishita Y, Pajak A, Bolar NA, Marobbio CM, Maffezzini C, Miniero DV, Monné M, Kohda M, Stranneheim H, Murayama K, Naess K, Lesko N, Bruhn H, Mourier A, Wibom R, Nennesmo I, Jesspers A, Govaert P, Ohtake A, Van Laer L, Loeys BL, Freyer C, Palmieri F, Wredenberg A, Okazaki Y, Wedell A. Intra-mitochondrial methylation deficiency due to mutations in SLC25A26. *Am J Hum Genet* 97: 761–768, 2015. doi:10.1016/j.ajhg.2015.09.013.
83. Klingenberg M. The ADP and ATP transport in mitochondria and its carrier. *Biochim Biophys Acta* 1778: 1978–2021, 2008. doi:10.1016/j.bbame.2008.04.011.
84. Klingenberg M. UCP1 - A sophisticated energy valve. *Biochimie* 134: 19–27, 2017. doi:10.1016/j.biochi.2016.10.012.
85. Knight SAB, Yoon H, Pandey AK, Pain J, Pain D, Dancis A. Splitting the functions of Rim2, a mitochondrial iron/pyrimidine carrier. *Mitochondrion* 47: 256–265, 2019. doi:10.1016/j.mito.2018.12.005.
86. Kobayashi K, Sinasac DS, Iijima M, Boright AP, Begum L, Lee JR, Yasuda T, Ikeda S, Hirano R, Terazono H, Crackower MA, Kondo I, Tsui LC, Scherer SW, Saheki T. The gene mutated in adult-onset type II citrullinaemia encodes a putative mitochondrial carrier protein. *Nat Genet* 22: 159–163, 1999. doi:10.1038/9667.
87. Kory N, Wyant GA, Prakash G, Uit de Bos J, Bottanelli F, Pacold ME, Chan SH, Lewis CA, Wang T, Keys HR, Guo YE, Sabatini DM. SFXN1 is a mitochondrial serine transporter required for one-carbon metabolism. *Science* 362: eaat9528, 2018. doi:10.1126/science.aat9528.
88. Ku DH, Kagan J, Chen ST, Chang CD, Baserga R, Wurzel J. The human fibroblast adenine nucleotide translocator gene. Molecular cloning and sequence. *J Biol Chem* 265: 16060–16063, 1990.
89. Kunji ERS. *Structural and Mechanistic Aspects of Mitochondrial Transport Proteins*. Amsterdam, The Netherlands: Elsevier, 2012, p. 174–205.
90. Kunji ERS, Aleksandrova A, King MS, Majd H, Ashton VL, Cerson E, Springett R, Kibalchenko M, Tavoulari S, Crichton PG, Ruprecht JJ. The transport mechanism of the mitochondrial ADP/ATP carrier. *Biochim Biophys Acta* 1863: 2379–2393, 2016. doi:10.1016/j.bbamcr.2016.03.015. A corrigendum for this article is available at <https://doi.org/10.1016/j.bbamcr.2016.09.021>.
91. Kunji ERS, Crichton PG. Mitochondrial carriers function as monomers. *Biochim Biophys Acta* 1797: 817–831, 2010. doi:10.1016/j.bbabi.2010.03.023.
92. Kunji ERS, Harding M. Projection structure of the atractyloside-inhibited mitochondrial ADP/ATP carrier of *Saccharomyces cerevisiae*. *J Biol Chem* 278: 36985–36988, 2003. doi:10.1074/jbc.C300304200.
93. Kunji ERS, Harding M, Butler PJG, Akamine P. Determination of the molecular mass and dimensions of membrane proteins by size exclusion chromatography. *Methods* 46: 62–72, 2008. doi:10.1016/j.jymeth.2008.10.020.
94. Kunji ERS, Robinson AJ. The conserved substrate binding site of mitochondrial carriers. *Biochim Biophys Acta* 1757: 1237–1248, 2006. doi:10.1016/j.bbabi.2006.03.021.
95. Kunji ERS, Robinson AJ. Coupling of proton and substrate translocation in the transport cycle of mitochondrial carriers. *Curr Opin Struct Biol* 20: 440–447, 2010. doi:10.1016/j.sbi.2010.06.004.
96. Lee Y, Willers C, Kunji ERS, Crichton PG. Uncoupling protein 1 binds one nucleotide per monomer and is stabilized by tightly bound cardiolipin. *Proc Natl Acad Sci USA* 112: 6973–6978, 2015. doi:10.1073/pnas.1503833112.
97. Lemattre C, Imbert-Bouteille M, Gatinois V, Benit P, Sanchez E, Guignard T, Tran Mau-Them F, Haquet E, Rivier F, Carme E, Roubertie A, Boland A, Lechner D, Meyer V, Thevenon J, Duffourd Y, Rivière JB, Deleuze JF, Wells C, Molinari F, Rustin P, Blanchet P, Geneviève D. Report on three additional patients and genotype-phenotype correlation in SLC25A22-related disorders group. *Eur J Hum Genet* 27: 1692–1700, 2019. doi:10.1038/s41431-019-0433-2.
98. Lill R, Kispal G. Mitochondrial ABC transporters. *Res Microbiol* 152: 331–340, 2001. doi:10.1016/S0923-2508(01)01204-9.
99. Lindhurst MJ, Fiermonte G, Song S, Struys E, De Leonhardt F, Schwartzberg PL, Chen A, Castegna A, Verhoeven N, Mathews CK, Palmieri F, Biesecker LG. Knockout of SLC25A19 causes mitochondrial thiamine pyrophosphate depletion, embryonic lethality, CNS malformations, and anemia. *Proc Natl Acad Sci USA* 103: 15927–15932, 2006. doi:10.1073/pnas.0607661103.
100. Liu Y, Wang X, Chen XJ. Misfolding of mutant adenine nucleotide translocase in yeast supports a novel mechanism of Ant1-induced muscle diseases. *Mol Biol Cell* 26: 1985–1994, 2015. doi:10.1091/mbc.E15-01-0030.
101. Liu Y, Wang X, Coyne LP, Yang Y, Qi Y, Middleton FA, Chen XJ. Mitochondrial carrier protein overloading and misfolding induce aggregates and proteostatic adaptations in the cytosol. *Mol Biol Cell* 30: 1272–1284, 2019. doi:10.1091/mbc.E19-01-0046.
102. Lunetti P, Damiano F, De Benedetto G, Siculella L, Pennetta A, Muto L, Paradies E, Marobbio CM, Dolce V, Capobianco L. Characterization of human and yeast mitochondrial glycine carriers with implications for heme biosynthesis and anemia. *J Biol Chem* 291: 19746–19759, 2016. doi:10.1074/jbc.M116.736876.
103. Majd H, King MS, Smith AC, Kunji ERS. Pathogenic mutations of the human mitochondrial citrate carrier SLC25A1 lead to impaired citrate export required for lipid, dolichol, ubiquinone and sterol synthesis. *Biochim Biophys Acta Bioenerg* 1859: 1–7, 2018. doi:10.1016/j.bbabi.2017.10.002.
104. Mayr JA, Haack TB, Graf E, Zimmermann FA, Wieland T, Haberberger B, Superti-Furga A, Kirschner J, Steinmann B, Baumgartner MG, Moroni I, Lamantea E, Zeviani M, Rodenburg RJ, Smeitink J, Strom TM, Meitinger T, Sperl W, Prokisch H. Lack of the mitochondrial protein acylglycerol kinase causes Sengers syndrome. *Am J Hum Genet* 90: 314–320, 2012. doi:10.1016/j.ajhg.2011.12.005.
105. Mayr JA, Merkel O, Kohlwein SD, Gebhardt BR, Böhles H, Fötschl U, Koch J, Jaksch M, Lochmüller H, Horváth R, Freisinger P, Sperl W. Mitochondrial phosphate-carrier deficiency: a novel disorder of oxidative phosphorylation. *Am J Hum Genet* 80: 478–484, 2007. doi:10.1086/511788.
106. Mayr JA, Zimmermann FA, Horváth R, Schneider HC, Schoser B, Holinski-Feder E, Czermin B, Freisinger P, Sperl W. Deficiency of the mitochondrial phosphate carrier presenting as myopathy and cardiomyopathy in a family with three affected children. *Neuromuscul Disord* 21: 803–808, 2011. doi:10.1016/j.nmd.2011.06.005.
107. Mifsud J, Ravaut S, Krammer EM, Chipot C, Kunji ERS, Pebay-Peyroula E, Dehez F. The substrate specificity of the human ADP/ATP carrier AAC1. *Mol Membr Biol* 30: 160–168, 2013. doi:10.3109/09687688.2012.745175.
108. Miniero DV, Cappello AR, Curcio R, Ludovico A, Daddabbo L, Stipani I, Robinson AJ, Kunji ERS, Palmieri F. Functional and structural role of amino acid residues in the matrix alpha-helices, termini and cytosolic loops of the bovine mitochondrial oxoglutarate carrier. *Biochim Biophys Acta* 1807: 302–310, 2011. doi:10.1016/j.bbabi.2010.12.005.
109. Mizurari S, Miki S, Araki H, Takahashi K, Kotani H. Identification of dicarboxylate carrier SLC25A10 as malate transporter in de novo fatty acid synthesis. *J Biol Chem* 280: 32434–32441, 2005. doi:10.1074/jbc.M503152200.
110. Molinari F, Kaminska A, Fiermonte G, Boddart N, Raas-Rothschild A, Plouin P, Palmieri L, Brunelle F, Palmieri F, Dulac O, Munnich A, Colleaux L. Mutations in the mitochondrial glutamate carrier SLC25A22 in neonatal epileptic encephalopathy with suppression bursts. *Clin Genet* 76: 188–194, 2009. doi:10.1111/j.1399-0004.2009.01236.x.
111. Monné M, Miniero DV, Daddabbo L, Robinson AJ, Kunji ERS, Palmieri F. Substrate specificity of the two mitochondrial ornithine carriers can be swapped by single mutation in substrate binding site. *J Biol Chem* 287: 7925–7934, 2012. doi:10.1074/jbc.M111.324855.
112. Morciano P, Carrisi C, Capobianco L, Mannini L, Burgio G, Cestra G, De Benedetto GE, Corona DF, Musio A, Cenci G. A conserved role for the mitochondrial citrate transporter Sea/SLC25A1 in the maintenance of chromosome integrity. *Hum Mol Genet* 18: 4180–4188, 2009. doi:10.1093/hmg/ddp370.
113. Mühlhoff U, Stadler JA, Richhardt N, Seubert A, Eickhorst T, Schweyen RJ, Lill R, Wiesenberger G. A specific role of the yeast mitochondrial carriers MR53/4p in mitochondrial iron acquisition under iron-limiting conditions. *J Biol Chem* 278: 40612–40620, 2003. doi:10.1074/jbc.M307847200.
114. Mühlhausen C, Salomons GS, Lukacs Z, Struys EA, van der Knaap MS, Ullrich K, Santer R. Combined D2-/L2-hydroxyglutaric aciduria (SLC25A1 deficiency): clinical course and effects of citrate treatment. *J Inher Metab Dis* 37: 775–781, 2014. doi:10.1007/s10545-014-9702-y.
115. Neckelmann N, Li K, Wade RP, Shuster R, Wallace DC. cDNA sequence of a human skeletal muscle ADP/ATP translocator: lack of a leader peptide, divergence from a fibroblast translocator cDNA, and coevolution with mitochondrial DNA genes. *Proc Natl Acad Sci USA* 84: 7580–7584, 1987. doi:10.1073/pnas.84.21.7580.

116. Nedergaard J, Bengtsson T, Cannon B. Unexpected evidence for active brown adipose tissue in adult humans. *Am J Physiol Endocrinol Metab* 293: E444–E452, 2007. doi:10.1152/ajpendo.00691.2006.
117. Nelson DR, Felix CM, Swanson JM. Highly conserved charge-pair networks in the mitochondrial carrier family. *J Mol Biol* 277: 285–308, 1998. doi:10.1006/jmbi.1997.1594.
118. Nicholls DG. The bioenergetics of brown adipose tissue mitochondria. *FEBS Lett* 61: 103–110, 1976. doi:10.1016/0014-5793(76)81014-9.
119. Nicholls DG. The hunt for the molecular mechanism of brown fat thermogenesis. *Biochimie* 134: 9–18, 2017. doi:10.1016/j.biochi.2016.09.003.
120. Nicholls DG, Rial E. A history of the first uncoupling protein, UCP1. *J Bioenerg Biomembr* 31: 399–406, 1999. doi:10.1023/A:1005436121005.
121. Nota B, Struys EA, Pop A, Jansen EE, Fernandez Ojeda MR, Kanhai WA, Kranendijk M, van Dooren SJ, Bevoova MR, Sijstermans EA, Nieuwint AW, Barth M, Ben-Omran T, Hoffmann GF, de Lonlay P, McDonald MT, Meberg A, Muntax AC, Nuoffer JM, Parini R, Read MH, Renneberg A, Santer R, Strahleck T, van Schaftingen E, van der Knaap MS, Jakobs C, Salomons GS. Deficiency in SLC25A1, encoding the mitochondrial citrate carrier, causes combined D-2- and L-2-hydroxyglutaric aciduria. *Am J Hum Genet* 92: 627–631, 2013. doi:10.1016/j.ajhg.2013.03.009.
122. Okano Y, Ohura T, Sakamoto O, Inui A. Current treatment for citrin deficiency during NICCD and adaptation/compensation stages: strategy to prevent CTLN2. *Mol Genet Metab* 127: 175–183, 2019. doi:10.1016/j.ymgme.2019.06.004.
123. Palmieri F, Monné M. Discoveries, metabolic roles and diseases of mitochondrial carriers: A review. *Biochim Biophys Acta* 1863: 2362–2378, 2016. doi:10.1016/j.bbamcr.2016.03.007.
124. Palmieri F, Scarcia P, Monné M. Diseases caused by mutations in mitochondrial carrier genes SLC25: a review. *Biomolecules* 10: E655, 2020. doi:10.3390/biom10040655.
125. Palmieri L, Alberio S, Pisano I, Lodi T, Meznaric-Petrusa M, Zidar J, Santoro A, Scarcia P, Fontanesi F, Lamantea E, Ferrero I, Zeviani M. Complete loss-of-function of the heart/muscle-specific adenine nucleotide translocator is associated with mitochondrial myopathy and cardiomyopathy. *Hum Mol Genet* 14: 3079–3088, 2005. doi:10.1093/hmg/ddi341.
126. Palmieri L, Pardo B, Lasorsa FM, del Arco A, Kobayashi K, Iijima M, Runswick MJ, Walker JE, Saheki T, Satrustegui J, Palmieri F. Citrin and aralar1 are Ca(2+)-stimulated aspartate/glutamate transporters in mitochondria. *EMBO J* 20: 5060–5069, 2001. doi:10.1093/emboj/20.18.5060.
127. Paul BT, Tesfay L, Winkler CR, Torti FM, Torti SV. Sideroflexin 4 affects Fe-S cluster biogenesis, iron metabolism, mitochondrial respiration and heme biosynthetic enzymes. *Sci Rep* 9: 19634, 2019. doi:10.1038/s41598-019-55907-z.
128. Pebay-Peyroula E, Dahout-Gonzalez C, Kahn R, Trézéguet V, Lauquin GJ, Brandolin G. Structure of mitochondrial ADP/ATP carrier in complex with carboxyatractyloside. *Nature* 426: 39–44, 2003. doi:10.1038/nature02056.
129. Poduri A, Heinzen EL, Chitsazzadeh V, Lasorsa FM, Elhosary PC, LaCoursiere CM, Martin E, Yuskaitis CJ, Hill RS, Atabay KD, Barry B, Partlow JN, Bashiri FA, Zeidan RM, Elmalik SA, Kabiraj MM, Kothare S, Stöckberg T, McTague A, Kurian MA, Scheffer IE, Barkovich AJ, Palmieri F, Salih MA, Walsh CA. SLC25A22 is a novel gene for migrating partial seizures in infancy. *Ann Neurol* 74: 873–882, 2013. doi:10.1002/ana.23998.
130. Pop A, Williams M, Struys EA, Monné M, Jansen EEW, De Grassi A, Kanhai WA, Scarcia P, Ojeda MRF, Porcelli V, van Dooren SJM, Lennertz P, Nota B, Abdenur JE, Coman D, Das AM, El-Gharbawy A, Nuoffer JM, Polic B, Santer R, Weinhönd N, Zuccarelli B, Palmieri F, Palmieri L, Salomons GS. An overview of combined D-2- and L-2-hydroxyglutaric aciduria: functional analysis of CIC variants. *J Inher Metab Dis* 41: 169–180, 2018. doi:10.1007/s10545-017-0106-7.
131. Porcelli V, Fiermonte G, Longo A, Palmieri F. The human gene SLC25A29, of solute carrier family 25, encodes a mitochondrial transporter of basic amino acids. *J Biol Chem* 289: 13374–13384, 2014. doi:10.1074/jbc.M114.547448.
132. Prasun P, Young S, Salomons G, Werneke A, Jiang YH, Struys E, Paige M, Avantagegiati ML, McDonald M. Expanding the clinical spectrum of mitochondrial citrate carrier (SLC25A1) deficiency: facial dysmorphism in siblings with epileptic encephalopathy and combined D,L-2-hydroxyglutaric aciduria. *JIMD Rep* 19: 111–115, 2015. doi:10.1007/8904_2014_378.
133. Prohl C, Pelzer W, Diekert K, Kmita H, Bedekovics T, Kispal G, Lill R. The yeast mitochondrial carrier Leu5p and its human homologue Graves' disease protein are required for accumulation of coenzyme A in the matrix. *Mol Cell Biol* 21: 1089–1097, 2001. doi:10.1128/MCB.21.4.1089-1097.2001.
134. Rial E, Poustie A, Nicholls DG. Brown-adipose-tissue mitochondria: the regulation of the 32000-Mr uncoupling protein by fatty acids and purine nucleotides. *Eur J Biochem* 137: 197–203, 1983. doi:10.1111/j.1432-1033.1983.tb07815.x.
135. Robinson AJ, Kunji ER, Gross A. Mitochondrial carrier homolog 2 (MTC2): the recruitment and evolution of a mitochondrial carrier protein to a critical player in apoptosis. *Exp Cell Res* 318: 1316–1323, 2012. doi:10.1016/j.yexcr.2012.01.026.
136. Robinson AJ, Kunji ERS. Mitochondrial carriers in the cytoplasmic state have a common substrate binding site. *Proc Natl Acad Sci USA* 103: 2617–2622, 2006. doi:10.1073/pnas.0509994103.
137. Robinson AJ, Overy C, Kunji ERS. The mechanism of transport by mitochondrial carriers based on analysis of symmetry. *Proc Natl Acad Sci USA* 105: 17766–17771, 2008. doi:10.1073/pnas.0809580105.
138. Rodríguez-García ME, Cotrina-Vinagre FJ, Cruz-Rojo J, Garzón-Lorenzo L, Carnicero-Rodríguez P, Pozo JS, Martínez-Azorín F. A rare male patient with Fontaine progeroid syndrome caused by p.R217H de novo mutation in SLC25A24. *Am J Med Genet A* 176: 2479–2486, 2018. doi:10.1002/ajmg.a.40496.
139. Rottiers V, Francisco A, Platov M, Zaltsman Y, Ruggiero A, Lee SS, Gross A, Libert S. MTC2 is a conserved regulator of lipid homeostasis. *Obesity (Silver Spring)* 25: 616–625, 2017. doi:10.1002/oby.21751.
140. Runswick MJ, Powell SJ, Nyren P, Walker JE. Sequence of the bovine mitochondrial phosphate carrier protein: structural relationship to ADP/ATP translocase and the brown fat mitochondria uncoupling protein. *EMBO J* 6: 1367–1373, 1987. doi:10.1002/j.1460-2075.1987.tb02377.x.
141. Ruprecht JJ, Hellawell AM, Harding M, Crichton PG, McCoy AJ, Kunji ERS. Structures of yeast mitochondrial ADP/ATP carriers support a domain-based alternating-access transport mechanism. *Proc Natl Acad Sci USA* 111: E426–E434, 2014. doi:10.1073/pnas.1320692111.
142. Ruprecht JJ, King MS, Zögg T, Aleksandrova AA, Pardon E, Crichton PG, Steyaert J, Kunji ERS. The molecular mechanism of transport by the mitochondrial ADP/ATP carrier. *Cell* 176: 435–447.e15, 2019. doi:10.1016/j.cell.2018.11.025.
143. Ruprecht JJ, Kunji ERS. The SLC25 mitochondrial carrier family: structure and mechanism. *Trends Biochem Sci* 45: 244–258, 2020. doi:10.1016/j.tibs.2019.11.001.
144. Ruprecht JJ, Kunji ERS. Structural changes in the transport cycle of the mitochondrial ADP/ATP carrier. *Curr Opin Struct Biol* 57: 135–144, 2019. doi:10.1016/j.sbi.2019.03.029.
145. Ryu J, Ko JM, Shin CHA. A 9-year-old Korean girl with Fontaine progeroid syndrome: a case report with further phenotypical delineation and description of clinical course during long-term follow-up. *BMC Med Genet* 20: 188, 2019. doi:10.1186/s12881-019-0921-9.
146. Saheki T, Inoue K, Tushima A, Mutoh K, Kobayashi K. Citrin deficiency and current treatment concepts. *Mol Genet Metab* 100, Suppl 1: S59–S64, 2010. doi:10.1016/j.ymgme.2010.02.014.
147. Saraste M, Walker JE. Internal sequence repeats and the path of polypeptide in mitochondrial ADP/ATP translocase. *FEBS Lett* 144: 250–254, 1982. doi:10.1016/0014-5793(82)80648-0.
148. Schaedler TA, Faust B, Shintre CA, Carpenter EP, Srinivasan V, van Veen HW, Balk J. Structures and functions of mitochondrial ABC transporters. *Biochem Soc Trans* 43: 943–951, 2015. doi:10.1042/BST20150118.
149. Schiff M, Veauville-Merlié A, Su CH, Tzagoloff A, Rak M, Ogier de Baulny H, Boutron A, Smedts-Walters H, Romero NB, Rigal O, Rustin P, Vianey-Saban C, Acquaviva-Bourdain C. SLC25A32 mutations and riboflavin-responsive exercise intolerance. *N Engl J Med* 374: 795–797, 2016. doi:10.1056/NEJMc1513610.
150. Shamseldin HE, Smith LL, Kentab A, Alkhalidi H, Summers B, Alsedairy H, Xiong Y, Gupta VA, Alkuray FS. Mutation of the mitochondrial carrier SLC25A42 causes a novel form of mitochondrial myopathy in humans. *Hum Genet* 135: 21–30, 2016. doi:10.1007/s00439-015-1608-8.
151. Shaw GC, Cope JJ, Li L, Corson K, Hersey C, Ackermann GE, Gwynn B, Lambert AJ, Wingert RA, Traver D, Trede NS, Barut BA, Zhou Y, Minet E, Donovan A, Brownlie A, Balzan R, Weiss MJ, Peters LL, Kaplan J, Zon LI, Paw BH. Mitoferrin is essential for erythroid iron assimilation. *Nature* 440: 96–100, 2006. doi:10.1038/nature04512.
152. Smith A, McBride S, Marcadier JL, Michaud J, Al-Dirbashi OY, Schwartzentruber J, Beaulieu CL, Katz SL, Majewski J, Bulman DE, Geraghty MT, Harper ME, Chakraborty P, Lines MA; FORGE Canada Consortium. Severe neonatal presentation of mitochondrial citrate carrier (SLC25A1) deficiency. *JIMD Rep* 30: 73–79, 2016. doi:10.1007/8904_2016_536.
153. Smith VR, Fearnley IM, Walker JE. Altered chromatographic behaviour of mitochondrial ADP/ATP translocase induced by stabilization of the protein by binding of 6'-O-fluorescein-atractyloside. *Biochem J* 376: 757–763, 2003. doi:10.1042/bj20030942.
154. Spaan AN, Ijlst L, van Roermund CW, Wijburg FA, Wanders RJ, Waterham HR. Identification of the human mitochondrial FAD transporter and its potential role in multiple acyl-CoA dehydrogenase deficiency. *Mol Genet Metab* 86: 441–447, 2005. doi:10.1016/j.ymgme.2005.07.014.
155. Spiegel R, Shaag A, Edvardson S, Mandel H, Stepensky P, Shalev SA, Horovitz Y, Pines O, Elpeleg O. SLC25A19 mutation as a cause of neuropathy and bilateral striatal necrosis. *Ann Neurol* 66: 419–424, 2009. doi:10.1002/ana.21752.
156. Springett R, King MS, Crichton PG, Kunji ERS. Modelling the free energy profile of the mitochondrial ADP/ATP carrier. *Biochim Biophys Acta Bioenerg* 1858: 906–914, 2017. doi:10.1016/j.bbabi.2017.05.006.

157. Thangaratnarajah C, Ruprecht JJ, Kunji ERS. Calcium-induced conformational changes of the regulatory domain of human mitochondrial aspartate/glutamate carriers. *Nat Commun* 5: 5491, 2014. doi:10.1038/ncomms6491.
158. Thompson K, Majd H, Dallabona C, Reinson K, King MS, Alston CL, He L, Lodi T, Jones SA, Fattal-Valevski A, Fraenkel ND, Saada A, Haham A, Isohanni P, Vara R, Barbosa IA, Simpson MA, Deshpande C, Puusepp S, Bonnen PE, Rodenburg RJ, Suomalainen A, Öunap K, Elpeleg O, Ferrero I, McFarland R, Kunji ERS, Taylor RW. Recurrent de novo dominant mutations in SLC25A4 cause severe early-onset mitochondrial disease and loss of mitochondrial DNA copy number. *Am J Hum Genet* 99: 860–876, 2016. doi:10.1016/j.ajhg.2016.08.014. A correction for this article is available at <https://doi.org/10.1016/j.ajhg.2016.11.001>.
159. Titus SA, Moran RG. Retrovirally mediated complementation of the glyB phenotype. Cloning of a human gene encoding the carrier for entry of folates into mitochondria. *J Biol Chem* 275: 36811–36817, 2000. doi:10.1074/jbc.M005163200.
160. Traba J, Satrustegui J, Del Arco A. Characterization of SCaMC-3-Like/slc25a41 a novel calcium-independent mitochondrial ATP-Mg/Pi carrier. *Biochem J* 418: 125–133, 2008. doi:10.1042/BJ20081262.
161. Visser WF, van Roermund CW, Waterham HR, Wanders RJ. Identification of human PMP34 as a peroxisomal ATP transporter. *Biochem Biophys Res Commun* 299: 494–497, 2002. doi:10.1016/S0006-291X(02)02663-3.
162. Vozza A, Blanco E, Palmieri L, Palmieri F. Identification of the mitochondrial GTP/GDP transporter in *Saccharomyces cerevisiae*. *J Biol Chem* 279: 20850–20857, 2004. doi:10.1074/jbc.M313610200.
163. Vozza A, Parisi G, De Leonardis F, Lasorsa FM, Castegna A, Amorese D, Marmo R, Calcagnile VM, Palmieri L, Ricquier D, Paradies E, Scarcia P, Palmieri F, Bouillaud F, Fiermonte G. UCP2 transports C4 metabolites out of mitochondria, regulating glucose and glutamine oxidation. *Proc Natl Acad Sci USA* 111: 960–965, 2014. doi:10.1073/pnas.1317400111.
164. Wan J, Steffen J, Yourshaw M, Mamsa H, Andersen E, Rudnik-Schöneborn S, Pope K, Howell KB, McLean CA, Kornberg AJ, Joseph J, Lockhart PJ, Zerres K, Ryan MM, Nelson SF, Koehler CM, Jen JC. Loss of function of SLC25A46 causes lethal congenital pontocerebellar hypoplasia. *Brain* 139: 2877–2890, 2016. doi:10.1093/brain/aww212.
165. Wang Y, Langer NB, Shaw GC, Yang G, Li L, Kaplan J, Paw BH, Bloomer JR. Abnormal mitochondrial ferritin-1 expression in patients with erythropoietic protoporphyria. *Exp Hematol* 39: 784–794, 2011. doi:10.1016/j.exphem.2011.05.003.
166. Wang Y, Tajkhorshid E. Electrostatic funneling of substrate in mitochondrial inner membrane carriers. *Proc Natl Acad Sci USA* 105: 9598–9603, 2008. doi:10.1073/pnas.0801786105.
167. Wibom R, Lasorsa FM, Töhönen V, Barbaro M, Sterky FH, Kucinski T, Naess K, Jonsson M, Pierré CL, Palmieri F, Wedell A. AGC1 deficiency associated with global cerebral hypomyelination. *N Engl J Med* 361: 489–495, 2009. doi:10.1056/NEJMoa0900591.
168. Writzl K, Maver A, Kovačič L, Martínez-Valero P, Contreras L, Satrustegui J, Castori M, Faivre L, Lapunzina P, van Kuilenburg ABP, Radović S, Thauvin-Robinet C, Peterlin B, Del Arco A, Hennekam RC. De novo mutations in SLC25A24 cause a disorder characterized by early aging, bone dysplasia, characteristic face, and early demise. *Am J Hum Genet* 101: 844–855, 2017. doi:10.1016/j.ajhg.2017.09.017.
169. Yang Q, Brüscheweiler S, Chou JJ. A self-sequestered calmodulin-like Ca²⁺ sensor of mitochondrial SCaMC carrier and its implication to Ca²⁺-dependent ATP-Mg/P(i) transport. *Structure* 22: 209–217, 2014. doi:10.1016/j.str.2013.10.018.
170. Ye X, Xu J, Cheng C, Yin G, Zeng L, Ji C, Gu S, Xie Y, Mao Y. Isolation and characterization of a novel human putative anemia-related gene homologous to mouse sideroflexin. *Biochem Genet* 41: 119–125, 2003. doi:10.1023/A:1022026001114.
171. Yoneshiro T, Wang Q, Tajima K, Matsushita M, Maki H, Igarashi K, Dai Z, White PJ, McGarrah RW, Ilkayeva OR, Deleye Y, Oguri Y, Kuroda M, Ikeda K, Li H, Ueno A, Ohishi M, Ishikawa T, Kim K, Chen Y, Sponton CH, Pradhan RN, Majd H, Greiner VJ, Yoneshiro M, Brown Z, Chondronikola M, Takahashi H, Goto T, Kawada T, Sidossis L, Szoka FC, McManus MT, Saito M, Soga T, Kajimura S. BCAA catabolism in brown fat controls energy homeostasis through SLC25A44. *Nature* 572: 614–619, 2019. doi:10.1038/s41586-019-1503-x.
172. Zaltsman Y, Shachnai L, Yivgi-Ohana N, Schwarz M, Maryanovich M, Houtkooper RH, Vaz FM, De Leonardis F, Fiermonte G, Palmieri F, Gillissen B, Daniel PT, Jimenez E, Walsh S, Koehler CM, Roy SS, Walter L, Hajnóczky G, Gross A. MTCH2/MIMP is a major facilitator of tBID recruitment to mitochondria. *Nat Cell Biol* 12: 553–562, 2010. doi:10.1038/ncb2057.
173. Zarrilli R, Oates EL, McBride OW, Lerman MI, Chan JY, Santisteban P, Ursini MV, Notkins AL, Kohn LD. Sequence and chromosomal assignment of a novel cDNA identified by immunoscreening of a thyroid expression library: similarity to a family of mitochondrial solute carrier proteins. *Mol Endocrinol* 3: 1498–1505, 1989. doi:10.1210/mend-3-9-1498.