

University of Groningen

Methods of enhancing translocation of charged analytes through transmembrane protein pores

Maglia, Giovanni; Bayley, John Hagan Pryce

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2010

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Maglia, G., & Bayley, J. H. P. (2010). Methods of enhancing translocation of charged analytes through transmembrane protein pores. (Patent No. *WO2010055307*).

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 May 2010 (20.05.2010)

PCT

(10) International Publication Number
WO 2010/055307 A1

(51) International Patent Classification:

C07K 14/31 (2006.01) *G01N 33/68* (2006.01)
C12N 15/31 (2006.01) *G01N 27/06* (2006.01)
C12Q 1/68 (2006.01)

(21) International Application Number:

PCT/GB2009/002666

(22) International Filing Date:

13 November 2009 (13.11.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0820927.2 14 November 2008 (14.11.2008) GB

(71) Applicant (for all designated States except US): **ISIS INNOVATION LIMITED** [GB/GB]; Ewert House, Ewert Place, Summertown, Oxford OX2 7SG (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MAGLIA, Giovanni** [IT/GB]; University of Oxford, Department of Chemistry, South Parks Road, Oxford OX1 3QZ (GB). **BAYLEY, John, Hagan, Pryce** [GB/GB]; University of Oxford, Department of Chemistry, South Parks Road, Oxford OX1 3QZ (GB).

(74) Agent: **CHAPMAN, Lee, Phillip**; J.A. Kemp & Co., 14 South Square, Gray's Inn, London WC1R 5JJ (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2010/055307 A1

(54) Title: METHODS OF ENHANCING TRANSLOCATION OF CHARGED ANALYTES THROUGH TRANSMEMBRANE PROTEIN PORES

(57) Abstract: The invention relates to enhancing translocation of a charged analyte through a transmembrane protein pore. Translocation is enhanced by increasing the net opposing charge of the barrel or channel and/or entrance of the pore. The invention also relates to pores enhanced in accordance with the invention.

METHODS OF ENHANCING TRANSLOCATION OF CHARGED ANALYTES THROUGH TRANSMEMBRANE PROTEIN PORES

Field of the invention

The invention relates to enhancing translocation of a charged analyte through a transmembrane protein pore. Translocation is enhanced by increasing the net opposing charge of the barrel or channel and/or entrance of the pore. The invention also relates to a pore enhanced in accordance with the invention.

Background of the invention

Stochastic detection is an approach to sensing that relies on the observation of individual binding events between analyte molecules and a receptor. Stochastic sensors can be created by placing a single pore of nanometer dimensions in an insulating membrane and measuring voltage-driven ionic transport through the pore in the presence of analyte molecules. The frequency of occurrence of fluctuations in the current reveals the concentration of an analyte that binds within the pore. The identity of an analyte is revealed through its distinctive current signature, notably the duration and extent of current block (Braha, O., Walker, B., Cheley, S., Kasianowicz, J. J., Song, L., Gouaux, J. E., and Bayley, H. (1997) *Chem.Biol.* **4**, 497-505; and Bayley, H., and Cremer, P. S. (2001) *Nature* **413**, 226-230).

Engineered versions of the bacterial pore-forming toxin α -hemolysin (α -HL) have been used for stochastic sensing of many classes of molecules (Bayley, H., and Cremer, P. S. (2001) *Nature* **413**, 226-230; Shin, S.-H., Luchian, T., Cheley, S., Braha, O., and Bayley, H. (2002) *Angew.Chem.Int.Ed.* **41**, 3707-3709; and Guan, X., Gu, L.-Q., Cheley, S., Braha, O., and Bayley, H. (2005) *ChemBioChem* **6**, 1875-1881). In the course of these studies, it was found that attempts to engineer α -HL to bind small organic analytes directly can prove taxing, with rare examples of success (Guan, X., Gu, L.-Q., Cheley, S., Braha, O., and Bayley, H. (2005) *ChemBioChem* **6**, 1875-1881).

Summary of the invention

When analytes translocate through a transmembrane protein pore, they are capable of affecting the current flowing through the pore in a manner specific for that analyte. This allows the analyte to be detected using stochastic sensing. The analyte usually affects the current flowing through a pore by interacting with the pore. A distinctive current flows through the pore whenever the analyte interacts with the pore.

The inventors have surprisingly demonstrated that increasing the net opposing charge of the barrel or channel and/or entrance of a pore enhances the translocation of charged analytes through

the pore. For instance, increasing the net positive charge of the barrel or channel and/or entrance of a pore enhances the translocation of negatively charged analytes through the pore. Increasing the net opposing charge of the barrel or channel and/or entrance increases the frequency of translocation of the analyte through the pore. It also decreases the threshold voltage for translocation of the analyte through the pore. It also decreases the translocation speed of the analyte through the pore. Further, events in which the analyte enters the pore, but is not translocated, are almost eliminated. Pores enhanced in this way are useful tools for stochastic sensing of charged analytes, especially for sequencing nucleic acids.

The invention therefore provides a method of enhancing translocation of a charged analyte through a transmembrane protein pore, comprising:

- (a) increasing the net opposing charge of the barrel or channel and/or entrance of the pore; and
- (b) determining whether or not translocation of the analyte through the resulting pore is enhanced.

The invention also provides:

- use of an increase in net opposing charge of the barrel or channel and/or entrance of a transmembrane protein pore to enhance translocation of a charged analyte through the pore;
- a transmembrane protein pore enhanced by a method of the invention;
- a transmembrane protein pore in which the net opposing charge of its barrel or channel and/or entrance has been increased to enhance the translocation of a charged analyte through the pore;
- a polynucleotide encoding a transmembrane protein pore of the invention;
- a polynucleotide encoding a transmembrane protein pore subunit having the sequence of SEQ ID NO: 6, 8 or 10 or a variant thereof;
- a method of determining presence or absence of an analyte in a sample, comprising:
 - (a) contacting the sample with a transmembrane protein pore of the invention under conditions that allow the analyte, if present, to translocate through and interact with the pore; and
 - (b) measuring the current passing through the pore during the interaction and thereby determining the presence or absence of the analyte;
- a method of sequencing a target nucleic acid sequence, comprising:
 - (a) pushing or pulling the target sequence through a transmembrane protein pore of the invention so that a proportion of the nucleotides in the target sequence interact with the pore; and

(b) measuring the current passing through the pore during each interaction and thereby determining the sequence of the target sequence; and

- a kit for sequencing a nucleic acid, comprising a transmembrane protein pore of the invention and a nucleic acid handling enzyme.

5

Description of the Figures

Fig. 1 shows sections through the α -hemolysin (α -HL) nanopore (PDB: 7AHL). (a) Charge distribution in the WT α -HL nanopore. Positively charged amino acids of α -HL are coloured in blue and negatively charged amino acids in red. (b) Sites in the α -HL nanopore modified in this work. Lys-8 is in purple. The constriction formed by the ion pair Glu-111/Lys-147 is coloured in green. Met-113, Thr-115, Thr-117, Gly-119, Asn-121, Asn-123 and Thr-125 are in orange.

10

Fig. 2 shows DNA translocation through the α -HL pore at +120 mV. (a) The five types of interaction of DNA with the WT α -HL pore. Type A event: a simple low-amplitude event (interpreted as the passage of DNA straight through the pore); type B: a simple mid-amplitude event (DNA is captured in the vestibule and exits the pore from the side of entry, the *cis* side); type C: a mid-amplitude signal, followed by a low-amplitude signal within the same event (DNA is detained in the vestibule before it passes through the β barrel to exit at the *trans* side); type D: a low-amplitude signal, followed by a mid-amplitude signal within the same event (DNA enters the β barrel directly and then retracts into the vestibule to exit the pore from the *cis* side); type E: a mid-amplitude signal, followed by a low-amplitude signal, followed by a second mid-amplitude signal within the same event (DNA explores the vestibule, threads into the β barrel, but retracts back into the vestibule and exits the pore from *cis* side). (b) Events histogram showing mean residual currents for 2,500 events expressed as fractions of the open pore current through the WT α -HL pore. The peaks representing type A and type B events are labelled. The type C, D and E events, characterized by their mean event currents, appear between the two major peaks. (c) Histogram of the DNA translocation times for the type A events. DNA (0.95 μ M) was presented from the *cis* chamber in 1 M KCl, 100 μ M EDTA, 25 mM Tris.HCl at pH 8.0.

15

20

25

30

Fig. 3 shows the effects of charged side-chains, introduced or removed by mutagenesis, on the frequency of DNA translocation at +120 mV. (a) M113R-WT (0.46 μ M DNA, *cis*), (b) E111N-WT (0.47 μ M), (c) WT (0.53 μ M), (d) M113R-RL2 (0.74 μ M), (e) RL2 (0.60 μ M) and (f) M113D-RL2 (2.0 μ M). M113D-RL2 showed no current blockades at up to +300 mV. Additional conditions are described in the legend to Fig. 2.

Fig. 4 shows the effects of arginine substitution in the β barrel of α -HL on the frequency of DNA translocation through the pore +120 mV. The frequency of DNA translocation is normalized

to 1 μM DNA. The line is a single exponential fit of the frequency of DNA translocation versus the distance of the arginine substitution from the constriction as represented by residue number. The mutants are in the RL2 background. Additional conditions are described in the legend to Fig. 2.

Fig. 5 shows the voltage dependence of the DNA capture rates. The DNA translocation frequency is plotted on a logarithmic scale and normalized to 1 μM DNA. Only the translocation events (i.e. type A and C) are included in the analysis. M113R-WT (red circles), E111N-WT (blue diamonds), M113R-RL2 (black squares), WT (green triangles down) and RL2 (blue triangles up). The lines show fits of the data with equation 1.

Fig. 6 shows the model for the translocation of DNA through the α -HL pore. To pass through the pore (V), DNA must first collide with the *cis* entrance of the pore (II), be transported through the vestibule (III) and enter the β barrel (IV). The DNA can prolong its visit within the vestibule and produce mid-amplitude and low-amplitude events corresponding to VI and VII. The processes that can be observed experimentally are labelled as events and are shown with slow inter-event rates. All steps are reversible with exception of translocation. In type A events, DNA collides with the pore, enters the vestibule and then directly translocates through the β barrel to exit on the trans side of the pore. Sampling of the barrel (III \leftrightarrow IV) during this process is too fast to observe. In type B events, DNA enters the pore but instead of penetrating the β barrel it has a prolonged interaction with the vestibule to produce the mid-amplitude event (I \rightarrow II \rightarrow III \rightarrow VI). In this case, DNA exits the pore from the same *cis* entrance from which it entered (VI \rightarrow III \rightarrow II \rightarrow I). Like type B events, type C events first show a mid-amplitude current (I \rightarrow II \rightarrow III \rightarrow VI), but then one end of the DNA enters the β barrel (VII) and a translocation event occurs (VII \rightarrow V \rightarrow I). In type D events, the DNA most likely enters the β barrel (I \rightarrow II \rightarrow III \rightarrow IV) but then stops translocating (VII), retracts back into the vestibule (VI) and exits the pore from the *cis* side (VI \rightarrow III \rightarrow II \rightarrow I). Finally, in type E events, the DNA begins to translocate through the β barrel as in type C events (I \rightarrow II \rightarrow III \rightarrow VI \rightarrow VII \rightarrow V), but then retracts into the vestibule (V \rightarrow VII \rightarrow VI) and exits the pore from the *cis* side as in type D events (VI \rightarrow III \rightarrow II \rightarrow I). Intermediates expected to give mid-amplitude events are numbered in yellow, although II and III would be too short for observation. Intermediates expected to give low-amplitude events are numbered in red, although IV would be too short for observation.

Fig. 7 shows DNA blockades for the homoheptameric mutant α -HL pores E111N-WT and M113R-WT. (a) Currents through E111N-WT during DNA-induced blockades at +200 mV, expressed as a fraction of the open-pore current, versus the event duration displayed on a semilogarithmic scale. (b) Corresponding plot for M113R-WT. (c) Long blockades for E111N-WT (diamonds) and M113R-WT (circles) as a percentage of the total for applied potentials from +100

to +200 mV. The DNA was added to the *cis* chamber at $\sim 0.5 \mu\text{M}$; the exact concentration was determined after each experiment. For further conditions, see Fig. 2, legend.

Fig. 8 shows amplitude histograms of mean event currents for DNA blockades of various α -HL pores. The main peak represents the type A events (DNA translocation). Each plot comprises 1000 events. The conditions are described in the legend to Fig. 2.

Fig. 9 shows current blockades after DNA addition ($1.0 \mu\text{M}$) to the *trans* chamber for: (a) M113R-RL2 and (b) N121R-RL2. The applied potential was -120 mV. For these mutants the currents through the open pores are much noisier at negative potentials than at positive potentials. Further conditions are described in the legend to Fig. 2.

Fig. 10 shows rare (<1%) current blockade events for DNA translocation through WT- α -HL nanopores at +120 mV showing multiple transitions between the mid-amplitude and low-amplitude states. The conditions are described in the legend to Fig. 2.

Fig. 11 shows: a) Section through a 7R nanopore. The amino acids at position 113, 115, 117, 119, 121, 123 and 125 were replaced by arginine using PyMOL (DeLano Scientific LLC, v 1.0) software from the WT nanopore (PDB:7AHL). Negative charged residues are coloured in red and positively charged residues in blue. b) Single channel recordings of 7R (top) and RL2- α -HL nanopores (bottom) at +120 mV after addition of $1 \mu\text{M}$ of ssDNA in the CIS side of the chamber. Experiments were performed at 1M KCl 25 mM Tris.HCl containing 100uM EDTA at pH 8.

Fig. 12 shows DNA translocation through 7R nanopores. a) Event histogram showing the translocation time distribution through a 7R pore upon addition of $0.7 \mu\text{M}$ of ssDNA (92mer) to the CIS side of the chamber. Events <5 ms are attributed to the transient gating of 7R nanopore (see SI). b) 2-D distribution plot of DNA translocation events ($\tau > 5 \text{ ms}$) versus the residual current during DNA translocation. The colour represents the density of the events as indicated in the inset. c) Dependence of the most likely DNA translocation speed of DNA through 7R nanopores on the applied voltage. The red line shows a single exponential fit.

Fig. 13 shows residual current through WT and modified α -HL nanopores during ssDNA translocation at +120mV. Errors express the standard deviation. All nanopores, except from WT, and 2N are in RL2 background. The fourth to last columns from the left concern the nanopores with additional positive charges.

Fig. 14 shows the dependence of the DNA translocation speed on the internal charge of 7R heteromers. The data are fitted to a single-exponential curve (upper panel). Heteromers nanopores were obtained by mixing monomers of RL2 (open circles) with monomer of 7R (purple circles). In this way, eight different classes of heptamers (7RmRL2n-m ; $n=7$, total number of subunits, $m=0\dots 7$) containing from 7 to 49 extra positive charges can be created (lower panel).

Fig. 15 shows the potential uses of 7R nanopores. a) Single strands are sequenced by reading the tunneling current. The 7R channel is incorporated into solid state nanopore functionalized with tunneling probes. Transverse tunneling currents through a ssDNA that is translocated through the pore are measured for recognition of single bases. 7R will control the time that each nucleotide remains between the tunneling probes. b) Strand sequencing using electrophysiological techniques. A β -barrel containing intra-positive charges is attached to the CIS entrance of the α -HL nanopore for controlling the speed of ssDNA. Differences between single nucleotides are measured by changes in conductance of the nanopore generated by the translocation of DNA.

10 **Description of the Sequence Listing**

SEQ ID NO: 1 shows the polynucleotide sequence that encodes one subunit of wild-type (WT) α -hemolysin.

SEQ ID NO: 2 shows the amino acid sequence of one subunit of WT α -hemolysin. Amino acids 2 to 6, 73 to 75, 207 to 209, 214 to 216 and 219 to 222 form α -helices. Amino acids 22 to 30, 35 to 44, 52 to 62, 67 to 71, 76 to 91, 98 to 103, 112 to 123, 137 to 148, 154 to 159, 165 to 172, 229 to 235, 243 to 261, 266 to 271, 285 to 286 and 291 to 293 form β -strands. All the other non-terminal amino acids, namely 7 to 21, 31 to 34, 45 to 51, 63 to 66, 72, 92 to 97, 104 to 111, 124 to 136, 149 to 153, 160 to 164, 173 to 206, 210 to 213, 217, 218, 223 to 228, 236 to 242, 262 to 265, 272 to 274 and 287 to 290 form loop regions. Amino acids 1 and 294 are terminal amino acids.

SEQ ID NO: 3 shows the polynucleotide sequence that encodes one subunit of α -hemolysin RL2. RL2 is the product of a semisynthetic gene that was devised to permit cassette mutagenesis of the sequence encoding the transmembrane β barrel (Cheley, S., Braha, O., Lu, X., Conlan, S., and Bayley, H. (1999) *Protein Sci.* **8**, 1257-1267). It contains six silent restriction sites and five altered amino acids in the encoded polypeptide sequence (K8A, V124L, G130S, N139Q and I142L).

D8RL2 is RL2 with an octa-aspartate tail.

SEQ ID NO: 4 shows the amino acid sequence of one subunit of α -hemolysin RL2. The same amino acids discussed above for SEQ ID NO: 2 form α -helices, β -strands, loop regions and terminal amino acids.

SEQ ID NO: 5 shows the polynucleotide sequence that encodes one subunit of α -hemolysin M113R-WT.

SEQ ID NO: 6 shows the amino acid sequence of one subunit of α -hemolysin M113R-WT.

SEQ ID NO: 7 shows the polynucleotide sequence that encodes one subunit of α -hemolysin E111N-WT.

SEQ ID NO: 8 shows the amino acid sequence of one subunit of α -hemolysin E111N-WT.

SEQ ID NO: 9 shows the polynucleotide sequence that encodes one subunit of α -hemolysin A8R-RL2 used in the Example.

SEQ ID NO: 10 shows the amino acid sequence of one subunit of α -hemolysin A8R-RL2 used in the Example.

5 SEQ ID NO: 11 shows the polynucleotide sequence that encodes one subunit of α -hemolysin M113K-RL2 used in the Example.

SEQ ID NO: 12 shows the amino acid sequence of one subunit of α -hemolysin M113K-RL2 used in the Example.

10 SEQ ID NO: 13 shows the polynucleotide sequence that encodes one subunit of α -hemolysin M113R-RL2 used in the Example.

SEQ ID NO: 14 shows the amino acid sequence of one subunit of α -hemolysin M113R-RL2 used in the Example.

SEQ ID NO: 15 shows the DNA sequence used in the Example.

15 **Detailed description of the invention**

It is to be understood that different applications of the disclosed products and methods may be tailored to the specific needs in the art. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting.

20 In addition as used in this specification and the appended claims, the singular forms “a”, “an”, and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to “an analyte” includes “analytes”, reference to “a transmembrane protein pore” includes two or more such pores, reference to “nucleic acid sequence” includes two or more such sequences, and the like.

25 All publications, patents and patent applications cited herein, whether supra or infra, are hereby incorporated by reference in their entirety.

Method of enhancing translocation

30 The present invention provides a method of enhancing translocation of a charged analyte through a transmembrane protein pore. The transmembrane protein pore is typically for use in detecting the analyte via stochastic sensing. As the analyte translocates through the pore, it typically interacts with the pore and affects the current flowing through the pore in a manner specific for that analyte. A distinctive current flows through the pore whenever the analyte interacts with the pore. This allows the analyte to be detected using stochastic sensing.

The method may enhance the translocation of the analyte in any way. The method preferably (1) increases the frequency of translocation of the analyte through the pore, (2) decreases the threshold voltage for translocation of the analyte through the pore, (3) decreases the speed of translocation of the analyte through the pore or (4) decreases the number of non-translocation interactions between the analyte and the pore. Non-translocation interactions are events where the analyte enters the pore and interacts with the pore, but does not translocate through the pore. In some instance, the non-translocation interactions are eliminated. The method more preferably increases (1) and (2); (1) and (3); (1) and (4); (2) and (3); (2) and (4); (3) and (4); (1), (2) and (3); (1), (3) and (4); (1), (2) and (4); or (2), (3) and (4). The method most preferably increases (1), (2), (3) and (4). In other words, the method most preferably increases the frequency of translocation of the analyte through the pore, decreases the threshold voltage for translocation of the analyte through the pore, decreases the speed of translocation of the analyte through the pore and decreases the number of non-translocation interactions between the analyte and the pore.

It is straightforward for a person skilled in the art to determine whether or not the increase in net opposing charge enhances translocation of the analyte through the pore. For instance, the analyte may be contacted with a non-enhanced or wild-type pore and the same type of pore in which the net opposing charge of its barrel or channel and/or entrance has been increased and any difference in translocation of the analyte between the two pores may be determined. Analyte translocation may be measured using any method known in the art. One method is described in the Examples.

By enhancing translocation of the analyte through the pore, the method of the invention provides an improved pore for stochastic sensing of the analyte. An enhanced translocation through the pore results in a more sensitive system that: (1) allows the detection of the analyte at both high and low concentrations; (2) allows the detection of the analyte even amongst impurities; (3) allows more rapid detection of the analyte; (4) reduces the background noise produced by non-translocation interactions and (5) allows the detection of the analyte at a lower voltage. Most importantly for the sequencing of nucleic acid sequences, an enhanced translocation, and in particular an enhanced frequency of translocation, results in a reduced deadtime between reads of the consecutive nucleotides in the sequence. This allows nucleic acids to be sequenced more quickly. A decreased translocation speed through the pore increases the read time of the consecutive nucleotides in the nucleic acid sequence and so allows the sequence of the nucleic acid to be determined more accurately.

Analyte

The analyte can be any charged substance. An analyte is charged if it has a net charge. The

analyte may be negatively charged or positively charged. An analyte is negatively charged if it has a net negative charge. An analyte is positively charged if it has a net positive charge.

Suitable analytes include, but are not limited to, metal ions, inorganic salts, polymers, such as a polymeric acids or bases, dyes, bleaches, pharmaceuticals, diagnostic agents, recreational
5 drugs, explosives and environmental pollutants.

The analyte can be an analyte that is secreted from cells. Alternatively, the analyte can be an analyte that is present inside cells such that the analyte must be extracted from cells.

The analyte is preferably a polymer. The polymer is preferably a nucleic acid sequence. Nucleic acids are negatively charged. A nucleic acid is a macromolecule comprising two or more
10 nucleotides. The nucleic acid handled by the enzyme may comprise any combination of any nucleotides. The nucleotides can be naturally occurring or artificial. A nucleotide typically contains a nucleobase, a sugar and at least one phosphate group. The nucleobase is typically heterocyclic. Nucleobases include, but are not limited to, purines and pyrimidines and more specifically adenine, guanine, thymine, uracil and cytosine. The sugar is typically a pentose sugar.

15 Nucleotide sugars include, but are not limited to, ribose and deoxyribose. The nucleotide is typically a ribonucleotide or deoxyribonucleotide. The nucleotide typically contains a monophosphate, diphosphate or triphosphate. Phosphates may be attached on the 5' or 3' side of a nucleotide.

Suitable nucleotides include, but are not limited to, adenosine monophosphate (AMP),
20 adenosine diphosphate (ADP), adenosine triphosphate (ATP), guanosine monophosphate (GMP), guanosine diphosphate (GDP), guanosine triphosphate (GTP), thymidine monophosphate (TMP), thymidine diphosphate (TDP), thymidine triphosphate (TTP), uridine monophosphate (UMP), uridine diphosphate (UDP), uridine triphosphate (UTP), cytidine monophosphate (CMP), cytidine diphosphate (CDP), cytidine triphosphate (CTP), cyclic adenosine monophosphate (cAMP), cyclic
25 guanosine monophosphate (cGMP), deoxyadenosine monophosphate (dAMP), deoxyadenosine diphosphate (dADP), deoxyadenosine triphosphate (dATP), deoxyguanosine monophosphate (dGMP), deoxyguanosine diphosphate (dGDP), deoxyguanosine triphosphate (dGTP), deoxythymidine monophosphate (dTMP), deoxythymidine diphosphate (dTDP), deoxythymidine triphosphate (dTTP), deoxyuridine monophosphate (dUMP), deoxyuridine diphosphate (dUDP),
30 deoxyuridine triphosphate (dUTP), deoxycytidine monophosphate (dCMP), deoxycytidine diphosphate (dCDP) and deoxycytidine triphosphate (dCTP). The nucleotides are preferably selected from AMP, TMP, GMP, UMP, dAMP, dTMP, dGMP or dCMP.

The nucleic acid is preferably double stranded, such as DNA. The nucleic acid may be single stranded, such as cDNA or RNA.

The polymer can be a peptide, polypeptide or a protein. The peptide, polypeptide or protein can be naturally-occurring or non-naturally-occurring. The polypeptide or protein can include within it synthetic or modified amino acids. A number of different types of modification to amino acids are well known in the art. For the purposes of the invention, it is to be understood that the
5 analyte can be modified by any method available in the art.

The protein can be an enzyme, antibody, hormone, growth factor or growth regulatory protein, such as a cytokine. The cytokine may be selected from an interleukin, preferably IFN-1, IL-2, IL-4, IL-5, IL-6, IL-10, IL-12 or IL-13, an interferon, preferably IL- γ or other cytokines such as TNF- α . The protein may be a bacterial protein, fungal protein, virus protein or parasite-derived
10 protein. Before it is contacted with the pore, the protein may be unfolded to form a polypeptide chain and thereby allow it to enter the barrel or channel of the pore and interact with the pore.

The analyte may be an amino acid. Charged amino acids are well known in the art and are discussed in more detail below. The amino acid can be naturally-occurring or non-naturally-occurring. The amino acid may be synthetic or modified. A number of different types of
15 modification to amino acids are well known in the art.

The analyte may be an individual nucleotide or a single nucleotide. An individual nucleotide is one which is not bound to another polynucleotide by a nucleotide bond. A nucleotide bond involves one of the phosphate groups of a nucleotide being bound to the sugar group of another nucleotide. An individual nucleotide is typically one which is not bound by a nucleotide
20 bond to another polynucleotide sequence of at least 5, at least 10, at least 20, at least 50, at least 100, at least 200, at least 500, at least 1000 or at least 5000 nucleotides. For example, the individual nucleotide has been digested from a target polynucleotide sequence, such as a DNA or RNA strand. Nucleotides are negatively charged. The nucleotide may be any of those discussed above.

The nucleotide may be derived from the digestion of a nucleic acid sequence such as
25 ribonucleic acid (RNA) or deoxyribonucleic acid. Nucleic acid sequences can be digested using any method known in the art. Suitable methods include, but are not limited to, those using enzymes or catalysts. Catalytic digestion of nucleic acids is disclosed in Deck *et al.*, *Inorg. Chem.*, 2002; 41: 669-677.

Individual nucleotides from a single nucleic acid sequence may be contacted with the pore
30 in a sequential manner in order to sequence the whole or part of the nucleic acid. Sequencing nucleic acids in accordance with the second embodiment of the invention is discussed in more detail below.

The nucleotide is typically unmodified, such as when the nucleotide is derived from the digestion of a nucleic acid sequence. Alternatively, the nucleotide may be modified or damaged.

35 The nucleotide is typically methylated or oxidised. The nucleotide may be labelled with a revealing

label. The revealing label may be any suitable label which allows the nucleotide to be detected. Suitable labels include fluorescent molecules, radioisotopes, e.g. ^{125}I , ^{35}S , and linkers such as biotin.

The nucleotide is typically present in any suitable biological sample. Suitable biological samples are discussed below.

5

Transmembrane protein pore

A transmembrane protein pore is a polypeptide that permits ions to flow from one side of the membrane to the other along an applied potential. The pore preferably permits the analyte to flow from one side of the membrane to the other along the applied potential.

10 The pore is typically an oligomer. The pore is preferably made up of several repeating subunits, such as 6, 7 or 8 subunits. The pore is more preferably heptameric. The pore typically comprises a barrel or channel through which the ions may flow. The subunits of the pore typically surround a central axis and contribute strands to a transmembrane β barrel or channel or a transmembrane α -helix bundle or channel.

15 Pores for use in accordance with the invention can be β -barrel pores or α -helix bundle pores. β -barrel pores comprise a barrel or channel that is formed from β -sheets. Suitable β -barrel pores include, but are not limited to, β -toxins, such as α -hemolysin and leukocidins, and outer membrane proteins/porins of bacteria, such as *Mycobacterium smegmatis* porin A (MspA), outer membrane porin F (OmpF), outer membrane porin G (OmpG), outer membrane phospholipase A and *Neisseria* autotransporter lipoprotein (NalP). α -helix bundle pores comprise a barrel or channel
20 that is formed from α -helices. Suitable α -helix bundle pores include, but are not limited to, inner membrane proteins and α outer membrane proteins, such as WZA.

The most preferred pore for use in the invention is α -hemolysin or a variant thereof. The α -hemolysin pore is formed of seven identical subunits (i.e. it is heptameric). The sequence of one
25 subunit of wild-type α -hemolysin is shown in SEQ ID NO: 2. The sequence of one subunit of α -hemolysin RL2 is shown in SEQ ID NO: 4. A variant is a heptameric pore in which one or more of the seven subunits has an amino acid sequence which varies from that of SEQ ID NO: 2 or 4 and which retains pore activity. A variant may include modifications that facilitate interaction with the analyte.

30 1, 2, 3, 4, 5, 6 or 7 of the subunits in a variant α -hemolysin may have an amino acid sequence that varies from that of SEQ ID NO: 2 or 4. All seven subunits within a variant pore are typically identical but may be different, particularly if one or more of the subunits has been modified to facilitate interaction with the analyte.

The variant may be a naturally-occurring variant which is expressed by an organism, for instance by a *Staphylococcus* bacterium. Variants also include non-naturally occurring variants produced by recombinant technology. Over the entire length of the amino acid sequence of SEQ ID NO: 2 or 4, a subunit of a variant will preferably be at least 50% homologous to that sequence based on amino acid identity. More preferably, the subunit polypeptide may be at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90% and more preferably at least 95%, 97% or 99% homologous based on amino acid identity to the amino acid sequence of SEQ ID NO: 2 or 4 over the entire sequence. There may be at least 80%, for example at least 85%, 90% or 95%, amino acid identity over a stretch of 200 or more, for example 230, 250, 270 or 280 or more, contiguous amino acids (“hard homology”). Preferred variants are shown in SEQ ID NOs: 6, 8, 10, 12 and 14.

Amino acid substitutions may be made to the amino acid sequence of SEQ ID NO: 2 or 4, for example up to 1, 2, 3, 4, 5, 10, 20 or 30 substitutions. Conservative substitutions may be made, for example, according to Table 1 below.

15

Table 1 – Conservative substitutions

Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other.

NON-AROMATIC	Non-polar	G A P
		I L V
	Polar – uncharged	C S T M
		N Q
	Polar – charged	D E
		H K R
AROMATIC		H F W Y

20

One or more amino acid residues of the amino acid sequence of SEQ ID NO: 2 or 4 may alternatively or additionally be deleted. Up to 1, 2, 3, 4, 5, 10, 20 or 30 residues may be deleted, or more.

Variants may include subunits made of fragments of SEQ ID NO: 2 or 4. Such fragments retain pore forming activity. Fragments may be at least 50, 100, 200 or 250 amino acids in length. Such fragments may be used to produce chimeric pores. A fragment preferably comprises the pore forming domain of SEQ ID NO: 2 or 4.

25

Variants include chimeric protein pores comprising fragments or portions of SEQ ID NO: 2 or 4. Chimeric protein pores are formed from subunits each comprising fragments or portions of SEQ ID NO: 2 or 4. The pore or channel part of a chimeric protein pore is typically formed by the fragments or portions of SEQ ID NO: 2 or 4.

5 One or more amino acids may be alternatively or additionally added to the polypeptides described above. An extension may be provided at the N-terminus or C-terminus of the amino acid sequence of SEQ ID NO: 2 or 4 or polypeptide variant or fragment thereof. The extension may be quite short, for example from 1 to 10 amino acids in length. Alternatively, the extension may be longer, for example up to 50 or 100 amino acids. A carrier protein may be fused to an amino acid
10 sequence according to the invention.

As discussed above, a variant of SEQ ID NO: 2 or 4 is a subunit that has an amino acid sequence which varies from that of SEQ ID NO: 2 or 4 and which retains its ability to form a pore. A variant typically contains the regions of SEQ ID NO: 2 or 4 that are responsible for pore formation. The pore forming ability of α -HL, which contains a β -barrel, is provided by β -sheets in
15 each subunit. A variant of SEQ ID NO: 2 or 4 typically comprises the regions in SEQ ID NO: 2 or 4 that form β -sheets. Amino acids 22 to 30, 35 to 44, 52 to 62, 67 to 71, 76 to 91, 98 to 103, 112 to 123, 137 to 148, 154 to 159, 165 to 172, 229 to 235, 243 to 261, 266 to 271, 285 to 286 and 291 to 293 of SEQ ID NO: 2 and 4 form β -sheets. One or more modifications can be made to the regions of SEQ ID NO: 2 or 4 that form β -sheets as long as the resulting variant retains its ability to form a
20 pore. Specific modifications that can be made to the β -sheet regions of SEQ ID NO: 2 or 4 in accordance with the invention are discussed below.

A variant of SEQ ID NO: 2 or 4 preferably includes one or more modifications, such as substitutions, additions or deletions, within its α -helices and/or loop regions. Amino acids 2 to 6,
25 73 to 75, 207 to 209, 214 to 216 and 219 to 222 of SEQ ID NO: 2 and 4 form α -helices. Amino acids 7 to 21, 31 to 34, 45 to 51, 63 to 66, 72, 92 to 97, 104 to 111, 124 to 136, 149 to 153, 160 to 164, 173 to 206, 210 to 213, 217, 218, 223 to 228, 236 to 242, 262 to 265, 272 to 274 and 287 to 290 of SEQ ID NO: 2 and 4 form loops. Amino acids 1 and 294 are terminal amino acids. Specific modifications that can be made to the loop regions of SEQ ID NO: 2 or 4 in accordance with the invention are discussed below.

30 Standard methods in the art may be used to determine homology. For example the UWGCG Package provides the BESTFIT program which can be used to calculate homology, for example used on its default settings (Devereux *et al* (1984) *Nucleic Acids Research* **12**, p387-395). The PILEUP and BLAST algorithms can be used to calculate homology or line up sequences (such as identifying equivalent residues or corresponding sequences (typically on their default settings)), for

example as described in Altschul S. F. (1993) *J Mol Evol* 36:290-300; Altschul, S.F *et al* (1990) *J Mol Biol* 215:403-10.

Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first
5 identifying high scoring sequence pair (HSPs) by identifying short words of length W in the query sequence that either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighbourhood word score threshold (Altschul *et al*, supra). These initial neighbourhood word hits act as seeds for initiating searches to find HSP's containing them. The word hits are extended in both directions along each
10 sequence for as far as the cumulative alignment score can be increased. Extensions for the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the
15 alignment. The BLAST program uses as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1992) *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

The BLAST algorithm performs a statistical analysis of the similarity between two sequences; see e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90: 5873-5787. One
20 measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two amino acid sequences would occur by chance. For example, a sequence is considered similar to another sequence if the smallest sum probability in comparison of the first sequence to the second sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most
25 preferably less than about 0.001.

A transmembrane protein pore may be modified for example by the addition of histidine or aspartic acid residues to assist their identification or purification or by the addition of a signal sequence to promote their secretion from a cell where the polypeptide does not naturally contain such a sequence.

30 A pore may be labelled with a revealing label. The revealing label may be any suitable label which allows the pore to be detected. Suitable labels include, but are not limited to, fluorescent molecules, radioisotopes, e.g. ¹²⁵I, ³⁵S, enzymes, antibodies, antigens, polynucleotides and ligands such as biotin.

The pore may be derived from a pore producing organism, such as *Staphylococcus aureus*,
35 or made synthetically or by recombinant means. For example, the pore may be synthesized by *in*

vitro transcription and translation (IVTT). In addition to the modifications discussed in detail below that increase the opposing charge of the barrel or channel and/or entrance of the pore, the amino acid sequence of the pore may be modified to include non-naturally occurring amino acids, for instance to increase the stability of the pore. When the pore is produced by synthetic means, such amino acids may be introduced during production. Methods for including non-naturally-occurring amino acids in the pore are well known in the art and are discussed in more detail below with reference to increasing the net opposing charge.

The pore may also be produced using D-amino acids. For instance, the pore may comprise a mixture of L-amino acids and D-amino acids. This is conventional in the art for producing such proteins or peptides.

The pore can also be altered following either synthetic or recombinant production. The pore may contain other non-specific modifications as long as they do not interfere with translocation of the analyte. A number of non-specific side chain modifications are well known in the art and may be made to the side chains of the pore. Such modifications include, for example, reductive alkylation of amino acids by reaction with an aldehyde followed by reduction with NaBH₄, amidination with methylacetimidate and thiol-specific modifications using ethyleneamine.

The pore can be produced using standard methods known in the art. Polynucleotide sequences encoding a pore or a pore subunit may be derived and replicated using standard methods in the art. Such sequences are discussed in more detail below. Polynucleotide sequences encoding a pore or a pore subunit may be expressed in a bacterial host cell using standard techniques in the art. The pore or pore subunit may be produced in a cell by *in situ* expression of the polypeptide from a recombinant expression vector. The expression vector optionally carries an inducible promoter to control the expression of the polypeptide.

A pore subunit may be produced in large scale following purification by any protein liquid chromatography system from pore producing organisms or after recombinant expression as described below. Typical protein liquid chromatography systems include FPLC, AKTA systems, the Bio-Cad system, the Bio-Rad BioLogic system and the Gilson HPLC system. The naturally occurring or recombinantly-produced pore or pore subunit may then be inserted into a naturally occurring or artificial membrane for use in accordance with the invention. Methods for inserting pore into membranes are discussed below.

The transmembrane pore may comprise a molecular adaptor, such as a cyclodextrin, a cyclic peptide or a cucurbituril, that facilitates the interaction between the pore and the analyte by improving the host-guest chemistry of the pore and the analyte. Suitable adaptors are well known in the art. The molecular adaptor may be covalently attached to the pore.

Increasing the net opposing charge

The method of the invention comprises increasing the net opposing charge of the barrel or channel and/or entrance of the pore. The opposing charge is the opposite charge to the charge of the analyte. If the analyte is negatively charged, the method involves increasing the net positive charge of the barrel or channel and/or entrance of the pore. If the analyte is positively charged, the method comprises increasing the net negative charge of the barrel or channel and/or entrance of the pore. The net opposing charge may be increased by any means known in the art.

The net opposing charge is increased in a manner that does not interfere with translocation of the analyte through the pore. The net opposing charge is increased in a manner that does not interfere with the interaction between the analyte and the pore.

In a preferred embodiment, the analyte is negatively charged and the method involves increasing the net positive charge of the barrel or channel and/or entrance of the pore. The net positive charge is preferably increased by introducing one or more positively charged amino acids into the barrel or channel and/or entrance of the pore. The one or more positively charged amino acids may be introduced by addition. The one or more positively charged amino acids are preferably introduced by substitution.

A positively charged amino acid is an amino acid with a net positive charge. The positively charged amino acid(s) can be naturally-occurring or non-naturally-occurring. The positively charged amino acids may be synthetic or modified. For instance, modified amino acids with a net positive charge may be specifically designed for use in the invention. A number of different types of modification to amino acids are well known in the art.

Preferred naturally-occurring positively charged amino acids include, but are not limited to, histidine (H), lysine (K) and arginine (R). Any number and combination of H, K and/or R may be introduced into the barrel or channel and/or entrance of the pore.

Methods for adding or substituting naturally-occurring amino acids are well known in the art. For instance, methionine (M) may be substituted with arginine (R) by replacing the codon for methionine (ATG) with a codon for arginine (AGA) at the relevant position in a polynucleotide encoding the pore. The polynucleotide can then be expressed as discussed below.

Methods for adding or substituting non-naturally-occurring amino acids are also well known in the art. For instance, non-naturally-occurring amino acids may be introduced by including synthetic aminoacyl-tRNAs in the IVTT system used to express the pore. Alternatively, they may be introduced by expressing the pore in *E. coli* that are auxotrophic for specific amino acids in the presence of synthetic (i.e. non-naturally-occurring) analogues of those specific amino acids. They may also be produced by naked ligation if the pore is produced using partial peptide synthesis.

Any amino acid may be substituted with a positively charged amino acid. One or more uncharged amino acids, non-polar amino acids and/or aromatic amino acids may be substituted with one or more positively charged amino acids. Uncharged amino acids have no net charge. Suitable uncharged amino acids include, but are not limited to, cysteine (C), serine (S), threonine (T), methionine (M), asparagine (N) and glutamine (Q). Non-polar amino acids have non-polar side chains. Suitable non-polar amino acids include, but are not limited to, glycine (G), alanine (A), proline (P), isoleucine (I), leucine (L) and valine (V). Aromatic amino acids have an aromatic side chain. Suitable aromatic amino acids include, but are not limited to, histidine (H), phenylalanine (F), tryptophan (W) and tyrosine (Y). Preferably, one or more negatively charged amino acids are substituted with one or more positively charged amino acids. Suitable negatively charged amino acids include, but are not limited to, aspartic acid (D) and glutamic acid (E).

Preferred introductions include, but are not limited to, substitution of M with R, substitution of M with H, substitution of M with K, substitution of D with R, substitution of D with H, substitution of D with K, substitution of E with R, substitution of E with H, substitution of E with K, substitution of N with R, substitution of T with R and substitution of G with R.

Any number of positively charged amino acids may be introduced. For instance, 1, 2, 5, 10, 15, 20, 25 or more positively charged amino acids may be introduced. If the pore is oligomeric, the one or more positively charged amino acids may be introduced into one, some or all of the subunits of the pore. In the case of α -HL (i.e. SEQ ID NO: 2 and 4 and variants thereof discussed above), the one or more positively charged amino acids may be introduced into 1, 2, 3, 4, 5, 6 or 7 of the subunits in the pore. In each of the seven subunits, the one or more positively charged amino acids may be introduced at the same or different positions. Preferably, the pore is a homoheptamer and one or more positive amino acids are introduced at the same position(s) in each subunit.

The net positive charge may also be increased by replacing by substitution one or more negatively charged amino acids with one or more uncharged amino acids, non-polar amino acids and/or aromatic amino acids. The removal of negative charge from the barrel or channel and/or entrance increases the net positive charge. The uncharged amino acids, non-polar amino acids and/or aromatic amino acids can be naturally-occurring or non-naturally-occurring. They may be synthetic or modified. Suitable uncharged amino acids, non-polar amino acids and aromatic amino acids are discussed above. Preferred substitutions include, but are not limited to, substitution of E with N, substitution of D with N, substitution of E with T, substitution of D with T, substitution of E with G and substitution of D with G.

Any number and combination of uncharged amino acids, non-polar amino acids and/or aromatic amino acids may substituted into the barrel or channel and/or entrance. For instance, 1, 2, 5, 10, 15, 20, 25 or more uncharged amino acids, non-polar amino acids and/or aromatic amino

acids may be substituted into the barrel or channel and/or entrance. If the pore is oligomeric, the uncharged amino acids, non-polar amino acids and/or aromatic amino acids may be substituted into one, some or all of the subunits of the pore. In the case of α -HL (i.e. SEQ ID NO: 2 and 4 and variants thereof discussed above), the uncharged amino acids, non-polar amino acids and/or aromatic amino acids may be substituted into 1, 2, 3, 4, 5, 6 or 7 of the subunits in the pore. In each of the seven subunits, the one or more uncharged amino acids, non-polar amino acids and/or aromatic amino acids may be substituted into the same or different positions. Preferably, the pore is a homoheptamer and uncharged amino acids, non-polar amino acids and/or aromatic amino acids are substituted into the same position(s) in each subunit. Negatively charged amino acids may be substituted with (1) uncharged amino acids; (2) non-polar amino acids; (3) aromatic amino acids; (4) uncharged amino acids and non-polar amino acids; (5) uncharged amino acids and aromatic amino acids; and (6) uncharged amino acids, non-polar amino acids and aromatic amino acids.

In another embodiment, the analyte is positively charged and the method involves increasing the net negative charge of the barrel or channel and/or entrance of the pore. The net negative charge is preferably increased by introducing one or more negatively charged amino acids into the barrel or channel and/or entrance of the pore. The one or more negatively charged amino acids may be introduced by addition. The one or more negatively charged amino acids are preferably introduced by substitution. Methods for adding and substituting amino acids are well known in the art.

Suitable negatively charged amino acids are discussed above. The negatively charged amino acid(s) can be naturally-occurring or non-naturally-occurring. The negatively charged amino acids may be synthetic or modified.

Any amino acid may be substituted with a negatively charged amino acid. One or more uncharged amino acids, non-polar amino acids and/or aromatic amino acids may be substituted with one or more negatively charged amino acids. Preferably, one or more positively charged amino acids are substituted with one or more negatively charged amino acids. Any number of negatively charged amino acids may be introduced as discussed above.

The net negative charge may also be increased by replacing by substitution one or more positively charged amino acids with one or more uncharged amino acids, non-polar amino acids and/or aromatic amino acids. A preferred substitution is substitution of K with N. The removal of positive charge from the barrel or channel and/or entrance increases the net negative charge. The uncharged amino acids, non-polar amino acids and/or aromatic amino acids can be naturally-occurring or non-naturally-occurring. They may be synthetic or modified.

Any number and combination of uncharged amino acids, non-polar amino acids and/or aromatic amino acids may be substituted into the barrel or channel and/or entrance as discussed above.

Barrel or channel and/or entrance

In one embodiment, the net opposing charge of the barrel or channel of the pore is increased. The barrel or the channel of the pore is the portion of the pore through which ions travel through the pore across the membrane. The analyte also translocates through the barrel or the
5 channel. The barrel or channel may be formed from α -helices or β -strands depending on the type of pore. Preferably, the net opposing charge of the lumen of the barrel or channel is increased.

As described above, the net opposing charge is increased by introducing or replacing by substitution one or more amino acids. Amino acid(s) may be introduced or substituted at any position in the barrel or channel as long as translocation of the analyte is enhanced. Amino acid(s)
10 are preferably introduced or substituted at position(s) that form part of the lumen of the barrel or channel. In other words, it is preferred that amino acid(s) are not introduced or substituted at positions in the barrel or channel that are buried in the pore or hidden from the lumen.

Amino acid(s) are preferably introduced or substituted towards the end of the barrel or channel through which the analyte enters. The end of the pore through which the analyte enters
15 may be the *cis* end or the *trans* end. The end is preferably the *cis* end. Amino acid(s) are preferably introduced or substituted near a constriction of the barrel or channel. Amino acid(s) are more preferably introduced or substituted near a constriction of the lumen of the barrel or channel.

In another embodiment, the net opposing charge of the entrance of the pore is increased. An entrance of the pore is the portion of the pore through which the analyte enters into the barrel or
20 channel. Entrances are typically formed primarily from loop regions in the pore.

As described above, the net opposing charge is increased by introducing or replacing by substitution one or more amino acids. Amino acid(s) may be introduced or substituted at any position in the entrance as long as translocation of the analyte is enhanced. Preferably, amino acid(s) are introduced or substituted at position(s) that form part of the lumen of the entrance. In
25 other words, it is preferred that amino acid(s) are not introduced or substituted at positions in the entrance that are buried in the pore or hidden from the lumen.

Amino acid(s) may be introduced or substituted in one or both entrances. Amino acid(s) are typically introduced or substituted near the entrance through which the analyte enters the pore. Amino acid(s) may be introduced or substituted in the barrel or channel and in one or both
30 entrance(s).

In SEQ ID NOs: 2 and 4, amino acid(s) may be introduced or substituted at any of positions 1 to 22, 35 to 41, 54 to 59, 100 to 155 and 218 to 229 to increase the net opposing charge. All of these positions form part of the lumen of the barrel or channel or part of the lumen of an entrance of
35 SEQ ID NOs: 2 and 4. Positions 111 to 147 form part of the lumen of the barrel or channel.

Positions 1 to 22, 35 to 41, 54 to 59, 100 to 110, 148 to 155 and 218 to 229 form part of the lumen

of an entrance. Amino acid(s) are preferably introduced or substituted at positions of SEQ ID NO: 2 or 4 located near the constriction of the barrel or channel. Such positions include, but are not limited to, positions 111, 113, 115, 117, 119, 121, 123, 125, 127, 145 and 147. A preferred position in the entrance of SEQ ID NO: 2 is position 8.

5 Preferred substitutions located near the constriction of the barrel or channel of SEQ ID NO: 2 or 4 include, but are not limited to, M113R, E111N, M113K, T115R, T117R, G119R, N121R, N123R, M113D, K147N, D127R, T125R and T145R. Preferred substitutions located in the entrance of SEQ ID NO: 4 include, but are not limited to, A8R and A8K.

10 Transmembrane protein pore of the invention

The present invention also provides a transmembrane protein pore enhanced in accordance with the invention. Such a pore will have an increased net opposing charge in its barrel or channel and/or entrance. An increased net positive charge enhances translocation of a negatively charged analyte through the pore. An increased net negative charge enhances translocation of a positively
15 charged analyte through the pore. The pore may be enhanced using any method described above.

A pore of the invention may be isolated, substantially isolated, purified or substantially purified. A pore of the invention is isolated or purified if it is completely free of any other components, such as lipids or other pores. A pore is substantially isolated if it is mixed with carriers or diluents which will not interfere with its intended use. For instance, a pore is
20 substantially isolated or substantially purified if it present in a form that comprises less than 10%, less than 5%, less than 2% or less than 1% of other components, such as lipids or other pores. Alternatively, a pore of the invention may be present in a lipid bilayer.

A pore of the invention may be present as an individual or single pore. Alternatively, a pore of the invention may be present in a homologous or heterologous population of two or more pores.

25 The transmembrane pore preferably comprises at least one subunit comprising SEQ ID NO: 6, 8, 10 or a variant thereof. The transmembrane pore is more preferably a homoheptamer comprising seven subunits comprising SEQ ID NO: 6, 8 or 10 or a variant thereof. Variants of SEQ ID NO: 6, 8 and 10 may differ from SEQ ID NO: 6, 8 and 10 to the same extent discussed above for variants of SEQ ID NOs: 2 and 4.

30 Variants of SEQ ID NO: 6 must comprise arginine at position 113 (M113R). The variant of SEQ ID NO: 6 is preferably not M113R-RL2 (SEQ ID NO: 14). Variants of SEQ ID NO: 8 must comprise asparagine at position 111 (E111N). Variants of SEQ ID NO: 10 must comprise arginine at position 8 (A8R-RL2).

Polynucleotide

The present invention also provides a polynucleotide sequence which encodes a transmembrane protein pore of the invention. The polynucleotide will encode a pore having an increased net opposing charge in its barrel or channel and/or entrance. An increased net positive charge enhances translocation of a negatively charged analyte through the pore. An increased net negative charge enhances translocation of a positively charged analyte through the pore. The pore may be enhanced using any method described above.

The present invention also provides a polynucleotide sequence encoding a transmembrane protein pore subunit having the sequence of SEQ ID NO: 6, 8 or 10 or a variant thereof. The polynucleotide sequence preferably comprises the sequence shown in SEQ ID NO: 5, 7 or 9 or a variant thereof. A variant of SEQ ID NO: 5, 7 or 9 is a sequence at least 50%, 60%, 70%, 80%, 90% or 95% homologous based on nucleotide identity to sequence of SEQ ID NO: 5, 7 or 9 over the entire sequence. There may be at least 80%, for example at least 85%, 90% or 95% nucleotide identity over a stretch of 600 or more, for example 700, 750, 850 or 900 or more, contiguous nucleotides ("hard homology"). Homology may be calculated as described above. The variant may comprise a sequence that differs from SEQ ID NO: 5, 7 or 9 on the basis of the degeneracy of the genetic code.

A variant of SEQ ID NO: 5 must encode a pore subunit comprising an arginine at position 113 (M113R) of SEQ ID NO: 6 or at the position equivalent to position 113 (M113R) of SEQ ID NO: 6. A variant of SEQ ID NO: 5 preferably does not encode M113R-RL2 (SEQ ID NO: 14).

A variant of SEQ ID NO: 7 must encode a pore subunit comprising asparagine at position 111 (E111N) of SEQ ID NO: 8 or at the position equivalent to position 111 (E111N) of SEQ ID NO: 8.

A variant of SEQ ID NO: 9 must encode a pore subunit comprising arginine at the position 8 (A8R) of SEQ ID NO: 10 or at the position equivalent to position 8 (A8R) of SEQ ID NO: 10.

Polynucleotide sequences may be derived and replicated using standard methods in the art. Chromosomal DNA may be extracted from a pore producing organism, such as *Staphylococcus aureus*. The gene encoding the pore subunit may be amplified using PCR involving specific primers. The amplified sequence may then be incorporated into a recombinant replicable vector such as a cloning vector. The vector may be used to replicate the polynucleotide in a compatible host cell. Thus polynucleotide sequences encoding a pore subunit may be made by introducing a polynucleotide encoding a pore subunit into a replicable vector, introducing the vector into a compatible host cell, and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell. Suitable host cells for cloning of

polynucleotides encoding a pore subunit are well known in the art and described in more detail below.

The polynucleotide sequence encoding a pore subunit may be cloned into suitable expression vector. In an expression vector, the polynucleotide sequence encoding a pore subunit is typically operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell. Such expression vectors can be used to express a pore subunit.

The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. Multiple copies of the same or different pore subunit sequences may be introduced into the vector.

The expression vector may then be introduced into a suitable host cell. Thus, a pore subunit can be produced by inserting a polynucleotide sequence encoding a pore subunit into an expression vector, introducing the vector into a compatible bacterial host cell, and growing the host cell under conditions which bring about expression of the polynucleotide sequence encoding the pore subunit. The recombinantly-expressed pore subunit may self-assemble into a pore in the host cell membrane. Alternatively, the recombinant pore produced in this manner may be removed from the host cell and inserted into another membrane. When producing a heptameric pore comprising at least two different subunits, the different subunits may be expressed separately in different host cells as described above, removed from the host cells and assembled into a pore in a separate membrane, such as a rabbit cell membrane.

The vectors may be for example, plasmid, virus or phage vectors provided with an origin of replication, optionally a promoter for the expression of the said polynucleotide sequence and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example a tetracycline resistance gene. Promoters and other expression regulation signals may be selected to be compatible with the host cell for which the expression vector is designed. A T7, *trc*, *lac*, *ara* or λ_L promoter is typically used.

The host cell typically expresses the pore subunit at a high level. Host cells transformed with a polynucleotide sequence encoding a pore subunit will be chosen to be compatible with the expression vector used to transform the cell. The host cell is typically bacterial and preferably *Escherichia coli*. Any cell with a λ DE3 lysogen, for example C41 (DE3), BL21 (DE3), JM109 (DE3), B834 (DE3), TUNER, Origami and Origami B, can express a vector comprising the T7 promoter.

Method of detecting analytes

The present invention also relates to a method of determining the presence or absence of an analyte in a sample. The analyte is charged. The analyte may be any of those discussed above. The method comprises (1) contacting the sample with a transmembrane protein pore of the invention under conditions that allow the analyte, if present, to translocate through and interact with the pore and (2) measuring the current passing through the pore during the interaction and thereby determining the presence or absence of the analyte. Any transmembrane protein pore of the invention can be used. The benefits associated with using a transmembrane protein pore of the invention to detect an analyte are discussed above. In a preferred embodiment, the method further comprises enhancing translocation of the analyte through the pore as described above. The enhancement is of course carried out before the analyte is detected.

The analyte is present if the current flows through the pore in a manner specific for the analyte (i.e. if a distinctive current associated with the analyte is detected flowing through the pore). The analyte is absent if the current does not flow through the pore in a manner specific for the analyte.

The invention therefore involves stochastic sensing of an analyte. The invention can be used to differentiate analytes of similar structure on the basis of the different effects they have on the current passing through a transmembrane protein pore. The invention can also be used to measure the concentration of a particular analyte in a sample.

The invention may also be used in a sensor that uses many or thousands of pores of the invention in bulk sensing applications.

The method may be carried out using any suitable membrane/pore system in which a transmembrane protein pore of the invention is inserted into a membrane. The method is typically carried out using (i) an artificial membrane comprising a transmembrane protein pore of the invention, (ii) an isolated, naturally-occurring membrane comprising a transmembrane protein pore of the invention, or (iii) a cell expressing a transmembrane protein pore that has been modified in accordance with the invention. The method is preferably carried out using an artificial membrane. The membrane may comprise other transmembrane and/or intramembrane proteins as well as other molecules in addition to the transmembrane protein pore of the invention.

The method of the invention is typically carried out *in vitro*.

Membrane

The membrane forms a barrier to the flow of ions, nucleotides and nucleic acids. The membrane is preferably a lipid bilayer. Lipid bilayers suitable for use in accordance with the invention can be made using methods known in the art. For example, lipid bilayer membranes can

be formed using the method of Montal and Mueller (1972). Lipid bilayers can also be formed using the method described in International Application No. PCT/GB08/000563.

The method of the invention may be carried out using lipid bilayers formed from any membrane lipid including, but not limited to, phospholipids, glycolipids, cholesterol and mixtures thereof. Any of the lipids described in International Application No. PCT/GB08/000563 may be used.

Methods are known in the art for inserting pores into membranes, such as lipid bilayers. For example, the pore may be suspended in a purified form in a solution containing a lipid bilayer such that it diffuses to the lipid bilayer and is inserted by binding to the lipid bilayer and assembling into a functional state. Alternatively, the pore may be directly inserted into the membrane using the “pick and place” method described in M.A. Holden, H. Bayley. J. Am. Chem. Soc. 2005, 127, 6502-6503 and International Application No. PCT/GB2006/001057 (published as WO 2006/100484).

The method of the invention is typically carried out *in vitro*.

Interaction between the pore and analyte

The analyte may be contacted with the pore on either side of the membrane. The analyte may be introduced to the pore on either side of the membrane. The analyte may be contacted with the side of the membrane that allows the analyte to pass through the pore to the other side of the membrane. For example, the analyte is contacted with an end of the pore, which in its native environment allows the entry of ions or small molecules, such as analytes, into the barrel or channel of the pore such that the analyte may pass through the pore. In such cases, the analyte interacts with the pore as it passes across the membrane through the barrel or channel of the pore. Alternatively, the analyte may be contacted with the side of the membrane that allows the analyte to interact with the pore, dissociate from the pore and remain on the same side of the membrane.

The analyte may interact with the pore in any manner and at any site. The analyte may reversibly bind to the pore.

During the interaction between the analyte and the pore, the analyte affects the current flowing through the pore in a manner specific for that analyte. For example, a particular analyte will reduce the current flowing through the pore for a particular mean time period and to a particular extent. In other words, the current flowing through the pore is distinctive for a particular analyte. Control experiments may be carried out to determine the effect a particular analyte has on the current flowing through the pore. Results from carrying out the method of the invention on a test sample can then be compared with those derived from such a control experiment in order to identify a particular analyte in the sample or determine whether a particular analyte is present in the sample.

The frequency at which the current flowing through the pore is affected in a manner indicative of a particular analyte can be used to determine the concentration of that analyte in the sample.

Apparatus

5 The method may be carried out using any apparatus that is suitable for investigating a membrane/pore system in which a transmembrane protein pore is inserted into a membrane. The method may be carried out using any apparatus that is suitable for stochastic sensing. For example, the apparatus comprises a chamber comprising an aqueous solution and a barrier that separates the chamber into two sections. The barrier has an aperture in which the membrane containing the pore
10 is formed. The analyte may be contacted with the pore by introducing the analyte into the chamber. The analyte may be introduced into either of the two sections of the chamber.

The method may be carried out using the apparatus described in International Application No. PCT/GB08/000562.

15 The method of the invention involves measuring the current passing through the pore during interaction with the analyte. Therefore the apparatus also comprises an electrical circuit capable of applying a potential and measuring an electrical signal across the membrane and pore. The method may be carried out using a patch clamp or a voltage clamp. The method preferably involves the use of a voltage clamp. The Example discloses one way to carry out a voltage clamp method.

20 *Sample*

The analyte is present in any suitable sample. The invention is typically carried out on a sample that is known to contain or suspected to contain the analyte. The invention may be carried out on a sample that contains one or more analytes whose identity is unknown. Alternatively, the invention may be carried out on a sample to confirm the identity of one or more analytes whose
25 presence in the sample is known or expected.

The sample may be a biological sample. The invention may be carried out *in vitro* on a sample obtained from or extracted from any organism or microorganism. The organism or microorganism is typically prokaryotic or eukaryotic and typically belongs to one the five kingdoms: plantae, animalia, fungi, monera and protista. The invention may be carried out *in vitro*
30 on a sample obtained from or extracted from any virus. The sample is preferably a fluid sample. The sample typically comprises a body fluid of the patient. The sample may be urine, lymph, saliva, mucus or amniotic fluid but is preferably blood, plasma or serum. Typically, the sample is human in origin, but alternatively it may be from another mammal animal such as from commercially
35 farmed animals such as horses, cattle, sheep or pigs or may alternatively be pets such as cats or dogs.

The sample may be a non-biological sample. The non-biological sample is preferably a fluid sample. Examples of a non-biological sample include surgical fluids, water such as drinking water, sea water or river water, and reagents for laboratory tests.

5 The sample is typically processed prior to being assayed, for example by centrifugation or by passage through a membrane that filters out unwanted molecules or cells, such as red blood cells. The sample may be measured immediately upon being taken. The sample may also be typically stored prior to assay, preferably below -70°C .

Conditions

10 The method of the invention involves the measuring of a current passing through the pore during interaction with the analyte. Suitable conditions for measuring ionic currents through transmembrane protein pores are well known in the art and disclosed in the Example. The method is carried out with a voltage applied across the membrane and pore. The voltage used is typically from -400mV to $+400\text{mV}$. The voltage used is preferably in a range having a lower limit selected
15 from -400 mV , -300mV , -200 mV , -150 mV , -100 mV , -50 mV , -20mV and 0 mV and an upper limit independently selected from $+10\text{ mV}$, $+20\text{ mV}$, $+50\text{ mV}$, $+100\text{ mV}$, $+150\text{ mV}$, $+200\text{ mV}$, $+300\text{ mV}$ and $+400\text{ mV}$. The voltage used is more preferably in the range 120mV to 170mV . Enhancing translocation of the analyte through the pore in accordance with the invention allows lower voltages to be used.

20 The method is carried out in the presence of any alkali metal chloride salt. In the exemplary apparatus discussed above, the salt is present in the aqueous solution in the chamber. Potassium chloride (KCl), sodium chloride (NaCl) or caesium chloride (CsCl) is typically used. KCl is preferred. The salt concentration is typically from 0.1 to 2.5M , from 0.3 to 1.9M , from 0.5 to 1.8M , from 0.7 to 1.7M , from 0.9 to 1.6M or from 1M to 1.4M . The salt concentration is preferably from
25 150 to 500mM . High salt concentrations provide a high signal to noise ratio and allow for currents indicative of the presence of a analyte to be identified against the background of normal current fluctuations. Lower salt concentrations must be used if analyte detection is carried out in the presence of an enzyme, such as when sequencing nucleic acids. The salt concentration is preferably from 150 to 500mM . Good nucleotide discrimination at these low salt concentrations can be
30 achieved by carrying out the method at temperatures above room temperature, such as from 30°C to 40°C .

The method is typically carried out in the presence of a buffer. In the exemplary apparatus discussed above, the buffer is present in the aqueous solution in the chamber. Any buffer may be used in the method of the invention. One suitable buffer is Tris-HCl buffer. The method is

typically carried out at a pH of from 4.0 to 10.0, from 4.5 to 9.5, from 5.0 to 9.0, from 5.5 to 8.8, from 6.0 to 8.7 or from 7.0 to 8.8 or 7.5 to 8.5. The pH used is preferably about 7.5.

The method is typically carried out at from 0°C to 100°C, from 15°C to 95°C, from 16°C to 90°C, from 17°C to 85°C, from 18°C to 80°C, 19°C to 70°C, or from 20°C to 60°C. The method
5 may be carried out at room temperature. The method is preferably carried out at a temperature that supports enzyme function, such as about 37°C. Good analyte discrimination can be achieved at low salt concentrations if the temperature is increased.

In addition to increasing the solution temperature, there are a number of other strategies that can be employed to increase the conductance of the solution, while maintaining conditions that are
10 suitable for enzyme activity, for example when sequencing nucleic acids. One such strategy is to use the lipid bilayer to divide two different concentrations of salt solution, a low salt concentration of salt on the enzyme side and a higher concentration on the opposite side. One example of this approach is to use 200 mM of KCl on the *cis* side of the membrane and 500 mM KCl in the *trans*
15 chamber. At these conditions, the conductance through the pore is expected to be roughly equivalent to 400 mM KCl under normal conditions, and the enzyme only experiences 200 mM if placed on the *cis* side. Another possible benefit of using asymmetric salt conditions is the osmotic gradient induced across the pore. This net flow of water could be used to pull nucleotides into the pore for detection. A similar effect can be achieved using a neutral osmolyte, such as sucrose, glycerol or PEG. Another possibility is to use a solution with relatively low levels of KCl and rely
20 on an additional charge carrying species that is less disruptive to enzyme activity.

Method of sequencing nucleic acids

The present invention also relates to a method of sequencing a target nucleic acid sequence. The method comprises (a) pushing or pulling the target sequence through a transmembrane pore of
25 the invention so that a proportion of the nucleotides in the target sequence interacts with the pore and (b) measuring the current passing through the pore during each interaction and thereby determining the sequence of the target sequence. Hence, the method involves stochastic sensing of a proportion of the nucleotides in a target nucleic acid sequence as the nucleotides pass through the barrel or channel in a successive manner in order to sequence the target sequence.

Any pore of the invention may be used. A pore of the invention in which the net positive charge of the barrel or channel and/or entrance has been increased is particularly suited to this
30 method. For instance, the enhanced translocation of the negatively charged nucleic acid through the pore reduces the non-translocation events and deadtime between reads of each nucleotide in the sequence and/or increases the read time for each nucleotide in the sequence. In a preferred
35 embodiment, the method further comprises enhancing enhancing translocation of the analyte

through the pore as described above. The enhancement is of course carried out before the nucleic acid is sequenced.

The whole or only part of the target nucleic acid sequence may be sequenced using this method. The nucleic acid sequence can be any length. For example, the nucleic acid sequence can
5 be at least 10, at least 50, at least 100, at least 150, at least 200, at least 250, at least 300, at least 400 or at least 500 nucleotides in length. The nucleic acid sequence is preferably DNA or RNA. The nucleic acid sequence can be naturally occurring or artificial. For instance, the method may be used to verify the sequence of a manufactured oligonucleotide. The method is typically carried out *in vitro*.

10 The method may be carried out using any of the samples, membrane/pore systems and conditions discussed above.

Interaction between the pore and nucleotides

The nucleic acid may be contacted with the pore on either side of the membrane. The
15 nucleic acid may be introduced to the pore on either side of the membrane. The nucleotide or nucleic acid is typically contacted with the side of the membrane on which an enzyme is attached to the pore. This allows the enzyme to handle the nucleic acid during the method as discussed below.

A proportion of the nucleotides of the target nucleic acid sequence interact with the pore as it passes across the membrane through the barrel or channel of the pore.

20 The nucleotides may interact with the pore in any manner and at any site. As discussed above, the nucleotides preferably reversibly bind to the pore.

During the interaction between the nucleotides and the pore, the nucleotide affects the current flowing through the pore in a manner specific for that nucleotide. For example, a particular nucleotide will reduce the current flowing through the pore for a particular mean time period and to
25 a particular extent. In other words, the current flowing through the pore is distinctive for a particular nucleotide. Control experiments may be carried out to determine the effect a particular nucleotide has on the current flowing through the pore. Results from carrying out the method of the invention on a test sample can then be compared with those derived from such a control experiment in order to identify a particular nucleotide.

30

Pushing or pulling the nucleic acid through the pore

Strand sequencing in accordance with the invention involves the controlled and stepwise translocation of nucleic acid polymers through a pore. The nucleic acid may be pushed or pulled through the pore in any manner known to a person skilled in the art. For instance, the nucleic acid

could be pushed or pulled through the pore using an electroosmotic effect. The nucleic acid is preferably pushed or pulled through the pore using a DNA handling enzyme.

The majority of nucleic acid handling enzymes are suitable for use in this application provided they hydrolyse, polymerise or process single stranded nucleic acids, such as DNA or
5 RNA. Preferred enzymes that are capable of pushing or pulling the target nucleic acid sequence through the pore include polymerases, exonucleases, helicases and topoisomerases, such as gyrases. The polymerase is preferably a member of any of the Enzyme Classification (EC) groups 2.7.7.6, 2.7.7.7, 2.7.7.19, 2.7.7.48 and 2.7.7.49. The polymerase is preferably a DNA-dependent DNA
10 polymerase, an RNA-dependent DNA polymerase, a DNA-dependent RNA polymerase or an RNA-dependent RNA polymerase. The helicase is preferably a member of any of the Enzyme Classification (EC) groups 3.6.1.- and 2.7.7.-. The helicase is preferably an ATP-dependent DNA helicase (EC group 3.6.1.8), an ATP-dependent RNA helicase (EC group 3.6.1.8) or an ATP-independent RNA helicase. The topoisomerase is preferably a member of any of the Enzyme
15 Classification (EC) groups 5.99.1.2 and 5.99.1.3.

The two strategies for single strand DNA sequencing are the translocation of the DNA through the nanopore, both *cis* to *trans* and *trans* to *cis*, either with or against an applied potential. The most advantageous mechanism for strand sequencing is the controlled translocation of single
20 strand DNA through the nanopore with an applied potential. Exonucleases that act progressively or processively on double stranded DNA can be used on the *cis* side of the pore to feed the remaining single strand through under an applied potential or the *trans* side under a reverse potential. Likewise, a helicase that unwinds the double stranded DNA can also be used in a similar manner. There are also possibilities for sequencing applications that require strand translocation against an
25 applied potential, but the DNA must be first "caught" by the enzyme under a reverse or no potential. With the potential then switched back following binding the strand will pass *cis* to *trans* through the pore and be held in an extended conformation by the current flow. The single strand DNA exonucleases or single strand DNA dependent polymerases can act as molecular motors to pull the recently translocated single strand back through the pore in a controlled stepwise manner, *trans* to
30 *cis*, against the applied potential.

Kit

The present invention also relates to a kit that may be used to carry out the method of sequencing a target nucleic acid sequence. The kit is therefore suitable for sequencing nucleic acids. The kit comprises a transmembrane pore of the invention and a nucleic acid handling enzyme. The pore and the enzyme may be any of those discussed above.

The kit may additionally comprise one or more other reagents or instruments which enable any of the embodiments of the method mentioned above to be carried out. Such reagents or instruments include one or more of the following: suitable buffer(s) (aqueous solutions), means to obtain a sample from a subject (such as a vessel or an instrument comprising a needle), means to amplify nucleic acid sequences, a membrane as defined above or voltage or patch clamp apparatus. Reagents may be present in the kit in a dry state such that a fluid sample resuspends the reagents. The kit may also, optionally, comprise instructions to enable the kit to be used in the method of the invention or details regarding which patients the method may be used for. The kit may, optionally, comprise nucleotides.

The following Examples illustrate the invention:

Example 1

Here, by augmenting the internal positive charge within the α -hemolysin pore and altering its distribution, we increase the capture rate of a 92-nt single-stranded DNA by ~ 10 -fold over the wild-type pore at +120 mV and dramatically lower the voltage threshold at which the DNA is translocated through the pore, e.g. by 50 mV for 1 event $s^{-1}mM^{-1}$. Further, events in which DNA enters the pore, but is not translocated, are almost eliminated. These engineered nanopores provide a basis for improved nucleic acid analysis and may assist in implementing ultrarapid nanopore sequencing.

1. Materials and methods

1.1 Protein preparation

α -HL was produced as described elsewhere (Cheley S, Braha O, Lu X, Conlan S, Bayley H: A functional protein pore with a "retro" transmembrane domain. *Protein Sci.* 1999, 8:1257-1267). The sequence of one subunit of α -HL is shown in SEQ ID NO: 2. In short, the protein was expressed in the presence of [35 S]methionine in an *E. coli in vitro* transcription and translation (IVTT) system (*E. coli* T7 S30 Extract System for Circular DNA, Cat. # L1130, Promega, Madison, WI). IVTT reactions (100 μ l) containing α -HL monomers were incubated with rabbit red blood cell membranes for 1 h at 37°C. The solution was spun at 25,000 Xg and the pellet containing α -HL heptamers was loaded onto a 5% SDS-polyacrylamide gel. The gel was run for 4 h at 100 V and vacuum dried for 3 to 4 h without heating onto Whatman 3M filter paper. The dried gel was exposed to photographic film for 2 h and the developed film was used to locate the position of the heptameric protein in the gel. This region of the gel was cut out, rehydrated and crushed in 400 μ L

of 10 mM Tris.HCl, pH 8.0, containing 100 μ M EDTA. After 20 min at room temperature, the polyacrylamide was removed by centrifuging the suspension at 25,000 Xg for 7 min at room temperature through a cellulose micro spin column (Microfilterfuge tubes, Cat. # 7016-024, Rainin, Oakland, CA). Aliquots of the purified protein were stored at -80°C.

5 Mutants α -HL genes were prepared either by using a site-directed mutagenesis kit (QuickChange® II XL, Cat. # 200522-5, Stratagene, Cedar Creek, TX) or by cassette mutagenesis (Cheley S, Gu L-Q, Bayley H: Stochastic sensing of nanomolar inositol 1,4,5-trisphosphate with an engineered pore. *Chem.Biol.* 2002, 9:829-838). The RL2 gene encodes four amino acid replacements in the barrel domain (Val-124→Leu, Gly-130→Ser, Asn-139→Gln, Ile-142→Leu) and one in the amino latch region of the pore (Lys-8→Ala). The sequence of one subunit of α -HL-RL2 is shown in SEQ ID NO: 4. The DNA sequence of each new gene was verified.

1.2 Planar bilayer recordings

A bilayer of 1,2-diphytanoyl-*sn*-glycero-3-phosphocholine (Avanti Polar Lipids) was formed on an aperture (~100 μ m in diameter) in a 25- μ m thick polytetrafluoroethylene film (Goodfellow Cambridge Limited, Cat. no. FP301200/10, Huntingdon, UK) that divided a chamber into *cis* and *trans* compartments. Both compartments contained 0.4 mL of 1 M KCl, 25 mM Tris.HCl, pH 8, containing 100 μ M EDTA. Planar bilayer current recordings were performed with a patch clamp amplifier (Axopatch 200B, Axon Instruments, Foster City, CA). The protein and subsequently the DNA were added to the *cis* compartment, which was connected to ground. The DNA was prepared and PAGE-purified by Sigma Genosys, UK. The concentration of the DNA was determined by measuring the absorbance at 260 nm of the solution in the *cis* chamber at the end of each experiment. The DNA sequence was: 5'-
 AAAAAAAAAAAAAAAAAAAAAAAAAATTCccccccccccccccccccccctTAAAAAAAAAAATT
 25 CccccccccctTAAAAAAAAAAATTCcccccccc-3' (SEQ ID NO: 15).

At applied potentials above +120 mV, the amplified signal arising from the ionic current passing through the α -HL pore was low-pass filtered at 50 kHz and sampled at 250 kHz with a computer equipped with a Digidata 1440A digitizer (Molecular Devices, Sunnyvale, CA). At applied potentials of +100 mV and +80 mV the signal was filtered at 40 kHz, while at +60 mV and +40 mV the signal was filtered at 30 kHz and 20 kHz, respectively. The sampling rate was at least five times the filtering rate. The data used in Table 2 were analyzed with pClamp 10.1 software (Axon Instruments), by using three different current levels. Level 0 corresponded to the open channel current, level 1 to the mid-amplitude event current and level 2 to the low-amplitude event current. The assignment to each level was checked manually.

Table 2 - Distribution of the types of event as a percentage of the total. The error is the standard error of the mean for four (WT, E111N-WT and M113R-WT) or five (M113R-RL2 and RL2) experiments.

5

	Type A	Type B	Type C	Type D	Type E
WT	67.6±2.9	19.6±2.7	10.2±1.2	2.0±0.5	0.6±0.3
M113R-WT	94.1±1.7	1.2±0.3	3.3±1.2	1.4±0.2	0
E111N-WT	97.5±0.5	0.6±0.1	1.1±0.4	0.8±0.2	0
RL2	84.0±3.2	7.3±2.3	4.8±0.7	3.5±0.7	0.4±0.2
M113R-RL2	91.4±1.3	1.7±0.5	4.1±0.8	2.6±0.2	0.2±0.1

The data used in Fig. 2b were analyzed and prepared for presentation by using only two levels: level 0 was set to the open channel current, and level 1 was placed between the mid-amplitude and low-amplitude event currents. The frequency of DNA translocations was calculated by fitting the interevent intervals to a single exponential (logarithmic binning, 10 bin/decade) and normalizing the mean value of the duration to a DNA concentration of 1 μ M. The DNA translocation time was calculated from the peak of a histogram of the distribution of the translocation times (Meller A, Nivon L, Brandin E, Golovchenko J, Branton D: Rapid nanopore discrimination between single polynucleotide molecules. *Proc.Natl.Acad.Sci.USA* 2000, 97:1079-1084).

1.3 Curve fitting

The values of f_{\max} , b and V_0 (Table 3) were determined by fitting the voltage dependence of the frequency of DNA translocation for the WT and mutated α -HL nanopores to equation 1 (Fig. 5) by using Origin data analysis software (OriginLab, Northampton, MA). The errors in these values are expressed as standard errors. The curve fittings were adjusted for the experimental error of each point by using the instrumental weight.

Table 3 - Values of ' f_{\max} ', ' b ' and ' V_0 ' for various α -HL pores. The values were derived by fitting the data to equation 1 (see Methods). Additional conditions are described in the legend to Fig. 2.

	f_{max}	33 b	V_0	V_1^a
M113R-WT	7,300±2,200	1200±100	-90±7	45
E111N-WT	1,600±200	780±40	-56±3	50
M113R-RL2	990±300	990±90	-67±7	77
WT	300±60	420±40	25±6	100
RL2	300±150	980±130	20±17	151

2. Results and discussion

2.1 Electrical recording of DNA translocation through the WT α -HL nanopore

5 Single-channel electrical recordings in planar lipid bilayers were carried out in 1 M KCl, 25 mM Tris.HCl at pH 8.0, containing 100 μ M EDTA. Under these conditions, the WT α -HL pore had a unitary conductance of 1.04 ± 0.01 nS (+120 mV, n = 23). After the addition of a 92-nucleotide synthetic ssDNA to the *cis* compartment (0.5 to 2.5 μ M), the current through a single pore was interrupted by short-lived blockades owing to the translocation of the ssDNA

10 (Kasianowicz JJ, Brandin E, Branton D, Deamer DW: Characterization of individual polynucleotide molecules using a membrane channel. *Proc.Natl.Acad.Sci.USA* 1996, 93:13770-13773). At +120 mV, the interactions of the DNA molecules with the WT nanopore produced five different current patterns (type A, B, C, D and E) that are combinations of three conductance levels: the open state, a mid-amplitude state and a low-amplitude state (Fig. 2a) (Butler TZ, Gundlach JH, Troll MA: Determination of RNA orientation during translocation through a biological nanopore. *Biophys J* 15 2006, 90:190-199). Events with a duration of shorter than 10 μ s were judged to represent the transient interaction of DNA with the mouth of the pore and were ignored in our analysis. As substantiated below, in all five types of event, DNA is captured by the pore, but only in type A and C events is the DNA translocated through the pore, from the *cis* to the trans side.

20 In a histogram displaying the mean current associated with each event (Fig. 2b), the current levels are separated into two peaks centered at 0.11 times (low amplitude) and 0.58 times (mid amplitude) the open-pore current. The type A events characterized by a fractional current of 0.11 (Fig. 2b) had a most likely duration of 0.14 ± 0.01 ms (n = 7) as determined from the peak of a Gaussian fit to a histogram of the event durations (Fig. 2c) (Meller A, Nivon L, Brandin E, Golovchenko J, Branton D: Rapid nanopore discrimination between single polynucleotide molecules. *Proc.Natl.Acad.Sci.USA* 2000, 97:1079-1084). The peak in the events histogram at a

25

fractional current of 0.58 comprises type B current blockades, which show a mid-state current level only (Fig. 2b). Type C, D and E events contain both mid-amplitude and low-amplitude current levels. Type C events corresponded to 10% of the total number of events, type D events 2% of the total and type E events, which displayed a mid-low-mid pattern, were only 0.6% of the total. The type C, D and E events are scattered between the two main peaks in the histogram, reflecting their wide distribution of mean currents.

2.2 DNA interactions with engineered nanopores

We made a variety of homoheptameric α -HL pores in which the charge distribution within the lumen was altered. The pores examined were homoheptamers, so the mutations appear in all seven subunits, e.g. M113R has a ring of seven positively charged side chains at position 113. The mutants were made by using the WT or RL2 genes as templates. The RL2 gene encodes four amino acid replacements in the barrel domain (Val-124→Leu, Gly-130→Ser, Asn-139→Gln, Ile-142→Leu) and one in the *cis* entrance of the pore (Lys-8→Ala). The properties of the mutants differed significantly from the WT pore with respect to the open-pore current and the frequency of occurrence of DNA translocation events (Table 4).

Table 4 - DNA translocation through the α HL nanopore. τ is the most probable translocation time; f is the frequency of translocation (type A and C events) normalized to 1 μ M DNA; I_B is the fractional residual current during a type A event and I_{+120mV} is the open-pore current at +120mV. The errors are expressed as the standard error of the mean and the number of experiments is in parentheses. For I_B the errors were all within ± 0.01 .

	τ (ms)	f (s ⁻¹ μ M ⁻¹)	I_B	I_{+120mV} (pA)
WT	0.141±0.004 (n=6)	3.0±0.2 (n=12)	0.11 (n=7)	125±1 (n=23)
RL2	0.147±0.003 (n=4)	0.28±0.04 (n=5)	0.10 (n=4)	124±2 (n=16)
A8R-RL2	0.15±0.01 (n=7)	7.6±0.7 (n=5)	0.10 (n=7)	125±3 (n=10)
A8K-RL2	0.16±0.03 (n=3)	3.0±0.3 (n=3)	0.11 (n=3)	123±3 (n=7)
M113R-WT	0.138±0.003 (n=5)	23±2 (n=9)	0.04 (n=5)	140±2 (n=13)
E111N-WT	0.28±0.04 (n=10)	18±1 (n=9)	0.11 (n=10)	137±3 (n=15)
M113K-RL2	0.092±0.002 (n=3)	4.0±0.3 (n=3)	0.05 (n=3)	129±2 (n=10)
M113R-RL2	0.141±0.004 (n=4)	4.4±0.6 (n=7)	0.03 (n=5)	131±2 (n=18)
T115R-RL2	0.23±0.02 (n=3)	2.2±0.1 (n=7)	0.02 (n=3)	122±2 (n=16)
T117R-RL2	0.35±0.04 (n=8)	1.6±0.1 (n=7)	0.03 (n=4)	113±6 (n=9)
G119R-RL2	0.292±0.002 (n=4)	1.3±0.1 (n=7)	0.02 (n=4)	111±1 (n=8)
N121R-RL2	0.30±0.01 (n=9)	0.43±0.03 (n=10)	0.03 (n=6)	96±2 (n=13)
N123R-RL2	0.22±0.02 (n=5)	0.43±0.04 (n=8)	0.03 (n=4)	77±2 (n=10)
M113D-RL2	not observed (n=3)	not observed (n=3)	not observed (n=3)	112±3 (n=3)

The homoheptamers formed from M113R-WT (seven subunits of SEQ ID NO: 6) and E111N-WT (seven subunits of SEQ ID NO: 8) showed the largest increases in ionic current at +120 mV with respect to the WT pore (11% and 9%, respectively), while N123R-RL2 showed the largest

current decrease (38%, Table 4). After the addition of DNA to the *cis* chamber at +120 mV, current blockades were observed for all the mutants tested except for M113D-RL2, in which case the addition of 2 μ M DNA did not produce any blockades at up to +300 mV ($n = 3$).

The distribution of the five types of interaction of DNA with the nanopore (Fig. 2a) depended on the mutant (see also Fig. 7). For pores other than WT, RL2, A8R-RL2 and A8K-RL2, the numbers of captures without translocation were negligible and therefore the term "translocation" is used henceforth. Nevertheless, in all cases, the number of type A and type C events was used to determine the frequency of DNA translocation.

Substitution of Lys-8 with an amino acid with a neutral side chain (mutant RL2) increased the relative proportion of type A events (84% versus 68% in WT) and decreased the type B and C events (7% and 5% in RL2, respectively, compared to 20% and 10% in WT). The elimination of the charge at the central constriction or the addition of a positive charge to the interior of the β barrel also increased the percentage of the type A events, and again the type B and C events were less well represented by comparison with the WT pore (Table 2). Recordings from E111N-WT, M113R-WT and M113R-RL2 (seven subunits of SEQ ID NO: 14) showed more than 90% of type A events (98%, 94% and 91%, respectively) and less than 6% of type B and C events (2%, 5% and 6%, respectively). The fraction of the minor type D and E events was similar in WT, RL2 and M113R-RL2, but lower in M113R-WT and E111N-RL2 (Table 2).

The translocation times of DNA through the α -HL nanopores were not changed considerably by the point mutations. The mutant with the slowest mean translocation time was N117R-RL2 at 0.35 ± 0.04 ms, which is only twice the mean translocation time observed for the WT pore (Table 4). The various mutants showed different extents of current block during the translocation of DNA through the pore (as determined from the type A events or the lowest level of type C events). WT, RL2, A8K-RL2 and E111N-WT showed fractional residual currents of 0.10 of the open channel current, while the introduction of positive charges in the β barrel produced higher extents of block (Table 4).

2.3 Effects of point mutations on the DNA capture rate

The effects of charged amino acid side chains on the passage of DNA through the α -HL pore were examined at +120 mV (Fig. 3). Lys-8 is the charged residue in the closest proximity to entering DNA. The substitution of Lys-8 with Ala (in the mutant RL2) decreased the number of translocation events per unit time by almost an order of magnitude, while substitution with Arg (in A8R-RL2 i.e. seven subunits of SEQ ID NO: 10) produced a three-fold increase in the translocation rate (Table 4). The additional mutations in RL2 (with respect to WT) did not affect DNA

translocation, as confirmed by reintroducing Lys-8 into the RL2 background (A8K-RL2, Table 4 and Fig. 8).

Removal of the positive charge from the *cis* entrance of the pore combined with the introduction of a positive charge just below the constriction restored (M113K-RL2 i.e. seven
 5 subunits of SEQ ID NO: 12) or even slightly enhanced (M113R-RL2) the frequency of translocation (Fig. 3 and Table 4). The effect of charge is cumulative as showed by the ten-fold increase in translocation frequency of M113R-WT compared to WT and the increase of two orders of magnitude when M113R-WT is compared to RL2 (Fig. 3 and Table 4). By contrast with the effects of positive charge, the placement of negative charge just below the constriction (M113D-
 10 RL2) eliminated DNA translocation at all potentials tested (+100 to +300 mV). In accord with these findings, removal of the negative charge at the constriction (E111N-WT) increased the frequency of translocation six-fold.

Amino acid substitutions within the β barrel of α -HL also affected the frequency of DNA translocation. Individual replacements of the amino acids at positions 111, 113, 115, 117, 119, 121
 15 and 123 with Arg (in the RL2 background) produced α -HL pores that showed a progressive decrease in the translocation frequency as the Arg residue became located further from the *cis* entrance (Fig. 4 and Table 4). By contrast, when DNA was added to the *trans* side of the protein, the effect was reversed; the translocation of DNA molecules through the pore was enhanced by a ring of arginines close to the *trans* entrance when compared with one near the constriction site (Fig.
 20 9).

2.4 Voltage dependence of DNA translocation and reduction of the voltage threshold for translocation

We found that the frequency of DNA translocation $f(V)$ was described well by:

$$25 \quad f(V) = f_{\max} e^{\frac{b}{V_0 - V}} \quad \text{for } V > V_0, \quad (1)$$

where ' f_{\max} ' is the rate of DNA translocation at a hypothetical infinite applied potential, ' V_0 ' is the threshold for DNA translocation, and a larger value of 'b' gives a faster approach to ' f_{\max} ' (Fig. 5). More positive charge produced pores with a greatly reduced threshold and a steeper approach to f_{\max} (Table 3). Further, pores with no additional charge at the constriction had lower values of f_{\max}
 30 ($300 \pm 60 \text{ s}^{-1} \mu\text{M}^{-1}$ and $300 \pm 150 \text{ s}^{-1} \mu\text{M}^{-1}$ for WT and RL2, respectively), while additional positive charge at both the constriction and the *cis* entrance gave high values of f_{\max} ($7,300 \pm 2,200 \text{ s}^{-1} \mu\text{M}^{-1}$ and $1,600 \pm 200 \text{ s}^{-1} \mu\text{M}^{-1}$ for M113R-WT and E111N-WT, respectively). A pore formed with a ring of additional charge at the constriction, but no charge at the *cis* entrance (M113R-RL2), displayed an intermediate value of f_{\max} ($990 \pm 300 \text{ s}^{-1} \mu\text{M}^{-1}$).

2.5 Dwell-time / amplitude distributions of the events for M113R-WT and E111N-WT

Dwell-time / amplitude distributions of the events for the M113R-WT and E111N-WT homoheptamers (but not the other mutants) yielded two distinct clusters of current blockades at 100 to 200 mV (Fig. 7). The relative intensities of the two peaks in the distributions were voltage dependent, with the longer events increasing with the applied potential (Fig. 7c). The first peak comprised blockades with a fast translocation time and a fractional residual current at +120 mV of 0.96 for M113R-WT and 0.89 for E111N-WT. The second cluster of current blockades included events with a longer dwell time and almost full current block. It is unlikely that the two clusters can be accounted for by 5' versus 3' threading (Mathé J, Aksimentiev A, Nelson DR, Schulten K, Meller A: Orientation discrimination of single-stranded DNA inside the alpha-hemolysin membrane channel. *Proc Natl Acad Sci USA* 2005, 102:12377-12382), because the differences in dwell times are too great (~100-fold).

In this study, the blockades at +120 mV are considered. At this potential, the short-lived events dominate in the dwell-time / amplitude distributions for M113R-WT and E111N-WT and it is these events, which are predominantly type A (Table 2), that are considered. To determine the frequency of DNA translocation both short and long events were included, but the contribution of the latter is minor.

2.6 Assignment of intermediate V

The assignment of intermediate V is supported by the findings of Kasianowicz and co-workers, who showed that, under a positive applied potential, the cis addition of ssDNA produces short lived current blockades of 85-100% of the open pore current. PCR analysis of the trans chamber revealed that DNA had passed through the α -HL nanopore (Kasianowicz JJ, Brandin E, Branton D, Deamer DW: Characterization of individual polynucleotide molecules using a membrane channel. *Proc.Natl.Acad.Sci.USA* 1996, 93:13770-13773).

Further, Movileanu and colleagues observed that individual PEG molecules covalently attached inside the β barrel produce a much higher reduction of the open pore current (70%) than when they are attached within the vestibule (20%) (Movileanu L, Cheley S, Howorka S, Braha O, Bayley H: Location of a constriction in the lumen of a transmembrane pore by targeted covalent attachment of polymer molecules. *J.Gen.Physiol.* 2001, 117:239-251).

Howorka and Bayley found that DNA duplexes covalently attached near the cis entrance with overhangs that protrude into the β barrel produce a current reduction of ~90% of the open pore current (Howorka S, Bayley H: Probing distance and electrical potential within a protein pore with tethered DNA. *Biophys.J.* 2002, 83:3202-3210). Finally, DNA immobilized inside the β barrel as part of a DNA- α -HL rotaxane structure causes a 85% block of the pore current (Sánchez-Quesada J,

Saghatelian A, Cheley S, Bayley H, Ghadiri MR: Single molecule DNA rotaxanes of a transmembrane pore protein. *Angew.Chem.Int.Ed.Engl.* 2004, 43:3063-3067).

2.7 Assignment of the mid-level events

5 Occupancy of the vestibule is associated with a reduced conductance (but not to the extent caused by occupancy of the barrel). For example, when high molecular weight PEG molecules (Movileanu L, Cheley S, Howorka S, Braha O, Bayley H: Location of a constriction in the lumen of a transmembrane pore by targeted covalent attachment of polymer molecules. *J.Gen.Physiol.* 2001, 117:239-251) or PAMAM dendrimers with a hydrodynamic diameter of 4.1 nm (Martin H, Kinns
10 H, Mitchell N, Astier Y, Madathil R, Howorka S: Nanoscale protein pores modified with PAMAM dendrimers. *J Am Chem Soc* 2007, 129:9640-9649) are tethered within the vestibule, the current is reduced by 20% and 45%, respectively. Second, when DNA duplexes are tethered within the vestibule, a 38% reduction of the pore current is observed (Howorka S, Bayley H: Probing distance and electrical potential within a protein pore with tethered DNA. *Biophys.J.* 2002, 83:3202-3210).
15 Finally, when the α -HL pore was genetically engineered so that the vestibule was filled with flexible polypeptides chains, the current was reduced by up to 70%, the reduction depending on the number of amino acids inserted (Jung Y, Cheley S, Braha O, Bayley H: The internal cavity of the staphylococcal α -hemolysin pore accommodates \sim 175 exogenous amino acid residues. *Biochemistry* 2005, 44:8919-8929).

20

2.8 Discussion

We have enhanced the frequency of translocation of DNA molecules by the α HL pore by using site-directed mutagenesis. The introduction or removal of charged groups within the lumen had a profound effect on DNA translocation. Increasing the positive charge near the internal
25 constriction increased the DNA translocation frequency at +120 mV by almost an order of magnitude (Fig. 3), and greatly decreased the threshold voltage for DNA translocation (Fig. 5). By contrast, eliminating the positive charge at the *cis* entrance decreased the frequency of DNA translocation by ten-fold. The effect of charge at the constriction on the DNA translocation frequency is surprising, as the *cis* entrance of the pore is 5 nm away and under the experimental
30 conditions of 1 M KCl the Debye length is 0.3 nm. Therefore, charged residues at the constriction are not expected to affect the capture of DNA through electrostatic interactions that extend into the bulk solution. Rather some form of "action at a distance" must be proposed, the most likely being rapid, reversible sampling of the lumen of the pore by DNA from the *cis* compartment or electroosmosis, both of which are discussed here.

35

A scheme for the interaction of DNA with the α -HL pore

A scheme for DNA capture and translocation that incorporates reversible sampling of the lumen is depicted (Fig. 6). Prior to a typical translocation event (via intermediate V, Fig. 6), a DNA molecule first collides with the pore (II). Upon visiting the vestibule (III), the DNA is transported
5 by electrophoresis into the narrow β barrel (IV). Because the pores used here are anion selective, electroosmotic solvent flow will occur in the same direction (cis to trans) and assist DNA transport. At this point, the effect of the applied field on the DNA molecule is strongest and the polymer is pulled through the pore into the *trans* compartment (V \rightarrow I), the only irreversible step in our scheme. The translocation intermediate (V) was observed experimentally by the abrupt reduction of
10 the current to 90% of the open pore value (during type A and C events), an assignment that is supported by several observations in the literature. For example, Kasianowicz and co-workers showed that the *cis* addition of ssDNA produces short-lived current blockades of 85-100% of the open pore current. PCR analysis of the *trans* chamber revealed that DNA had passed through the α HL pore (Kasianowicz, J. J., Brandin, E., Branton, D., & Deamer, D. W. (1996) Characterization
15 of individual polynucleotide molecules using a membrane channel. *Proc.Natl.Acad.Sci.USA* **93**, 13770-13773).

In the case of the WT- α HL pore, not all the current blockades were due to straightforward DNA translocation. In one-third of the events, a mid-amplitude event was observed (Fig. 2a). This current signature is interpreted to represent DNA molecules during a prolonged occupancy of the α -
20 HL vestibule (intermediate VI, Fig. 6), as supported by the fact that occupancy of the vestibule is associated with a lower conductance, but not one reduced to the extent caused by occupancy of the barrel. For example, when DNA duplexes are tethered within the vestibule, a 38% reduction of the pore current is observed (Howorka, S. & Bayley, H. (2002) Probing distance and electrical
25 potential within a protein pore with tethered DNA. *Biophys.J.* **83**, 3202-3210). Capture within the vestibule is similar to the phenomenon of entropic trapping described by Han and Craighead (Han, J. & Craighead, H. G. (2002) Characterization and optimization of an entropic trap for DNA
separation. *Anal Chem* **74**, 394-401).

DNA molecules captured in the mid-amplitude configuration (VI, Fig. 6) can either exit the pore from the *cis* compartment (type B event, Fig. 2a), or proceed into the β barrel producing a low-
30 amplitude signal (intermediate VII). From this configuration, the DNA can either complete translocation through the pore and exit on the *trans* side of the bilayer (type C event, Fig. 2a, via intermediate V, Fig. 6) or retract back into the vestibule and exit on the *cis* side (type E event, Fig. 2a). Finally, DNA can visit the vestibule, move at once into the β barrel to produce a low-amplitude signal, but instead of proceeding through the pore (type A event) reverse into the

vestibule to produce a mid-amplitude signal, before exiting on the *cis* side (type D, Fig. 2a). Butler and colleagues recently observed the same phenomena and came to similar conclusions (Butler TZ, Gundlach JH, Troll MA: Determination of RNA orientation during translocation through a biological nanopore. *Biophys J* 2006, 90:190-199). In further support of the scheme, we have
5 observed a few events with multiple mid-to-low (VI ↔ VII) steps (Fig. 10). In addition, we have observed type A-E events with additional DNAs, both with and without weak secondary structure (D. Japrun, unpublished work). The 92-mer used in the present work is unlikely to possess stem-loop structures, but the existence of helical structure within single-stranded nucleic acids remains a contentious area (Seol, Y., Skinner, G. M., Visscher, K., Buhot, A., & Halperin, A. (2007)
10 Stretching of homopolymeric RNA reveals single-stranded helices and base-stacking. *Phys Rev Lett* 98, 158103).

The effects of charge in the lumen of the pore

A ring of seven positive charges introduced by site-directed mutagenesis near the internal
15 constriction increased the frequency of translocation of DNA molecules (Fig. 3), and reduced the fraction of events with mid-amplitude components (Fig. 2ab). However, according to our proposed scheme, the DNA is within a Debye length of the residues forming the constriction only as intermediate IV, V or VII (Fig. 6) and cannot sense these charged groups from bulk solution. Therefore, to explain the effect of charge at the constriction, we propose that a DNA molecule
20 enters and quickly exits the top end of the β barrel (sampling, III ↔ IV, Fig. 6) many times before either retracting back into the *cis* compartment or proceeding to intermediate V, VI or VII. An increase in positive charge at the constriction (Glu-111→Asn or Met-113→Arg) increases the affinity of DNA for the pore at this location. Therefore, in these mutants the number of molecules that return to the vestibule after sampling the β barrel is reduced and the rate of occurrence of direct
25 DNA translocation events (type A) is increased. In addition, since the fraction of time spent as intermediate III is decreased, the probability of observing a mid-amplitude event (VI) is reduced and fewer type B to E events occur. The sampling of the β barrel by ssDNA was observed experimentally at +100 mV for short DNA overhangs protruding from a duplex DNA tethered near the *cis* entrance of the pore (Howorka S, Bayley H: Probing distance and electrical potential within
30 a protein pore with tethered DNA. *Biophys.J.* 2002, 83:3202-3210). For untethered DNA molecules, we propose that the sampling of the barrel is much quicker and not observed because of bandwidth limitations.

The elimination of a charge at the *cis* entrance of the pore in the RL2 mutant (Lys-8 Ala) reduced the fraction of type B and C events (Table 2), and decreased the frequency of DNA
35 translocation by an order of magnitude (Fig. 3). We attribute these effects to a reduced affinity of

DNA for the pore. A weaker interaction with the pore during collisions with the entrance and visits to the vestibule (intermediates II and III) reduces the overall number of observable captures.

Importantly, we have observed an increased number of capture events with α HL pores with increased internal positive charge with several additional DNAs and RNAs, as described here for the 92-mer, Therefore, the phenomena we describe are not peculiar to a specific DNA sequence.

Role of the β barrel

The depth to which DNA samples the β barrel (step IV) was investigated by introducing a ring of arginines at various positions. When DNA was added from the *cis* side, we found that the DNA translocation frequency decreased roughly exponentially as the ring was moved from the constriction (M113R-RL2) towards the *trans* entrance (N123R-RL2) of the pore (Fig. 4), suggesting that DNA molecules can sample the β barrel deeply. When presented from the *cis* side, the DNA must occasionally come close to residue 119, which is 2.3 nm away from the constriction. Accordingly, when the DNA was added on the *trans* side of the pore, more translocation events were observed (at -120 mV) when the ring of arginines was closer to the *trans* entrance (Fig. 9). As the ring of charges is moved towards the *trans* entrance, the unitary conductance of the pore is decreased by about twofold (Table 4), but this is unlikely to be fully related to the dramatic change in DNA translocation frequency of more than ten-fold. Further, in other cases, the trend is not seen. For example, RL2 has a similar conductance to the WT pore, but a far lower translocation frequency.

Electroosmotic effect

In the preceding discussion, we have ignored the differences in electroosmotic solvent flow through the various α -HL pores. For example, the WT α -HL pore is slightly anion selective ($P_+/P_- = 0.78$, in 1 M NaCl *cis* and 0.2 M NaCl *trans*) and, therefore, there will be a net flow of water from *cis* to *trans* under a positive potential (Gu L-Q, Cheley S, Bayley H: Electroosmotic enhancement of the binding of a neutral molecule to a transmembrane pore. *Proc.Natl.Acad.Sci.USA* 2003, 100:15498-15503). The M113R-WT pore is more anion selective than the WT pore ($P_+/P_- = 0.38$, in 1 M NaCl *cis* and 0.2 M NaCl *trans*) and the increased frequency of DNA translocation with this mutant could be due, at least in part, to enhanced electroosmotic flow. A simple calculation, based on equation 2 in Gu, L.-Q., Cheley, S., & Bayley, H. (2003) Electroosmotic enhancement of the binding of a neutral molecule to a transmembrane pore. *Proc.Natl.Acad.Sci.USA* 100, 15498-15503, assuming that five molecules of water are carried per ion, suggests that with a weakly anion selective pore ($P_+/P_- = 0.9$) only two additional molecules are transported to the mouth of the pore per second in a 1 μ M DNA solution. With a moderately selective pore ($P_+/P_- = 0.5$), 23 additional

molecules arrive per second and with a more strongly selective pore ($P_+/P_- = 0.1$) an additional 53. These numbers are of a similar order of magnitude to the effects seen here.

However, several lines of evidence suggest that electroosmosis does not have a dominant effect on the frequency of DNA translocation through the α -HL pore at +120mV. For example, E111N-WT shows a higher DNA translocation frequency than M113R-WT, but it is much less anion selective ($P_+/P_- = 0.71$, in 1 M NaCl *cis* and 0.2 M NaCl *trans*) (Gu L-Q, Cheley S, Bayley H: Electroosmotic enhancement of the binding of a neutral molecule to a transmembrane pore. *Proc.Natl.Acad.Sci.USA* 2003, 100:15498-15503). The mutant RL2 has a similar ion selectivity to WT α -HL ($P_+/P_- = 0.68$, in 0.9 M KCl *cis* and 0.3 M KCl, *trans* R. Madathil, unpublished), but the DNA translocation frequency with RL2 is ten-fold lower (Table 4). RL2 also shows the same DNA translocation frequency as N123R-RL2 (Fig. 5), which has a higher anion selectivity ($P_+/P_- = 0.24$, in 0.3 M NaCl *cis* and 1.0 M NaCl *trans*, R. Madathil, unpublished). Finally, if the trend in DNA translocation frequencies observed when a ring of arginines is placed at different positions within the β barrel were due to electroosmosis, we would not expect the trend to reverse when DNA is threaded from the *trans* rather than the *cis* side. Previous workers have recognized the possibility that electroosmosis might affect the rate of DNA translocation through nanopores (Chen P, Mitsui T, Farmer DB, Golovchenko JA, Gordon RG, Branton D: Atomic layer deposition to fine-tune the surface properties and diameters of fabricated nanopores. *Nano Letters* 2004, 4:1333-1337) while others have ignored the effect (Meller A: Dynamics of polynucleotide transport through nanometre-scale pores. *J.Phys.: Condens.Matter* 2003, 15:R581-R607) or considered it an unlikely contributor under prevailing conditions (Gershow M, Golovchenko JA: Recapturing and trapping single molecules with a solid-state nanopore. *Nat Nanotechnol* 2007, 2:775-779). It is also possible that electroosmosis has a role in decreasing the type B to E events by promoting laminar flow through the vestibule.

Movileanu and colleagues have explored the passage of positively charged peptides through α -HL pores with additional negative charge in the lumen (Wolfe, A. J., Mohammad, M. M., Cheley, S., Bayley, H., & Movileanu, L. (2007) Catalyzing the translocation of polypeptides through attractive interactions. *J Am Chem Soc* 129, 14034-14041; Mohammad, M. M., Prakash, S., Matouschek, A., & Movileanu, L. (2008) Controlling a single protein in a nanopore through electrostatic traps. *J Am Chem Soc* 130, 4081-4088). Both the rate of capture and the translocation time were increased, which they explained in terms of reduced barriers to transit. The differences between nucleic acids, for which the translocation time is not greatly altered (τ , Table 4), and peptides will require further experimentation to resolve, but one obvious distinction is the higher charge density on nucleic acids.

Voltage dependence of DNA translocation.

The frequency of DNA translocation events through α -HL pores depends strongly on the applied potential in a non-linear fashion (Fig 5a). At the low applied potentials of our experiments, the oligonucleotides that visit the vestibule (intermediate III, Fig. 6) do not sense a strong potential drop (<10 mV at an applied potential of +100 mV; Howorka S, Bayley H: Probing distance and electrical potential within a protein pore with tethered DNA. *Biophys.J.* 2002, 83:3202-3210). Further, the contributions of local potentials from fixed charges in the protein are expected to be small, given the high ionic strength. Therefore, only intermediates IV and V (Fig. 6) in which the DNA is inside the β barrel and subjected to a strong electric field are expected to be significantly influenced by the applied potential. The existence of a voltage threshold for DNA translocation and the steep dependence of the frequency of translocation on potential at <150 mV (Fig. 5) suggests that the energy barrier for DNA translocation is large and only a fraction of the DNA molecules that reach the β barrel (intermediate IV) pass through the entire pore. Increasing the charge near the constriction lowers the energy barrier and more DNA molecules can pass through the β barrel (see above). The lower energy barrier is also associated with a lower threshold for DNA translocation (Table 3). Other groups have observed the threshold and the steep voltage dependence with WT α -HL pores and fitted their observations to exponential functions (Henrickson SE, Misakian M, Robertson B, Kasianowicz JJ: Driven DNA transport into an asymmetric nanometer-scale pore. *Phys.Rev.Lett.* 2000, 85:3057-3060).

At higher potentials ($V > 150$ mV), the voltage dependence of DNA translocation is less steep (Fig. 5). Other groups, working with the WT α -HL pore, have described a linear dependence on voltage at high potentials (Meller A: Dynamics of polynucleotide transport through nanometre-scale pores. *J.Phys.: Condens.Matter* 2003, 15:R581-R607), but in our case the dependence is stronger. Interestingly, the maximum frequency of DNA translocation (f_{max}) for α -HL WT and the mutant pores varies considerably (Fig. 5), the hallmark of electroosmosis (Wong CT, Muthukumar M: Polymer capture by electro-osmotic flow of oppositely charged nanopores. *J Chem Phys* 2007, 126:164903), which may play a greater role under these conditions.

3. Conclusion

We have used site-directed mutagenesis to increase the frequency of ssDNA translocation through α -HL pores, and at the same time the threshold for DNA translocation has been lowered. The most likely mechanisms for the increased rate of translocation are rapid sampling of the interior of the pore by the DNA (Fig. 6), increased electroosmotic flow (Chen P, Mitsui T, Farmer DB, Golovchenko JA, Gordon RG, Branton D: Atomic layer deposition to fine-tune the surface

properties and diameters of fabricated nanopores. *Nano Letters* 2004, 4:1333-1337) or a combination of the two that depends on the applied potential. The new engineered pores or derivatives of them will be useful for enhancing the sensitivity of α -HL as a biosensor of nucleic acids. An additional potential application is in nanopore sequencing, where the in a single DNA strand are read off one by one during translocation. An increase in the frequency of ssDNA translocation through α -HL pores, as demonstrated here, might, for example, be used to reduce the deadtime between reads. By analogy, engineered pores might also be used to increase the capture efficiency of individual bases during exonuclease sequencing, which will be vital for accurate reads.

10 **Example 2**

Here, we show that the translocation of DNA molecules through the α -HL biological nanopore can be reduced by more than two orders of magnitude by introducing positive charges in the lumen of the pore. Although the ionic current during DNA translocation was almost completely suppressed, we suggest that these nanopores can be used to control the speed of DNA translocation, which, in combination with emerging technologies for DNA analysis, could lead to the development of a robust platform for the sequencing of nucleic acids.

1. **Materials and methods**

The same Materials and Methods used in Example 1 were also used in this Example.

2. **Results and discussion**

2.1. **Ionic current through α -HL nanopores**

Individual channels were examined by single-channel current recordings in planar lipid bilayers in 1M KCl, 25 mM Tris.HCl containing 100 μ M EDTA at pH 8.0. At +120 mV, all channels displayed stable open pore currents (+120 mV), however, some of the pores showed isolated, short-lived current blockade events, attributed to opening and closing of the pore (Kasianowickz *et al.*, *Proc.Natl.Acad.Sci.USA* 1996, *supra*). The unitary conductance values (g) varied greatly among nanopores (Table 5).

Table 5 - DNA translocation through the α -HL nanopores at +120mV. All experiment were carried out in 1M KCL, 25 mM Tris.HCl containing 100 μ M of EDTA. τ_G is the most likely DNA translocation speed, g is the unitary conductance of the nanopores and I_{RES} is the residual current percent during DNA translocation. All mutants, except for WT and 2N, are in RL2 background (SEQ ID NOs: 3 and 4). Errors are shown in standard deviation.

Pore	τ_G (μ s/base)	I_{RES}	g , nS (+120mV)
WT*	1.52±0.00 (n=6)	9.5 (n=5)	1.04±0.01 (n=23)
RL2*	1.63±0.00 (n=4)	10.5 (n=4)	1.03±0.02 (n=16)
2N (E111N-K147N)	2.01±0.07 (n=4)	26.5 (n=4)	1.07±0.05 (n=4)
M113R-RL2*	1.53±0.10 (n=4)	3.0 (n=5)	1.19±0.06 (n=8)
N123R-RL2*	2.39±0.22 (n=5)	3.0 (n=4)	0.64±0.02 (n=10)
2R \emptyset (M113R-N123R)	2.99±0.15 (n=5)	2.0 (n=4)	0.98±0.05 (n=6)
2R \bullet (N123R-D127R)	4.43±0.23 (n=5)	2.5 (n=4)	0.57±0.03 (n=5)
2R \dagger (M113R-T145R)	4.49±0.69 (n=5)	2.0 (n=5)	1.06±0.06 (n=5)
3R (T115R-G119R-123R)	16.0±4.0 (n=7)	0.5 (n=6)	0.87±0.06 (n=7)
4R (T115R-G119R- N123R-D127R)	29.0±5.5 (n=4)	0.5 (n=4)	0.70±0.09 (n=6)
7R (M113R-T115R-T117R- G119R-N121R-N123R- T125R)	269 ±14 (n=6)	0.5 (n=6)	0.87±0.02 (n=6)
7R $_6$ RL2 $_1$	100±35 (n=7)	0.5 (n=5)	0.91±0.02 (n=7)
7R $_5$ RL2 $_2$	66.6±10.1 (n=5)	0.5 (n=5)	0.88±0.09 (n=5)
7R $_4$ RL2 $_3$	51.2±6.1 (n=5)	0.5 (n=5)	0.86±0.05 (n=5)
7R $_3$ RL2 $_4$	21.4±8.8 (n=7)	1.0 (n=5)	0.94±0.05 (n=7)
7R $_2$ RL2 $_5$	9.89±2.15 (n=5)	1.5 (n=5)	0.93±0.07 (n=5)
7R $_1$ RL2 $_6$	5.95±1.10 (n=5)	4.0 (n=4)	0.99±0.05 (n=5)

* - data from Example 1 and Maglia, *Proc.Natl.Acad.Sci.USA*, 2008, *supra*

All nanopores in this study, with exception of WT and 2N (E111N-K147N), were made by
5 using the RL2 gene (SEQ ID NO: 3) as template, which encodes four amino acid mutations in the
 β -barrel domain (Val-124→Leu, Gly-130→Ser, Asn-139→Gln, Ile-142→Leu) and one in the cis
entrance of the pore (Lys-8→Ala) (SEQ ID NO: 4; Cheley, *Protein Sci.*, 1999, *supra*). The

introduction of positive charges close to the constriction site increased the ionic current of the nanopores with respect to the WT pore, as observed in the homoheptamers formed by M113R and M113R-T145R. By contrast, ionic currents were reduced significantly when positive charges were introduced close to the trans exit of the α -HL pore (Maglia *et al.*, (2008) *Proc.Natl.Acad.Sci.USA*, 5 **105(50)**: 19720-19725). N123R and N123R-D127R pores displayed the highest decrease in ionic current, with unitary conductance values of 0.64 ± 0.02 nS ($n=10, +120$ mV) and 0.57 ± 0.03 ($n=5, +120$ mV), respectively (Table 5). The current rectification ratios, calculated as the value of ionic current at +50 divided by the current at -50, also varied depending on the position of the charges. Nanopores with additional positive charges introduced near the trans exit (after position 121) 10 showed current rectification lower than 1 (e.g. they show higher current at negative applied potentials than at positive applied potentials), while mutants with positive charges introduced close to the cis side displayed similar rectifying behaviour to WT- α -HL (higher than 1). On the other hand, nanopores with no charges at the central constriction displayed almost non-rectifying behaviour ($R\sim 1$ in 2N)). These results suggest, in agreement with previous results (Wolfe, *J Am* 15 *Chem Soc*, 2007, *supra*; Maglia *et al.*, *Proc.Natl.Acad.Sci.USA*, 2008, *supra*), that the transport of molecules (e.g. ions) through the relatively small dimensions of the α -HL nanopore can be strongly altered by the introduction of charged residues in the β -barrel region of the pore.

2.2 DNA translocation through α -HL nanopores

20 The addition short ssDNA (92 bases nucleotide in this work) to the CIS side of the chamber under positive applied potential provokes current blockade, which are due to the translocation of single DNA molecules (Kasianowicz, *Proc.Natl.Acad.Sci.USA*, 1996, *supra*). The most likely translocation time, τ_G , is defined as the peak of the Gaussian fit the histogram of events duration (Figure 11a). For the WT nanopore, at +120mV of DNA through the pore τ_G is of 0.141 ± 0.004 ms, 25 which corresponds to a translocation speed of 1.53 ± 0.00 μ s/nt, ($n=6$). Manipulation of charges in the barrel of the pore altered the interaction between DNA and the pore. The exponential decrease of the dwell time of the current blockades induced by DNA, as shown for example for 7R nanopores (Figure 12b) indicates, however, that DNA also translocate through the altered nanopores.

30 The removal of the charges at the central constriction (E111N/147N) or the introduction of single (1R) and double (2R) arginine rings in the pore lumen had a negligible effect on the translocation speeds of DNA through the pore (the most likely translocation speeds in the 2R- α -HL mutants were just 2-3 fold lower relatively to WT- α -HL). The introduction of three (3R- α -HL) and

four (4R) rings of arginines in the barrel of the nanopore reduced the most likely translocation speed ~10 and ~20 folds, respectively. The strongest effect was observed upon addition of seven rings of arginines residues in the lumen of the pore (7R), which decreased the translocation speeds of DNA through the pore by more than two orders of magnitude relative to the WT and RL2 α -HL pores (269 \pm 14 μ s/nt Table 5).

2.3 Residual current through α -HL nanopores

The residual current during the translocation of DNA molecules through α -HL nanopores (I_{RES}) is defined as the blocked pore current (I_B) divided by the open pore current (I_O) expressed in percent. WT nanopores showed a fractional residual current of 11%. In DNA nanopore sequencing, I_{RES} is important because it is the most likely mean to which single bases will be discriminated as they are translocated through the pore. Unfortunately, while introductions of arginine residues allow the control of DNA translocation through α -HL nanopores, the residual current while DNA is translocating is greatly reduced (Table 5). Single and double arginines substitutions in the lumen of the pore reduced the residual current to 4% and 2%, respectively (Maglia, *Proc.Natl.Acad.Sci.USA*, 2008, *supra* and Table 5), while three or more arginine rings in the barrel reduced I_{RES} to 0.5% (Table 5). By contrast, the replacement of the charged residues at the constriction site with smaller non-charged asparagine residues (E111N-K147N), increased the fractional residual current to 26%.

The effect of amino acids modification in the barrel of the pore on the residual current through the α -HL pore while DNA is in the pore is most likely due to a combination of steric and electrostatic effect. Arginine has a relatively large volume (173.4 \AA^3) and when replace threonine (116.1 \AA^3) or glycine (60.1 \AA^3) residues will reduce the diameter of the pore and thus I_{RES} . Similarly, substitution of glutamate (138.4 \AA^3) and lysine (166.7 \AA^3) residues at the constriction with asparagine (117.4 \AA^3) will slightly increase the diameter of the constriction and thus I_{RES} . However, electrostatic interactions are likely to play a large role. In the WT and mutants, a reduction of the volume alone can not account for the that positive charges in the barrel of the pore will also provide an additional energy barrier for the passage of positive ion, which provide most of the ionic current during DNA translocation.

2.4 DNA Translocation through α -HL hetero-heptamers

Hetero-heptamers α -HL pores were prepared by mixing 7R monomers containing a D8 tail at the C termini with RL2 monomers on rabbit red blood cells, and mono-derivatized heptamers were separated by SDS-PAGE (SI). The modified pores contained, therefore, lines of single

arginine residues along the barrel of the pore stretching from the constriction to the trans exit of the pore, forming a slide to guide DNA translocation through the α -HL pore. It was hoped that the arginine residues would reduce DNA translocation by interacting strongly with the phosphate groups of DNA, while allowing K⁺ ions still to pass through the pore, thus allowing base recognition. Derivatized heptamers, however, showed very similar residual current and DNA translocation speeds as homo-heptamers with comparable positive charges in the barrel (Table 5). Interestingly, the translocation speed of DNA through the pore depended exponentially on the internal charge of the barrel (Figure 14), suggesting a strong polymer-pore interactions (Wanunu, M. Sutin, J. McNally, B. Chow, A. and A. Meller (2008), *Biophys. J.*, **95**, 4716-25; Luo *et al.*, (2007), *Phys.Rev.Lett*, **99**(14): 148102).

Tables 6 to 13 below set out the sequence information.

<p>Table 6 – SEQ ID NOs: 1 and 2 - Wild-type α-hemolysin from <i>Staphylococcus aureus</i></p> <p>ATGGCAGATTCTGATATTAATATTAAAACCGGTA CTACAGATATTGGAAGCAATACTACAGTAA AAACAGGTGATTTAGTCACTTATGATAAAGAAAATGGCATGCACAAAAAAGTATTTTATAGTTT TATCGATGATAAAAATCACAATAAAAAACTGCTAGTTATTAGAACAAAAGGTACCATTGCTGGT CAATATAGAGTTTATAGCGAAGAAGGTGCTAACAAAAGTGGTTTAGCCTGGCCTTCAGCCTTTA AGGTACAGTTGCAACTACCTGATAATGAAGTAGCTCAAATATCTGATTACTATCCAAGAAATTC GATTGATACAAAAGAGTATATGAGTACTTTAACTTATGGATTCAACGGTAATGTTACTGGTGAT GATACAGGAAAAATTGGCGGCCTTATTGGTGCAAATGTTTCGATTGGTCATACACTGAAATATG TTCAACCTGATTTCAAACAATTTTAGAGAGCCCAACTGATAAAAAAGTAGGCTGGAAAGTGAT ATTTAACAATATGGTGAATCAAATTTGGGGACCATACGATCGAGATTCTTGAACCCGGTATAT GGCAATCAACTTTTCATGAAAAGTAGAAATGGTCTATGAAAGCAGCAGATAACTTCCTTGATC CTAACAAAGCAAGTCTCTATTATCTTCAGGGTTTTCACCAGACTTCGCTACAGTTATTACTAT GGATAGAAAAGCATCCAAACAACAACAATATAGATGTAATATACGAACGAGTTCGTGATGAT TACCAATTGCATTGGACTTCAACAATTTGGAAAGGTACCAATACTAAAGATAAATGGACAGATC GTTCTTCAGAAAGATATAAAATCGATTGGGAAAAAGAAGAAATGACAAAT</p> <p>ADSDINIKTGTTDIGSNTTVKTGDLVITYDKENGMHKVVFYSFIDDKNHNKLLVIRTKGTIAGQ YRVYSEEGANKSGLAWPSAFKVQLQLPDNEVAQISDYPRNSIDTKEYMSTLTYGFNGNVTGDD TGKIGGLIGANVSIGHTLKYVQPDFKTI LESPTDKKVGWKVIFNNMVNQNWGPYDRDSWNPVYG NQLFMKTRNGSMKAADNFLDPNKASSLLSSGFS PDFATVITMDRKASKQQTNI DVIYERVRDDY QLHWTSTNWKGTNTKDKWTD RSSERYKIDWEKEEMTN</p>
--

<p>Table 7 – SEQ ID NOs: 3 and 4 - α-hemolysin RL2</p> <p>ATGGCAGATTCTGATATTAATATTGCAACCGGTA CTACAGATATTGGAAGCAATACTACAGTAA AAACAGGTGATTTAGTCACTTATGATAAAGAAAATGGCATGCACAAAAAAGTATTTTATAGTTT TATCGATGATAAAAATCACAATAAAAAACTGCTAGTTATTAGAACAAAAGGTACCATTGCTGGT CAATATAGAGTTTATAGCGAAGAAGGTGCTAACAAAAGTGGTTTAGCCTGGCCTTCAGCCTTTA AGGTACAGTTGCAACTACCTGATAATGAAGTAGCTCAAATATCTGATTACTATCCGCGGAATTC GATTGATACAAAAGAGTATATGAGT</p> <p>ACGTTAACGTACGGATTCAACGGTAACCTTACTGGTGATGATACTAGTAAAATTTGGAGGCCTTA TTGGGGCCAGGTTTCCCTAGGTCATACACTTAAGTATGTTCAACCTGATTTCAAACAATTTCT CGAGAGCCCAACTGATAAAAAAGTAGGCTGGAAAGTGATATTTAACAATATGGTGAATCAAAT TGGGGACCATACGATCGAGATTCTTGGAAACCCGGTATATGGCAATCAACTTTTCATGAAAAC TA GAAATGGTTCTATGAAAGCAGCAGATAACTTCCCTTGATCCTAACAAAGCAAGTTCCTCTATTATC</p>
--

TTCAGGGTTTTACCAGACTTCGCTACAGTTATTACTATGGATAGAAAAGCATCCAAACAACAA
 ACAAATATAGATGTAATATACGAACGAGTTCGTGATGATTACCAATTGCATTGGACTTCAACAA
 ATTGGAAAGGTACCAATACTAAAGATAAATGGACAGATCGTTCTTCAGAAAGATATAAAATCGA
 TTGGGAAAAAGAAGAAATGACAAATTAA

ADSDINI**A**TGTTDIGSNTTVKTGDLVITYDKENGMHKKVFYSFIDDKNHNKLLVIRTKGTIAGQ
 YRVYSEEGANKSGLAWPSAFKVQLQLPDNEVAQISDYYPNSIDTKEY**Y****M****S****T****L****T****Y****G****F****N****G****N****L****T**GDD
 TSKIGGLIGAQVSLGHTLKYVQPDFKTI LESPTDKKVGWKVI FNNMVNQNWGPYDRDSWNPVYG
 NQLFMKTRNGSMKAADNFLDPNKASSLLSSGFS PDFATVITMDRKASKQQTNI DVIYERVRDDY
 QLHWTSTNWKGTNTKDKWTD RSSERYKIDWEKEEMTN

Table 8 – SEQ ID NOs: 5 and 6 - α -hemolysin M113R-WT

ATGGCAGATTCTGATATTAATATTTAAAACCGGTACTACAGATATTGGAAGCAATACTACAGTAA
 AACAGGTGATTTAGTCACTTATGATAAAGAAAATGGCATGCACAAAAAAGTATTTTATAGTTT
 TATCGATGATAAAAATCACAATAAAAACTGCTAGTTATTAGAACAAAAGGTACCATTGCTGGT
 CAATATAGAGTTTATAGCGAAGAAGGTGCTAACAAAAGTGGTTTAGCCTGGCCTTCAGCCTTTA
 AGGTACAGTTGCAACTACCTGATAATGAAGTAGCTCAAATATCTGATTACTATCCAAGAAATTC
 GATTGATACAAAAGAGTATAGGAGTACTTTAACTTATGGATTCAACGGTAATGTTACTGGTGAT
 GATACAGGAAAAATGGCGGCCTTATTGGTGCAAATGTTTCGATTGGTCATACACTGAAATATG
 TTCAACCTGATTTCAAACAATTTTAGAGAGCCCAACTGATAAAAAAGTAGGCTGGAAAGTGAT
 ATTTAACAATATGGTGAATCAAATTTGGGACCATACGATCGAGATTCCTTGAACCCGGTATAT
 GGCAATCAACTTTTCATGAAAAGTAGAAATGGTCTATGAAAGCAGCAGATAACTTCCTTGATC
 CTAACAAAGCAAGTTCTCTATTATCTTCAGGGTTTTACCAGACTTCGCTACAGTTATTACTAT
 GGATAGAAAAGCATCCAAACAACAACAATATAGATGTAATATACGAACGAGTTCGTGATGAT
 TACCAATTGCATTGGACTTCAACAATTTGGAAAGGTACCAATACTAAAGATAAATGGACAGATC
 GTTCTTCAGAAAGATATAAAATCGATTGGGAAAAAGAAGAAATGACAAATTAA

ADSDINI**K**TGTTDIGSNTTVKTGDLVITYDKENGMHKKVFYSFIDDKNHNKLLVIRTKGTIAGQ
 YRVYSEEGANKSGLAWPSAFKVQLQLPDNEVAQISDYYPNSIDTKEY**K****S****T****L****T****Y****G****F****N****G****N****V****T**GDD
 TGKIGGLIGANVSIGHTLKYVQPDFKTI LESPTDKKVGWKVI FNNMVNQNWGPYDRDSWNPVYG
 NQLFMKTRNGSMKAADNFLDPNKASSLLSSGFS PDFATVITMDRKASKQQTNI DVIYERVRDDY
 QLHWTSTNWKGTNTKDKWTD RSSERYKIDWEKEEMTN

Table 9 – SEQ ID NOs: 7 and 8 - α -hemolysin E111N-WT

ATGGCAGATTCTGATATTAATATTTAAAACCGGTACTACAGATATTGGAAGCAATACTACAGTAA
 AACAGGTGATTTAGTCACTTATGATAAAGAAAATGGCATGCACAAAAAAGTATTTTATAGTTT
 TATCGATGATAAAAATCACAATAAAAACTGCTAGTTATTAGAACAAAAGGTACCATTGCTGGT
 CAATATAGAGTTTATAGCGAAGAAGGTGCTAACAAAAGTGGTTTAGCCTGGCCTTCAGCCTTTA
 AGGTACAGTTGCAACTACCTGATAATGAAGTAGCTCAAATATCTGATTACTATCCAAGAAATTC
 GATTGATACAAAAACTATATGAGTACTTTAACTTATGGATTCAACGGTAATGTTACTGGTGAT
 GATACAGGAAAAATGGCGGCCTTATTGGTGCAAATGTTTCGATTGGTCATACACTGAAATATG
 TTCAACCTGATTTCAAACAATTTTAGAGAGCCCAACTGATAAAAAAGTAGGCTGGAAAGTGAT
 ATTTAACAATATGGTGAATCAAATTTGGGACCATACGATCGAGATTCCTTGAACCCGGTATAT
 GGCAATCAACTTTTCATGAAAAGTAGAAATGGTCTATGAAAGCAGCAGATAACTTCCTTGATC
 CTAACAAAGCAAGTTCTCTATTATCTTCAGGGTTTTACCAGACTTCGCTACAGTTATTACTAT
 GGATAGAAAAGCATCCAAACAACAACAATATAGATGTAATATACGAACGAGTTCGTGATGAT
 TACCAATTGCATTGGACTTCAACAATTTGGAAAGGTACCAATACTAAAGATAAATGGACAGATC
 GTTCTTCAGAAAGATATAAAATCGATTGGGAAAAAGAAGAAATGACAAATTAA

ADSDINI**K**TGTTDIGSNTTVKTGDLVITYDKENGMHKKVFYSFIDDKNHNKLLVIRTKGTIAGQ
 YRVYSEEGANKSGLAWPSAFKVQLQLPDNEVAQISDYYPNSIDTK**K****Y****M****S****T****L****T****Y****G****F****N****G****N****V****T**GDD

TGKIGGLIGANVSIGHTLKYVQPDFKTI LESPTDKKVGWKVIFNNMVNQNWGPYDRDSWNPVYG
 NQLFMKTRNGSMKAADNFLDPNKASSLLSSGFSPDFATVITMDRKASKQQTNI DVIYERVRDDY
 QLHWTSTNWKGTNTKDKWTD RSSERYKIDWEKEEMTN

Table 10 – SEQ ID NOs: 9 and 10 - α -hemolysin A8R-RL2

ATGGCAGATTCTGATATTAATATTTAAAACCGGTACTACAGATATTGGAAGCAATACTACAGTAA
 AACAGGTGATTTAGTCACTTATGATAAAGAAAATGGCATGCACAAAAAAGTATTTTATAGTTT
 TATCGATGATAAAAATCACAATAAAAAACTGCTAGTTATTAGAACAAAAGGTACCATTGCTGGT
 CAATATAGAGTTTATAGCGAAGAAGGTGCTAACAAAAGTGGTTTAGCCTGGCCTTCAGCCTTTA
 AGGTACAGTTGCAACTACCTGATAATGAAGTAGCTCAAATATCTGATTACTATCCGCGGAATTC
 GATTGATACAAAAGAGTATATGAGTACGTTAACGTACGGATTCAACGGTAACCTTACTGGTGAT
 GATACTAGTAAAAATTGGAGGCCTTATTGGGGCCCAGGTTCCCTAGGTCATACACTTAAGTATG
 TTCAACCTGATTTCAAACAATTCTCGAGAGCCCAACTGATAAAAAAGTAGGCTGGAAAGTGAT
 ATTTAACAATATGGTGAATCAAATTTGGGACCATAACGATCGAGATTCCTTGAACCCGGTATAT
 GGCAATCAACTTTTCATGAAAAC TAGAAATGGTTCTATGAAAGCAGCAGATAACTTCCTTGATC
 CTAACAAAGCAAGTTCTCTATTATCTTCAGGGTTTTACCAGACTTCGCTACAGTTATTACTAT
 GGATAGAAAAGCATCCAAACAACAACAATATAGATGTAATATACGAACGAGTTCGTGATGAT
 TACCAATTGCATTGGACTTCAACAAATTTGGAAAGGTACCAATACTAAAGATAAATGGACAGATC
 GTTCTTCAGAAAGATATAAAATCGATTGGGAAAAAGAAGAAATGACAAATTAA

ADSDINI **I**TGTTDIGSNTTVKTGDLV TYDKENGMHKKVFYSFIDDKNHNKLLVIRTKGTIAGQ
 YRVYSEEGANKSGLAWPSAFKVQLQLPDNEVAQISDYYPNSIDTKEY**MSTLTYGFNGNL**TGDD
 TSKIGGLIGAQVSLGHTLKYVQPDFKTI LESPTDKKVGWKVIFNNMVNQNWGPYDRDSWNPVYG
 NQLFMKTRNGSMKAADNFLDPNKASSLLSSGFSPDFATVITMDRKASKQQTNI DVIYERVRDDY
 QLHWTSTNWKGTNTKDKWTD RSSERYKIDWEKEEMTN

Table 11 – SEQ ID NOs: 11 and 12 - α -hemolysin M113K-RL2

ATGGCAGATTCTGATATTAATATTGCAACCGGTACTACAGATATTGGAAGCAATACTACAGTAA
 AACAGGTGATTTAGTCACTTATGATAAAGAAAATGGCATGCACAAAAAAGTATTTTATAGTTT
 TATCGATGATAAAAATCACAATAAAAAACTGCTAGTTATTAGAACAAAAGGTACCATTGCTGGT
 CAATATAGAGTTTATAGCGAAGAAGGTGCTAACAAAAGTGGTTTAGCCTGGCCTTCAGCCTTTA
 AGGTACAGTTGCAACTACCTGATAATGAAGTAGCTCAAATATCTGATTACTATCCGCGGAATTC
 GATTGATACAAAAGAGTATAAGAGTACGTTAACGTACGGATTCAACGGTAACCTTACTGGTGAT
 GATACTAGTAAAAATTGGAGGCCTTATTGGGGCCCAGGTTCCCTAGGTCATACACTTAAGTATG
 TTCAACCTGATTTCAAACAATTCTCGAGAGCCCAACTGATAAAAAAGTAGGCTGGAAAGTGAT
 ATTTAACAATATGGTGAATCAAATTTGGGACCATAACGATCGAGATTCCTTGAACCCGGTATAT
 GGCAATCAACTTTTCATGAAAAC TAGAAATGGTTCTATGAAAGCAGCAGATAACTTCCTTGATC
 CTAACAAAGCAAGTTCTCTATTATCTTCAGGGTTTTACCAGACTTCGCTACAGTTATTACTAT
 GGATAGAAAAGCATCCAAACAACAACAATATAGATGTAATATACGAACGAGTTCGTGATGAT
 TACCAATTGCATTGGACTTCAACAAATTTGGAAAGGTACCAATACTAAAGATAAATGGACAGATC
 GTTCTTCAGAAAGATATAAAATCGATTGGGAAAAAGAAGAAATGACAAATTAA

ADSDINI **A**TGTTDIGSNTTVKTGDLV TYDKENGMHKKVFYSFIDDKNHNKLLVIRTKGTIAGQ
 YRVYSEEGANKSGLAWPSAFKVQLQLPDNEVAQISDYYPNSIDTKEY **I****S****T****L****T****Y****G****F****N****G****N****L****T**GDD
 TSKIGGLIGAQVSLGHTLKYVQPDFKTI LESPTDKKVGWKVIFNNMVNQNWGPYDRDSWNPVYG
 NQLFMKTRNGSMKAADNFLDPNKASSLLSSGFSPDFATVITMDRKASKQQTNI DVIYERVRDDY
 QLHWTSTNWKGTNTKDKWTD RSSERYKIDWEKEEMTN

Table 12 – SEQ ID NOs: 13 and 14 - α -hemolysin M113R-RL2

ATGGCAGATTCTGATATTAATATTGCAACCGGTACTACAGATATTGGAAGCAATACTACAGTAA
 AACAGGTGATTTAGTCACTTATGATAAAGAAAATGGCATGCACAAAAAAGTATTTTATAGTTT

TATCGATGATAAAAATCACAAATAAAAAACTGCTAGTTATTAGAACAAAAGGTACCATTGCTGGT
 CAATATAGAGTTTATAGCGAAGAAGGTGCTAACAAAAGTGGTTTAGCCTGGCCTTCAGCCTTTA
 AGGTACAGTTGCAACTACCTGATAATGAAGTAGCTCAAATATCTGATTACTATCCGCGGAATTC
 GATTGATACAAAAGAGTATAGAAGTACGTTAACGTACGGATTCAACGGTAACCTTACTGGTGAT
 GATACTAGTAAAATTTGGAGGCCTTATTGGGGCCCAGGTTTCCCTAGGTCATACACTTAAGTATG
 TTCAACCTGATTTCAAACAATTCTCGAGAGCCCAACTGATAAAAAAGTAGGCTGGAAAGTGAT
 ATTTAACAATATGGTGAATCAAATTTGGGGACCATACGATCGAGATTCTTGAACCCGGTATAT
 GGCAATCAACTTTTCATGAAAAGTAGAAATGGTTCTATGAAAGCAGCAGATAACTTCCCTGATC
 CTAACAAAGCAAGTTCTCTATTATCTTCAGGGTTTTACCAGACTTCGCTACAGTTATTACTAT
 GGATAGAAAAGCATCCAAACAACAACAATATAGATGTAATATACGAACGAGTTCGTGATGAT
 TACCAATTGCATTGGACTTCAACAATTTGGAAAGGTACCAATACTAAAGATAAATGGACAGATC
 GTTCTTCAGAAAGATATAAAATCGATTGGGAAAAGAAGAAATGACAAATTAA

ADSDINIATGTTDIGSNTTVKTGDLVITYDKENGMHKKVFYSFIDDKNHNKLLVIRTKGTIAGQ
 YRVYSEEGANKSGLAWPSAFKVQLQLPDNEVAQISDYYPNSIDTKEYSTLTYGFNGNLTGDD
 TSKIGGLIGAQVSLGHTLKYVQPDFKTILESPTDKKVGWKVIFNNMVNQNWGPYDRDSWNPVYG
 NQLFMKTRNGSMKAADNFLDPNKASSLLSSGFSPDFATVITMDRKASKQQTNI DVIYERVRDDY
 QLHWTSTNWKGTNTKDKWTD RSSERYKIDWEKEEMTN

Table 13 – SEQ ID NO: 15 – DNA sequence used in the Example
AAAAAAAAAAAAAAAAAAAAAAAAATCCCCCCCCCCCCCCCCCTAAAAAAAAAATCCCCC CCCCTTAAAAAAAAAATCCCCCCCCC

CLAIMS

1. A method of enhancing translocation of a charged analyte through a transmembrane protein pore, comprising:
 - (a) increasing the net opposing charge of the barrel or channel and/or entrance of the pore; and
 - (b) determining whether or not translocation of the analyte through the resulting pore is enhanced.
2. A method according to claim 1, wherein increasing the net opposing charge increases the frequency of translocation of the analyte through the pore.
3. A method according to claim 1 or 2, wherein increasing the net opposing charge decreases the threshold voltage for translocation of the analyte through the pore.
4. A method according to any one of the preceding claims, wherein increasing the net opposing charge decreases the translocation speed of the analyte through the pore.
5. A method according to any one of the preceding claims, wherein increasing the net opposing charge decreases the number of non-translocation interactions between the analyte and the pore
6. A method according to any one of the preceding claims, wherein the analyte is a polymer.
7. A method according to any one of the preceding claims, wherein the analyte is a nucleic acid sequence.
8. A method according to any one of the preceding claims, wherein the pore comprises seven subunits each comprising SEQ ID NO: 2 or a variant thereof.
9. A method according to any one of claims 1 to 7, wherein the pore comprises seven subunits each comprising SEQ ID NO: 4 or a variant thereof.

10. A method according to any one of the preceding claims, wherein the analyte is negatively charged and step (a) involve increasing the net positive charge.
11. A method according to claim 10, wherein the net positive charge is increased by introducing one or more positively charged amino acids into the barrel or channel and/or entrance of the pore.
12. A method according to claim 11, wherein the introducing is carried out by substitution.
13. A method according to claim 12, wherein the one or more positively-charge amino acids are histidine (H), lysine (K) and/or arginine (R).
14. A method according to claim 10, wherein the net positive charge is increased substituting one or more negatively charged amino acids in the barrel or channel and/or entrance of the pore with one or more uncharged amino acids, non-polar amino acids and/or aromatic amino acids.
15. A method according to any one of the preceding claims, wherein the entrance is the portion of the pore through which the analyte enters the barrel or channel.
16. Use of an increase in net opposing charge of the barrel or channel and/or entrance of a transmembrane protein pore to enhance translocation of a charged analyte through the pore.
17. A transmembrane protein pore enhanced by a method according to any one of claims 1 to 15.
18. A transmembrane protein pore in which the net opposing charge of its barrel or channel and/or entrance has been increased to enhance the translocation of a charged analyte through the pore.
19. A transmembrane protein pore according to claim 17 or 18, wherein the pore comprises:
 - (a) at least one subunit comprising SEQ ID NO: 6 or a variant thereof;
 - (b) at least one subunit comprising SEQ ID NO: 8 or a variant thereof; or
 - (c) at least one subunit comprising SEQ ID NO: 10 or a variant thereof.

20. A transmembrane protein pore according to claim 19, wherein the pore is a homoheptamer comprising seven subunits of SEQ ID NO: 6, 8 or 10 or a variant thereof.
21. A polynucleotide encoding a transmembrane protein pore according to any one of claims 17 to 20.
22. A polynucleotide encoding a transmembrane protein pore subunit having the sequence of SEQ ID NO: 6, 8 or 10 or a variant thereof.
23. A polynucleotide according to claim 22, wherein the polynucleotide comprises the sequence shown in SEQ ID NO: 5, 7 or 9 or a variant thereof
24. A method of determining presence or absence of an analyte in a sample, comprising:
(a) contacting the sample with a transmembrane protein pore according to any one of claims 17 to 20 under conditions that allow the analyte, if present, to translocate through and interact with the pore; and
(b) measuring the current passing through the pore during the interaction and thereby determining the presence or absence of the analyte.
25. A method of sequencing a target nucleic acid sequence, comprising:
(a) pushing or pulling the target sequence through a transmembrane protein pore according to any one of claims 17 to 20 so that a proportion of the nucleotides in the target sequence interact with the pore; and
(b) measuring the current passing through the pore during each interaction and thereby determining the sequence of the target sequence.
26. A kit for sequencing a nucleic acid, comprising a transmembrane protein pore according to any one of claims 17 to 20 and a nucleic acid handling enzyme.




Figure 2




Figure 3




Figure 4





Figure 5






Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14





Figure 15



INTERNATIONAL SEARCH REPORT

International application No
PCT/GB2009/002666

A. CLASSIFICATION OF SUBJECT MATTER
 INV. C07K14/31 C12N15/31 C12Q1/68 G01N33/68 G01N27/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 C07K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
 EPO-Internal, WPI Data, BIOSIS, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GU LI-QUN ET AL: "Prolonged residence time of a noncovalent molecular adapter, beta-Cyclodextrin, within the lumen of mutant alpha-hemolysin pores" JOURNAL OF GENERAL PHYSIOLOGY, vol. 118, no. 5, November 2001 (2001-11), pages 481-493, XP002568389 ISSN: 0022-1295 the whole document <div style="text-align: center; margin-top: 10px;">----- -/--</div>	17-23

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family
---	---

Date of the actual completion of the international search 15 February 2010	Date of mailing of the international search report 26/02/2010
--	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Kools, Patrick
--	---

INTERNATIONAL SEARCH REPORT

International application No.

PCT/GB2009/002666

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.b of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application and necessary to the claimed invention, the international search was carried out on the basis of:
 - a. type of material
 - a sequence listing
 - table(s) related to the sequence listing
 - b. format of material
 - on paper
 - in electronic form
 - c. time of filing/furnishing
 - contained in the international application as filed
 - filed together with the international application in electronic form
 - furnished subsequently to this Authority for the purpose of search
2. In addition, in the case that more than one version or copy of a sequence listing and/or table relating thereto has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that in the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No

PCT/GB2009/002666

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>HENRICKSON S E ET AL: "Driven DNA transport into an asymmetric nanometer-scale pore." PHYSICAL REVIEW LETTERS 2 OCT 2000, vol. 85, no. 14, 2 October 2000 (2000-10-02), pages 3057-3060, XP002568385 ISSN: 0031-9007 page 3059, column 1, paragraph 2 - column 2, paragraph 1</p>	1-26
X	<p>MERZLYAK PETR G ET AL: "Conductance and ion selectivity of a mesoscopic protein nanopore probed with cysteine scanning mutagenesis" BIOPHYSICAL JOURNAL, vol. 89, no. 5, November 2005 (2005-11), pages 3059-3070, XP002568386 ISSN: 0006-3495 the whole document</p>	1,8-24
A	<p>ASTIER YANN ET AL: "Toward single molecule DNA sequencing: direct identification of ribonucleoside and deoxyribonucleoside 5'-monophosphates by using an engineered protein nanopore equipped with a molecular adapter" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, AMERICAN CHEMICAL SOCIETY, NEW YORK, USA, vol. 128, no. 5, 8 February 2006 (2006-02-08), pages 1705-1710, XP002503447 ISSN: 0002-7863 the whole document</p>	1-26
A	<p>WO 2007/057668 A1 (ISIS INNOVATION [GB]; BAYLEY HAGAN [GB]; ASTIER YANN [GB]; BRAHA ORIT) 24 May 2007 (2007-05-24) the whole document</p>	1-26
A	<p>WO 2007/084103 A2 (TEXAS A & M UNIV SYS [US]; KANG XIAOFENG [US]; GU LI QUN ANDREW [US];) 26 July 2007 (2007-07-26) the whole document</p>	1-26
A	<p>GU LI-QUN ET AL: "Electroosmotic enhancement of the binding of neutral molecule to a transmembrane pore." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 100, no. 26, 23 December 2003 (2003-12-23), pages 15498-15503, XP002568387 ISSN: 0027-8424 the whole document</p>	1-26

-/--

INTERNATIONAL SEARCH REPORT

International application No

PCT/GB2009/002666

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	COMAI MASSIMILIANO ET AL: "Protein engineering modulates the transport properties and ion selectivity of the pores formed by staphylococcal gamma-haemolysins in lipid membranes" MOLECULAR MICROBIOLOGY, vol. 44, no. 5, June 2002 (2002-06), pages 1251-1267, XP002568388 ISSN: 0950-382X the whole document -----	1-26
X,P	MAGLIA GIOVANNI ET AL: "Enhanced translocation of single DNA molecules through alpha-hemolysin nanopores by manipulation of internal charge" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 105, no. 50, December 2008 (2008-12), pages 19720-19725, XP002568390 ISSN: 0027-8424 the whole document -----	1-26

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/GB2009/002666

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007057668 A1	24-05-2007	AU 2006314296 A1	24-05-2007
		CA 2629009 A1	24-05-2007
		CN 101356288 A	28-01-2009
		EP 1951898 A1	06-08-2008
		JP 2009519705 T	21-05-2009
		KR 20080083281 A	17-09-2008
		US 2008311582 A1	18-12-2008
WO 2007084103 A2	26-07-2007	EP 1842061 A2	10-10-2007
		US 2008101988 A1	01-05-2008