

University of Groningen

Corrigendum

Campos, Ana Carolina C; Andrade, Nathália L; Ferdous, Mithila; Chlebowicz, Monika A; Santos, Carla C; Correal, Julio C D; Lo Ten Foe, Jerome R; Rosa, Ana Cláudia P; Damasco, Paulo V; Friedrich, Alex W

Published in:
Frontiers in Microbiology

DOI:
[10.3389/fmicb.2020.599031](https://doi.org/10.3389/fmicb.2020.599031)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Campos, A. C. C., Andrade, N. L., Ferdous, M., Chlebowicz, M. A., Santos, C. C., Correal, J. C. D., Lo Ten Foe, J. R., Rosa, A. C. P., Damasco, P. V., Friedrich, A. W., & Rossen, J. W. A. (2020). Corrigendum: Comprehensive Molecular Characterization of Escherichia coli Isolates from Urine Samples of Hospitalized Patients in Rio de Janeiro, Brazil (Frontiers in Microbiology, (2018), 9, (243), 10.3389/fmicb.2018.00243). *Frontiers in Microbiology*, 11, 1-4. [599031]. <https://doi.org/10.3389/fmicb.2020.599031>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Corrigendum: Comprehensive Molecular Characterization of *Escherichia coli* Isolates from Urine Samples of Hospitalized Patients in Rio de Janeiro, Brazil

Ana Carolina C. Campos^{1,2}, Nathália L. Andrade¹, Mithila Ferdous², Monika A. Chlebowicz², Carla C. Santos³, Julio C. D. Correal^{1,3}, Jerome R. Lo Ten Foe², Ana Cláudia P. Rosa¹, Paulo V. Damasco^{4,5}, Alex W. Friedrich² and John W. A. Rossen^{2*}

¹ Departamento de Microbiologia, Imunologia e Parasitologia, Faculdade de Ciências Médicas, Universidade Do Estado Do Rio de Janeiro, Rio de Janeiro, Brazil, ² Department of Medical Microbiology, University of Groningen, University Medical Center Groningen, Groningen, Netherlands, ³ Departamento de Controle de Infecções, Hospital Rio Laranjeiras, Rio de Janeiro, Brazil, ⁴ Departamento de Doenças Infecciosas e Parasitárias, Universidade Federal Do Estado Do Rio de Janeiro, Rio de Janeiro, Brazil, ⁵ Departamento de Doenças Infecciosas e Parasitárias, Universidade Do Estado Do Rio de Janeiro, Rio de Janeiro, Brazil

OPEN ACCESS

Edited and reviewed by:

Jorge Blanco,
University of Santiago de Compostela,
Spain

*Correspondence:

John W. A. Rossen
j.w.a.rossen@rug.nl

Specialty section:

This article was submitted to
Infectious Diseases,
a section of the journal
Frontiers in Microbiology

Received: 26 August 2020

Accepted: 15 September 2020

Published: 30 October 2020

Citation:

Campos ACC, Andrade NL, Ferdous M, Chlebowicz MA, Santos CC, Correal JCD, Lo Ten Foe JR, Rosa ACP, Damasco PV, Friedrich AW and Rossen JWA (2020) Corrigendum: Comprehensive Molecular Characterization of *Escherichia coli* Isolates from Urine Samples of Hospitalized Patients in Rio de Janeiro, Brazil. *Front. Microbiol.* 11:599031. doi: 10.3389/fmicb.2020.599031

Keywords: *Escherichia coli*, urinary tract infections, Brazil, ST131, antibiotic resistance, virulence genes, whole genome sequencing, diagnostic stewardship

A Corrigendum on

Comprehensive Molecular Characterization of *Escherichia coli* Isolates from Urine Samples of Hospitalized Patients in Rio de Janeiro, Brazil

by Campos, A. C. C., Andrade, N. L., Ferdous, M., Chlebowicz, M. A., Santos, C. C., Correal, J. C. D., et al. (2018). *Front. Microbiol.* 9:243. doi: 10.3389/fmicb.2018.00243

In the original article, there was a mistake in **Table 2** and **Supplementary Data Sheet 1, S.6** as published. The virotype of the isolate 4233 is not typeable. The corrected **Table 2** and **Supplementary Data Sheet 1, S.6** appear below.

In the original article, there was an error in the presented percentage of the virotype D among ST131 isolates.

A correction has been made to *Results, Escherichia coli* ST131, Paragraph 1

UPEC strains produce different adhesins and fimbriae, including type 1 fimbriae. The FimH protein is the adhesive subunit of type 1 fimbriae that is used for epidemiological typing of UPEC. In this study, three *fimH* types were identified among the ST131 isolates, two O25:H4/ST131 isolates belonged to *fimH22*, two O16:H5/ST131 isolates to *fimH41* while the majority of O25:H4/ST131 isolates ($n = 22$) belonged to *fimH30* (**Table 2**). The virulence genes (*afa/draBC*, *iroN*, *sat*, *ibeA*, *papGII*, *papGIII*, *cnf-1*, *hlyA*, *cdtB*, *neuC-K1*, *kpsMIIK2*, *kpsmII-K5*) were used to determine the virotype of ST131 isolates based on the virulence profile. O25:H4/ST131 isolates belonged to different virotypes, i.e., 7 (26.92%) to virotype A, 1 (3.84%) to virotype B, 14 (53.84%) to virotype C, and 3 (11.53%) to virotype D. Isolates belonging to virotype C could be divided into subtypes C2 ($n = 6$) or C3 ($n = 3$), whereas

five isolates could not be further subtyped. The only two isolates with serotype O16:H5/ST131 were classified as virotype A (see Data Sheet S6). Almost all O25:H4/ST131 isolates were resistant to fluoroquinolones, whereas the O16:H5/ST131 isolates were susceptible to this antibiotic. The *bla*_{CTX-M} gene was most prevalent in O25:H4/ST131 *fimH*30 fluoroquinolones resistant (O25:H4/ST131-H30-R) isolates belonging to virotype C. Within ST131, *bla*_{CTX-M15} was confined to the H30-R sub-clone known as O25:H4/ST131-H30-Rx, represented by 9 (34.61%) isolates (**Table 2**).

The authors apologize for these errors and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Campos, Andrade, Ferdous, Chlebowicz, Santos, Correal, Lo Ten Foe, Rosa, Damasco, Friedrich and Rossen. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

TABLE 2 | Distribution of *fimH* types among ST131 *Escherichia coli* isolates.

Isolates	Phylogenetic group	<i>FimH</i> type	Serotype	Virotype	ESBL genes	Fluoroquinolone resistant ^a
5332	B2	<i>fimH22</i>	O25:H4	D	<i>bla_{CMY-2}</i>	Pos
7018	B2	<i>fimH30</i>	O25:H4	A	<i>bla_{OXA-1}</i>	Pos
7104	B2	<i>fimH30</i>	O25:H4	C2	<i>bla_{KPC-2}</i>	Pos
9260	B2	<i>fimH30</i>	O25:H4	C	<i>bla_{CTX-M-15}</i>	Pos
3218	B2	<i>fimH30</i>	O25:H4	C2	<i>bla_{KPC-2}</i>	Pos
9581A	B2	<i>fimH30</i>	O25:H4	C	<i>bla_{CTX-M-15}</i>	Pos
x5770d	B2	<i>fimH30</i>	O25:H4	C	<i>bla_{CTX-M-15}</i>	Pos
x6638	B2	<i>fimH30</i>	O25:H4	A	<i>bla_{CTX-M-15}</i>	Pos
1294D	B2	<i>fimH30</i>	O25:H4	B	<i>bla_{KPC-2}</i>	Pos
2102	B2	<i>fimH30</i>	O25:H4	A	<i>bla_{KPC-2}</i>	Pos
1710D	B2	<i>fimH30</i>	O25:H4	C	<i>bla_{CTX-M-15}</i>	Pos
9533D	B2	<i>fimH30</i>	O25:H4	C	<i>bla_{CTX-M-15}</i>	Pos
3528	B2	<i>fimH30</i>	O25:H4	C2	<i>bla_{CTX-M-15}</i>	Neg
7078	B2	<i>fimH30</i>	O25:H4	C3	<i>bla_{TEM-1B}</i>	Pos
9893	B2	<i>fimH30</i>	O25:H4	C2	<i>bla_{KPC-2}</i>	Neg
7974	B2	<i>fimH30</i>	O25:H4	D	<i>bla_{CTX-M-2}</i>	Pos
4233	B2	<i>fimH30</i>	O25:H4	NT	<i>bla_{KPC-2}</i>	Pos
5420	B2	<i>fimH30</i>	O25:H4	A	<i>bla_{CTX-M-15}</i>	Neg
2478	B2	<i>fimH41</i>	O16:H5	A	<i>bla_{TEM-1B}</i>	Neg
4006	B2	<i>fimH41</i>	O16:H5	A	<i>bla_{TEM-1B}</i>	Pos
5976	B2	<i>fimH30</i>	O25:H4	C3	<i>bla_{TEM-1B}</i>	Pos
2206	B2	<i>fimH30</i>	O25:H4	A	<i>bla_{CTX-M-15}</i>	Pos
8565	B2	<i>fimH30</i>	O25:H4	C3	<i>bla_{TEM-1B}</i>	Pos
x2724	B2	<i>fimH30</i>	O25:H4	C2	<i>bla_{TEM-1B}</i>	Pos
6202	B2	<i>fimH30</i>	O25:H4	C2	<i>bla_{TEM-1B}</i>	Pos
5848	B2	<i>fimH22</i>	O25:H4	D	<i>bla_{CMY-2}</i>	Neg

^aNeg. indicates susceptible to fluoroquinolones and Pos. indicates resistant to fluoroquinolones. NT, not typeable.

Supplementary data S.6. | Virotypes distribution in ST131 isolates.

Isolates	FimH type	Serotype	<i>afa/draBC</i>	<i>afaoperon FM955459</i>	<i>iroN</i>	<i>sat</i>	<i>ibeA</i>	<i>papG II</i>	<i>papG III</i>	<i>cnf1</i>	<i>hlyA</i>	<i>cdtB</i>	<i>neuC-K1</i>	<i>kpsM II-K2</i>	<i>kpsM II-K5</i>	Virotypes
5332	<i>fimH22</i>	O25:H4	neg	neg	pos	neg	pos	neg	neg	neg	neg	neg	pos	pos	neg	D
7018	<i>fimH30</i>	O25:H4	pos	pos	neg	pos	neg	neg	neg	neg	neg	neg	neg	pos	neg	A
7104	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	pos	C2
9260	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	pos	neg	pos	neg	neg	neg	pos	neg	C
3218	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	pos	C2
9581A	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	pos	neg	pos	neg	neg	neg	pos	neg	C
x5770d	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	pos	neg	pos	neg	neg	neg	pos	neg	C
x6638	<i>fimH30</i>	O25:H4	pos	pos	neg	pos	neg	neg	neg	neg	neg	neg	neg	pos	neg	A
1294D	<i>fimH30</i>	O25:H4	neg	neg	pos	pos	neg	neg	neg	neg	neg	neg	neg	neg	pos	B
2102	<i>fimH30</i>	O25:H4	pos	pos	neg	pos	neg	neg	neg	neg	neg	neg	neg	pos	neg	A
1710D	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	pos	neg	pos	neg	neg	neg	neg	pos	C
9533D	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	pos	neg	pos	neg	neg	neg	pos	neg	C
3528	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	pos	C2
7078	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	neg	C3
9893	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	pos	C2
7974	<i>fimH30</i>	O25:H4	neg	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	pos	neg	D
4233	<i>fimH30</i>	O25:H4	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	NT
5420	<i>fimH30</i>	O25:H4	pos	pos	neg	pos	neg	neg	neg	neg	neg	neg	neg	pos	neg	A
2478	<i>fimH41</i>	O16:H5	pos	pos	neg	neg	neg	neg	neg	neg	neg	neg	neg	pos	neg	A
4006	<i>fimH41</i>	O16:H5	pos	pos	neg	neg	neg	neg	neg	neg	neg	neg	neg	pos	neg	A
5976	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	neg	C3
2206	<i>fimH30</i>	O25:H4	pos	pos	neg	pos	neg	neg	neg	neg	neg	neg	neg	pos	neg	A
8565	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	neg	C3
x2724	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	pos	C2
6202	<i>fimH30</i>	O25:H4	neg	neg	neg	pos	neg	neg	neg	neg	neg	neg	neg	neg	pos	C2
5848	<i>fimH22</i>	O25:H4	neg	neg	pos	neg	pos	neg	neg	neg	neg	neg	pos	pos	neg	D

pos (positive) and neg (negative); Some isolates were not possible to subtyping according with Dahbi et al., 2014 were just typing according Blanco et al., 2014, and other were not possible to type being classified as NT, not typeable.