

## University of Groningen

### The disembodied self

Mertens, Yoki Linn

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Publication date:*  
2019

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Mertens, Y. L. (2019). *The disembodied self: Using an experimental approach to study the neural correlates of dissociation*. Poster session presented at 35th Annual Meeting of the International Society for Traumatic Stress Studies (ISTSS), Boston, United States.

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# The Disembodied Self – Using an experimental approach to investigate objective markers of dissociation


Yoki L. Mertens<sup>1,2</sup>, Judith K. Daniels<sup>1</sup>, André Aleman<sup>1,2</sup>

<sup>1</sup>University of Groningen

<sup>2</sup>Cognitive Neuroimaging Centre, University Medical Centre Groningen

## Background

- Individuals with childhood trauma report elevated levels of **dissociation**, e.g. depersonalization symptoms<sup>1</sup>.
- Lack of objective markers of (somatoform) dissociation, e.g. **out-of-body experiences (OBEs)**; Picture 2).
- Reduced sense of self and diminished body awareness could relate to brain deficits in integrative capacity.
- The **temporoparietal junction (TPJ)** (Picture 1), a critical hub for multisensory integration, is a proposed key region for establishing a coherent sense of (bodily) self<sup>2,3</sup>.
- Lesion and **perturbation studies** indicate that stimulation of the TPJ affects own-body perceptions<sup>4,5</sup> and decreases depersonalization symptoms<sup>6</sup>.
- **Connectivity** between right TPJ and right (posterior) insula depicted the strongest activity associated with changes in self-location and first-person perspective<sup>7</sup>.




Picture 1. The temporoparietal junction.<sup>4</sup>

## Research Question

How does a temporary perturbation of the right temporoparietal junction (rTPJ) affect

- 1) Dissociative experiences in individuals with (and without) a history of childhood trauma
- 2) Experimentally-induced OBEs<sup>8</sup>?
- 3) The brain network subserving body awareness?



Picture 2. Out-of-Body Experience<sup>3</sup>

## Methods

**Study Design:** Sham-controlled, randomized, pre-post TMS study

**Study Population:** Total  $N = 52$  right-handed, female, young adults\*

**Trauma Group:**  $N = 26$  subjects with former childhood trauma

**Control Group:**  $N = 26$  subjects without former childhood trauma

\*Exclusion Criteria

- 1) Neurological disorder, e.g. epilepsy
- 2) Psychiatric disorder
- 3) Presence of metallic devices
- 4) Psychotropic medication intake for the last 6 months
- 5) Claustrophobia
- 6) Alcohol or drug abuse
- 7) Tinnitus

## Questionnaires

**Childhood Trauma.** Childhood Trauma Questionnaire (CTQ; Bernstein & Fink, 1998)

**State Dissociation.** Dissociation Tension Scale – Short (DSS-4; Stiglmayr et al., 2009)

**Trait Dissociation.** Dissociative Experiences Scale (DES-II; Carlson & Putnam, 2000)


**Depersonalization.** Cambridge Depersonalization Scale (CDS; Sierra & Berrios, 2000)

**Vestibular Symptoms.** Vertigo Symptom Scale-Short Form (VSS-SF; Wilhelmssen et al., 2008)

**Sleep Quality.** Pittsburgh Sleep Quality Index (PSQI; Buysse et al., 1989)




TMS Coil (Figure-8; Cool B65; MagVenture)




**Brain network PRE-TMS**

- Anatomical scan
- Resting-state fMRI
- Functional connectivity analyses (CONN toolbox, SPM12; ROIs: TPJ, insula, pSTG, S1, mPFC,...)




**Out-of-Body Illusion Paradigm<sup>8</sup> PRE-TMS**

- VR goggles connected to camera filming subject
- Synchronous tapping → OBE




**Transcranial Magnetic Stimulation (TMS)**

- cTBS protocol, 40s, 600 pulses
- Inhibitory effect
- Neuronavigated rTPJ stimulation (Brain Voyager TMS neuronavigation)



**Out-of-Body Illusion Paradigm<sup>8</sup> POST-TMS**

- VR goggles connected to camera filming subject
- Synchronous tapping → OBE



**Brain network POST-TMS**

- Resting-state fMRI
- Functional connectivity analyses (CONN toolbox, SPM12; ROIs: TPJ, insula, pSTG, S1, mPFC,...)


### Infobox 1: Heart rate as objective marker of disembodiment

- Measured with wireless C3 Corrium heart rate (HR) monitor
- Threat-evoked: Fake hammer attack on illusory body
- Before: Increased emotional distress measured via skin conductance response after OBE induction<sup>8</sup>
- Hypothesis: OBE induction leads to increased heart rate (=distress) when illusory body attacked.



### Infobox 2: Spatial map as objective marker of self-localization

- Subjects indicate shift in self-localization on map
- Hypothesis: OBE induction leads to shift in self-localization from original position (smiley) towards camera (red triangle)



## Contact

Yoki Linn Mertens, PhD student  
Department: Clinical Psychology & Experimental  
Psychopathology; University of Groningen, Netherlands

E-mail: [y.l.mertens@rug.nl](mailto:y.l.mertens@rug.nl)  
Phone: +31 50363 7373

## References

- <sup>1</sup>Vonderlin, R., Kleindienst, N., Alpers, G. W., Bohus, M., Lyssenko, L., & Schmahl, C. (2018). Dissociation in victims of childhood abuse or neglect: a meta-analytic review. *Psychological medicine*, 48(15), 2467-2476.
- <sup>2</sup>Eddy, C. M. (2016). The junction between self and other? Temporo-parietal dysfunction in neuropsychiatry. *Neuropsychologia*, 89, 465-477.
- <sup>3</sup>Blanke, O. (2012). Multisensory brain mechanisms of bodily self-consciousness. *Nature Reviews Neuroscience*, 13(8), 556.
- <sup>4</sup>Donaldson, P. H., Rinehart, N. J., & Enticott, P. G. (2015). Noninvasive stimulation of the temporoparietal junction: A systematic review. *Neuroscience & Biobehavioral Reviews*, 55, 547-572.
- <sup>5</sup>Grivaz, P., Blanke, O., & Serino, A. (2017). Common and distinct brain regions processing multisensory bodily signals for peripersonal space and body ownership. *NeuroImage*, 147, 602-618.
- <sup>6</sup>Jay, E. L., Sierra, M., Van den Eynde, F., Rothwell, J. C., & David, A. S. (2014). Testing a neurobiological model of depersonalization disorder using repetitive transcranial magnetic stimulation. *Brain stimulation*, 7(2), 252-259.
- <sup>7</sup>Ionta, S., Martuzzi, R., Salomon, R., & Blanke, O. (2014). The brain network reflecting bodily self-consciousness: a functional connectivity study. *Social cognitive and affective neuroscience*, 9(12), 1904-1913.
- <sup>8</sup>Ehrsson, H. H. (2007). The experimental induction of out-of-body experiences. *Science*, 317(5841), 1048-1048.