

University of Groningen

**Addendum to ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI expert consensus recommendations for multimodality imaging in cardiac amyloidosis**

Dorbala, Sharmila; Ando, Yukio; Bokhari, Sabahat; Dispenzieri, Angela; Falk, Rodney H.; Ferrari, Victor A.; Fontana, Marianna; Gheysens, Olivier; Gillmore, Julian D.; Glaudemans, Andor W. J. M.

*Published in:*  
Journal of Nuclear Cardiology

*DOI:*  
[10.1007/s12350-020-02455-z](https://doi.org/10.1007/s12350-020-02455-z)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2021

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Dorbala, S., Ando, Y., Bokhari, S., Dispenzieri, A., Falk, R. H., Ferrari, V. A., Fontana, M., Gheysens, O., Gillmore, J. D., Glaudemans, A. W. J. M., Hanna, M. A., Hazenberg, B. P. C., Kristen, A., Kwong, R. Y., Maurer, M. S., Merlini, G., Miller, E. J., Moon, J. C., Murthy, V. L., ... Bourque, J. M. (2021). Addendum to ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI expert consensus recommendations for multimodality imaging in cardiac amyloidosis: Part 1 of 2-evidence base and standardized methods of imaging. *Journal of Nuclear Cardiology*. <https://doi.org/10.1007/s12350-020-02455-z>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Addendum to ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI expert consensus recommendations for multimodality imaging in cardiac amyloidosis: Part 1 of 2—evidence base and standardized methods of imaging

Sharmila Dorbala, MD, MPH, FASNC,<sup>a</sup> Yukio Ando, MD, PhD,<sup>b</sup> Sabahat Bokhari, MD,<sup>c</sup> Angela Dispenzieri, MD,<sup>d</sup> Rodney H. Falk, MD,<sup>a</sup> Victor A. Ferrari, MD,<sup>e</sup> Marianna Fontana, PhD,<sup>f</sup> Olivier Gheysens, MD, PhD,<sup>g</sup> Julian D. Gillmore, MD, PhD,<sup>f</sup> Andor W. J. M. Glaudemans, MD, PhD,<sup>h</sup> Mazen A. Hanna, MD,<sup>i</sup> Bouke P. C. Hazenberg, MD, PhD,<sup>j</sup> Arnt V. Kristen, MD,<sup>k</sup> Raymond Y. Kwong, MD, MPH,<sup>a</sup> Mathew S. Maurer, MD,<sup>c</sup> Giampaolo Merlini, MD,<sup>l,m</sup> Edward J. Miller, MD, PhD,<sup>n</sup> James C. Moon, MD,<sup>f</sup> Venkatesh L. Murthy, MD, PhD,<sup>o</sup> C. Cristina Quarta, MD, PhD,<sup>f</sup> Claudio Rapezzi, MD,<sup>p</sup> Frederick L. Ruberg, MD,<sup>q</sup> Sanjiv J. Shah, MD,<sup>r</sup> Riemer H. J. A. Slart, MD,<sup>h</sup> Hein J. Verberne, MD, PhD,<sup>s</sup> and Jamieson M. Bourque, MD, MHS, FASNC<sup>t</sup>

<sup>a</sup> Cardiac Amyloidosis Program, Cardiovascular Imaging Programs, Departments of Medicine and Radiology, Harvard Medical School, Brigham and Women's Hospital, Boston, MA

<sup>b</sup> Department of Neurology, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

<sup>c</sup> Columbia University Medical Center, New York Presbyterian Hospital, Columbia University, New York, NY

<sup>d</sup> Division of Hematology, Division of Cardiovascular Diseases, Department of Radiology, Division of Nuclear Medicine, Department of Medicine, Mayo Clinic, Rochester, MN

<sup>e</sup> Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA

<sup>f</sup> Division of Medicine, National Amyloidosis Centre, University College London, London, UK

<sup>g</sup> University Hospitals Leuven, Nuclear Medicine and Molecular Imaging, Leuven, Belgium

<sup>h</sup> Department of Nuclear Medicine and Molecular Imaging, Medical Imaging Center, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

<sup>i</sup> Department of Cardiovascular Medicine, Cleveland Clinic, Cleveland, OH

<sup>j</sup> Department of Rheumatology & Clinical Immunology, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

<sup>k</sup> Department of Cardiology, University of Heidelberg, Heidelberg, Germany

<sup>l</sup> Fondazione Istituto di Ricovero e Cura a Carattere Scientifico Policlinico San Matteo, Amyloidosis Research and Treatment Center, Pavia, Italy

<sup>m</sup> Department of Molecular Medicine, University of Pavia, Pavia, Italy

<sup>n</sup> Cardiovascular Medicine, Yale University School of Medicine, New Haven, CT

<sup>o</sup> Frankel Cardiovascular Center, Michigan Medicine, Ann Arbor, MI

Refer to the related article (<https://doi.org/10.1007/s12350-019-01760-6>).

Reprint requests: Jamieson M. Bourque, MD, MHS, FASNC (Co-Chair), Departments of Medicine and Radiology, Cardiovascular Imaging Center, University of Virginia, Charlottesville, VA; [JMB8T@hscmail.mcc.virginia.edu](mailto:JMB8T@hscmail.mcc.virginia.edu)

This article is being jointly published in *Circulation: Cardiovascular Imaging* (<https://doi.org/10.1161/HCI.0000000000000029>), the

*Journal of Cardiac Failure* (<https://doi.org/10.1016/j.cardfail.2019.08.002>) and the *Journal of Nuclear Cardiology* (<https://doi.org/10.1007/s12350-020-02455-z>).

*J Nucl Cardiol*  
1071-3581/\$34.00

Copyright © 2021 American Society of Nuclear Cardiology, American Heart Association, Inc., and Heart Failure Society of America.

doi:10.1007/s12350-020-02455-z

- <sup>p</sup> Cardiology Unit, Department of Experimental, Diagnostic and Specialty Medicine, Alma Mater-University of Bologna, Bologna, Italy
- <sup>q</sup> Department of Medicine, Amyloidosis Center and Section of Cardiovascular Medicine, Boston Medical Center, Boston University School of Medicine, Boston, MA
- <sup>r</sup> Feinberg School of Medicine, Northwestern University, Chicago, IL
- <sup>s</sup> Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam, The Netherlands
- <sup>t</sup> Departments of Medicine and Radiology, Cardiovascular Imaging Center, University of Virginia, Charlottesville, VA

Received Nov 23, 2020; accepted Nov 24, 2020

doi:10.1007/s12350-020-02455-z

## THE NEED FOR AN ADDENDUM

There are two primary reasons for an addendum. The first is that the document reviewer list is being updated to include Dr. Richard Cheng and Dr. Roy John, who have critically reviewed the document, but were inadvertently not listed as reviewers. In addition, since the publication of this document and introduction of approved therapies for transthyretin cardiac amyloidosis (ATTR-CA), the clinical use of bone tracer cardiac scintigraphy has been extended to populations with lower prevalence of ATTR-CA. Numerous observations have raised concerns about (1) incorrect diagnosis of ATTR-CA based on <sup>99m</sup>Tc-pyrophosphate (PYP) planar imaging and heart-to-contralateral lung (H/CL) ratio without confirmation of diffuse myocardial uptake on SPECT imaging at some sites; (2) excess blood pool activity on the 1-hour planar and SPECT images being interpreted as positive scans; and (3) missed diagnosis of light chain (AL) amyloidosis, as serum-free light chain studies and serum and urine immunofixation electrophoresis studies may not be recommended in the <sup>99m</sup>Tc-PYP/-3,3-diphosphono-1,2-propanodicarboxylic acid/hydroxymethylene diphosphonate (<sup>99m</sup>Tc-PYP/DPD/HMDP) report. Incorrect diagnosis leads to inappropriate therapy and worse patient outcomes. SPECT and planar imaging performed at 3 hour maximize specificity.<sup>1-3</sup> Additionally, technical parameters have been updated.

Accordingly, we are issuing this addendum to clarify the protocols, interpretation, and reporting of <sup>99m</sup>Tc-PYP imaging:

### 1. Acquisition (Table 4):

- a. The time between injection of <sup>99m</sup>Tc-PYP and scan is revised: 2- or 3-hour imaging is recommended, and 1-hour imaging is optional (Table 4).

If excess blood pool activity is noted, 3-hour imaging is recommended. The timing between injection and scanning is now consistent for <sup>99m</sup>Tc-PYP, -DPD, and -HMDP. We recognize some experienced centers that have become proficient at 1-hour scanning; the recommendation for 2- or 3-hour imaging is particularly important for centers starting new <sup>99m</sup>Tc-PYP programs.

- b. SPECT imaging is required in all studies (irrespective of time between injection and scan) to highlight the importance of directly visualizing tracer uptake in the myocardium.
- c. 1-hour planar-only imaging is not recommended.
- d. Emerging literature suggests that cadmium zinc telluride (CZT) SPECT can also be used for <sup>99m</sup>Tc-PYP/DPD/HMDP imaging.<sup>4,5</sup>

### 2. Interpretation (Table 5):

- a. Planar imaging and H/CL ratio alone are insufficient for diagnosis of ATTR cardiac amyloidosis. SPECT imaging is necessary to identify myocardial uptake of <sup>99m</sup>Tc-PYP/DPD/HMDP.
- b. Repeat imaging is recommended at 3 hours if excess blood pool activity is noted.
- c. The steps in Table 5 clarify that visual grading on planar and SPECT imaging is the primary method for diagnosis of ATTR cardiac amyloidosis.
- d. Recommendations are clarified for ease of interpretation.

### 3. Reporting (Table 6):

- a. Diffuse myocardial uptake should be visualized to report a positive scan.
- b. The criterion for H/CL ratio > 1.5 as strongly positive has been removed (consistent with diagnostic criteria listed in the ASNC/AHA/ASE/

**Table 4.** Recommendations for standardized acquisition of  $^{99m}\text{Tc}$ -PYP/DPD/HMDP for cardiac amyloidosis. Adapted from Reference (207)

Imaging procedures	Parameters	Recommendation
Preparation	No specific preparation. No fasting required	Required
Scan	Rest scan	Required
Dose	$^{99m}\text{Tc}$ -PYP: 10-20 mCi (370-740 MBq) intravenously	Recommended
	$^{99m}\text{Tc}$ -DPD: 10-20 mCi (370-740 MBq) intravenously	
	$^{99m}\text{Tc}$ -HMDP: 10-20 mCi (370-740 MBq) intravenously	
Time between injection and acquisition: $^{99m}\text{Tc}$ -PYP/DPD/HMDP	2 or 3 h	Recommended
Time between injection and acquisition: $^{99m}\text{Tc}$ -PYP only	1 h	Optional. If excess blood pool activity noted on 1-h images, 3-h imaging is recommended. See below regarding image type.
General imaging parameters <sup>†</sup>		
Field of view	Heart	Required
	Chest	Optional for planar
	Heart	Recommended
CT attenuation correction		SPECT/CT fusion images helpful to localize tracer uptake to the myocardium
Image type: planar	Chest	Recommended
	2 or 3 h	1-hour planar-only imaging is not recommended
Image type: SPECT	Heart	Required
Position	Supine	Required
	Upright	Optional
Energy window	140 keV, 15-20%	Required
Collimators	Low energy, high resolution	Recommended
Matrix-Planar	256 × 256	Recommended
Matrix-SPECT	128 × 128	Recommended
Pixel size	(at least 64 by 64 is required) 2.3-6.5 mm	Recommended
Planar imaging specific parameters <sup>†</sup>		
Number of views*	Anterior and lateral	Required

**Table 4** continued

<b>Imaging procedures</b>	<b>Parameters</b>	<b>Recommendation</b>
Detector configuration	90°	Recommended
Image duration (count based)	750,000 counts	Recommended
Magnification	1.46 for large field of view systems 1.0 for small field of view systems	Recommended with goal of achieving recommended pixel size Recommended
SPECT imaging specific parameters <sup>†</sup>		
Angular range/detector configuration	180°/90° 360°/180°	Minimum Required Optional, recommended if large FOV camera is available
ECG gating	Off; Non-gated imaging	Recommended
Number of views/detector	40/32	Recommended
Time per stop	20 s/25 s	Recommended
Magnification	1.46 (180° angular range) 1.0 (360° angular range)	Recommended

ECG electrocardiogram, *PYP* pyrophosphate<sup>\*</sup>Anterior and lateral views are obtained at the same time; lateral planar views or SPECT imaging may help separate sternal from myocardial uptake<sup>†</sup>Parameters for NaI SPECT scanners

**Table 5.** Recommendations for interpretation of  $^{99m}\text{Tc}$ -PYP/DPD/HMDP for cardiac amyloidosis. Adapted from Reference (207)

---

Step 1: Visual interpretation

Evaluate planar and SPECT images to confirm diffuse radiotracer uptake in the myocardium

Differentiate myocardial radiotracer uptake from residual blood pool activity, focal myocardial infarct, and overlapping bone (e.g., from rib hot spots from fractures) on SPECT images. If excess blood pool activity is noted, recommend repeat SPECT imaging at 3 h

If myocardial tracer uptake is visually present on SPECT, proceed to step 2, semi-quantitative visual grading. If no myocardial tracer uptake is present on SPECT, the visual grade is 0

Step 2: Semi-quantitative visual grading to diagnose ATTR cardiac amyloidosis

Examine planar and SPECT images for relative tracer uptake in the myocardium relative to ribs and grade using the following scale:

Grade 0	No myocardial uptake and normal bone uptake
Grade 1	Myocardial uptake less than rib uptake
Grade 2	Myocardial uptake equal to rib uptake
Grade 3	Myocardial uptake greater than rib uptake with mild/absent rib uptake

Step 3: Heart/contralateral lung uptake ratio assessment (when applicable)

A circular ROI should be drawn over the heart on the anterior planar images with care to avoid sternal overlap and with size adjusted to maximize coverage of the heart without inclusion of adjacent lung. This ROI (same size) should be mirrored over the contralateral chest without inclusion of the right ventricle, to adjust for background and rib uptake (see Fig. 6\*). The heart and contralateral ROIs should be drawn above the diaphragm

An H/CL ratio is calculated as the fraction of heart ROI mean counts to contralateral lung ROI mean counts.

H/CL ratios of  $\geq 1.5$  at 1 h can accurately identify ATTR cardiac amyloidosis if myocardial PYP uptake is visually confirmed on SPECT and systemic AL amyloidosis is excluded (114). An H/CL ratio of  $\geq 1.3$  at 3 h can identify ATTR cardiac amyloidosis

NOTE: Diagnosis of ATTR cardiac amyloidosis cannot be made solely based on H/CL ratio alone with PYP. H/CL ratio is not recommended if there is absence of myocardial uptake on SPECT. Additionally, if the visual grade is 2 or 3, diagnosis is confirmed and H/CL ratio assessment is not necessary. H/CL ratio is typically concordant with visual grade. If discordant or the visual grade is equivocal, H/CL ratio may be helpful to classify equivocal visual grade 1 versus 2 as positive or negative

See Fig. 7.\* Grade 2 or Grade 3 uptake is consistent with ATTR cardiac amyloidosis if a monoclonal plasma cell dyscrasia is excluded, as this degree of uptake can be seen in  $> 20\%$  of patients with AL cardiac amyloidosis.(3) Grade 0 and Grade 1 uptake may be observed in AL cardiac amyloidosis and warrants further evaluation to exclude AL amyloidosis.(3) The writing group would like to emphasize the importance of excluding a monoclonal process with serum/urine immunofixation and a serum-free light-chains assay in all patients with suspected amyloidosis

Of note:  $^{99m}\text{Tc}$ -PYP/DPD/HMDP uptake could be seen in other causes of myocardial injury, including pericarditis, myocardial infarction (regional uptake), and chemotherapy or drug associated myocardial toxicity

---

\*Fig. 6 and 7 refer to figures in the original document

AL amyloid light chain, ATTR amyloid transthyretin, H/CL heart/contralateral lung, ROI region of interest

**Table 6.** Recommendations for standardized reporting of  $^{99m}\text{Tc}$ -PYP/DPD/HMDP imaging for cardiac amyloidosis. Adapted from Reference (207)

Parameters	Elements
Demographics	Patient name, age, sex, reason for the test, date of study, prior imaging procedures, biopsy results if available (required)
Methods	Imaging technique, radiotracer dose and mode of administration, interval between injection and scan, scan technique (planar and SPECT) (required)
Findings	Image quality Visual scan interpretation (required) Semi-quantitative interpretation in relation to rib uptake (required) Quantitative findings H/CL lung ratio (optional; recommended for positive scans)
Ancillary findings	Whole-body imaging if planar whole-body images are acquired (optional) Interpret CT for attenuation correction if SPECT/CT scanners are used (recommended)
Conclusions	1. An overall interpretation of the findings into categories of (1) not suggestive of ATTR cardiac amyloidosis; (2) strongly suggestive of ATTR cardiac amyloidosis; or (3) equivocal for ATTR cardiac amyloidosis after exclusion of a systemic plasma cell dyscrasia (required) (a) <b>Not suggestive:</b> A semi-quantitative visual Grade of 0 (b) <b>Equivocal:</b> If diffuse myocardial uptake of $^{99m}\text{Tc}$ -PYP/DPD/HMDP is visually confirmed and the semi-quantitative visual grade is 1 or there is interpretive uncertainty of grade 1 versus grade 2 on visual grading (c) <b>Strongly suggestive:</b> If diffuse myocardial uptake of $^{99m}\text{Tc}$ -PYP/DPD/HMDP is visually confirmed, a semi-quantitative visual grade of 2 or 3 2. Statement that evaluation for AL amyloidosis by serum FLCs, serum, and urine immunofixation is recommended in all patients undergoing $^{99m}\text{Tc}$ -PYP/DPD/HMDP scans for cardiac amyloidosis (required) 3. Statement that results should be interpreted in the context of prior evaluation and referral to a hematologist or amyloidosis expert is recommended if either: (a) Recommended echo/CMR is strongly suggestive of cardiac amyloidosis and $^{99m}\text{Tc}$ -PYP/DPD/HMDP is not suggestive or equivocal and/or (b) FLCs are abnormal or equivocal (recommended)

AL amyloid light chain, ATTR amyloid transthyretin, CMR cardiovascular magnetic resonance, echo echocardiography, FLC free light chain, H/C heart-to-contralateral lung ratio

EANM/HFSA/ISA/SCMR/SNMMI Expert Consensus Recommendations for Multimodality Imaging in Cardiac Amyloidosis: Part 2 of 2—Diagnostic Criteria and Appropriate Utilization, where H/CL ratio was not listed).

c. Conclusions have been clarified.

## References

- Sperry BW, Burgett E, Bybee KA, McGhie AI, O'Keefe JH, Saeed IM, et al. Technetium pyrophosphate nuclear scintigraphy for cardiac amyloidosis: Imaging at 1 vs 3 hours and planar vs SPECT/CT. *J Nucl Cardiol* 2020;27(5):1802–7. <https://doi.org/10.1007/s12350-020-02139-8>.
- Masri A, Bukhari S, Ahmad S, Nieves R, Eisele YS, Follansbee W, et al. Efficient 1-hour Technetium-99m pyrophosphate imaging protocol for the diagnosis of Transthyretin Cardiac Amyloidosis. *Circ Cardiovasc Imaging* 2020;13:e010249. <https://doi.org/10.1161/CIRCIMAGING.119.010249>.
- Castano A, Haq M, Narotsky DL, Goldsmith J, Weinberg RL, Morgenstern R, et al. Multicenter study of planar Technetium 99m pyrophosphate cardiac imaging: Predicting survival for patients With ATTR Cardiac Amyloidosis. *JAMA Cardiol* 2016;1(8):880–9. <https://doi.org/10.1001/jamacardio.2016.2839>.
- Tamarappoo B, Otaki Y, Manabe O, Hyun M, Cantu S, Armon Y, et al. Simultaneous Tc-99m PYP/Tl-201 dual-isotope SPECT myocardial imaging in patients with suspected cardiac amyloidosis. *J Nucl Cardiol* 2020;27:28–37.
- Manrique A, Dudoignon D, Brun S, N'Ganoa C, Cassol E, Legallois D, et al. Quantification of myocardial ( $^{99m}\text{Tc}$ )-labeled bisphosphonate uptake with cadmium zinc telluride camera in patients with transthyretin-related cardiac amyloidosis. *EJNMMI Res* 2019;9:117.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.