

University of Groningen

## Salpivory by colonial reef corals at Curaçao, Southern Caribbean

Ter Horst, Lars J. V.; Hoeksema, Bert

*Published in:*  
Diversity

*DOI:*  
[10.3390/d13110560](https://doi.org/10.3390/d13110560)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2021

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Ter Horst, L. J. V., & Hoeksema, B. (2021). Salpivory by colonial reef corals at Curaçao, Southern Caribbean. *Diversity*, 13(11), [560]. <https://doi.org/10.3390/d13110560>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Salpivory by Colonial Reef Corals at Curaçao, Southern Caribbean

Lars J. V. ter Horst<sup>1</sup> and Bert W. Hoeksema<sup>2,3,\*</sup> 

<sup>1</sup> Department of Freshwater and Marine Ecology, Institute for Biodiversity and Ecosystem Dynamics, University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, The Netherlands; larzsth@gmail.com

<sup>2</sup> Taxonomy, Systematics and Geodiversity Group, Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands

<sup>3</sup> Groningen Institute for Evolutionary Life Sciences, University of Groningen, P.O. Box 11103, 9700 CC Groningen, The Netherlands

\* Correspondence: bert.hoeksema@naturalis.nl

Salps (Phylum Tunicata, Family Salpidae) are transparent, gelatinous zooplankton that may occur in large densities (swarms) in the open ocean, where they form a food source for a variety of predators. Most salpivores are fishes, but other pelagic animals (e.g., heteropod snails) and penguins have also been reported to eat salps [1–3]. Salps may be accessible to benthic predators after they reach the bottom through currents and vertical migration [4,5]. When they die and sink to the ocean floor, their dead bodies may also be eaten by benthos [6]. It is only recently that detailed information has started to become available on corals (Phylum Cnidaria, Class Anthozoa) as salp predators [5,7,8].

These salpivorous corals include reef-dwelling scleractinians [5,8], which implies that salps ingested by stony corals contribute to the food web of coral reefs. Since salps are relatively large planktonic animals (either as solitary zooids or colonial, fusiform ones) and most of the observed predatory corals had big mouths (>1 cm wide), the capability to eat salps was linked to the predator's large gape [5,8]. This is consistent with observations of large-mouthed solitary corals and sea anemones eating jellyfish, which are also large gelatinous plankton [9,10]. Based on observations in the Mediterranean, however, it is known that the colonial scleractinians *Phyllangia americana* and *Astroides calycularis*, both with relatively small mouths (<1 cm wide), are also able to catch and consume jellyfish [11,12]. Interestingly, in the latter coral species, the polyps were observed to cooperate in order to catch large prey [12].

So far, only limited information is available on Caribbean reef coral species preying on large gelatinous plankton, and to our knowledge there is no photographic evidence available for this. In Panama (southwestern Caribbean), the massive scleractinian *Montastraea cavernosa* was observed to eat salps [13], whereas in Curaçao (Southern Caribbean), the foliose scleractinian *Agaricia agaricites* was observed to act as a salpivore (R.P.M. Bak, pers. comm.). Both coral species are colonial and prominent reef builders and widespread and common in the tropical West Atlantic.

During recent surveys at Curaçao (10 July 2021), one of us (LJVtH) observed a salp swarm in Director's Bay (12°03'59" N, 68°51'38" W) at depths of 5–10 m, where they were wave-swept or swimming slowly and aggregating near the sea floor (Video footage as Supplementary Material). The timing is consistent with February–August as the usual period of salp swarms off the southwestern part of the USA [14]. The salps are translucent in water and the in situ photographs reveal very little about the morphological characters that are used to distinguish salp taxa (Figure 1a,b). Since several salp species are known to occur in the Caribbean [15], it is impossible to identify the animals accurately and therefore it is most reliable to refer to them as Salpidae sp. It is noteworthy, though, that one salp species in particular, the widespread *Salpa fusiformis*, has been reported in Curaçao [16].



**Citation:** ter Horst, L.J.V.; Hoeksema, B.W. Salpivory by Colonial Reef Corals at Curaçao, Southern Caribbean. *Diversity* **2021**, *13*, 560. <https://doi.org/10.3390/d13110560>

Academic Editor: Michael Wink

Received: 19 October 2021


Accepted: 2 November 2021

Published: 3 November 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



**Figure 1.** Salpivory by scleractinian corals at the coral reef in Director’s Bay, Curaçao. (a,b) Solitary zooids of Salpidae sp. above the reef; test length: ca. 5 cm. (c) A salp (marked by yellow ellipse) trapped by several polyps in a branching colony of *Madracis auretenra*. (d) A large number of salps (some marked by yellow ellipses) caught by a massive coral colony of *Meandrina meandrites*. (e) A few salps (yellow ellipses) captured by a massive colony of *Montastraea cavernosa*. Photo credits: (a,b): Jason Baer; (c): LJVT; (d): Mark Little; (e): Antoni Luque.

Three common reef-dwelling scleractinians had their tentacles extended in the daytime, trying to catch the salps: *Madracis auretenra*, *Meandrina meandrites*, and *Montastraea cavernosa* (Figure 1c–e). The latter was also reported earlier [13], but the first two were newly recorded as salpivores. The bodies of the salps that were captured by the coral polyps showed signs of early decomposition (Figure 1d,e), a process that we assume to be associated with ingestion. When polyp size is taken into account, the branching *Madracis auretenra* is the most remarkable, because its polyps are very small (ca. 0.3 cm wide when extended), whereas those of the other species are larger, with *M. cavernosa* reported as having extended polyps that are ca. 1.3 cm in diameter [17].

The present observations include new results: (1) two scleractinian species were newly recorded as salpivores (*Madracis auretenra* and *Meandrina meandrites*); (2) since salp swarms occur frequently, salpivory by reef corals is probably more common and widespread than previously assumed; (3) large coral colonies are able to catch and digest several salp zooids simultaneously (Figure 1d,e), in contrast to solitary corals that can capture one or two zooids at a time [5,8]; and (4) predation on salps does not appear to be limited by polyp size. Although *Madracis auretenra* with small polyps was supposed to be more selective in its diet by preying on smaller copepods than *Montastraea cavernosa* with large polyps [17], a restriction by polyp size does not appear to apply to salpivory.

In conclusion, both solitary and colonial corals can be salpivorous and since salp swarms appear to be occurring frequently on reefs, it is probably just a matter of time before additional new records of salpivorous coral species will be added. Furthermore, aquarium experiments may help to study the timing of capture and ingestion of salps by corals more precisely, since it is not always clear whether corals eat large prey entirely or whether they expel some parts [18]. Moreover, some large prey species (such as sea slugs)

are preferred over others and consumed more quickly by corals than others [19], which indicates that more studies should be done on the ingestion of large gelatinous plankton by corals [5,7–9,11–13]. This plankton should also include ctenophores, which are also gelatinous but so far have not been reported as prey of corals.

**Supplementary Materials:** The following are available online at <https://www.mdpi.com/article/10.3390/d13110560/s1>, Video footage (two clips) of a salp swarm on the reef of Director’s Bay, Curaçao.

**Author Contributions:** Conceptualization and supervision, L.J.V.t.H. and B.W.H.; methodology and illustrations, L.J.V.t.H.; investigation, L.J.V.t.H.; writing—original draft preparation, B.W.H. and L.J.V.t.H.; writing—review and editing, B.W.H. and L.J.V.t.H. All authors have read and agreed to the published version of the manuscript.

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Acknowledgments:** We would like to express our appreciation to the staff of Carmabi and the Diveshop at Curaçao for their hospitality. A special thanks to Cynthia B. Silveira, Natascha S. Varona, Antoni Luque, Jason Baer, and Mark Little and the rest of the Rohwer lab for letting us use the pictures and the good company on the island of Curaçao. We are grateful to two reviewers for their constructive comments, which helped us to improve the manuscript.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Seapy, R.R. Predation by the epipelagic heteropod mollusk *Carinaria cristata* forma *japonica*. *Mar. Biol.* **1980**, *60*, 137–146. [[CrossRef](#)]
- Henschke, N.; Everett, J.D.; Richardson, A.J.; Suthers, I.M. Rethinking the role of salps in the ocean. *Trends. Ecol. Evol.* **2016**, *31*, 720–733. [[CrossRef](#)] [[PubMed](#)]
- Cavallo, C.; Chiaradia, A.; Deagle, B.E.; McInnes, J.C.; Sánchez, S.; Hays, G.C.; Reina, R.D. Molecular analysis of predator scats reveals role of salps in temperate inshore food webs. *Front. Mar. Sci.* **2018**, *5*, 381. [[CrossRef](#)]
- Madin, L.P.; Kremer, P.; Hacker, S. Distribution and vertical migration of salps (Tunicata, Thaliacea) near Bermuda. *J. Plankton Res.* **1996**, *18*, 747–755. [[CrossRef](#)]
- Hoeksema, B.W.; Waheed, Z. It pays to have a big mouth: Mushroom corals ingesting salps at northwest Borneo. *Mar. Biodivers.* **2012**, *42*, 297–302. [[CrossRef](#)]
- Henschke, N.; Bowden, D.A.; Everett, J.D.; Holmes, S.P.; Kloser, R.J.; Lee, R.W.; Suthers, I.M. Salp-falls in the Tasman Sea: A major food input to deep-sea benthos. *Mar. Ecol. Prog. Ser.* **2013**, *491*, 165–175. [[CrossRef](#)]
- Gili, J.M.; Rossi, S.; Pagès, F.; Orejas, C.; Teixidó, N.; López-González, P.J.; Arntz, W.E. A new trophic link between the pelagic and benthic systems on the Antarctic shelf. *Mar. Ecol. Prog. Ser.* **2006**, *322*, 43–49. [[CrossRef](#)]
- Mehrotra, R.; Scott, C.M.; Hoeksema, B.W. A large gape facilitates predation on salps by *Heteropsammia* corals. *Mar. Biodivers.* **2016**, *46*, 323–324. [[CrossRef](#)]
- Alamaru, A.; Bronstein, O.; Loya, Y.; Dishon, G. Opportunistic feeding by the fungiid coral *Fungia scruposa* on the moon jellyfish *Aurelia aurita*. *Coral Reefs* **2009**, *28*, 865. [[CrossRef](#)]
- Hoeksema, B.W.; Tuti, Y.; Becking, L.E. Mixed medusivory by the sea anemone *Entacmaea medusivora* (Anthozoa: Actiniaria) in Kakaban Lake, Indonesia. *Mar. Biodivers.* **2015**, *45*, 141–142. [[CrossRef](#)]
- Aguilar, R. *The Corals of the Mediterranean*; Fondazione Zegna & Oceana: Biella, Italy, 2007; p. 85.
- Musco, L.; Vega Fernández, T.; Caroselli, E.; Roberts, J.M.; Badalamenti, F. Proto-cooperation among small polyps allows the coral *Astroides calycularis* to prey on large jellyfish. *Ecology* **2018**, *99*, 2400–2401. [[CrossRef](#)] [[PubMed](#)]
- Porter, J.W. Zooplankton feeding by the Caribbean reef-building coral *Montastrea cavernosa*. In Proceedings of the Second International Symposium on Coral Reefs, Brisbane, Australia, 22 June–2 July 1974; Volume 1, pp. 111–125.
- Deibel, D.; Paffenhöfer, G.A. Predictability of patches of neritic salps and doliolids (Tunicata, Thaliacea). *J. Plankton Res.* **2009**, *31*, 1571–1579. [[CrossRef](#)]
- Hereu, C.M.; Suarez-Morales, E. Checklist of the salps (Tunicata, Thaliacea) from the western Caribbean Sea with a key for their identification and comments on other North Atlantic salps. *Zootaxa* **2012**, *32*, 50–60. [[CrossRef](#)]
- van Soest, R.M.W. A revision of the genera *Salpa* Forskål, 1775, *Pegea* Savigny, 1816, and *Ritteriella* Metcalf, 1919 (Tunicata, Thaliacea). *Beaufortia* **1974**, *22*, 153–191.
- Sebens, K.P.; Vandersall, K.S.; Savina, L.A. Zooplankton capture by two scleractinian corals, *Madracis mirabilis* and *Montastrea cavernosa*, in a field enclosure. *Mar. Biol.* **1996**, *127*, 303–317. [[CrossRef](#)]

- 
18. Mehrotra, R.; Scott, C.M.; Rohrer, J.M.; Hoeksema, B.W. Predation on a sacoglossan gastropod by a mushroom coral. *Coral Reefs* **2015**, *34*, 517. [[CrossRef](#)]
  19. Mehrotra, R.; Monchanin, C.; Scott, C.M.; Phongsuwan, N.; Caballer Gutierrez, M.C.; Chavanich, S.; Hoeksema, B.W. Selective consumption of sacoglossan sea slugs (Mollusca: Gastropoda) by scleractinian corals (Cnidaria: Anthozoa). *PLoS ONE* **2019**, *14*, e0215063. [[CrossRef](#)] [[PubMed](#)]