

University of Groningen

The long-term course of cognition in bipolar disorder

Samamé, Cecilia; Cattaneo, Brenda Lucía; Richaud, María Cristina; Strejilevich, Sergio; Arahamian, Ivan

Published in:
 Psychological Medicine

DOI:
[10.1017/S0033291721004517](https://doi.org/10.1017/S0033291721004517)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Samamé, C., Cattaneo, B. L., Richaud, M. C., Strejilevich, S., & Arahamian, I. (2022). The long-term course of cognition in bipolar disorder: A systematic review and meta-analysis of patient-control differences in test-score changes. *Psychological Medicine*, *52*(2), 217 - 228.
<https://doi.org/10.1017/S0033291721004517>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Review Article

Cite this article: Samamé C, Cattaneo BL, Richaud MC, Strejilevich S, Aprahamian I (2021). The long-term course of cognition in bipolar disorder: a systematic review and meta-analysis of patient-control differences in test-score changes. *Psychological Medicine* 1–12. <https://doi.org/10.1017/S0033291721004517>

Received: 11 August 2021

Revised: 14 October 2021

Accepted: 15 October 2021


Key words:

Bipolar disorder; cognition; long-term; meta-analysis

Author for correspondence:

Cecilia Samamé,
E-mail: ceciliasamame@gmail.com

The long-term course of cognition in bipolar disorder: a systematic review and meta-analysis of patient-control differences in test-score changes

Cecilia Samamé^{1,2} , Brenda Lucía Cattaneo², María Cristina Richaud¹, Sergio Strejilevich^{2,3} and Ivan Aprahamian^{4,5}

¹National Scientific and Technical Research Council (CONICET), Buenos Aires, Argentina; ²Favaloro University, Buenos Aires, Argentina; ³AREA, Assistance and Research in Affective Disorders, Buenos Aires, Argentina; ⁴Group of Investigation on Multimorbidity and Mental Health in Aging (GIMMA), Geriatrics Division, Internal Medicine Department, Jundiaí Medical School, Jundiaí, Brazil and ⁵Department of Psychiatry, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

Abstract

Neuropsychological impairment represents a key aspect of bipolar disorder (BD) that is evident even in early-course patients and is a strong predictor of functional outcomes among those affected. Previous meta-analyses of longitudinal studies suggest that BD-related cognitive deficits may not progress along the course of the disorder. However, short test-retest periods were used in most primary studies and comparisons with healthy controls were limited. The aim of this review was to synthesize the findings of research reports comparing long-term neurocognitive trajectories between BD patients and healthy individuals. PubMed, PsycINFO, and Scopus databases were searched from inception through July 2021. Publications were considered for inclusion if they reported cognitive test scores of BD patients and healthy controls at two different time points, with a minimum test-retest interval of 5 years. Fifteen studies compared the long-term course of cognition in BD patients with that of healthy controls. Ten of these were included in the quantitative analysis and involved 540 BD patients and 644 healthy individuals (mean follow-up period: 8.9 years). Patient-control effect sizes (standardized mean differences) were calculated for test-score changes in 24 neuropsychological variables and combined by means of meta-analytic procedures. No significant differences were found between patients and controls regarding long-term cognitive outcomes. These findings are consistent with previous shorter-term longitudinal meta-analyses and do not provide evidence for progressive cognitive deterioration in most bipolar individuals. Future studies should address the longitudinal course of cognition in different subgroups of BD patients and its prognostic and therapeutic value.

Introduction

Bipolar disorder (BD) comprises a group of chronic and recurrent neuropsychiatric conditions characterized by pathological mood instability, which finds its maximum expression in the full-blown manic and depressive episodes that many affected individuals experience throughout their lifetime. At present, it has been widely documented that a substantial proportion of BD patients exhibit measurable neuropsychological impairments, which persist beyond acute mood episodes (Bourne et al., 2013; Kjærstad, Eikeseth, Vinberg, Kessing, & Miskowiak, 2019; Varo et al., 2020) and have been shown to be highly predictive of suboptimal outcomes in different aspects of real-world functioning (Ehrminger et al., 2019; Gilbert & Marwaha, 2013; Gitlin & Miklowitz, 2017). Consequently, cognitive impairments are currently acknowledged as key aspects of BD that should be targeted in the clinical management of affected individuals (e.g. controlling variables known to worsen cognitive outcomes) and, in turn, in our theoretical and research considerations, as these neuropsychological features could shed light on the underlying mechanisms of the disorder. However, a large knowledge gap exists about the causes, onset, correlates, and longitudinal course of cognitive dysfunction in BD.

In contrast to research findings in the field of psychotic disorders, most population-based studies of premorbid neuropsychological functioning in BD have reported good and even outstanding cognitive outcomes (Koenen et al., 2009; MacCabe et al., 2010; Reichenberg et al., 2002; Smith et al., 2015). However, these studies explored neurocognitive performance by means of intelligence quotient scales, fluid intelligence tests, or proxies for cognitive outcomes (i.e. scholastic achievement) and therefore did not provide any information regarding individuals' functioning in those circumscribed domains typically found to be impaired in BD

patients (i.e. verbal episodic memory, processing speed, attention, and executive processes). Consequently, the possibility of neuropsychological dysfunction occurring in the premorbid phases of BD cannot be excluded. Indeed, studies of young offspring of bipolar patients have reported the presence of deficits in the same cognitive domains found to be impaired in well-established BD and with similar inter-individual heterogeneity (Bora *et al.*, 2019; Bora & Özerdem, 2017a). In addition, prospective studies of adolescents and young adults at high risk for BD, though scant, have reported impaired cognitive outcomes in individuals with subsequent development of the disorder (Meyer *et al.*, 2004; Ratheesh *et al.*, 2013). Altogether, these pieces of evidence suggest that cognitive impairments may be present prior to illness onset (despite being unclear when exactly their emergence occurs), and they may be selective and milder than those observed in premorbid psychotic disorders. Thus, neurodevelopmental abnormalities may play a role in BD-related cognitive dysfunction (Martino, Samamé, Ibañez, & Strejilevich, 2015).

In well-established BD, both state- and trait-related neuropsychological impairments represent a consistent finding even in early-course patients (i.e. following the first episode of mania) (Bora & Pantelis, 2015; Bourne *et al.*, 2013; Chakrabarty *et al.*, 2021; Kjærstad *et al.*, 2020). In recent years, a widespread hypothesis known as ‘neuroprogression’ has proposed that such impairments develop during the long-term course of the disorder as a result of illness-related neurodegenerative changes (Carvalho, Firth, & Vieta, 2020; Kapczynski *et al.*, 2014; Velosa *et al.*, 2020). This assumption was initially supported by findings of cross-sectional studies that reported a relationship between a larger number of previous affective episodes (especially of mania) and severity of cognitive deficits (López-Jaramillo *et al.*, 2010; Robinson & Ferrier, 2006). However, early follow-up studies of cognition in BD have not supported neuroprogression, and alternative explanations have been proposed for the association between number of manic episodes and cognitive deficits (for a review, see Strejilevich, Samamé, & Martino, 2015). For instance, cognitive dysfunction in BD might be a severity marker associated with an increased number of recurrences and a poorer clinical course rather than being the consequence of cumulative effects of multiple mood episodes (Martino *et al.*, 2013).

It is evident that only findings from longitudinal research might provide direct evidence to answer the question about cognitive stability or decline in BD. At present, three meta-analyses have been conducted to synthesize the results of investigations on the longitudinal course of cognition in bipolar individuals. A pioneering meta-analysis by Samamé, Martino, and Strejilevich (2014), which combined the results of studies including only euthymic participants on both test and retest occasions, did not find any significant differences between BD patients’ performance at baseline and after a mean follow-up period of 4.62 years for 14 neuropsychological variables. However, data from controls were scant and only made it possible to perform meta-analyses for four cognitive variables, for which no significant patient-control differences were found regarding test-retest effect sizes in a mean interval of 2.2 years. In a subsequent meta-analysis including a larger number of studies (Bora & Özerdem, 2017b), similar results were found for BD individuals’ test-retest differences (mean follow-up period: 3.7 years). In addition, this meta-analysis compared longitudinal cognitive outcomes between short-term (mean duration: <3 years) and long-term (mean duration: ≥ 3 years) follow-up studies and reported similar results between subgroups for seven cognitive

variables. When patients’ test-retest neuropsychological differences were compared with those of healthy controls, similar trajectories were observed. However, this analysis included only seven neuropsychological variables, and the follow-up periods used at the primary study level ranged from 1 to 5 years (mean follow-up period: 3.3 years). More recently, a small review by Szmulewicz, Valerio, and Martino (2020) analyzed the longitudinal course of neurocognitive functioning in recent-onset BD (mean follow-up period: 1.4 years) and late-life BD (mean follow-up period: 2.8 years) and found no evidence of cognitive deterioration. However, only a limited number of cognitive domains were analyzed besides using short follow-up periods, which may have been insufficient to detect changes in cognitive function.

In this context, further evidence from controlled long-term follow-up studies is needed to gain insight into the course of neuropsychological functioning in BD and determine whether progressive decline indeed occurs. Broadening our knowledge on the longitudinal trajectory of cognition in BD would contribute to a better understanding of the pathophysiological mechanisms of the disorder, the establishment of possible bipolar subtypes, the identification of targets for treatment, and the development of more efficacious therapeutic strategies. The aim of the current study was to conduct a systematic review and meta-analysis of studies exploring the long-term neurocognitive course (≥ 5 years) of BD individuals as compared with healthy controls.

Material and methods

Search strategy

The present study was conducted in accordance with the PRISMA 2020 Statement guidelines (Page *et al.*, 2021). The review protocol was registered in the International Prospective Register of Systematic Reviews (PROSPERO, CRD42020198367), and can be accessed at https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42020198367.

PubMed, PsycINFO, and Scopus databases were searched from inception through 1 July 2021 to retrieve publications available in English. At the first step of the search, combinations of keywords were used as follows: ‘bipolar disorder’ AND (‘cognition’ OR ‘neuropsychol*’) AND (‘longitudinal’ OR ‘follow-up’). Additionally, in order not to miss potentially relevant literature not covered by the aforementioned databases, a search was performed through Google Scholar using the same algorithm. First, titles and abstracts retrieved using this strategy were screened to identify relevant studies. Full texts of the articles identified in this initial screening were thoroughly assessed to confirm or reject their inclusion based on prespecified criteria. As a second step, the reference lists of the articles identified for inclusion and other relevant studies on the topic (e.g. systematic reviews) were checked for additional eligible reports. The different steps of the literature search were conducted independently by two reviewers (CS, BLC). Disagreements were resolved by consensus-based discussion.

Study selection criteria

Studies were considered for the current meta-analysis if they met the following criteria: (1) included a patient group with the diagnosis of BD according to standardized criteria; (2) included a healthy control group; (3) involved longitudinal study design

with neuropsychological assessment at baseline and after a follow-up period of at least 5 years; (4) provided data to estimate patient-control effect sizes for neuropsychological change; (5) included at least one behavioral measure of cognitive functioning that was used in a minimum of three independent studies.

In addition, if a study was based on a group of patients with different diagnoses including BD (e.g. 'first-episode psychoses', 'affective psychoses'), we contacted the original authors to request separate data for BD individuals. If there were studies with overlapping content based on the same patient sample, we considered the data from the study with the largest sample size. Two studies on the same patient group were only included if they reported different cognitive measures and were therefore not meta-analyzed together.

Data extraction and quality assessment

Two reviewers (CS, BLC) independently extracted data from full texts of the studies selected for inclusion using a standardized pre-coded spreadsheet. Data regarding age, education, sample size, and neuropsychological functioning at both assessment time points were extracted for both BD and control groups. When available, the following data were also extracted for the BD group: mood state at both assessment time points, subtype of the disorder, number of previous depressive/(hypo)manic episodes, duration of illness, and medication use. A consensus meeting was held to resolve any disparities between the two reviewers.

To appraise study quality, the National Institutes of Health Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies (NIH, 2014) was used, with some questions adapted to the current review. Quality assessment was performed in duplicate. Studies were considered 'good quality' if they achieved at least a score of eight (i.e. eight 'yes' answers). Disagreements regarding quality scores for each individual study were resolved by consensus.

Meta-analytic procedure

Meta-analyses were performed using the Comprehensive Meta-Analysis software version 3.0 (Borenstein, Hedges, Higgins, & Rothstein, 2013). Patient-control effect sizes for test-score changes (i.e. changes in scores of neuropsychological tests from baseline to a follow-up time point) were calculated as standardized mean differences (Hedges' g). To compute the effect sizes, a conservative value of 0.5 was assumed for pre-post correlations. For each neuropsychological test or cognitive domain composite score, the sign of patient-control effect sizes for test-retest change was adjusted so that positive effect sizes reflected greater decrease in neuropsychological performance in the BD group. Effect sizes were weighted using the inverse variance method (DerSimonian & Laird, 2015). A random-effects model was used in all the analyses performed. According to this model, the studies in the analysis are assumed to be a random sample in the universe of studies (Borenstein et al., 2013).

The Q -test for heterogeneity was used to estimate the homogeneity of the resulting mean weighted effect size for each variable. The I^2 index was calculated to describe the percentage of total variation across reports due to heterogeneity rather than chance. I^2 values of 25, 50, and 75% indicate low, moderate, and high heterogeneity, respectively (Higgins, Thompson, Deeks, & Altman, 2003). In analyses including at least five studies, publication bias (the tendency of small studies to show larger

effects) was assessed using Egger's test (Egger, Davey Smith, Schneider, & Minder, 1997).

In addition, subgroup meta-analyses were performed considering only those studies including homogeneous samples of euthymic or stable (not in an acute mood episode) BD patients at both assessment time points.

A significance level of $p < 0.05$ was set for all the analyses performed.

Cognitive variables

For the purposes of this study, the results of investigations utilizing the same test or tapping approximately the same neuropsychological construct were combined into a single summary measure. Twenty-four overall neuropsychological measures were obtained (Table 1, online Supplementary material).

Results

The selection process of included studies is summarized in Fig. 1. Fifteen studies compared the long-term course (≥ 5 years) of cognition in BD patients with that of healthy individuals and are synthesized in Table 1. Five studies were not included in the meta-analysis due to overlapping sample with other reports (Jiménez-López et al., 2019; Mora, Portella, Forcada, Vieta, & Mur, 2016; Ryan et al., 2016, 2017; Santos et al., 2014). Finally, 10 studies were included in the quantitative analysis and involved 540 BD patients and 644 healthy controls (mean follow-up period: 8.9 years). Two studies based on the same sample were included (Correa-Ghisays et al., 2017, 2019) as they explored different neuropsychological variables, but only that with the largest sample size was considered for calculating the total number of individuals included in the meta-analysis. The authors of these two reports were contacted by e-mail and requested to provide separate data corresponding to the same group of patients who underwent cognitive assessment both at baseline and after a 5-year follow-up period, as that information was not available from the publication. The studies by Camprodon-Boadas et al. (2021), Fett et al. (2020), and Zanelli et al. (2019) explored the longitudinal course of cognition in a group of patients with 'psychoses', which included both patients diagnosed with psychotic disorders and patients suffering from affective disorders with psychotic features. The authors of these studies were contacted by e-mail and requested to provide separate data for the individuals diagnosed with BD. The study by Fett et al. (2020) provided a cross-sectional comparison for healthy controls only at the end of the follow-up period but was included in the meta-analysis as it used a reference for normative cognitive change, based on recent data from a longitudinal study that included people of the same age group (in their 30s and 40s) from the same country (Hughes, Agrigoroaei, Jeon, Bruzzese, & Lachman, 2018).

Out of the total studies reviewed, most reports were based on samples of middle-aged adults. One report included only late-life individuals (Schouws, Comijs, Dols, Beekman, & Stek, 2016) and one report included only pediatric patients at baseline (Camprodon-Boadas et al., 2021). Three studies included BD individuals followed up after a first episode of psychosis (Camprodon-Boadas et al., 2021; Fett et al., 2020; Zanelli et al., 2019) (Table 1). Overall, the quality of the studies included in this review was good (Table 2, online Supplementary material).

There were no statistically significant differences for age or baseline years of education between the group of BD individuals

Table 1. Long-term (≥ 5 years) follow-up studies of neuropsychological functioning in BD patients and healthy individuals

Study	Sample ^a BD (type)/HC	Patients' age (baseline) Mean (SD)	Patients' race/ethnicity	No. of previous mood episodes (baseline) Mean (SD)	Study characteristics	Follow-up	BD patients' mood state (baseline)	BD patients' mood state (follow-up)	Cognitive variables ^b	Results ^c
Camprodon-Boadas et al. (2021)*	21 (NR)/37	16.1 (1.26)	NR	NR	All pediatric patients (age <18) at baseline, followed after a first episode of psychosis	5 years	NR	NR	-Executive functions (cognitive flexibility, response inhibition, working memory). -Episodic verbal memory -Attention	Evidence of decline was found for cognitive flexibility
Correa-Ghisays et al. (2017)*	54 (NR)/23	43 (NR)	NR	NR	All adult patients (age >18 at baseline).	5 years	Different phases of illness	Different phases of illness	-Psychomotor speed	No evidence of decline
Correa-Ghisays et al. (2019)*	65 (NR)/25	43 (NR)	NR	NR	All adult patients (age >18 at baseline).	5 years	Different phases of illness	Different phases of illness	-Episodic visual memory	No evidence of decline
Fett et al. (2020)*	46 (NR)/258	28.67 (9.55)	88% white Full racial breakdown NR	NR	Age 15–60 at baseline. History of psychotic symptoms	18 years	NR	NR	-Verbal knowledge -Episodic memory (verbal, visual) -Attention/ Processing speed -Executive functions (set-shifting, verbal fluency) -Visuoconstructive abilities	No evidence of decline
Hinrichs et al. (2017)*	159 (I-II-NOS)/54	40.70 (12.03)	NR	NR	NR	5 years	Different phases of illness	Different phases of illness	-Attention/processing speed -Episodic memory (verbal, visual) -Executive functions (response inhibition, set-shifting, flexibility) -Visuoconstructive abilities	No evidence of decline
Jiménez-López et al. (2019)	76 (I)// 40	41 (11.3)	NR	ME: 4.95 (5.1) DE: 8.98 (8.7)	All adult patients (age 18–55 at baseline)	5 years	All euthymic HAM-D < 7 YMRS < 6	All euthymic HAM-D < 7 YMRS < 6	-Processing speed -Attention/vigilance -Verbal learning and memory -Visual memory -Executive functions (working memory, set-shifting, response inhibition, verbal fluency) -Visuospatial abilities	No evidence of decline

Mora et al. (2013)*	28 (I-II)/26	41.71 (12.4)	NR	ME: 2.54 (2.1)	All adult patients (age 18–65 at baseline) on lithium (monotherapy or combined treatment)	6.1 years	All euthymic HDRS < 8 YMRS < 6	All euthymic HDRS < 8 YMRS < 6	-Executive functions (response inhibition, flexibility, set-shifting) -Attention/ processing speed -Episodic memory (verbal, visual)-Visuoconstructive abilities	No evidence of decline
Mora et al. (2016)	10 (I-II)/10	45.6 (10.9)	NR	ME: 2.2 (1.6) DE: 1.6 (1.0)	All adult patients (age 18–65 at baseline) with excellent response to lithium	6 years	All euthymic HAM-D < 8 YMRS < 6	All euthymic HAM-D < 8 YMRS < 6	-Executive functions (response inhibition, cognitive flexibility, set-shifting) -Attention/ processing speed -Episodic memory (verbal, visual) -Visuoconstructive abilities	No evidence of decline
Ryan et al. (2016)	91 (I-II-NOS)/17	42.06 (11.30)	NR	NR	NR	5 years	All symptomatic	All symptomatic	-Executive functions (response inhibition, set-shifting, flexibility)	No evidence of decline
Ryan et al. (2017)	91 (I-II-NOS)/17	42.06 (11.30)	NR	NR	NR	5 years	All symptomatic	All symptomatic	-Episodic memory (verbal, visual) -Emotion processing -Psychomotor speed	No evidence of decline.
Sánchez-Morla et al. (2019)*	76 (I-II)/40	41.4 (11.0)	NR	ME: 5.5 (5.2) DE: 6.8 (7.0)	All adult patients (age 18–55 at baseline) Predominance of type I patients	5 years	All euthymic HDRS < 8 YMRS < 6	All euthymic HDRS < 8 YMRS < 6	-Processing speed -Attention/vigilance -Verbal learning and memory -Visual memory -Executive functions (set-shifting, response inhibition, working memory, flexibility) -Visuospatial abilities	No evidence of decline. Improvement was observed in a measure of cognitive flexibility
Santos et al. (2014)	62 (I-II)/40	44.4 (10.5)	NR	ME: 5.8 (5.4) DE: 7.5 (7.1)	All adult patients (age 18–55 at baseline)	5 years	All euthymic HDRS < 7 YMRS < 6	All euthymic HDRS < 7 YMRS < 6	-Processing speed -Attention/vigilance -Verbal learning and memory -Visual memory -Executive functions (response inhibition, working memory)	Only a measure from the verbal memory domain (delayed free recall) worsened more in BD patients. No evidence of decline was observed for other variables

(Continued)

Table 1. (Continued.)

Study	Sample ^a BD (type)/HC	Patients' age (baseline) Mean (SD)	Patients' race/ethnicity	No. of previous mood episodes (baseline) Mean (SD)	Study characteristics	Follow-up	BD patients' mood state (baseline)	BD patients' mood state (follow-up)	Cognitive variables ^b	Results ^c
Schouws et al. (2016)*	56 (I-II)/44	68.16 (7.0)	NR	ME: 2.96 (1.7) DE: 4.25 (3.0)	All elderly patients (age >60 at baseline).	5 years	All euthymic CES-D < 15 YMRS < 7	All euthymic CES-D < 15 YMRS < 7	-Attention -Episodic verbal memory -Executive functions (set-shifting, verbal fluency, planning)	No evidence of decline
Sparding et al. (2021)*	72 (I-II)/59	37.4 (12.1)	Ethnically diverse sample from northern Stockholm Full racial breakdown NR	NR (only no. of episodes occurring during follow-up are available)	All adult patients (Age >18 at baseline)	6 years	All stable (not suffering from an acute mood episode: euthymic or with subsyndromal symptoms)	All stable (not suffering from an acute mood episode: euthymic or with subsyndromal symptoms)	-Attention/processing speed -Executive functions (inhibition, set-shifting, planning, verbal fluency, working memory) -Reasoning/ concept formation -Verbal learning -Visual memory	No evidence of decline
Zanelli et al. (2019)*	17 (NR)/101	29.82 (9.42)	58% white Full racial breakdown NR	NR	All patients followed after a first episode of psychosis. Age 15-65 at baseline.	9.1 years	NR	NR	-Verbal and executive IQ -Attention -Visuoconstructive abilities	No evidence of decline

BD, bipolar disorder; CES-D, Centre for Epidemiologic Studies Depression Scale; DE, depressive episodes; HC, healthy controls; HDRS/HAM-D/HRDS, Hamilton Depression Rating Scale; IQ, intelligence quotient; ME, (hypo)manic episodes; NR, not reported; NOS, not otherwise specified; s.d., standard deviation; YMRS, Young Mania Rating Scale.

Studies included in the meta-analysis.

^aSome studies included other clinical samples besides BD (e.g. psychotic disorders) or a sample of first-degree relatives of bipolar patients, but the corresponding data are not reported here as they fall beyond the scope of the current review. The sample size reported for each study corresponds to the number of participants (BD/HC) who underwent cognitive assessment on two occasions separated by a minimum interval of 5 years. The study by Fett et al. (2020) did not provide cognitive outcomes for HC at baseline, but an estimation of cognitive change was made from longitudinal normative data.

^bNeuropsychological variables are reported here only if both test and retest outcomes were available.

^cThe results reported here correspond to the long-term cognitive outcomes of individuals with a BD diagnosis as compared with healthy individuals. Given that the studies by Camprodon-Boadas et al. (2021), Fett et al. (2020), and Zanelli et al. (2019) were based on mixed samples of individuals with different diagnoses, separate data for bipolar individuals were provided by the authors of these studies. Hence, the results here reported correspond to individuals with a BD diagnosis exclusively.




Fig. 1. PRISMA 2020 flow diagram for systematic reviews.

and the healthy control group neither in the total sample nor in any of the meta-analyses performed ($p > 0.05$).

Meta-analytic results for patient-control differences in neuropsychological change

No significant patient-control differences were observed for longitudinal cognitive outcomes in any of the 24 variables explored (Table 2). Large significant heterogeneity was observed for five neuropsychological variables. Figure 2 provides forest plots depicting individual and pooled patient-control effect sizes for changes in category fluency and the Wisconsin Card Sorting Test (WCST), the two executive measures with the least consistent effects across studies. In the WCST analysis (Fig. 2), there was one clear outlier (Camprodon-Boadas et al., 2021). When removing this study, the distribution of effect sizes became largely homogeneous in the absence of significant patient-control differences [before: Hedges' $g = 0.09$, $p = 0.68$, $Q(p) = 0.006$, $I^2 = 75.76\%$; after: Hedges' $g = -0.15$, $p = 0.19$, $Q(p) = 0.47$, $I^2 = 00.00\%$]. No evidence of publication bias was observed (Table 3, online Supplementary material).

Subgroup meta-analysis of euthymic patients

Subgroup meta-analyses were performed including only studies of euthymic/stable individuals at both assessment time points. Four studies (Mora, Portella, Forcada, Vieta, & Mur, 2013; Sánchez-Morla et al., 2019; Schouws et al., 2016; Sparding et al., 2021) compared longitudinal neuropsychological outcomes between 232 euthymic/stable BD patients and 169 healthy controls (mean follow-up: 5.5 years) and were included in this analysis. No significant between-group differences were found for 11 of

the 12 variables analyzed (Table 3). Homogeneous distributions of effect sizes were observed for most of the analyses performed.

Discussion

This meta-analysis aimed to compare the long-term cognitive trajectory of BD patients with that of healthy individuals. Meta-analytic findings for 24 variables yielded no significant patient-control differences in test-retest neuropsychological change (mean follow-up: 8.9 years). In a subanalysis considering studies of euthymic individuals only (mean follow-up: 5.5 years), no significant patient-control differences were found for 11 out of 12 cognitive variables analyzed. In addition, at the primary study level, most studies reported similar long-term cognitive trajectories between bipolar patients and healthy individuals.

Our results are in keeping with previous meta-analyses of longitudinal neuropsychological outcomes in BD (Bora & Özerdem, 2017b; Samamé et al., 2014; Szmulewicz et al., 2020). However, unlike these meta-analyses, the current review includes primary studies with a long follow-up period and a healthy control group in addition to analyzing a much larger number of neurocognitive variables. Therefore, the results here reported provide more robust evidence not supportive of the concept of progressive cognitive deterioration of BD individuals as a group.

Furthermore, the results of our review and those from previous meta-analyses are in line with the classic concept of BD, according to which leaving aside the general impact of mood swings on the overall functioning of those affected, cognition and general intelligence are preserved along the course of the disorder. Despite these considerations, it is not possible to exclude the possibility that cognitive deterioration indeed occurs. Several issues should

Table 2. Pooled weighted effect sizes for patient-control differences in test-score changes.

Variable	K	BD	HC	ES	CI 95%	Z	p	Q test(p)	I ² (%)
TMT, part A	5	365	422	0.04	−0.12 to 0.20	0.51	0.61	0.88	0.00
Digit-symbol coding	4	210	457	0.06	−0.12 to 0.25	0.68	0.50	0.73	0.00
Sustained attention (CPT)	3	149	108	0.29	−0.08 to 0.65	1.52	0.13	0.13	51.61
Processing speed (composite score)	8	511	541	0.10	−0.03 to 0.24	1.49	0.14	0.84	0.00
Digit span forward	3	105	107	−0.06	−0.33 to 0.21	−0.40	0.69	0.61	0.00
Digit span backward	4	181	147	0.06	−0.16 to 0.28	0.56	0.58	0.70	0.00
TMT, part B	6	421	477	0.03	−0.15 to 0.20	0.30	0.77	0.26	22.69
Stroop	7	447	516	0.06	−0.08 to 0.20	0.82	0.41	0.82	0.00
WCST	4	284	157	0.09	−0.34 to 0.52	0.41	0.68	0.006	75.76
Category fluency	4	357	193	0.09	−0.18 to 0.36	0.65	0.51	0.08	55.56
Phonological fluency	5	365	422	−0.03	−0.22 to 0.16	−0.29	0.77	0.27	23.40
Working memory (composite score)	4	181	147	0.04	−0.18 to 0.26	0.36	0.72	0.73	0.00
VLT – list learning	6	374	251	0.10	−0.15 to 0.35	0.77	0.44	0.05	54.18
VLT – free immediate recall	5	340	201	0.05	−0.28 to 0.38	0.30	0.76	0.01	68.03
VLT – free delayed recall	5	340	201	−0.09	−0.31 to 0.13	−0.82	0.41	0.24	27.79
VLT – delayed recall (recognition)	3	160	110	−0.21	−0.45 to 0.03	−1.69	0.09	0.72	0.00
Short-term verbal memory (composite score)	6	386	459	0.09	−0.18 to 0.35	0.63	0.53	0.02	63.60
Long-term verbal memory (composite score)	6	386	459	−0.03	−0.23 to 0.16	−0.34	0.73	0.19	33.24
CRF (immediate recall)	4	330	177	−0.02	−0.21 to 0.16	−0.23	0.82	0.75	0.00
CRF (delayed recall)	4	328	145	−0.12	−0.32 to 0.08	−1.20	0.23	0.42	0.00
Short-term visual memory (composite score)	5	376	435	0.14	−0.20 to 0.48	0.82	0.41	0.002	77.13
Long-term visual memory (composite score)	5	374	403	0.05	−0.31 to 0.41	0.28	0.78	0.002	76.60
CRF (copy)	3	263	120	−0.10	−0.43 to 0.23	−0.58	0.56	0.13	51.75
Visuoconstructive abilities (composite score)	5	351	279	−0.10	−0.29 to −0.09	−1.03	0.30	0.33	13.81

CPT, Continuous Performance Tests; CRF, Complex Rey Figure; TMT, Trail Making Test; VLT, Verbal Learning Tests; WCST, Wisconsin Card Sorting Test; K, number of studies; BD, bipolar disorder patients; HC, healthy controls; ES, effect size (Hedges' *g*); CI, confidence interval.

be taken into account before drawing any firm conclusions about the long-term course of cognition in BD.

First, a number of investigations using cluster-analytic approaches have consistently demonstrated the existence of at least three different cognitive subgroups of BD individuals, including a 'cognitively intact group' in comparison with healthy individuals, a 'globally impaired group', with severe and generalized deficits, and a 'moderately impaired group' with moderate or selective impairment (Burdick et al., 2014; Chakrabarty et al., 2021; Varo et al., 2020). These distinct groups of BD individuals could have different long-term neurocognitive trajectories. However, it should be noted that cognitive subgroups have been found to be equally present in young (Frias et al., 2017) and late-life patients, with a remarkable similarity in their distribution compared to that found in middle-aged patients (Martino, Marengo, Igoa, & Strejilevich, 2018). These findings and the fact that cognitive subgroups are detectable even in first-episode BD and before illness onset (Bora et al., 2019; Chakrabarty et al., 2021) suggest that cognitive heterogeneity is not the result of neuroprogressive changes or solely attributable to differences in long-term exposure to pharmacological agents. It is likely that these subgroups reflect different underlying processes and cognitive reserve, which, in turn, may interact with other clinical

characteristics. Another issue that warrants consideration when interpreting the neurocognitive heterogeneity observed among bipolar individuals is the fact that some cases of late-onset BD (LOBD) may be behavioral manifestations of other conditions that mimic the disorder. LOBD, which is associated with more severe cognitive deficits (Samamé, Martino, & Strejilevich, 2013; Schouws et al., 2009) and a possible progressive course, has been proposed to be different from the 'primary' and most prevalent forms of the disorder as its etiology may be related to vascular illness or early manifestations of neurodegenerative diseases (Mendez, Parand, & Akhlaghipour, 2020; Subramaniam, Dennis, & Byrne, 2007). As evident, both illness-related processes and other clinical factors (e.g. drug treatment, severity of the course of illness) have an impact on the overall cognitive picture of BD, in which the relative weight of each variable remains to be determined. All this said, although progressive cognitive decline is not a general rule in BD, it is possible that a subgroup of patients presents with a neuroprogressive course. In line with this hypothesis, the study by Sánchez-Morla et al. (2019) reported that the number of previous (hypo)manic episodes was a significant predictor of long-term progression of neuropsychological impairments. Similarly, Schouws et al. (2016) found that more manic symptoms were significantly associated with a greater decline in




Fig. 2. Forest plots of individual and pooled patient-control effect sizes for changes in two executive measures. BD, bipolar disorder patients; HC, healthy controls.

Table 3. Pooled weighted effect sizes for patient-control differences in test-score changes (subanalysis of studies including euthymic patients on both assessment occasions)

Variable	<i>K</i>	BD	HC	ES	CI 95%	<i>Z</i>	<i>p</i>	Q test(<i>p</i>)	<i>I</i> ² (%)
TMT, part A	3	160	110	0.04	-0.21 to 0.28	0.29	0.77	0.81	0.00
Processing speed (composite score)	4	231	169	0.10	-0.10 to 0.30	0.99	0.32	0.76	0.00
TMT, part B	4	216	165	0.00	-0.28 to 0.28	-0.01	0.99	0.13	47.05
Category fluency	3	198	139	0.05	-0.34 to 0.44	0.26	0.80	0.04	68.04
Phonological fluency	3	160	110	-0.16	-0.41 to 0.09	-1.25	0.21	0.35	4.77
Digit span backward	3	160	110	0.04	-0.20 to 0.28	0.31	0.75	0.54	0.00
Stroop	4	221	167	0.11	-0.09 to 0.31	1.04	0.30	0.70	0.00
VLT – list learning	4	194	160	0.26	0.05–0.48	2.43	0.02	0.54	0.00
VLT – free immediate recall	3	160	110	0.26	-0.17 to 0.67	1.19	0.24	0.05	65.73
VLT – free delayed recall	3	160	110	0.04	-0.20 to 0.29	0.36	0.72	0.52	0.00
VLT – delayed recall (recognition)	3	160	110	-0.21	-0.45 to 0.04	-1.69	0.09	0.72	0.00
CRF (immediate recall)	3	171	123	0.02	-0.21 to 0.26	0.19	0.85	0.66	0.00

CRF, Complex Rey Figure; TMT, Trail Making Test; VLT, Verbal Learning Tests; WCST, Wisconsin Card Sorting Test; *K*, number of studies; BD, bipolar disorder patients; HC, healthy controls; ES, effect size (Hedges' *g*); CI, confidence interval.

memory. Further studies considering the neurocognitive heterogeneity of BD may contribute to better understanding of the expected cognitive evolution of the disorder and disentangling the correlates of neuropsychological dysfunction.

Second, it is possible that, in the most 'classic' forms of BD, cognitive deterioration occurs prior to the onset of the disorder as a result of illness-related processes and may remain quite stable during the course of illness. It is also possible that such cognitive decline occurs when the first mood manifestations of the disorder emerge in the absence of a BD diagnosis [taking into account that

depression occurs long before the first episode of (hypo)mania that is necessary for performing any diagnosis] or immediately after the first well-established manic episode and then remain stable over time. For instance, the only study reviewed here that documented a decline in executive functions (Camprodon-Boadas et al., 2021) was based on pediatric patients who were followed up after their first episode of psychosis, and the differences observed with other studies could be explained by the fact that this study included younger patients in the very early manifestations of BD. It is evident that neurodevelopmental factors do not

exclude the possibility of neurodegeneration. However, to prove a progressive neurodevelopmental nature of BD, more consistent evidence for progressive deterioration is needed. At present, the strongest empirical support for neuroprogression is drawn from studies showing progressive cortical thinning in bipolar individuals (Carvalho et al., 2020). However, these studies have not controlled for variables such as exposure to antipsychotic drugs, which have also been correlated with cortical thinning (Hibar et al., 2018; Strejilevich, Quiroz, & Bitran, 2020).

Third, it is possible that cognitive decline occurs in the very-long term and the studies reviewed here do not capture such changes. As is evident, follow-up periods of more than a decade are difficult to accomplish in research settings with accurate data. However, the findings of the only meta-analysis of neuropsychological outcomes in late-life bipolar individuals, with longstanding illness in most cases, showed a pattern and a magnitude of cognitive impairments similar to those observed among younger bipolar individuals (Samamé et al., 2013).

A number of limitations should be acknowledged when interpreting the results of this meta-analysis. Of note, very few reports were included in some analyses. However, most studies were rated as 'good quality' and yielded quite consistent results. It should also be noted that the studies reviewed were quite diverse as regards clinical and demographic characteristics of patient samples. In some individual studies, considerable between-patient heterogeneity was also observed regarding such variables. Furthermore, it is worthy of note that, in terms of race and ethnicity, there was very limited information on how diverse the samples were, as the vast majority of studies did not report these important variables. All these issues may hinder comprehension of the relationship between specific characteristics of bipolar patients and cognitive change. In addition, the small number of studies included prevented us from conducting meta-regression or further subgroup analyses to explore the relation between change-score effect sizes and characteristics of each study. Additional long-term follow-up studies are needed to help ascertain the generalizability of our results and gain insight into the influence of certain variables on the long-term cognitive outcomes of BD patients. Other shortcomings are essentially those of the primary studies reviewed. First, despite a large number of neuropsychological variables being included, long-term outcomes for emotion processing and 'hot' executive domains such as decision-making could not be analyzed given the lack of longitudinal primary studies exploring these domains. Second, differences in pharmacological variables between assessment occasions could not be controlled and may have influenced the results of this review. Maintenance of pharmacologic status over prolonged periods of time is hardly possible to accomplish in BD patients and may have an impact on cognition. Indeed, preliminary findings suggest that prolonged exposure to lithium may have a protective effect on the risk of cognitive deterioration (De-Paula, Gattaz, & Forlenza, 2016; Forlenza, Radanovic, Talib, & Gattaz, 2019; Won & Kim, 2017). However, the alleged long-term neurotrophic effects of this agent have yet to be confirmed in large, controlled studies of bipolar individuals. By contrast, other psychotropic drugs commonly prescribed to bipolar patients, such as antipsychotics, have been shown to be related to gray matter loss and cognitive impairment (Cullen et al., 2016; Torrent et al., 2011; Vita, De Peri, Deste, Barlati, & Sacchetti, 2015). Another important limitation of the current review involves the fact that some of the primary reports did not provide mean scores for mood rating scales both at baseline

and after the follow-up period. Therefore, it was not possible to explore differences in mood state between time points that could have influenced our results. Lack of information was also evident at the primary study level regarding race and ethnicity, and it was therefore not possible to explore how these variables may have influenced the observed outcomes. Finally, the potential influence of attrition on the results of this study should not be overlooked. As evident, the number of missing observations increases over time, with attrition rates ranging between 15% and 60% in the studies reviewed, thus raising concerns of drop-out bias. Therefore, although there is no evidence so far suggesting that patients who dropped out were those who would display worse cognitive outcomes, this hypothesis cannot be excluded.

To conclude, although it is not possible to assert that cognitive deficits are stable in BD, the strongest and most updated evidence from longitudinal studies, which is synthesized in the current meta-analysis, is not supportive of the hypothesis of neuroprogression in BD. Further studies should map longitudinal trajectories in different cognitive subgroups of bipolar individuals combining behavioral task with neuroimaging techniques and trying to control the effects of mood and medication variables on the observed outcomes.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0033291721004517>

Financial support. This study was supported by a grant from the National Scientific and Technical Research Council (CONICET, Buenos Aires, Argentina) awarded to Dr Samamé. The funding source played no role in the study design, collection, analysis, or interpretation of the data, in the writing of this study, or in the decision to submit this article for publication.

Conflict of interest. None.

References^{†*}

- Bora, E., Can, G., Ildiz, A., Ulas, G., Ongun, C. H., Inal, N. E., & Ozerdem, A. (2019). Neurocognitive heterogeneity in young offspring of patients with bipolar disorder: The effect of putative clinical stages. *Journal of Affective Disorders*, 257, 130–135. doi: 10.1016/j.jad.2019.07.015
- Bora, E., & Özerdem, A. (2017a). A meta-analysis of neurocognition in youth with familial high risk for bipolar disorder. *European Psychiatry: the Journal of the Association of European Psychiatrists*, 44, 17–23. doi: 10.1016/j.eurpsy.2017.02.483
- Bora, E., & Özerdem, A. (2017b). Meta-analysis of longitudinal studies of cognition in bipolar disorder: Comparison with healthy controls and schizophrenia. *Psychological Medicine*, 47(16), 2753–2766. doi: 10.1017/S0033291717001490
- Bora, E., & Pantelis, C. (2015). Meta-analysis of cognitive impairment in first-episode bipolar disorder: Comparison with first-episode schizophrenia and healthy controls. *Schizophrenia Bulletin*, 41(5), 1095–1104. doi: 10.1093/schbul/sbu198
- Borenstein, M., Hedges, L., Higgins, J., & Rothstein, H. (2013). *Comprehensive meta-analysis*, version 2. Englewood, NJ: Biostat.
- Bourne, C., Aydemir, Ö., Balanzá-Martínez, V., Bora, E., Brissos, S., Cavanagh, J. T., ... Goodwin, G. M. (2013). Neuropsychological testing of cognitive impairment in euthymic bipolar disorder: An individual patient data meta-analysis. *Acta Psychiatrica Scandinavica*, 128(3), 149–162. doi: 10.1111/acps.12133
- Burdick, K. E., Russo, M., Frangou, S., Mahon, K., Braga, R. J., Shanahan, M., & Malhotra, A. K. (2014). Empirical evidence for discrete neurocognitive subgroups in bipolar disorder: Clinical implications. *Psychological Medicine*, 44(14), 3083–3096. doi: 10.1017/S0033291714000439

[†]The notes appear after the main text.

*References indicated with an asterisk represent studies included in the meta-analysis

- *Camprodón-Boadas, P., de la Serna, E., Baeza, I., Puig, O., Ilzarbe, D., Sugranyes, G., ... Castro-Fornieles, J. (2021). Cognitive reserve in patients with first-episode psychosis as outcome predictor at 5-year follow-up. *European Child & Adolescent Psychiatry*, 30(12), 1959–1967. doi: 10.1007/s00787-020-01668-4
- Carvalho, A. F., Firth, J., & Vieta, E. (2020). Bipolar disorder. *The New England Journal of Medicine*, 383(1), 58–66. doi: 10.1056/NEJMra1906193
- Chakrabarty, T., Torres, I. J., Su, W. W., Sawatzky, R., Keramatian, K., & Yatham, L. N. (2021). Cognitive subgroups in first episode bipolar I disorder: Relation to clinical and brain volumetric variables. *Acta Psychiatrica Scandinavica*, 143(2), 151–161. doi: 10.1111/acps.13245
- *Correa-Ghisays, P., Balanzá-Martínez, V., Selva-Verá, G., Vila-Francés, J., Soria-Olivas, E., Vivas-Lalinde, J., ... Tabarés-Seisdedos, R. (2017). Manual motor speed dysfunction as a neurocognitive endophenotype in euthymic bipolar disorder patients and their healthy relatives. Evidence from a 5-year follow-up study. *Journal of Affective Disorders*, 215, 156–162. doi: 10.1016/j.jad.2017.03.041
- *Correa-Ghisays, P., Sánchez-Ortí, J. V., Ayesa-Arriola, R., Setién-Suero, E., Balanzá-Martínez, V., Selva-Verá, G., ... Tabarés-Seisdedos, R. (2019). Visual memory dysfunction as a neurocognitive endophenotype in bipolar disorder patients and their unaffected relatives. Evidence from a 5-year follow-up Valencia study. *Journal of Affective Disorders*, 257, 31–37. doi: 10.1016/j.jad.2019.06.059
- Cullen, B., Ward, J., Graham, N. A., Deary, I. J., Pell, J. P., Smith, D. J., & Evans, J. J. (2016). Prevalence and correlates of cognitive impairment in euthymic adults with bipolar disorder: A systematic review. *Journal of Affective Disorders*, 205, 165–181. doi: 10.1016/j.jad.2016.06.063
- De-Paula, V. J., Gattaz, W. F., & Forlenza, O. V. (2016). Long-term lithium treatment increases intracellular and extracellular brain-derived neurotrophic factor (BDNF) in cortical and hippocampal neurons at subtherapeutic concentrations. *Bipolar Disorders*, 18(8), 692–695. doi: 10.1111/bdi.12449
- DerSimonian, R., & Laird, N. (2015). Meta-analysis in clinical trials revisited. *Contemporary Clinical Trials*, 45, 139–145. doi: 10.1016/j.cct.2015.09.002
- Egger, M., Davey Smith, G., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple, graphical test. *BMJ (Clinical Research Ed.)*, 315(7109), 629–634. doi: 10.1136/bmj.315.7109.629
- Ehrminger, M., Brunet-Gouet, E., Cannavo, A. S., Aouizerate, B., Cussac, I., Azorin, J. M., ... Roux, P. (2019). Longitudinal relationships between cognition and functioning over 2 years in euthymic patients with bipolar disorder: A cross-lagged panel model approach with the FACE-BD cohort. *The British Journal of Psychiatry: the Journal of Mental Science*, 218(2), 80–87. doi: 10.1192/bjp.2019.180
- *Fett, A. J., Velthorst, E., Reichenberg, A., Ruggiero, C. J., Callahan, J. L., Fochtmann, L. J., ... Kotov, R. (2020). Long-term changes in cognitive functioning in individuals with psychotic disorders: Findings from the Suffolk county mental health project. *JAMA Psychiatry*, 77(4), 387–396. doi: 10.1001/jamapsychiatry.2019.3993
- Forlenza, O. V., Radanovic, M., Talib, L. L., & Gattaz, W. F. (2019). Clinical and biological effects of long-term lithium treatment in older adults with amnesic mild cognitive impairment: Randomised clinical trial. *The British Journal of Psychiatry: the Journal of Mental Science*, 215(5), 668–674. doi: 10.1192/bjp.2019.76
- Frías, Á., Dickstein, D. P., Merranko, J., Gill, M. K., Goldstein, T. R., Goldstein, B. I., ... Birmaher, B. (2017). Longitudinal cognitive trajectories and associated clinical variables in youth with bipolar disorder. *Bipolar Disorders*, 19(4), 273–284. doi: 10.1111/bdi.12510
- Gilbert, E., & Marwaha, S. (2013). Predictors of employment in bipolar disorder: A systematic review. *Journal of Affective Disorders*, 145(2), 156–164. doi: 10.1016/j.jad.2012.07.009
- Gitlin, M. J., & Miklowitz, D. J. (2017). The difficult lives of individuals with bipolar disorder: A review of functional outcomes and their implications for treatment. *Journal of Affective Disorders*, 209, 147–154. doi: 10.1016/j.jad.2016.11.021
- Hibar, D. P., Westlye, L. T., Doan, N. T., Jahanshad, N., Cheung, J. W., Ching, C., ... Andreassen, O. A. (2018). Cortical abnormalities in bipolar disorder: An MRI analysis of 6503 individuals from the ENIGMA Bipolar Disorder Working Group. *Molecular Psychiatry*, 23(4), 932–942. doi: 10.1038/mp.2017.73
- Higgins, J. P., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *British Medical Journal*, 327(7414), 557–560. doi: 10.1136/bmj.327.7414.557
- *Hinrichs, K. H., Easter, R. E., Angers, K., Pester, B., Lai, Z., Marshall, D. F., ... Ryan, K. A. (2017). Influence of cognitive reserve on neuropsychological functioning in bipolar disorder: Findings from a 5-year longitudinal study. *Bipolar Disorders*, 19(1), 50–59. doi: 10.1111/bdi.12470
- Hughes, M. L., Agrigoroaei, S., Jeon, M., Bruzzese, M., & Lachman, M. E. (2018). Change in cognitive performance from midlife into old age: Findings from the Midlife in the United States (MIDUS) study. *Journal of the International Neuropsychological Society: JINS*, 24(8), 805–820. doi: 10.1017/S1355617718000425
- Jiménez-López, E., Sánchez-Morla, E. M., López-Villarreal, A., Aparicio, A. I., Martínez-Vizcaíno, V., Vieta, E., ... Santos, J. L. (2019). Neurocognition and functional outcome in patients with psychotic, non-psychotic bipolar I disorder, and schizophrenia. A five-year follow-up. *European Psychiatry*, 56, 60–68. doi: 10.1016/j.eurpsy.2018.11.008
- Kapczinski, F., Magalhães, P. V., Balanzá-Martínez, V., Dias, V. V., Frangou, S., Gama, C. S., ... Berk, M. (2014). Staging systems in bipolar disorder: An International Society for Bipolar Disorders Task Force Report. *Acta Psychiatrica Scandinavica*, 130(5), 354–363. doi: 10.1111/acps.12305
- Kjærstad, H. L., Eikeseth, F. F., Vinberg, M., Kessing, L. V., & Miskowiak, K. (2019). Neurocognitive heterogeneity in patients with bipolar disorder and their unaffected relatives: Associations with emotional cognition. *Psychological Medicine*, 51(4), 668–679. doi: 10.1017/S0033291719003738
- Kjærstad, H. L., Mistarz, N., Coello, K., Stanislaus, S., Melbye, S. A., Harmer, C. J., ... Kessing, L. V. (2020). Aberrant cognition in newly diagnosed patients with bipolar disorder and their unaffected relatives. *Psychological Medicine*, 50(11), 1808–1819. doi: 10.1017/S0033291719001867
- Koenen, K. C., Moffitt, T. E., Roberts, A. L., Martin, L. T., Kubzansky, L., Harrington, H., ... Caspi, A. (2009). Childhood IQ and adult mental disorders: A test of the cognitive reserve hypothesis. *American Journal of Psychiatry*, 166(1), 50–57. doi: 10.1176/appi.ajp.2008.08030343
- López-Jaramillo, C., Lopera-Vásquez, J., Gallo, A., Ospina-Duque, J., Bell, V., Torrent, C., ... Vieta, E. (2010). Effects of recurrence on the cognitive performance of patients with bipolar I disorder: Implications for relapse prevention and treatment adherence. *Bipolar Disorders*, 12(5), 557–567. doi: 10.1111/j.1399-5618.2010.00835.x
- MacCabe, J. H., Lambe, M. P., Cnattingius, S., Sham, P. C., David, A. S., Reichenberg, A., ... Hultman, C. M. (2010). Excellent school performance at age 16 and risk of adult bipolar disorder: National cohort study. *British Journal of Psychiatry*, 196(2), 109–115. doi: 10.1192/bjp.bp.108.060368
- Martino, D. J., Marengo, E., Igoa, A., & Strejilevich, S. A. (2018). Neurocognitive heterogeneity in older adults with bipolar disorders. *Psychiatry Research*, 262, 510–512. doi: 10.1016/j.psychres.2017.09.035
- Martino, D. J., Samamé, C., Ibañez, A., & Strejilevich, S. A. (2015). Neurocognitive functioning in the premorbid stage and in the first episode of bipolar disorder: A systematic review. *Psychiatry Research*, 226(1), 23–30. doi: 10.1016/j.psychres.2014.12.044
- Martino, D. J., Strejilevich, S. A., Marengo, E., Igoa, A., Fassi, G., Teitelbaum, J., & Caravotta, P. (2013). Relationship between neurocognitive functioning and episode recurrences in bipolar disorder. *Journal of Affective Disorders*, 147(1–3), 345–351. doi: 10.1016/j.jad.2012.11.037
- Mendez, M. F., Parand, L., & Akhlaghiour, G. (2020). Bipolar disorder among patients diagnosed with frontotemporal dementia. *The Journal of Neuropsychiatry and Clinical Neurosciences*, 32(4), 376–384. doi: 10.1176/appi.neuropsych.20010003
- Meyer, S. E., Carlson, G. A., Wiggs, E. A., Martinez, P. E., Ronsaville, D. S., Klimes-Dougan, B., ... Radke-Yarrow, M. (2004). A prospective study of the association among impaired executive functioning, childhood attentional problems, and the development of bipolar disorder. *Development and Psychopathology*, 16(2), 461–476. doi: 10.1017/s095457940404461x
- *Mora, E., Portella, M. J., Forcada, I., Vieta, E., & Mur, M. (2013). Persistence of cognitive impairment and its negative impact on psychosocial functioning in lithium-treated, euthymic bipolar patients: A 6-year follow-up study. *Psychological Medicine*, 43(6), 1187–1196. doi: 10.1017/S0033291712001948

- Mora, E., Portella, M. J., Forcada, I., Vieta, E., & Mur, M. (2016). A preliminary longitudinal study on the cognitive and functional outcome of bipolar excellent lithium responders. *Comprehensive Psychiatry*, *71*, 25–32. doi: 10.1016/j.comppsy.2016.07.008
- National Institutes of Health (2014). Quality assessment tool for observational cohort and cross-sectional studies. Available online at: <https://www.nhlbi.nih.gov/health-pro/guidelines/in-develop/cardiovascular-risk-reduction/tools/cohort> (Accessed 2 January 2021).
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., ... Moher, D. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ (Clinical Research Ed.)*, *372*, n71. doi: 10.1136/bmj.n71
- Ratheesh, A., Lin, A., Nelson, B., Wood, S. J., Brewer, W., Betts, J., ... Bechdorf, A. (2013). Neurocognitive functioning in the prodrome of mania – an exploratory study. *Journal of Affective Disorders*, *147*(1–3), 441–445. doi: 10.1016/j.jad.2012.09.017
- Reichenberg, A., Weiser, M., Rabinowitz, J., Caspi, A., Schmeidler, J., Mark, M., ... Davidson, M. (2002). A population-based cohort study of premorbid intellectual, language, and behavioral functioning in patients with schizophrenia, schizoaffective disorder, and nonpsychotic bipolar disorder. *American Journal of Psychiatry*, *159*(12), 2027–2035. doi: 10.1176/appi.ajp.159.12.2027
- Robinson, L. J., & Ferrier, I. N. (2006). Evolution of cognitive impairment in bipolar disorder: A systematic review of cross-sectional evidence. *Bipolar Disorders*, *8*(2), 103–116. doi: 10.1111/j.1399-5618.2006.00277.x
- Ryan, K. A., Assari, S., Angers, K., Marshall, D. F., Hinrichs, K., Easter, R., ... McInnis, M. G. (2017). Equivalent linear change in cognition between individuals with bipolar disorder and healthy controls over 5 years. *Bipolar Disorders*, *19*(8), 689–697. doi: 10.1111/bdi.12532
- Ryan, K. A., Assari, S., Pester, B. D., Hinrichs, K., Angers, K., Baker, A., ... Langenecker, S. A. (2016). Similar trajectory of executive functioning performance over 5 years among individuals with bipolar disorder and unaffected controls using latent growth modeling. *Journal of Affective Disorders*, *199*, 87–94. doi: 10.1016/j.jad.2016.04.016
- Samamé, C., Martino, D. J., & Strejilevich, S. A. (2013). A quantitative review of neurocognition in euthymic late-life bipolar disorder. *Bipolar Disorders*, *15*(6), 633–644. doi: 10.1111/bdi.12077
- Samamé, C., Martino, D. J., & Strejilevich, S. A. (2014). Longitudinal course of cognitive deficits in bipolar disorder: A meta-analytic study. *Journal of Affective Disorders*, *164*, 130–138. doi: 10.1016/j.jad.2014.04.028
- *Sánchez-Morla, E. M., López-Villarreal, A., Jiménez-López, E., Aparicio, A. I., Martínez-Vizcaino, V., Roberto, R. J., ... Santos, J. L. (2019). Impact of number of episodes on neurocognitive trajectory in bipolar disorder patients: A 5-year follow-up study. *Psychological Medicine*, *49*(8), 1299–1307. doi: 10.1017/S0033291718001885
- Santos, J. L., Aparicio, A., Bagney, A., Sánchez-Morla, E. M., Rodríguez-Jiménez, R., Mateo, J., & Jiménez-Arriero, MÁ (2014). A five-year follow-up study of neurocognitive functioning in bipolar disorder. *Bipolar Disorders*, *16*(7), 722–731. doi: 10.1111/bdi.12215
- *Schouws, S. N., Comijs, H. C., Dols, A., Beekman, A. T., & Stek, M. L. (2016). Five-year follow-up of cognitive impairment in older adults with bipolar disorder. *Bipolar Disorders*, *18*(2), 148–154. doi: 10.1111/bdi.12374
- Schouws, S. N., Comijs, H. C., Stek, M. L., Dekker, J., Oostervink, F., Naarding, P., ... Beekman, A. T. (2009). Cognitive impairment in early and late bipolar disorder. *The American Journal of Geriatric Psychiatry: Official Journal of the American Association for Geriatric Psychiatry*, *17*(6), 508–515. doi: 10.1097/JGP.0b013e31819e2d50
- Smith, D. J., Anderson, J., Zammit, S., Meyer, T. D., Pell, J. P., & Mackay, D. (2015). Childhood IQ and risk of bipolar disorder in adulthood: Prospective birth cohort study. *British Journal of Psychiatry*, *1(1)*, 74–80. doi: 10.1192/bjpo.bp.115.000455
- *Sparding, T., Joas, E., Clements, C., Sellgren, C. M., Pålsson, E., & Landén, M. (2021). Long-term trajectory of cognitive performance in people with bipolar disorder and controls: 6-year longitudinal study. *BJPsych Open*, *7*(4), e115. doi: 10.1192/bjo.2021.66
- Strejilevich, S. A., Quiroz, D., & Bitran, J. A. (2020). Bipolar disorder. *The New England Journal of Medicine*, *383*(14), 1398. doi: 10.1056/NEJMc202646.2
- Strejilevich, S. A., Samamé, C., & Martino, D. J. (2015). The trajectory of neuropsychological dysfunctions in bipolar disorders: A critical examination of a hypothesis. *Journal of Affective Disorders*, *175*, 396–402. doi: 10.1016/j.jad.2015.01.018
- Subramaniam, H., Dennis, M. S., & Byrne, E. J. (2007). The role of vascular risk factors in late onset bipolar disorder. *International Journal of Geriatric Psychiatry*, *22*(8), 733–737. doi: 10.1002/gps.1730
- Szmulewicz, A., Valerio, M. P., & Martino, D. J. (2020). Longitudinal analysis of cognitive performances in recent-onset and late-life bipolar disorder: A systematic review and meta-analysis. *Bipolar Disorders*, *22*(1), 28–37. doi: 10.1111/bdi.12841
- Torrent, C., Martínez-Arán, A., Daban, C., Amann, B., Balanzá-Martínez, V., del Mar Bonnín, C., ... Vieta, E. (2011). Effects of atypical antipsychotics on neurocognition in euthymic bipolar patients. *Comprehensive Psychiatry*, *52*(6), 613–622. doi: 10.1016/j.comppsy.2010.12.009
- Varo, C., Solé, B., Jiménez, E., Bonnin, C. M., Torrent, C., Valls, E., ... Reinares, M. (2020). Identifying social cognition subgroups in euthymic patients with bipolar disorder: A cluster analytical approach. *Psychological Medicine*, 1–10. Advance online publication. doi: 10.1017/S0033291720001865
- Velosa, J., Delgado, A., Finger, E., Berk, M., Kapczinski, F., & de Azevedo Cardoso, T. (2020). Risk of dementia in bipolar disorder and the interplay of lithium: A systematic review and meta-analyses. *Acta Psychiatrica Scandinavica*, *141*(6), 510–521. doi: 10.1111/acps.13153
- Vita, A., De Peri, L., Deste, G., Barlati, S., & Sacchetti, E. (2015). The effect of antipsychotic treatment on cortical gray matter changes in schizophrenia: Does the class matter? A meta-analysis and meta-regression of longitudinal magnetic resonance imaging studies. *Biological Psychiatry*, *78*(6), 403–412. doi: 10.1016/j.biopsych.2015.02.008
- Won, E., & Kim, Y. K. (2017). An oldie but goodie: Lithium in the treatment of bipolar disorder through neuroprotective and neurotrophic mechanisms. *International Journal of Molecular Sciences*, *18*(12), 2679. doi: 10.3390/ijms18122679
- Zanelli, J., Mollon, J., Sandin, S., Morgan, C., Dazzan, P., Pilecka, I., ... Reichenberg, A. (2019). Cognitive change in schizophrenia and other psychoses in the decade following the first episode. *The American Journal of Psychiatry*, *176*(10), 811–819. doi: 10.1176/appi.ajp.2019.18091088