

University of Groningen

Corrigendum

Simoes, Jorge Piano; Daoud, Elza; Shabbir, Maryam; Amanat, Sana; Assouly, Kelly; Biswas, Roshni; Casolani, Chiara; Dode, Albi; Enzler, Falco; Jacquemin, Laure

Published in:
Frontiers in Aging Neuroscience

DOI:
[10.3389/fnagi.2021.730758](https://doi.org/10.3389/fnagi.2021.730758)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Simoes, J. P., Daoud, E., Shabbir, M., Amanat, S., Assouly, K., Biswas, R., Casolani, C., Dode, A., Enzler, F., Jacquemin, L., Joergensen, M., Kok, T., Liyanage, N., Lourenco, M., Makani, P., Mehdi, M., Ramadhani, A. L., Riha, C., Santacruz, J. L., ... Genitsaridi, E. (2021). Corrigendum: Multidisciplinary Tinnitus Research: Challenges and Future Directions From the Perspective of Early Stage Researchers (Front. Aging Neurosci., (2021), 13, (647285), 10.3389/fnagi.2021.647285). *Frontiers in Aging Neuroscience*, 13, [730758]. <https://doi.org/10.3389/fnagi.2021.730758>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Corrigendum: Multidisciplinary Tinnitus Research: Challenges and Future Directions From the Perspective of Early Stage Researchers

Jorge Piano Simoes^{1*†}, Elza Daoud^{2*†}, Maryam Shabbir^{3*†}, Sana Amanat⁴, Kelly Assouly^{5,6,7}, Roshni Biswas^{3,8}, Chiara Casolani^{9,10,11}, Albi Dode¹², Falco Enzler², Laure Jacquemin^{13,14}, Mie Joergensen^{9,15}, Tori Kok¹⁶, Nuwan Liyanage^{17,18}, Matheus Lourenco^{19,20}, Punitkumar Makani^{21,22}, Muntazir Mehdi²³, Anissa L. Ramadhani^{24,25}, Constanze Riha²⁶, Jose Lopez Santacruz^{21,22}, Axel Schiller¹, Stefan Schoisswohl¹, Natalia Trpchevska²⁷ and Eleni Genitsaridi^{3,28*†}

OPEN ACCESS

Edited and reviewed by:

Daniel Ortuño-Sahagún,
University of Guadalajara, Mexico

*Correspondence:

Eleni Genitsaridi
eleni.genitsaridi@nottingham.ac.uk
Jorge Piano Simoes
jorge.simoes@ukr.de
Maryam Shabbir
maryam.shabbir@nottingham.ac.uk
Elza Daoud
elzadaoud@gmail.com

[†]These authors have contributed equally to this work and share first and senior authorship

Received: 25 June 2021

Accepted: 02 July 2021

Published: 09 August 2021

Citation:

Simoes JP, Daoud E, Shabbir M, Amanat S, Assouly K, Biswas R, Casolani C, Dode A, Enzler F, Jacquemin L, Joergensen M, Kok T, Liyanage N, Lourenco M, Makani P, Mehdi M, Ramadhani AL, Riha C, Santacruz JL, Schiller A, Schoisswohl S, Trpchevska N and Genitsaridi E (2021) Corrigendum: Multidisciplinary Tinnitus Research: Challenges and Future Directions From the Perspective of Early Stage Researchers. *Front. Aging Neurosci.* 13:730758. doi: 10.3389/fnagi.2021.730758

¹ Department of Psychiatry and Psychotherapy, University of Regensburg, Regensburg, Germany, ² Centre National de la Recherche Scientifique, Aix-Marseille University, Marseille, France, ³ Hearing Sciences, Mental Health and Clinical Neurosciences, School of Medicine, University of Nottingham, Nottingham, United Kingdom, ⁴ Otology & Neurology Group CTS 495, Department of Genomic Medicine, GENYO - Centre for Genomics and Oncological Research Pfizer/University of Granada/Junta de Andalucía, PTS, Granada, Spain, ⁵ Department of Otorhinolaryngology and Head & Neck Surgery, University Medical Center Utrecht, Utrecht, Netherlands, ⁶ Department of Clinical and Experimental Neuroscience, University Medical Center Utrecht Brain Center, Utrecht University, Utrecht, Netherlands, ⁷ Cochlear Technology Centre, Mechelen, Belgium, ⁸ Laboratory of Lifestyle Epidemiology, Department of Environmental Health Sciences, Istituto di Ricerche Farmacologiche Mario Negri IRCCS, Milan, Italy, ⁹ Hearing Systems, Department of Health Technology, Technical University of Denmark, Lyngby, Denmark, ¹⁰ Oticon A/S, Smørum, Denmark, ¹¹ Interacoustics Research Unit, Lyngby, Denmark, ¹² Institute of Databases and Information Systems, Ulm University, Ulm, Germany, ¹³ Department of Otorhinolaryngology Head and Neck Surgery, Antwerp University Hospital, Edegem, Belgium, ¹⁴ Department of Translational Neurosciences, Faculty of Medicine and Health Sciences, Antwerp University, Wilrijk, Belgium, ¹⁵ WS Audiology, Lyngby, Denmark, ¹⁶ Ear Institute, University College London, London, United Kingdom, ¹⁷ University of Zurich, Zurich, Switzerland, ¹⁸ Department of Otorhinolaryngology, Head and Neck Surgery, University Hospital Zurich, Zurich, Switzerland, ¹⁹ Experimental Health Psychology, Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, Netherlands, ²⁰ Health Psychology Research Group, Faculty of Psychology and Educational Sciences, University of Leuven, Leuven, Belgium, ²¹ Department of Otorhinolaryngology, Head and Neck Surgery, University of Groningen, University Medical Center Groningen, Groningen, Netherlands, ²² Graduate School of Medical Sciences (Research School of Behavioral and Cognitive Neurosciences), University of Groningen, Groningen, Netherlands, ²³ Institute of Distributed Systems, Ulm University, Ulm, Germany, ²⁴ Radiological Sciences, Mental Health and Clinical Neurosciences, School of Medicine, University of Nottingham, Nottingham, United Kingdom, ²⁵ Sir Peter Mansfield Imaging Centre, School of Medicine, University of Nottingham, Nottingham, United Kingdom, ²⁶ Chair of Neuropsychology, Department of Psychology, University of Zurich, Zurich, Switzerland, ²⁷ Department of Physiology and Pharmacology, Experimental Audiology Laboratory, Karolinska Institutet, Stockholm, Sweden, ²⁸ Nottingham Biomedical Research Centre, National Institute for Health Research, Nottingham, United Kingdom

Keywords: tinnitus, review, heterogeneity, standardization, interdisciplinary collaborations, big data, treatment development

A Corrigendum on

Multidisciplinary Tinnitus Research: Challenges and Future Directions From the Perspective of Early Stage Researchers

by Simoes, J. P., Daoud, E., Shabbir, M., Amanat, S., Assouly, K., Biswas, R., et al. (2021). *Front. Aging Neurosci.* 13:647285. doi: 10.3389/fnagi.2021.647285

In the original article, there was an error. For the sentence “NMDA receptor antagonists (AM-101) have been discontinued in phase III for not meeting endpoints (van de Heyning et al., 2014)” there was a typographical error (phase III should have been phase II). In addition, it was brought to our attention that clinical trials for AM-101 are ongoing.

A correction has been made to section 6. Treatment Development, Subsection 6.4. Pharmacology-Based Interventions, paragraph 1. The corrected paragraph is below.

A wide variety of therapeutic drugs have been used to relieve tinnitus (Elgoyhen and Langguth, 2010). For acute tinnitus, a dose-dependent reduction in tinnitus intensity was observed with intravenous lidocaine (Trellakis et al., 2006). However, its use is controversial due to its short-lasting response, its potentially life threatening arrhythmogenic side effects, and the low bioavailability of its oral form (Israel et al., 1982; Trellakis et al., 2007; Gil-Gouveia and Goadsby, 2009). A potential goal of pharmacologic tinnitus research could be to identify the mechanism by which lidocaine interferes with tinnitus and mimic this effect using a drug with better tolerance that can be orally administered. For chronic tinnitus, the off-label use of medicines like betahistine (Hall et al., 2018d), anticonvulsants

(Hoekstra et al., 2011), and glutamate receptor antagonists have shown little or no effect in clinical trials. Prescription of antidepressants and benzodiazepines is limited to tinnitus-associated comorbidities such as depression, insomnia and anxiety (Langguth et al., 2019). Moreover, three clinical research programs, in the last few years, were discontinued in phase II and III. AMPA antagonist selurampanel (BGG492) has not resulted in a new compound (Cederroth et al., 2018). NMDA receptor antagonists (AM-101) did not meet the primary endpoint of improving minimum masking level in acute tinnitus in a phase II clinical trial but showed improvement for tinnitus loudness, annoyance, sleep difficulties, and tinnitus impact in patients with tinnitus after noise trauma or otitis media (van de Heyning et al., 2014). Many other treatments decreasing tinnitus percept or targeting central auditory processing pathways are at a preclinical phase (Schilder et al., 2019). The modulator of voltage-gated potassium channels (Kv3.1) (AUT00063) was not effective in alleviating tinnitus symptoms (Hall et al., 2019b).

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

REFERENCES

- Cederroth, C. R., Dyhrfeld-Johnsen, J., and Langguth, B. (2018). An update: emerging drugs for tinnitus. *Expert Opin. Emerg. Drugs* 23, 251?260. doi: 10.1080/14728214.2018.1555240
- Elgoyhen, A. B., and Langguth, B. (2010). Pharmacological approaches to the treatment of tinnitus. *Drug Discov. Today* 15, 300?305. doi: 10.1016/j.drudis.2009.11.003
- Gil-Gouveia, R., and Goadsby, P. (2009). Neuropsychiatric side-effects of lidocaine: examples from the treatment of headache and a review. *Cephalgia* 29, 496?508. doi: 10.1111/j.1468-2982.2008.01800.x
- Hall, D. A., Ray, J., Watson, J., Sharman, A., Hutchison, J., Harris, P., et al. (2019b). A balanced randomised placebo controlled blinded phase iia multi-centre study to investigate the efficacy and safety of aut00063 versus placebo in subjective tinnitus: the quiet-1 trial. *Hear. Res.* 377, 153?166. doi: 10.1016/j.heares.2019.03.018
- Hall, D. A., Wegner, I., Smit, A. L., McFerran, D., and Stegeman, I. (2018d). Betahistine for tinnitus. *Cochrane Database Syst. Rev.* 2018:CD013093. doi: 10.1002/14651858.CD013093.pub2
- Hoekstra, C. E., Rynja, S. P., van Zanten, G. A., and Rovers, M. M. (2011). Anticonvulsants for tinnitus. *Cochrane Database Syst. Rev.* 7:CD007960. doi: 10.1002/14651858.CD007960.pub2
- Israel, J. M., Connelly, J. S., McTigue, S. T., Brummett, R. E., and Brown, J. (1982). Lidocaine in the treatment of tinnitus aurium: a double-blind study. *Arch. Otolaryngol.* 108, 471?473.
- Langguth, B., Elgoyhen, A. B., and Cederroth, C. R. (2019). Therapeutic approaches to the treatment of tinnitus. *Annu. Rev. Pharmacol. Toxicol.* 59, 291?313. doi: 10.1146/annurev-pharmtox-010818-021556
- Schilder, A. G., Su, M. P., Blackshaw, H., Lustig, L., Staecker, H., Lenarz, T., et al. (2019). Hearing protection, restoration, and regeneration: an overview of emerging therapeutics for inner ear and central hearing disorders. *Otol. Neurotol.* 40, 559?570. doi: 10.1097/MAO.0000000000002194
- Trellakis, S., Benzenberg, D., Urban, B. W., and Friederich, P. (2006). Differential lidocaine sensitivity of human voltage-gated potassium channels relevant to the auditory system. *Otol. Neurotol.* 27, 117?123. doi: 10.1097/01.mao.0000186443.11832.8a
- Trellakis, S., Lautermann, J., and Lehnerdt, G. (2007). Lidocaine: neurobiological targets and effects on the auditory system. *Prog. Brain Res.* 166, 303?322. doi: 10.1016/S0079-6123(07)66028-2
- van de Heyning, P., Muehlmeier, G., Cox, T., Lisowska, G., Maier, H., Morawski, K., et al. (2014). Efficacy and safety of am-101 in the treatment of acute inner ear tinnitus a double-blind, randomized, placebo-controlled phase II study. *Otol. Neurotol.* 35:589. doi: 10.1097/MAO.0000000000000268

Publisher’s Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Simoes, Daoud, Shabbir, Amanat, Assouly, Biswas, Casolani, Dode, Enzler, Jacquemin, Joergensen, Kok, Liyanage, Lourenco, Makani, Mehdi, Ramadhani, Riha, Santacruz, Schiller, Schoiswohl, Trpchevska and Genitsaridi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.