

University of Groningen

Cross-education

Farthing, Jonathan P.; Zehr, E. Paul; Hendy, Ashlee M.; Andrushko, Justin W.; Manca, Andrea; Deriu, Franca; Loenneke, Jeremy; Minetto, Marco A.; Hortobagyi, Tibor

Published in:
Brazilian Journal of Motor Behavior

DOI:
[10.20338/bjmb.v15i1.215](https://doi.org/10.20338/bjmb.v15i1.215)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Farthing, J. P., Zehr, E. P., Hendy, A. M., Andrushko, J. W., Manca, A., Deriu, F., Loenneke, J., Minetto, M. A., & Hortobagyi, T. (2021). Cross-education: Is it a viable method for rehabilitation? *Brazilian Journal of Motor Behavior*, 15(1), 1–4. <https://doi.org/10.20338/bjmb.v15i1.215>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Cross-education: Is it a viable method for rehabilitation?

JONATHAN P. FARTHING¹ | E. PAUL ZEHR² | ASHLEE M. HENDY³ | JUSTIN W. ANDRUSHKO¹ | ANDREA MANCA⁴ |
FRANCA DERIU⁴ | JEREMY LOENNEKE⁵ | MARCO A. MINETTO⁶ | TIBOR HORTOBÁGYI⁷

¹ College of Kinesiology, University of Saskatchewan, Saskatoon, Canada.

² Division of Medical Sciences; Exercise Science, Physical & Health Education; Centre for Biomedical Research, University of Victoria, Victoria, Canada.

³ Institute for Physical Activity and Nutrition (IPAN), School of Exercise and Nutrition Sciences, Deakin University, Victoria, Australia.

⁴ Department of Biomedical Sciences, University of Sassari, Sassari, Italy.

⁵ Department of Health, Exercise Science, and Recreation Management, University of Mississippi, Oxford, USA.

⁶ Division of Physical Medicine and Rehabilitation, Department of Surgical Sciences, University of Turin, Torino, Italy.

⁷ Center for Human Movement Sciences, University Medical Center Groningen, The University of Groningen, Groningen, The Netherlands.

Correspondence to: Jonathan P. Farthing, College of Kinesiology, University of Saskatchewan, Saskatoon, Canada.

email: jon.farthing@usask.ca

<https://doi.org/10.20338/bjmb.v15i1.215>

ABBREVIATIONS

MS Multiple sclerosis
RCT Randomized controlled trial

PUBLICATION DATA

Received 03 12 2020

Accepted 02 01 2021

Published 01 03 2021

In 1894, psychologists Edward W. Scripture, Theodate L. Smith, and Emily M. Brown reported for the first time the curious observation that practicing a motor skill with one hand also dramatically improved the non-practiced hand, giving rise to the phenomenon now coined as *cross-education*. Cross-education is the increase in motor output (i.e., force generation, skill) of the opposite, untrained limb following a period of unilateral motor training.¹ The potential to exploit such inter-limb adaptations for the purposes of rehabilitation of unilateral neurological or orthopedic injuries has captured the attention of scientists and therapists for years.

The magnitude of cross-education varies greatly between muscles and participants. In healthy adults, resistance training improves maximal voluntary force of the untrained limb by up to ~20%, usually half of the trained limb's improvement, but mirror training,² non-invasive brain stimulation³ and neuromuscular electrical stimulation⁴ can augment the transfer effects. Although not completely unraveled, researchers agree that the effects are likely driven by neuroplasticity in the primary and supplementary motor brain regions.¹

Until recently, it has remained unclear if cross-education could aid rehabilitation of patients after a (unilateral) fracture, surgical intervention, a stroke, or multiple sclerosis (MS). During experimental arm immobilization of healthy adults, cross-education offset declines in strength and muscle cross-sectional area.⁵ Cross-education appears to be amplified in clinical settings, with evidence for improved grip strength and range of motion after wrist fracture,⁶ wrist and ankle strength in chronic post-stroke hemiparesis,⁷ and ankle strength and mobility in persons with MS.⁸ Cross-education effects in MS were similar to direct training of the more affected side;⁸ efficacious for scenarios where the more affected limb is unable to train or becomes fatigued.

Still, caution is needed in interpreting these data because the promising clinical studies have small samples,^{6,7,8} or lack a control or 'standard of care' comparison in a randomized controlled trial (RCT) design,⁷ do not show persistent effects at six months follow-up,^{5,8} or do not consistently show benefits beyond standard care for clinical-oriented outcomes.^{6,8} The largest RCT involving cross-education as an adjuvant post knee surgery rehabilitation intervention reported no effects beyond standard care for neuromuscular or functional assessments.⁹

One hypothesis is that contraction history of the muscle influences inter-limb outcomes after unilateral training, where improvements in the untrained muscles following high-intensity strength training of the contralateral limb exceed gains from low-load direct training of a limb. Cross-education may be lessened when both limbs are trained because the local contraction history of the muscle inhibits potential signaling from the opposite limb motor pathway.¹⁰ Cross-education could be best applied clinically in isolation, and not in combination with affected limb therapy - especially if it involves low load functional movement.

Amidst the limited evidence from large RCTs, can we say that cross-education is a viable method of rehabilitation? Indeed, most clinical studies are positive, and critically, none report negative outcomes. The risks of cross-education were based on the premise that it can worsen inter-limb asymmetry and exacerbate neglect of an impaired limb. But clinical studies so far suggest upper limb asymmetry was in fact reduced by cross-education^{6,7} and lower limb asymmetry was not worsened.^{7,9} Clinical emphasis on avoiding asymmetry may be short-sighted because it diverts focus from the absolute functional capacity of the impaired limb. If clinical function of the impaired limb is improved, and neuromuscular activation yields movements that before were not possible,⁷ the risk of enhancing asymmetry should be a secondary concern¹ that can be addressed by direct exercise once functional ability is restored. Caution is important where we lack clinical data on cross-education effects (e.g., acute and sub-acute stroke). Training the unaffected limb is somewhat incompatible with the widely used constraint-induced movement therapy, and the techniques are yet to be studied in conjunction. Perhaps the two are antithetical and target different patients. The evidence suggests cross-education is most usefully applied when the impaired or injured limb is very weak, immobilized, or unable to function unaided, whereas ideal candidates to constraint-induced movement therapy must have a minimum functional reservoir in order to benefit. Future studies need to assess whether adding cross-education prior to direct training is functionally and clinically relevant.

We suggest that contralateral training stimulates neuroplasticity in motor pathways of the injured or impaired side, serving to offset weakness and wasting and stimulate recovery after orthopedic injury or neurological impairment, or diminish effects of neurological disease. The possible implications of cross-education for management of contralateral musculoskeletal pain seem promising but require further investigation. Importantly, we lack evidence from large RCTs that explore the combination of cross-education with mirror training, electrical stimulation, and non-invasive brain stimulation. We endorse collaborative efforts for clinical studies to explore these novel avenues as we deepen our understanding of benefits and limitations of cross-education as a rehabilitation method.

REFERENCES

1. Manca A, Hortobágyi T, Carroll TJ et al. Contralateral Effects of Unilateral Strength and Skill Training: Modified Delphi Consensus to Establish Key Aspects of Cross-Education. *Sports Med.* 2021;51:11–20. 10.1007/s40279-020-01377-7.
2. Zult T, Goodall S, Thomas K, Solnik S, Hortobágyi T, Howatson G. Mirror Training Augments the Cross-education of Strength and Affects Inhibitory Paths. *Med Sci Sports Exerc.* 2016;48(6):1001-13. 10.1249/MSS.0000000000000871.
3. Hendy AM, Teo WP, Kidgell DJ. Anodal Transcranial Direct Current Stimulation Prolongs the Cross-education of Strength and Corticomotor Plasticity. *Med Sci Sports Exerc.* 2015;47(9):1788-97. 10.1249/MSS.0000000000000600..
4. Minetto MA, Botter A, Gambero G et al. Contralateral effect of short-duration unilateral neuromuscular electrical stimulation and focal vibration in healthy subjects. *Eur J Phys Rehabil Med.* 2018;54(6):911-920. 10.23736/S1973-9087.18.05004-9.
5. Andrushko JW, Lanovaz JL, Björkman KM, Kontulainen SA, Farthing JP. Unilateral strength training leads to muscle-specific sparing effects during opposite homologous limb immobilization. *J Appl Physiol.* 2018;124(4):866-876. 10.1152/jappphysiol.00971.2017.
6. Magnus CR, Arnold CM, Johnston G et al. Cross-education for improving strength and mobility after distal radius fractures: a randomized controlled trial. *Arch Phys Med Rehabil.* 2013;94(7):1247-55. 10.1016/j.apmr.2013.03.005.
7. Sun Y, Zehr EP. Training-Induced Neural Plasticity and Strength Are Amplified After Stroke. *Exerc Sport Sci Rev.* 2019;47(4):223-229. 10.1249/JES.000000000000199.
8. Manca A, Cabboi MP, Dragone D et al. Resistance Training for Muscle Weakness in Multiple Sclerosis: Direct Versus Contralateral Approach in Individuals With Ankle Dorsiflexors' Disparity in Strength. *Arch Phys Med Rehabil.* 2017;98(7):1348-1356.e1. 10.1016/j.apmr.2017.02.019.
9. Zult T, Gokeler A, van Raay JJAM et al. Cross-education does not accelerate the rehabilitation of neuromuscular functions after ACL reconstruction: a randomized controlled clinical trial. *Eur J Appl Physiol.* 2018;118(8):1609-1623. 10.1007/s00421-018-3892-1.
10. Bell ZW, Wong V, Spitz RW et al. The contraction history of the muscle and strength change: lessons learned from unilateral training models. *Physiol Meas.* 2020;41(1):01TR01. 10.1088/1361-6579/ab516c.

Citation: Farthing JP, Zehr EP, Hendy AM, Andrushko JW, Manca A, Deriu F, Loenneke J, Minetto MA, Hortobágyi T. Cross-education: Is it a viable method for rehabilitation?. BJMB. 2021;15(1):1-4.

Editors: Dr Fabio Augusto Barbieri - São Paulo State University (UNESP), Bauru, SP, Brazil; Dr José Angelo Barela - São Paulo State University (UNESP), Rio Claro, SP, Brazil; Dr Natalia Madalena Rinaldi - Federal University of Espírito Santo (UFES), Vitória, ES, Brazil.

Section Editors (Current Opinion): Dr Luis Augusto Teixeira - University of São Paulo (USP), São Paulo, SP, Brazil; Dr Tibor Hortobágyi - University of Groningen, The Netherlands; Dr Renato de Moraes - University of São Paulo (USP), Ribeirão Preto, SP, Brazil.

Copyright: © 2021 Farthing, Zehr, Hendy, Andrushko, Manca, Deriu, Loenneke, Minetto and Hortobágyi and BJMB. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International License which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: There was no funding for this study.

Competing interests: The authors have declared that no competing interests exist.

DOI: <https://doi.org/10.20338/bjmb.v15i1.215>