

University of Groningen

Role of transmembrane pH gradient and membrane binding in nisin pore formation

Moll, Gert N.; Clark, Jonathan; Chan, Weng C.; Bycroft, Barrie W.; Roberts, Gordon C.K.; Konings, Wil N.; Driessen, Arnold J.M.

Published in:
Journal of Bacteriology

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1997

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Moll, G. N., Clark, J., Chan, W. C., Bycroft, B. W., Roberts, G. C. K., Konings, W. N., & Driessen, A. J. M. (1997). Role of transmembrane pH gradient and membrane binding in nisin pore formation. *Journal of Bacteriology*, 179(1), 135-140.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Role of Transmembrane pH Gradient and Membrane Binding in Nisin Pore Formation

GERT N. MOLL,¹ JONATHAN CLARK,² WENG C. CHAN,² BARRIE W. BYCROFT,²
GORDON C. K. ROBERTS,³ WIL N. KONINGS,¹ AND ARNOLD J. M. DRIESSEN^{1*}

Department of Microbiology, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, 9751 NN Haren, The Netherlands,¹ and Department of Pharmaceutical Sciences, University of Nottingham, Nottingham NG7 2RD,² and Department of Biochemistry and Biological NMR Centre, University of Leicester, Leicester LE1 9HN,³ United Kingdom

Received 18 July 1996/Accepted 21 October 1996

Nisin is a cationic antimicrobial peptide that belongs to the group of lantibiotics. It is thought to form oligomeric pores in the target membrane by a mechanism that requires the transmembrane electrical potential ($\Delta\psi$) and that involves local perturbation of the lipid bilayer structure. Here we show that nisin does not form exclusively voltage-dependent pores: even in the absence of a $\Delta\psi$, nisin is able to dissipate the transmembrane pH gradient (ΔpH) in sensitive *Lactococcus lactis* cells and proteoliposomes. The rate of dissipation increases with the magnitude of the ΔpH . Nisin forms pores only when the ΔpH is inside alkaline. The efficiency of $\Delta\psi$ -induced pore formation is strongly affected by the external pH, whereas ΔpH -induced pore formation is rather insensitive to the external pH. Nisin¹⁻¹², an amino-terminal fragment of nisin, and (*des*- ΔAla_5)-nisin¹⁻³² amide have a strongly reduced capacity to dissipate the $\Delta\psi$ and ΔpH in cytochrome *c* oxidase proteoliposomes and *L. lactis* cells. Both variants bind with reduced efficiency to liposomes containing negatively charged phospholipids, suggesting that both ring A and rings C to E play a role in membrane binding. Nisin¹⁻¹² competes with nisin for membrane binding and antagonizes pore formation. These findings are consistent with the wedge model of nisin-induced pore formation (A. J. M. Driessen et al., *Biochemistry* 34:1606–1614, 1995).

Lantibiotics are posttranslationally modified peptide antibiotics that owe their name to the presence of cyclic structures formed by lanthionines or 3-methylanthionine residues. The lantibiotic nisin is produced by *Lactococcus lactis* subsp. *lactis* and has five such intramolecular rings, termed A, B, C, D, and E, in a total of 34 amino acids. Because of its bactericidal activity against a broad range of gram-positive bacteria, nisin is exploited as a food preservative. Its genetics (23) as well as the regulation of its synthesis (22) are known, and several mutants have been generated (24–26).

Nisin forms nonselective, transient, multistate pores in membranes (38–40). In addition to having bactericidal activity, it inhibits the outgrowth of spores (11, 18, 29, 33) and interferes with the activity of biosynthetic (36) and autolytic (2–5) enzymes. Nisin is able to form pores in cells, proteoliposomes, liposomes, and black lipid membranes. It has been reported that the pores are formed only when the transmembrane electrical potential ($\Delta\psi$), negative inside, is sufficiently high (15, 38, 39). In cells, pore formation induces the release of ions, amino acids, and ATP and causes the collapse of the proton motive force (Δp) (7, 16, 21, 35, 37, 39, 40). Nisin requires anionic phospholipids for membrane binding and pore formation (12, 15), while in the absence of anionic phospholipids, it acts as an anion carrier (15, 17).

The structure of nisin in solution (28, 44) and in the bound state to micelles of dodecylphosphocholine and of sodium dodecyl sulfate (42, 43) has been studied by nuclear magnetic resonance (NMR) analysis. Nisin is a flexible molecule in solution but contains two relatively structured domains: an amino-terminal domain (residues 3 to 19) containing rings A, B,

and C, in which rings A and B are particularly well structured; and a carboxyl-terminal domain (residues 22 to 28) containing rings D and E. Both domains are amphipathic, a property consistent with the ability of nisin to interact with phospholipid membranes. Based on structural and functional studies, we have proposed a wedge model for pore formation by nisin (15, 32a). A wedge-like pore composed of multiple nisin molecules may be formed once the $\Delta\psi$ drives the membrane insertion of the phospholipid surface-bound molecules. In this model, the structure of nisin changes only in the putative hinge region between rings C and D/E, while the orientation of the molecule relative to the lipid headgroups remains unchanged. The carboxyl-terminal residues, which have a net positive charge, together with the bound lipids, are drawn across the membrane in response to the $\Delta\psi$, causing the formation of a structural defect in the phospholipid bilayer.

The aim of this work was to obtain more insight into the mechanism of pore formation and to study the role of the structural domains of nisin in this process. For this purpose, we studied the role of the transmembrane pH gradient (ΔpH) in pore formation and the characteristics of two nisin variants: nisin¹⁻¹², a fragment of nisin containing only rings A and B (10), and (*des*- ΔAla_5)-nisin¹⁻³²amide (8), a form of nisin in which ring A is opened. The results demonstrate that nisin is not a voltage-dependent bacteriocin per se, but that it can be activated by the ΔpH as well. A possible mechanism of the ΔpH -induced pore formation is discussed.

MATERIALS AND METHODS

Materials. Nisin was obtained as a gift of Aplin & Barrett. Dioleoylphosphatidylglycerol (DOPG), egg phosphatidylcholine (PC), *N*-(7-nitro-2, 1,3-benzoxadiazol-4-yl) (NBD)-labelled phosphatidylethanolamine (PE), and PE from *Escherichia coli* were purchased from Avanti Polar Lipids Inc. (Birmingham, Ala.). 9-Amino-6-chloro-2-methoxyacridine (ACMA) was obtained from Molecular Probes. [¹⁴C]benzoic acid (22 mCi/mmol) was purchased from NEN. (*des*- ΔAla_5)-nisin¹⁻³²amide and nisin¹⁻¹² were obtained as described previously (8,

* Corresponding author. Mailing address: Department of Microbiology, University of Groningen, Kerklaan 30, 9751 NN Haren, The Netherlands. Phone: (31) (50) 3632164. Fax: (31) (50) 3632154. E-mail: A.J.M.DRIESSEN@BIOL.RUG.NL.




FIG. 1. (A) Nisin dissipates the ΔpH in sensitive *L. lactis* cells. Logarithmically grown cells loaded with BCECF (17 μg of protein/ml) in 50 mM KPi (pH 7.0) were energized with 0.5% glucose (arrow 1). Valinomycin (0.25 μM ; arrow 2) collapsed the total membrane potential and increased the pH gradient. Subsequently, either nisin (42 nM) (a), or (*des*- ΔAla_5)-nisin¹⁻³²amide (42 nM) (b), or nisin¹⁻¹² (42 nM) (c) was added (arrow 3). Nigericin (0.25 μM ; arrow 4) dissipated the remaining pH gradient. (B) pH dependence of the ΔpH -induced pore formation. Logarithmically grown cells (2.1 mg of protein/ml) in 50 mM KPi , at pH 7.0 (\circ), pH 6.5 (\square), or pH 6.0 (\blacktriangle) were energized with 0.5% glucose and incubated with radiolabelled benzoic acid. After 4 min, 0.25 μM valinomycin was added; 1 min later, nisin was added. Cells were incubated for 15 min and separated from the medium by centrifugation through silicon oil (6, 31). (C) pH dependence of the $\Delta\psi$ -induced pore formation. Logarithmically grown cells (2.1 mg of protein/ml) in 50 mM KPi , at pH 7.0 (\circ), pH 6.5 (\square), or pH 6.0 (\blacktriangle) were energized with 0.5% glucose and supplemented with nigericin (0.25 μM). The $\Delta\psi$ was measured by using a TPP^+ electrode as described in Materials and Methods.

10). *L. lactis* IL 1403 was used as a sensitive strain and grown as described previously (32).

Preparation of liposomes. Liposomes were made by injection of phospholipid-containing ethanol into buffer (1).

Binding of peptides to liposomes. Binding of nisin and the variants was measured by a fluorimetric assay that is based on the increase in the fluorescence quantum yield of NBD when the molecules bind to the liposomes containing 2 mol% NBD-PE (15). Experiments were performed in 50 mM KPi (pH 7.0) at 30°C. Fluorescence was measured with a Perkin-Elmer LS-50 spectrofluorimeter, using excitation and emission wavelengths of 460 and 536 nm, respectively. The slit widths were set to 4 nm.

Growth inhibition assay. Half-maximal inhibitory concentrations were determined as previously described (34), except that Tween 80 was excluded, using a microtiter assay system. Experiments were started at a cell density of 20 μg of cellular protein/ml.

Cytochrome *c* oxidase vesicles. Beef heart cytochrome *c* oxidase was reconstituted into liposomes composed of ether-acetone-washed *E. coli* lipids as described previously (14). After dialysis, proteoliposomes were subjected to five subsequent freeze-thaw cycles, by freezing them in liquid nitrogen and thawing them slowly. Thereafter they were extruded 11 times through 400-nm and subsequently 11 times through 200-nm-pore-size polycarbonate filters. Proteoliposomes (0.4 mg of lipid/ml) were energized with the electron donor cytochrome *c* (10 μM)-*N,N,N',N'*-tetramethylphenylenediamine (TMPD) (10 μM)-ascorbate (10 mM). The K^+/H^+ antiporter nigericin (0.25 μM) prevented the generation of a ΔpH , or alternatively, the K^+ carrier valinomycin (0.25 μM) prevented the generation of a $\Delta\psi$.

Bacteriorhodopsin vesicles and measurement of a ΔpH , inside acid. Bacteriorhodopsin was reconstituted into lipid vesicles (*E. coli* lipids/egg PC lipids, 3:1) as described previously (14) in 50 mM KPi . Bacteriorhodopsin-mediated proton pumping was initiated by illumination with white light (Xenon lamp, 450 W), filtered through a 495-nm-pore-size high-pass filter. Changes in the ΔpH were measured by recording the fluorescence of ACMA at 2.2 μM , using excitation and emission wavelengths of 409 and 474 nm, respectively.

Measurements of proton motive force. The $\Delta\psi$, inside negative, was calculated from the distribution of tetraphenylphosphonium ion (TPP^+) as measured with a TPP^+ -selective electrode (20). Cells were used at a concentration of 2.1 mg of protein/ml. The magnitude of the $\Delta\psi$ was calculated according to the Nernst equation. A correction for TPP^+ binding to the cells and proteoliposomes according to reference 30 was made.

The internal pH of cytochrome *c* oxidase vesicles was determined from the fluorescence of pyranine (excitation and emission wavelengths of 450 and 508 nm, respectively) as described previously (16). Entrapment was achieved by adding 1 mM pyranine to the vesicles before the sonication step. External

pyranine was removed by passing the vesicles through a Sephadex G-50-medium column (1 by 20 cm).

The internal pH of cells was measured with the fluorescent pH indicator 2',7'-bis-(2-carboxyethyl)-5-(and-6)-carboxyfluorescein (BCECF) as described previously (31). In short, two suspensions of 20 μl of 50 mM KPi (pH 7.0) containing washed cells (100 μg of protein) on ice were supplemented with 0.5 mM BCECF. To one suspension 2.0 μl , and to the other 2.5 μl , of 0.5 M HCl was added. Suspensions were kept for exactly 5 min at room temperature. External BCECF was removed by five washes in 1 ml of 50 mM KPi (pH 7.0). Cells loaded under the lowest HCl concentration were used for experiments. Experiments were performed in 50 mM KPi , at 30°C. Alternatively, the ΔpH was measured by monitoring the distribution of radiolabelled benzoic acid by the silicon oil centrifugation technique (6, 19).

RESULTS

Dissipation of the ΔpH and $\Delta\psi$ by nisin in intact cells.

Previously it has been shown that in the absence of a $\Delta\psi$, nisin is able to dissipate the ΔpH in cytochrome *c* oxidase vesicles composed of *E. coli* lipids and egg PC (9:1) (16). To determine if this activity of nisin is also relevant for intact cells of *L. lactis*, ΔpH measurements were performed with cells loaded with the fluorescent pH indicator BCECF (31). The presence of valinomycin and potassium ions prevented the generation of a $\Delta\psi$, as evident from measurements of the distribution of TPP^+ (data not shown). At a concentration of 42 nM, nisin completely collapsed the ΔpH in *L. lactis* cells (Fig. 1A, a). Similar results were obtained when the ΔpH measurements were performed by monitoring the distribution of radiolabelled benzoic acid (Fig. 1B). This concentration is about 250-fold lower than that needed to dissipate the ΔpH in proteoliposomes (see Fig. 3 and reference 16). These data demonstrate that nisin is not a voltage-dependent bacteriocin per se, but that a ΔpH suffices as well.

The efficiency of nisin-induced ΔpH and $\Delta\psi$ dissipation was studied as a function of the external pH in cells energized with glucose. In the presence of valinomycin, the efficiency of ΔpH




FIG. 2. Δ pH stimulates nisin action. To logarithmically grown cells loaded with BCECF (17 μ g of protein/ml), valinomycin (arrow 1), nisin (arrow 2), and glucose (0.5%; arrow 3) were added. Nisin was used at concentrations of 0 (a), 42 (b), 100 (c), and 400 (d) nM. Finally, the Δ pH was dissipated by the addition of nigericin (arrow 4).

dissipation by nisin is hardly affected by the external pH within the range pH 6.0 to 7.0 (Fig. 1B). In contrast, $\Delta\psi$ measurements in the presence of nigericin show a striking pH dependence of the nisin-induced $\Delta\psi$ dissipation (Fig. 1C). At a low pH, nisin is much less effective than at the higher pHs. Comparison of the dose-response curves shows that at pH 7.0, nisin is equally effective in dissipating the Δ pH and $\Delta\psi$. However, at pH 6.0, the Δ pH is more effectively dissipated than the $\Delta\psi$, implying an important role of Δ pH in nisin pore formation.

When nisin was added to the cells prior to energization, 400 nM nisin (Fig. 2, d) was needed to prevent the generation of a Δ pH upon the addition of glucose. This concentration is nearly 10-fold higher than needed to dissipate a preexisting Δ pH, i.e., when nisin is added after energization with glucose (Fig. 1A, a). Moreover, especially at the lower concentrations of nisin, the generation of a Δ pH appeared to be transient (Fig. 2, b and c), suggesting that in the absence of a $\Delta\psi$, a Δ pH is required for pore formation.

Dissipation of the Δ pH in proteoliposomes. It was previously shown that nisin dissipates only a $\Delta\psi$, inside negative (15, 38, 39). To establish whether a similar polarity effect exists with the Δ pH, nisin was added to proteoliposomes containing cytochrome *c* oxidase or bacteriorhodopsin. In the presence of reduced cytochrome *c*, cytochrome *c* oxidase liposomes gener-




FIG. 3. The outside-acid Δ pH induces nisin pores. Nisin was added to bacteriorhodopsin (BR) vesicles (●; inside-acid Δ pH) or cytochrome *c* oxidase vesicles (■; outside-acid Δ pH). BR vesicles (0.4 mg of lipid/ml, 1.7 nmol of BR/ml) were incubated in 50 mM KP_i (pH 7.0) in the presence 50 nM valinomycin, 2.5 mM $MgCl_2$, and 2.2 μ M ACMA and illuminated with a Xenon lamp (450 W) via fiber optics, using a 495-nm-pore-size high-pass filter. The inside-acid Δ pH was estimated from the ACMA fluorescence quenching. Cytochrome *c* oxidase vesicles (0.4 mg of lipid/ml) were loaded with 1 mM pyranine and incubated in 50 mM KP_i (pH 7.0) in the presence of valinomycin (0.25 μ M), ascorbate (10 mM), TMPD (10 μ M), and cytochrome *c* (10 μ M). The alkaline Δ pH was estimated from the pH changes recorded by pyranine.

ate a Δ pH, inside alkaline (14). On the other hand, upon illumination, bacteriorhodopsin liposomes generate a Δ pH, inside acid (14). Valinomycin was added to prevent the formation of a $\Delta\psi$. Nisin was much more effective in dissipating a Δ pH, inside alkaline, than a Δ pH, inside acid (Fig. 3). Only at very high concentrations (around 46 μ M) of nisin, dissipation of the Δ pH, inside acid, occurred (data not shown). Addition of 0.2 μ M nigericin caused in both cases complete dissipation of the Δ pH (data not shown). These data demonstrate that nisin dissipates the Δ pH only in an oriented manner.

Dissipation of the Δ pH and $\Delta\psi$ by nisin derivatives. Nisin¹⁻¹² is an amino-terminal fragment derived from nisin upon chymotrypsin cleavage (10), and (*des*- Δ Ala₅)-nisin¹⁻³²amide is a form of nisin with an open ring A resulting from the hydrolysis of the dehydroalanine at position 5 (10). Both derivatives exhibit a strongly reduced bactericidal activity with intact cells (8-10) (Table 1). To establish if the reduced bactericidal activity of nisin¹⁻¹² and (*des*- Δ Ala₅)-nisin¹⁻³²amide correlates with their pore-forming capacity, their effects on the Δ p in cells and cytochrome *c* oxidase vesicles were determined. Both de-

TABLE 1. Activity and membrane binding of nisin and variants^a

Lantibiotic	IC ₅₀ (μ M)		Membrane binding (%)
	Viability	$\Delta\psi$	
Nisin	0.0086 \pm 0.0016	0.94 \pm 0.06	100.0 \pm 1.7
(<i>des</i> - Δ Ala ₅)-nisin ¹⁻³² amide	>30.0	>30.0	74.0 \pm 1.9
Nisin ¹⁻¹²	15.3 \pm 1.4	>18.0	17.0 \pm 0.8

^a IC₅₀ represents the concentration of nisin or derivative needed for half-maximal killing of the indicator strain or dissipation of the $\Delta\psi$. Binding experiments were performed at 10.0 μ M nisin or nisin variant with DOPG vesicles (220 μ M lipid) containing 2 mol% NBD-PE. The relative levels of binding were estimated from the NBD fluorescence increase, with the value for nisin set to 100%.




FIG. 4. Nisin¹⁻¹² antagonizes nisin-induced $\Delta\psi$ dissipation in cytochrome *c* oxidase vesicles. The effects of nisin¹⁻¹² in the absence (■) and presence (●) of nisin (42 nM) on the initial rate of $\Delta\psi$ dissipation and the steady-state level of $\Delta\psi$ (inset) were determined. Experiments were performed in 50 mM KP₁ (pH 7.0).




FIG. 5. Competition of nisin¹⁻¹² with nisin for binding to liposomes. Nisin¹⁻¹² at the concentrations indicated, without (■) or with (●) 817 nM nisin, was added to DOPG liposomes (12.6 μ M) containing 4 mol% NBD-PE, and the increase in NBD fluorescence was recorded. a.u., arbitrary units.

derivatives showed a dramatically reduced ability to dissipate the Δ pH (Fig. 1A) and $\Delta\psi$ (Table 1) in cells. In cytochrome *c* oxidase vesicles, both nisin¹⁻¹² and (*des*- Δ Ala₅)-nisin¹⁻³²amide were nearly inactive in eliminating the Δ pH and $\Delta\psi$ (data not shown). On the other hand, increasing concentrations of nisin¹⁻¹² reduce the rate and extent of the nisin-induced $\Delta\psi$ dissipation in cytochrome *c* oxidase vesicles (Fig. 4). This antagonistic effect of nisin¹⁻¹² on the nisin activity was stronger on the initial rate of $\Delta\psi$ dissipation (Fig. 4) than on the steady-state level of the $\Delta\psi$ (Fig. 4, inset). These data suggest that nisin¹⁻¹² interferes with the formation of a functional nisin pore.

Membrane binding of nisin derivatives. The antagonistic effect of nisin¹⁻¹² on the action of nisin could be due to competition for membrane binding sites, to the formation of hetero-oligomeric pore complexes, or both. The ability of the nisin variants to interact with the membrane was analyzed by the use of phosphatidylglycerol liposomes containing 2 mol% NBD-PE (15). Binding of nisin to these membranes results in an increase in the fluorescence quantum yield of NBD, and this effect has been attributed to shielding of NBD from the aqueous environment (15). Nisin¹⁻¹² but more notably (*des*- Δ Ala₅)-nisin¹⁻³²amide retained the ability to bind to phosphatidylglycerol liposomes, but the binding activity was lower than that of nisin (Table 1). The membrane-bound nisin¹⁻¹² rapidly dissociates from the liposomes, as the addition of a 30-fold excess of phosphatidylglycerol liposomes without NBD-PE resulted in a complete loss of NBD fluorescence at a rate of about 50%/s, while only a very slow release was observed with nisin (0.8%/s). These results demonstrate that truncation of the carboxyl-terminal domain of nisin severely interferes with binding to phosphatidylglycerol.

A binding experiment was performed with increasing concentrations of nisin¹⁻¹² followed by addition of a fixed but a suboptimal amount of nisin (Fig. 5). The addition of increasing amounts of nisin¹⁻¹² results in a gradual increase in NBD fluorescence, but identical final fluorescence levels were ob-

tained after the subsequent addition of a fixed amount of nisin. These results demonstrate that nisin and nisin¹⁻¹² compete for the same binding sites, which can, at least in part, explain the antagonism of nisin¹⁻¹² on nisin action.

DISCUSSION

The present results show that even in the absence of a $\Delta\psi$, nisin is able to effectively dissipate the Δ pH in *L. lactis* cells. This finding demonstrates that, in contrast to previous suggestions, nisin does not obligatorily require a $\Delta\psi$, but that this function can also be fulfilled by the Δ pH. The characterization of nisin pores as voltage-gated channels is therefore not correct. The efficiency with which nisin dissipates the $\Delta\psi$ decreases markedly when the pH is lowered from pH 7.0 to 6.0. In contrast, the efficiency by which nisin collapses the Δ pH is only marginally lower at pH 6.0 than at pH 7.0.

At this stage, it is unclear why nisin becomes less effective in dissipating the $\Delta\psi$ when the pH is lowered. Proton conductance may involve protonable side chains in the nisin molecules. For instance, the histidine residues (His27 and His31) at the carboxyl-terminal part of nisin may play an essential role in the Δ pH-dissipating activity. In aqueous solution, both histidines have a pK_a of about 6.5 (41), and thus a significant fraction will be in the protonated state around neutral pH. When bound to micelles of sodium dodecyl sulfate, the pK_a increases to 7.5 (43); this phenomenon has been explained by the electrostatic effects of the adjacent negatively charged sulfate groups, implying that the histidine side chains reside at the micellar surface. A similar phenomenon may occur when nisin binds to liposomes containing acidic phospholipids. Under the influence of a pH gradient, the C-terminal part of nisin, which contains the protonated histidines, may insert into the membrane together with bound phospholipids, forming a wedged pore (15). The C-terminal domain harbors most of the cationic amino acid side chains that have been implicated in the $\Delta\psi$ -dependent formation of a nisin pore. It is important to note that nisin is highly active only against a Δ pH, inside alkaline, not a Δ pH, inside acid. This mechanism is consistent with the




FIG. 6. Wedge model of nisin pore formation. Nisin binds to the anionic membrane surface, leading to a high local concentration and disturbance of the lipid dynamics. Both components of the proton motive force are by themselves sufficient for inducing insertion of nisin into the membrane. Coinsertion of bound, anionic phospholipids results in bending of the lipid surface, giving rise to a wedge-like pore.

wedge model, which assumes an oriented interaction of nisin with the membrane relative to $\Delta\psi$.

The naturally occurring variant of nisin A, nisin Z, in which His27 has been replaced by Asp27 (24), has full bactericidal activity, as do the H27K (13) mutant of nisin A and H31K (26) mutant of nisin Z, showing that the histidine residues are apparently not essential for activity. Similarly, the fragment nisin¹⁻²⁹ shows only a 16-fold decrease in activity (10), and even nisin¹⁻²⁰, lacking the cationic C-terminal domain of the molecule, has detectable though low (~1%) activity (10). However, it is not clear if these variants are still able to form pores in the presence of only a ΔpH .

According to the wedge model of pore formation (Fig. 6), a hinge region may act as a twist that allows nisin to bend the lipid surface (15). Our model speculates that the ABC rings might bind to the membrane and remain surface bound, while the Δp drives the membrane insertion of the carboxyl-terminal domain containing the DE rings together with the bound lipids. The detectable activity of nisin¹⁻²⁰ and the importance of an intact ring C (10) suggest that ring C may also partly insert into the pore. NMR studies of nisin in solution (41, 44) indicate that the flexibility of the hinge between rings B and C is nearly as high as that of the hinge between rings C and D.

The wedge model predicts that contributions to membrane binding will not be confined to specific subdomains but will be found all along the nisin molecule. In line with this assumption is the observation that nisin¹⁻¹² exhibits a strongly reduced membrane binding activity, while the opening of ring A in (*des*- ΔAla_5)-nisin¹⁻³²amide has a smaller effect on membrane binding. NMR studies on micelle-bound nisin are also consistent with the idea that several parts along the nisin molecule contribute to membrane binding, as both dehydroalanines (residues 5 and 33) as well as residues Ile30 to Lys34 are shielded to some extent from the solvent. Spin label experiments indicate that the interactions of residues Ser29 to Lys34 with the micelles are the strongest (43). Absence of the latter interactions in the case of nisin¹⁻¹² and absence of the interaction of residues 33 and 34 in the case of (*des*- ΔAla_5)-nisin¹⁻

32amide might also contribute to the observed reduced membrane binding of both variants.

Both nisin¹⁻¹² and (*des*- ΔAla_5)-nisin¹⁻³²amide are strongly defective in $\Delta\psi$ and ΔpH dissipation in intact cells and cytochrome *c* oxidase proteoliposomes. This finding parallels the observations that these variants are devoid of bactericidal activity. It should be stressed that the loss of activity of (*des*- ΔAla_5)-nisin¹⁻³² must be the consequence of the opening of the A ring, since nisin¹⁻³²amide, which also lacks the two carboxyl-terminal residues, and the ΔAla_5 -Ala mutant have full bactericidal activity (8, 11). The largest conformational differences between soluble nisin and sodium dodecyl sulfate micelle-bound nisin are seen in ring A around Dha5 (41, 43). Studies with a Dha5Dhb mutant (25) and with [α -OH-Ala⁵]nisin further confirm the importance of the A ring. Both the bactericidal activity and the extent of the structural change of nisin upon micelle binding decrease in the following order: nisin A > nisinDha5Dhb > [α -OH-Ala⁵]nisin (42). This finding suggests that the micelle-bound conformation of ring A might indeed mimic that of active membrane-bound nisin and confirms the importance of the conformation of this ring for activity.

In summary, these studies have established that nisin is not a $\Delta\psi$ -dependent pore per se, but that pore formation can also be affected by the ΔpH . At an acidic pH, nisin dissipates the ΔpH more effectively than the $\Delta\psi$, indicating that under the conditions with which most bacterial fermentations in food manufacturing are performed, the ΔpH is the primary target for nisin action.

ACKNOWLEDGMENT

This work was supported by the European Community (Biotechnology Program contract BIOT-CT94-3055).

REFERENCES

- Batzri, S., and E. D. Korn. 1973. Single bilayer liposomes prepared without sonication. *Biochim. Biophys. Acta* **298**:1015-1019.
- Bierman, G., and H.-G. Sahl. 1985. Induction of autolysis of *Staphylococci* by the basic peptide antibiotics Pep5 and nisin and their influence on the activity of auto-lytic enzymes. *Arch. Microbiol.* **141**:249-254.
- Bierman, G., and H.-G. Sahl. 1987. Autolytic system of *Staphylococcus simulans* 22: influence of cationic peptides on activity of *N*-acetyl-muramoyl-L-alanine amidase. *J. Bacteriol.* **169**:5452-5458.
- Bierman, G., and H.-G. Sahl. 1988. Influence of cationic peptides on the activity of the autolytic endo- β -N-acetylglucosamidase of *Staphylococcus simulans* 22. *FEMS Microbiol. Lett.* **58**:223-228.
- Bierman, G., and H.-G. Sahl. 1991. Induction of autolysis of *Staphylococcus simulans* 22 by Pep5 and nisin and influence of the cationic peptides on the activity of the autolytic enzymes, p. 386-396. *In* G. Jung and H.-G. Sahl (ed.), *Nisin and novel lantibiotics*. ESCOM, Leiden, The Netherlands.
- Booth, J. R. 1985. Regulation of cytoplasmic pH in bacteria. *Microbiol. Rev.* **49**:359-378.
- Bruno, M. E. C., A. Kaiser, and T. J. Montville. 1992. Depletion of proton motive force by nisin in *Listeria monocytogenes* cells. *Appl. Environ. Microbiol.* **58**:2255-2259.
- Chan, W. C., B. W. Bycroft, L.-Y. Lian, and G. C. K. Roberts. 1989. Isolation and characterisation of two degradation products derived from the peptide antibiotic nisin. *FEBS Lett.* **252**:29-36.
- Chan, W. C., G. C. K. Roberts, and B. W. Bycroft. 1993. Structure-activity relationships of nisin and subtilin, p. 770-771. *In* C. H. Schneider and A. N. Eberle (ed.), *Peptides 1992*. ESCOM, Leiden, The Netherlands.
- Chan, W. C., M. Leyland, J. Clark, H. M. Dodd, L.-Y. Lian, M. J. Gasson, B. W. Bycroft, and G. C. K. Roberts. 1996. Structure-activity relationships in the peptide antibiotic nisin: antibacterial activity of fragments of nisin. *FEBS Lett.* **390**:129-132.
- Chan, W. C., H. M. Dodd, N. Horn, K. Maclean, L.-Y. Lian, B. W. Bycroft, M. J. Gasson, and G. C. K. Roberts. 1996. Structure-activity relationships in the peptide antibiotic nisin: the role of dehydroalanine 5. *Appl. Environ. Microbiol.* **62**:2966-2969.
- Demel, R. A., T. Peelen, R. J. Siezen, B. de Kruijff, and O. P. Kuipers. 1996. Nisin Z, mutant nisin Z and lactacin 481 interactions with anionic lipids correlate with antimicrobial activity. A monolayer study. *Eur. J. Biochem.* **235**:267-274.

13. Dodd, H. M., N. Horn, C. J. Giffard, and M. J. Gasson. 1996. A gene replacement strategy for engineering nisin. *Microbiology* **142**:47–55.
14. Driessen, A. J. M., and W. N. Konings. 1993. Insertion of lipids and proteins into bacterial membranes by fusion with liposomes. *Methods Enzymol.* **221**:394–408.
15. Driessen, A. J. M., H. W. van den Hooven, W. Kuiper, M. van de Kamp, H.-G. Sahl, R. N. H. Konings, and W. N. Konings. 1995. Mechanistic studies of lantibiotic-induced permeabilization of phospholipid vesicles. *Biochemistry* **34**:1606–1614.
16. Gao, F. H., T. Abee, and W. N. Konings. 1991. Mechanism of action of the peptide antibiotic nisin in liposomes and cytochrome *c* oxidase-containing proteoliposomes. *Appl. Environ. Microbiol.* **57**:2164–2170.
17. García Garcerá, M. J., M. G. L. Elferink, A. J. M. Driessen, and W. N. Konings. 1993. In vitro pore-forming activity of the lantibiotic nisin. *Eur. J. Biochem.* **212**:417–422.
18. Gross, E., and J. L. Morell. 1971. The presence of dehydroalanine in the lantibiotic nisin and its relationship to activity. *J. Am. Chem. Soc.* **93**:4634–4635.
19. Kashket, E. R. 1985. The proton motive force in bacteria: a critical assessment of methods. *Annu. Rev. Microbiol.* **39**:219–242.
20. Konings, W. N., B. Poolman, and A. J. M. Driessen. 1989. Bioenergetics and solute transport in *Lactococci*. *Crit. Rev. Microbiol.* **16**:419–476.
21. Kordel, M., and H.-G. Sahl. 1986. Susceptibility of bacterial eukaryotic and artificial membranes to the disruptive action of the cationic peptides Pep5 and nisin. *FEMS Microbiol. Lett.* **34**:139–144.
22. Kuipers, O. P., M. M. Beerthuizen, P. G. A. de Ruyter, E. J. Luesink, and W. M. de Vos. 1995. Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction. *J. Biol. Chem.* **45**:27299–27304.
23. Kuipers, O. P., M. M. Beerthuizen, R. J. Siezen, and W. M. de Vos. 1993. Characterization of the nisin gene cluster nis ABTCIPR of *Lactococcus lactis*. Requirement of expression of the nis A and nis I genes for the development of immunity. *Eur. J. Biochem.* **216**:281–291.
24. Kuipers, O. P., G. Bierbaum, B. Ottenwälder, H. M. Dodd, N. Horn, J. Metzger, T. Kupke, V. Gnau, R. Bongers, P. van den Bogaard, H. Kusters, H. S. Rollema, W. M. de Vos, R. J. Siezen, G. Jung, F. Götz, H.-G. Sahl, and M. J. Gasson. 1996. Protein engineering of lantibiotics. *Antonie Leeuwenhoek* **69**:161–170.
25. Kuipers, O. P., H. S. Rollema, W. M. G. J. Yap, H. J. Boot, R. J. Siezen, and W. M. de Vos. 1992. Engineering dehydrated amino acid residues in the antimicrobial peptide nisin. *J. Biol. Chem.* **267**:24340–24346.
26. Kuipers, O. P., W. M. G. J. Yap, H. S. Rollema, M. M. Beerthuizen, R. J. Siezen, and W. M. de Vos. 1991. Expression of wild-type and mutant nisin genes in *Lactococcus lactis*, p. 250–259. In G. Jung and H.-G. Sahl (ed.), *Nisin and novel lantibiotics*. ESCOM, Leiden, The Netherlands.
27. Lian, L.-Y., W. C. Chan, S. D. Morley, G. C. K. Roberts, B. W. Bycroft, and D. Jackson. 1991. NMR studies of the solution structure of nisin A and related peptides, p. 43–58. In G. Jung and H.-G. Sahl (ed.), *Nisin and novel lantibiotics*. ESCOM, Leiden, The Netherlands.
28. Lian, L.-Y., W. C. Chan, S. D. Morley, G. C. K. Roberts, B. W. Bycroft, and D. Jackson. 1992. Solution structures of nisin A and its two major degradation products determined by NMR. *Biochem. J.* **283**:413–420.
29. Liu, W., and J. N. Hansen. 1993. The antimicrobial effect of a structural variant of subtilin against outgrowing *Bacillus cereus* T spores and vegetative cells occurs by different mechanisms. *Appl. Environ. Microbiol.* **59**:648–651.
30. Lolkema, J. S., K. J. Hellingwerf, and W. N. Konings. 1982. The effect of “probe binding” on the quantitative determination of the proton-motive force in bacteria. *Biochim. Biophys. Acta* **681**:85–94.
31. Molenaar, D., T. Abee, and W. N. Konings. 1991. Measurement of intracellular pH in bacteria with a fluorescent probe. *Biochim. Biophys. Acta* **1115**:75–83.
32. Moll, G., T. Ubbink-Kok, H. Hildeng-Hauge, J. Nissen-Meyer, I. F. Nes, W. N. Konings, and A. J. M. Driessen. 1996. Lactococin G is a potassium ion-conducting two component bacteriocin. *J. Bacteriol.* **178**:600–605.
- 32a. Moll, G. N., G. C. K. Roberts, W. N. Konings, and A. J. M. Driessen. 1996. Mechanism of lantibiotic-induced pore-formation. *Antonie Leeuwenhoek* **69**:185–191.
33. Morris, S. L., R. C. Walsh, and J. N. Hansen. 1984. Identification and characterization of some bacterial membrane sulfhydryl groups which are targets of bacteriostatic and antibiotic action. *J. Biol. Chem.* **259**:13590–13594.
34. Nissen-Meyer, J., H. Holo, L. S. Havårstein, K. Sletten, and I. F. Nes. 1992. A novel lactococcal bacteriocin whose activity depends on the complementary action of two peptides. *J. Bacteriol.* **174**:5686–5692.
35. Okereke, A., and T. J. Montville. 1992. Nisin dissipates the proton motive force of the obligate anaerobe *Clostridium sporogenes* PA 3679. *Appl. Environ. Microbiol.* **58**:2463–2467.
36. Reisinger, P., H. Seidel, H. Tschesche, and W. P. Hammes. 1980. The effect of nisin on murein synthesis. *Arch. Microbiol.* **127**:187–193.
37. Ruhr, E., and H.-G. Sahl. 1985. Mode of action of the peptide antibiotic nisin and influence on the membrane potential of whole cells and on cytoplasmic and artificial membrane vesicles. *Antimicrob. Agents Chemother.* **27**:841–845.
38. Sahl, H.-G., M. Kordel, and R. Benz. 1987. Voltage dependent depolarization of bacterial membranes and artificial lipid bilayers by the peptide antibiotic nisin. *Arch. Microbiol.* **149**:120–124.
39. Sahl, H.-G. 1991. Pore formation in bacterial membranes by cationic lantibiotics, p. 347–358. In G. Jung and H.-G. Sahl (ed.), *Nisin and novel lantibiotics*. ESCOM, Leiden, The Netherlands.
40. Sahl, H.-G., R. W. Jack, and G. Bierman. 1995. Lantibiotics: biosynthesis and biological activities of peptides with unique post-translational modifications. *Eur. J. Biochem.* **230**:827–853.
41. Van den Hooven, H. W., F. Fogolari, H. S. Rollema, R. N. H. Konings, C. W. Hilbers, and F. J. M. van de Ven. 1993. NMR and circular dichroism studies of the lantibiotic nisin in non-aqueous environments. *FEBS Lett.* **319**:189–194.
42. Van den Hooven, H. W., C. C. M. Doeland, M. van de Kamp, R. N. H. Konings, C. W. Hilbers, and F. J. M. van de Ven. 1996. Three-dimensional structure of the lantibiotic nisin in the presence of membrane-mimetic micelles of dodecylphosphocholine and of sodium dodecylsulphate. *Eur. J. Biochem.* **235**:382–393.
43. Van den Hooven, H. W., C. A. E. M. Spronk, M. van de Kamp, R. N. H. Konings, C. W. Hilbers, and F. J. M. van de Ven. 1996. Surface location and orientation of the lantibiotic nisin bound to membrane-mimicking micelles of dodecylphosphocholine and of sodium dodecylsulphate. *Eur. J. Biochem.* **235**:394–403.
44. van de Ven, F. J. M., H. W. van den Hooven, R. N. H. Konings, and C. W. Hilbers. 1991. NMR studies of lantibiotics. The structure of nisin in aqueous solution. *Eur. J. Biochem.* **202**:1181–1188.