

University of Groningen

Arctic waders are not capital breeders

Klaassen, Marcel; Lindström, Åke; Meltofte, Hans; Piersma, Theunis

Published in:
 Nature

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2001

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Klaassen, M., Lindström, Å., Meltofte, H., & Piersma, T. (2001). Arctic waders are not capital breeders. *Nature*, 413, 794-794.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Ornithology

Arctic waders are not capital breeders

Birds prepare their eggs from recently ingested nutrients ('income' breeders) or from body stores ('capital' breeders)¹. As summers are short at Arctic latitudes, Arctic migrants have been presumed to bring nutrients for egg production from their previous habitats, so that they can start breeding immediately upon arrival¹⁻³. But we show here that eggs laid by 10 different wader species from 12 localities in northeast Greenland and Arctic Canada are produced from nutrients originating from tundra habitats, as inferred from carbon stable-isotope ratios in eggs, natal down, and juvenile and adult feathers.

During winter and migration, most Arctic-breeding waders eat estuarine invertebrates, shifting to terrestrial and limnic invertebrates on tundra breeding grounds. Invertebrates from estuarine and tundra habitats have distinctly different carbon-isotope ratios⁴. Diet-based differences in carbon-isotope ratios are expressed in bird tissues, including eggs and feathers^{4,5}. We investigated whether Arctic-breeding wader species that use estuarine habitats during the non-breeding season are capital breeders, producing eggs and hatchlings with carbon-isotope ratios that are typical of estuaries.

As expected, the flight feathers produced during early winter show ratios that

are typical of estuarine systems (Fig. 1). The same is true for shoulder feathers, which are among the last to be moulted before migration to breeding grounds⁶. However, the carbon-isotope signatures of eggs and natal down are typical of terrestrial and limnic systems. The eggs and resultant hatchlings therefore seem to be produced from local nutrients. This conclusion is supported by the strong resemblance of the carbon-isotope signatures of natal down and juvenile flight feathers, showing that eggs and post-hatch tissues have the same nutritional source.

Our results also indicate that the use of a mixed capital/income breeding strategy by these waders is unlikely. Egg-laying females use a single, local nutrient source, as indicated by the low within-clutch variation compared with the among-clutch variation in carbon-isotope ratios of natal down. (We used a hierarchically nested design in a mixed-model ANOVA, in which brood is nested within species as a random factor, and found significant effects of species ($F_{6,30} = 4.8, P < 0.002$) and brood ($F_{29,93} = 16.8, P < 0.001$) on the carbon stable-isotope ratios of natal down.)

We conclude that, with respect to egg production, the Arctic-breeding waders investigated here are income, rather than capital, breeders. For these waders, the fitness costs of transporting extra nutrient stores to breeding grounds⁷ outweigh the potential benefits¹⁻³. The capital strategy may still be used by large species of Arctic breeding migrants, such as geese^{1,2} — it is likely that larger species need relatively

smaller body stores for egg production. Also, larger species, although constrained by the same fixed period of opportunity as smaller birds, require longer to complete their breeding.

Marcel Klaassen*, **Åke Lindström†‡**, **Hans Meltofte§**, **Theunis Piersma‡**

*Centre for Limnology, Netherlands Institute of Ecology, PO Box 1299, 3600 BG Maarsse, The Netherlands

e-mail: klaassen@cl.nioo.knaw.nl

†Department of Animal Ecology, Ecology Building, Lund University, 22362 Lund, Sweden

‡Netherlands Institute for Sea Research, PO Box 59, 1790 AB Den Burg, Texel, and Centre for Ecological and Evolutionary Studies, University of Groningen, PO Box 14, 9750 AA Haren, The Netherlands

§Department of Arctic Environment, National Environmental Research Institute, PO Box 358, 4000 Roskilde, Denmark

1. Drent, R. H. & Daan, S. *Ardea* **68**, 225–252 (1980).
2. Newton, I. in *Evolutionary Ecology* (eds Stonehouse, B. & Perrins, C.) 113–126 (Macmillan, London, 1977).
3. Davidson, N. C. & Evans, P. R. in *Acta XIX Congressus Internationalis Ornithologici* (ed. Ouellet, H.) 342–352 (Univ. Ottawa Press, Ottawa, 1988).
4. Peterson, B. J. & Fry, B. *Annu. Rev. Ecol. Syst.* **18**, 293–320 (1987).
5. Hobson, K. A. *Oecologia* **120**, 314–326 (1999).
6. Cramp, S. (ed.) *Handbook of the Birds of Europe, the Middle East and North Africa* Vol. 3 (Oxford Univ. Press, Oxford, 1983).
7. Witter, M. S. & Cuthill, I. C. *Phil. Trans. R. Soc. Lond. B* **340**, 73–92 (1993).
8. Stettenheim, P. in *Avian Biology* Vol. 2 (eds Farner, D. S., King, J. R. & Parkes, K. C.) 1–63 (Academic, New York, 1972).

Intracellular signalling

Key enzyme in leptin-induced anorexia

Leptin is a key hormonal regulator of energy balance that acts upon hypothalamic neurons to reduce food intake, but the intracellular mechanisms involved are incompletely understood. Here we show that systemic administration of leptin in rats activates the enzyme phosphatidylinositol-3-OH kinase (PI(3)K) in the hypothalamus and that intracerebroventricular (i.c.v.) infusion of inhibitors of this enzyme prevents leptin-induced anorexia. Our results indicate that PI(3)K is a crucial enzyme in the signal-transduction pathway that links hypothalamic leptin to reduced food intake.

Leptin, a hormone derived from adipocytes (fat cells), is known to activate PI(3)K in non-neuronal cells and to elicit cellular responses that are blocked by PI(3)K inhibitors *in vitro*^{1,2}; it also activates the transcription factor STAT3 (ref. 3). To determine whether leptin activates PI(3)K, as well as STAT3, in the hypothalamus *in vivo*, we observed the effect of systemically administered leptin on their activity in male Wistar rats. As before³, leptin induced rapid tyrosine-phosphorylation of STAT3




Figure 1 Carbon stable-isotope ratios (parts per thousand difference from ¹³C/¹²C ratio in PeeDee Limestone⁴; δC) of eggs, natal down and feathers of different species of Arctic-breeding waders at different times during the year. All samples were collected in northeast Greenland and Arctic Canada in 1999 and 2000. Feathers collected from nest-attending adults were grown either in winter (adult flight feathers) or during spring migration (adult shoulder feathers); data are averages across individuals ± s.d. Eggs were collected from deserted nests (egg content); natal down was collected from hatchlings; data are averages across clutches ± s.d. per species. Data for natal down still attached to the tips of head and neck feathers of independent young⁸, and for secondary flight feathers from independent young are averages across individuals ± s.d.). Values of δC were determined using a Carlo Erba model 1106 elemental analyser which was coupled online, through a Finnigan conflow 2 interface, to a Finnigan Delta S mass spectrometer. Feathers were washed in chloroform before carbon-isotope analysis. Using a hierarchically nested design in a mixed model ANOVA, in which species was nested within tissue type, which in turn was nested within the tissue groups 'adult feathers' and 'egg, natal down and juvenile feathers', we found significant effects of tissue group ($F_{1,152} = 805.3, P < 0.001$), tissue type ($F_{4,152} = 6.7, P < 0.001$) and species ($F_{9,152} = 2.8, P < 0.005$) on δC.