

University of Groningen

Theoretical models for fluid thermodynamics based on the quasi-Gaussian entropy theory

Amadei, Andrea

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1998

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Amadei, A. (1998). *Theoretical models for fluid thermodynamics based on the quasi-Gaussian entropy theory*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Summary

The theoretical modeling of fluid thermodynamics is one of the most challenging fields in physical chemistry. In fact the fluid behavior, except at very low density conditions, is still extremely difficult to be modeled from a statistical mechanical point of view, as for any realistic model Hamiltonian the configurational part of the partition function cannot be evaluated, i.e., the corresponding high dimensional integral is far too complex to be solved. Hence once a molecular Hamiltonian has been modeled, often severe approximations are necessary in order to obtain a model for the thermodynamics.

The usual approximations, based on integral equations or on perturbation theory make use of different kinds of expansions in the particle correlation functions or energy fluctuations for a given molecular Hamiltonian model. Much has been done on this from the 50's on and in the literature many different versions of these two basic approaches can be found. However, these methods still cannot be used in general for polar fluids, and even in the case of the simpler apolar fluids, where the Lennard-Jones potential seems to be a good model, they do not always provide an accurate thermodynamic description. Direct molecular simulations of fluids are also severely limited by the difficulties connected to the still insufficient computer power to simulate large enough systems reaching a sufficient sampling of phase space. In addition, the modeling of molecular Hamiltonians for real fluids is difficult and often it can be achieved only with the use of approximations in the molecular description.

Recently a new theoretical approach, the quasi-Gaussian entropy (QGE) theory, has been proposed. In this theory the basic statistical mechanical relations

are rewritten in terms of the distribution of the fluctuations of a macroscopic property instead of the partition function. In fact, for each statistical mechanical ensemble it is possible to define a proper reference state such that the free energy difference between the actual condition and the reference one can be expressed exactly in terms of the moment generating function, defined by the distribution of a specific macroscopic fluctuation. Instead of modeling as usual a molecular Hamiltonian, in the QGE theory we model the distribution and hence its moment generating function, to describe the thermodynamics of the system. In this way we obtain for each model distribution a corresponding model system at the level of thermodynamic behavior.

In modeling the fluctuation distributions we can use basic physical principles to restrict the possible set of distributions. In fact considering that a (macroscopic) thermodynamic system can be always decomposed into a large set of identical virtually independent subsystems, defined as “elementary systems” in the QGE theory, we realize that the distribution for the fluctuations of a macroscopic property is the convolution of a large set of independent “elementary” distributions. Hence applying the central limit theorem we obtain that every distribution of a macroscopic property must be “quasi-Gaussian”, and thus can be modeled as a unimodal distribution. Note that the concept of elementary systems is purely a consequence of the thermodynamic principle of linear scalability of extensive properties and size independence of the intensive ones. The elementary system hence defines the “universal” minimum size required to obtain the thermodynamic behavior. Moreover, each type of model distribution provides a closure relation for a general thermodynamic equation, resulting in an ordinary differential equation, the thermodynamic master equation (TME). According to the combination of the ensemble and distribution chosen, the TME solution provides the temperature or density dependence of the thermodynamic properties. Each model distribution, combined with the input information at a single thermodynamic state point, defines a corresponding “statistical state” of the system, and hence for a given model distribution the corresponding exact thermodynamics is obtained.

It should be clear that modeling the moment generating function, which is directly connected to the entropy of the system, in terms of quasi-Gaussian distributions simplifies tremendously the statistical mechanical description, substituting a high dimensional integral, the partition function, with a one dimensional integral, the moment generating function. Many different types of molecular Hamiltonians could in principle result in the same type of distribution and thus the Hamiltonian fixes the distribution and not vice versa. It should also be noted that a good model distribution for a real fluid is likely to have a mathematical complexity which is still relatively low, although the corresponding model Hamiltonian can be rather complicated. Only in the critical point region and at phase transitions extra complications can arise and a proper choice of the ensemble and type of fluctuation used is essential to keep a relatively simple model distribution.

In this thesis the QGE theory has been developed and systematically applied to obtain general thermodynamic models for fluids. The theory was presented in a general and comprehensive way, describing in detail its derivation in different statistical mechanical ensembles and using different kinds of fluctuations. The basic theoretical models (statistical states) for the temperature and density dependence were used to describe the thermodynamics of different fluid systems, finding that the level of the QGE theory employed (the Gamma state level) can really provide an accurate general model for fluids. The theory was used in combination with both experimental and simulation data, and the derivations of general fluid equations of state, obtained from the theory, were described. Finally the use of the QGE theory to obtain the partial molar properties in a fluid mixture was addressed.