

University of Groningen

Autolysis of *Lactococcus lactis* caused by induced overproduction of its major autolysin, AcmA

Buist, Girbe; Karsens, H; Nauta, A; van Sinderen, D; Venema, G; Kok, J

Published in:
Applied and environmental microbiology

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1997

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Buist, G., Karsens, H., Nauta, A., van Sinderen, D., Venema, G., & Kok, J. (1997). Autolysis of *Lactococcus lactis* caused by induced overproduction of its major autolysin, AcmA. *Applied and environmental microbiology*, 63(7), 2722-2728.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Autolysis of *Lactococcus lactis* Caused by Induced Overproduction of Its Major Autolysin, AcmA

GIRBE BUIST, HARMA KARSENS, ARJEN NAUTA, DOUWE VAN SINDEREN,†
GERARD VENEMA,* AND JAN KOK

Department of Genetics, Groningen Biomolecular Sciences and Biotechnology Institute,
University of Groningen, 9751 NN Haren, The Netherlands

Received 16 December 1996/Accepted 22 April 1997

The optical density of a culture of *Lactococcus lactis* MG1363 was reduced more than 60% during prolonged stationary phase. Reduction in optical density (autolysis) was almost absent in a culture of an isogenic mutant containing a deletion in the major autolysin gene, *acmA*. An *acmA* mutant carrying multiple copies of a plasmid encoding AcmA lysed to a greater extent than the wild-type strain did. Intercellular action of AcmA was shown by mixing end-exponential-phase cultures of an *acmA* deletion mutant and a tripeptidase (*pepT*) deletion mutant. PepT, produced by the *acmA* mutant, was detected in the supernatant of the mixed culture, but no PepT was present in the culture supernatant of the *acmA* mutant. A plasmid was constructed in which *acmA*, lacking its own promoter, was placed downstream of the inducible promoter/operator region of the temperate lactococcal bacteriophage r1t. After mitomycin induction of an exponential-phase culture of *L. lactis* LL302 carrying this plasmid, the cells became subject to autolysis, resulting in the release of intracellular proteins.

The action of some of the bacterial peptidoglycan hydrolases (proteins degrading the peptidoglycan of bacterial cell walls) can result in cell lysis (30). Therefore, the potentially lethal enzymes causing this phenomenon can be referred to as autolysins. In the only paper to date on the genetics of autolysis of *Lactococcus lactis*, we have described the cloning of the major autolysin gene, *acmA*, of *L. lactis* subsp. *cremoris* MG1363 (3). AcmA is a lysozyme-like enzyme (muramidase) that hydrolyzes the *N*-acetylmuramyl-1,4- β -*N*-acetylglucosamine bonds in the peptidoglycan.

Autolysis and the subsequent release of intracellular substances from the cells of a number of lactococcal strains have been shown during growth in liquid media (2, 17, 28, 29, 44) as well as during cheese production (5–7, 18, 46). Various factors such as pH, temperature, carbon source, and salt concentration appear to be important for the autolytic process. The degree of autolysis is strain dependent, and the process starts after exponential growth has ceased. The proteolytic activities of lactococci are involved in ripening and in flavor development in fermented milk products, such as cheese (27, 45). Lactococci contain more than 10 different intracellular peptidases (14) whose action leads to the production of small peptides and free amino acids which are flavors and flavor precursors. The degree and rate of release of these peptidases into the cheese matrix after lysis of the cells is of great importance for cheese maturation and flavor development (5, 7, 45, 46). Cheese maturation is a slow and therefore costly process and may be accelerated by enhanced lysis of cells with concomitant quick release of intracellular peptidases.

In a first attempt to construct starters with enhanced autolytic properties, Feirtag and McKay (11) mutagenised *L. lactis* C2 and obtained thermolytic variants which lysed at 38 to 40°C

but grew normally at 32°C. Lysis was evidenced by the reduction in optical density of the culture and by the release of the intracellular enzyme phospho- β -galactosidase. Shearman et al. (35) have constructed a lactococcal strain containing the Φ vML3 lysin gene under the control of its own promoter. After growth in milk at 30°C and subsequent storage at 12°C, the number of viable cells dropped to zero within 28 days, whereas the control strain still contained more than 10⁶ viable cells per ml. Apparently, the lysin caused enhanced lysis of lactococcal cells, although this was not documented by showing a release of intracellular components.

In this study, we proved that AcmA is an autolysin involved in stationary-phase lysis of *L. lactis* and used this information to construct a system for *L. lactis* with which enhanced autolysis and release of intracellular proteins was obtained. This system is based on the recently characterized promoter/operator region of the temperate lactococcal bacteriophage r1t (25, 43).

MATERIALS AND METHODS

Bacteria, plasmids, and growth conditions. The strains and plasmids used in this study are listed in Table 1. *L. lactis* was grown at 30°C in 0.5× M17 broth (Difco, West Molesey, United Kingdom) containing 1.9% β -glycerophosphate (Sigma Chemical Co., St. Louis, Mo.), or in M17 when indicated. M17 agar plates contained 1.5% agar. All of these media were supplemented with 0.5% glucose. When needed, 5 μ g of erythromycin (Boehringer GmbH, Mannheim, Germany) per ml was added. *Escherichia coli* was grown in TY (Difco Laboratories, Detroit, Mich.) medium at 37°C with vigorous agitation or on TY agar plates containing 1.5% agar. Ampicillin (Sigma) and erythromycin were used at final concentrations of 100 μ g/ml.

General DNA techniques and transformation. Molecular cloning techniques were performed essentially as described by Sambrook et al. (31). Restriction enzymes, Klenow enzyme, T4 DNA polymerase and T4 DNA ligase were obtained from Boehringer and used as specified by the supplier. Deoxynucleotides were obtained from Pharmacia LKB Biotechnology AB, Uppsala, Sweden. *E. coli* and *L. lactis* were transformed by electroporation with a gene pulser (Bio-Rad Laboratories, Richmond, Calif.), as described by Zabarovsky and Winberg (47) and Leenhouts and Venema (20), respectively. Plasmid DNA was isolated from *E. coli* and *L. lactis* by the method of Birnboim and Doly, with minor modifications for *L. lactis* (34).

Primer extension analysis. RNA was isolated as previously described (39) from an exponentially growing *L. lactis* culture at an optical density at 600 nm (OD₆₀₀) of 0.5. Oligonucleotide pALA-26 (5'-CGCCAGCAAATTTGTGGC TGGTTTATAAAAAGCGAGTGG), synthesized with a 381A DNA synthesizer (Applied Biosystems Inc., Foster City, Calif.), was used for primer extension

* Corresponding author. Mailing address: Department of Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Kerklaan 30, 9751 NN Haren, The Netherlands. Phone: 31-50-3632093. Fax: 31-50-3632348. E-mail: VenemaG@BIOL.RUG.NL.

† Present address: Department of Microbiology, University College Cork, Cork, Ireland.

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Relevant phenotype or genotype	Source or reference
Strains		
<i>L. lactis</i> subsp. <i>cremoris</i>		
MG1363	Plasmid-free strain	12
MG1363 <i>acm</i> Δ <i>I</i>	MG1363 derivative carrying a deletion in the <i>acmA</i> gene	3
MG1363 <i>pepT</i>	MG1363 derivative containing deletion in the <i>pepT</i> gene	22
LL302	MG1363 carrying the pWV01 <i>repA</i> gene on the chromosome	19
<i>E. coli</i>		
NM522	<i>supE thi (lac-proAB) hsd-5 (r_K⁻ m_K⁻) [F' proAB lacI^rZΔM15]</i>	Stratagene, La Jolla, Calif.
MC1000	<i>araD139 lacX74 (ara, leu) 7697 galU galK strA</i>	4
Plasmids		
pAL01	Ap ^r , pUC19 containing 4,137-bp lactococcal chromosomal DNA insert carrying the <i>acmA</i> gene	3
pAL08	Ap ^r , pAL01 with <i>Sma</i> I- <i>Eco</i> RI deletion	This work
pAL10	Ap ^r , pAL08 containing 2,716-bp <i>Sac</i> I fragment of ORF1 of the <i>srfA</i> operon of <i>B. subtilis</i>	This work
pAL11	Em ^r , pIR1EF containing 4,520-bp <i>Sca</i> I- <i>Bam</i> HI fragment of pAL10	This work
pAL12	Em ^r , pIR12 containing <i>acmA</i> under control of the regulatory region of phage r1t	This work
pIR12	Em ^r , pWV01 derivative carrying the regulatory region of phage r1t	25
pEF+	Ap ^r , SK+ containing a 1,740-bp <i>Eco</i> RI fragment of phage r1t	43
pIR12EF	Em ^r , pIR12 containing a 1,785-bp <i>Sca</i> I- <i>Bam</i> HI fragment of pEF+	This work
pIR1EF	Em ^r , pIR12EF in which <i>Sac</i> I site was removed	This work
pGK13	Em ^r Cm ^r , pWV01-based lactococcal plasmid	13
pGKAL1	Em ^r Cm ^r , pGK13 containing 1,942-bp <i>Ssp</i> I- <i>Bam</i> HI fragment of pAL01	This work
pGKAL2	Em ^r Cm ^r , pGK13 containing 1,804-bp <i>Sca</i> I- <i>Bam</i> HI fragment of pAL01	This work

reactions. Nucleotide sequence reactions were done on plasmid pAL01 by the dideoxy chain termination method (33) with the T7 sequencing kit and protocol (Pharmacia). A 25-ng portion of primer was added to 3.5 μ g of RNA in a reaction mixture containing dCTP, dGTP, dTTP, and [α -³²S]dATP, and cDNA was synthesized with avian myeloblastosis virus reverse transcriptase (Boehringer). After 10 min of incubation at 42°C, an excess of cold dATP was added, and incubation was continued for another 10 min at 42°C. The products were analyzed on a 6% polyacrylamide (PAA) sequencing gel.

Plasmid constructions. Plasmids pGKAL1 and pGKAL2 (Fig. 1) were constructed by subcloning of the 1,943-bp *Ssp*I-*Bam*HI or the 1,804-bp *Sca*I-*Bam*HI fragment of pAL01 (3), respectively, into the *Eco*RV-*Bam*HI sites of lactococcal plasmid pGK13. Plasmid pAL01 is a pUC19 derivative containing a 4,137-bp chromosomal DNA fragment from *L. lactis* MG1363 encompassing *acmA*. The ligation mixtures were used to transform *L. lactis* MG1363*acm* Δ *I*.

All cloning steps for the construction of pAL12 (Fig. 1) were performed with *E. coli* MC1000 unless stated otherwise. The *Sac*I site present in the multiple-cloning site of pAL01 was removed by cutting with *Eco*RI and *Sma*I. The plasmid was treated with Klenow enzyme, ligated, and used to electrotransform *E. coli* NM522, resulting in plasmid pAL08. Because *E. coli* grows very poorly when it carries an intact *acmA* gene (3), *acmA* was disrupted by cloning into the unique *Sac*I site of pAL08 a 2,716-bp *Sac*I fragment originating from the *srfA* operon of *Bacillus subtilis* (42). This resulted in pAL10.

One of the two *Sac*I sites present in pIR12 (25) was deleted by replacing the 2,750-bp *Sca*I-*Xho*II fragment by a 1,785-bp *Sca*I-*Bam*HI fragment, taken from pEF+ (43). The remaining *Sac*I site in the resulting plasmid, pIR12EF, was removed by digestion with *Sac*I and treatment with T4 DNA polymerase. After self-ligation, pIR1EF was obtained. The 1,764-bp *Eco*RV-*Xho*II fragment of pIR1EF was replaced by the 4,520-bp *Sca*I-*Bam*HI fragment of pAL10 containing the interrupted *acmA* gene. The resulting plasmid, pAL11, was digested with *Sac*I to remove the DNA fragment interrupting *acmA*. After self-ligation, the mixture was used to transform *L. lactis* LL302 and plasmid pAL12 was obtained.

Mitomycin induction. An overnight culture of *L. lactis* was diluted 100-fold in GM17 and grown to an OD₆₀₀ of 0.2. The culture was divided into two portions, and mitomycin (Sigma) was added to one of them to a final concentration of 1 μ g/ml. Incubation was continued at 30°C. The OD₆₀₀ values were measured in a Philips PU8720 UV/VIS spectrophotometer (Pye Unicam Ltd., Cambridge, United Kingdom).

Sample preparation, SDS-PAGE, and detection of lytic activity. For the analysis of the intercellular action of AcmA, 2-ml samples of culture were subjected to centrifugation. A 1-ml volume of the supernatant fraction was dialyzed against several changes of demineralized water, lyophilized, and dissolved in 0.5 ml of denaturation buffer (1). The cell pellet was resuspended in 1 ml of denaturation buffer, and cell extracts were prepared as described by van de Guchte et al. (41). The samples were boiled for 2 min and centrifuged, and 30 μ l of the mid-exponential-phase samples and 15 μ l of the other samples were loaded onto sodium dodecyl sulfate (SDS)-PAA gels.

For the analysis of (induced) lysis, 1-ml samples were treated as described

above and the supernatant and cell fractions were dissolved in 0.2 ml of denaturation buffer. The amount of sample loaded was equalized according to the measured optical density. SDS-PAGE was carried out by the method of Laemmli (16) with the Protean II minigel system (Bio-Rad). The standard low-range and prestained low- and high-range SDS-PAGE molecular weight markers of Bio-Rad were used as references. SDS-PAA gels were stained with Coomassie brilliant blue (Bio-Rad).

Lytic activity was detected in situ by using SDS-12.5% PAA gels containing 0.15% autoclaved, lyophilized *Micrococcus lysodeikticus* ATCC 4698 cells (Sigma) as described previously (3). Protein renaturation was performed at room temperature for 14 h.

Western blotting and immunodetection. After SDS-PAGE, the proteins were transferred to BA85 nitrocellulose membranes (Schleicher & Schuell, Dassel, Germany) as described by Towbin et al. (38). Endopeptidase and tripeptidase antigens were detected with 1:8,000-diluted polyclonal anti-endopeptidase antibodies (23) and 1:4,000-diluted polyclonal anti-tripeptidase antibodies (22), respectively, and alkaline phosphatase-conjugated goat anti-rabbit antibodies (Promega Corp., Madison, Wis.) by using the Western-light chemiluminescent detection system and protocol (TROPIX Inc., Bedford, Mass.).

RESULTS

AcmA is required for autolysis of *L. lactis* during stationary phase. Overnight cultures of *L. lactis* MG1363 and its *acmA* deletion mutant MG1363*acm* Δ *I* (3) were diluted 200-fold in fresh prewarmed M17 broth. During the first 9 h of growth and hourly sampling, the cultures were gently shaken at 30°C to prevent settling of MG1363*acm* Δ *I*, which grows as long filaments due to improper cell separation (3). Further incubation was carried out without shaking, but the cultures were briefly shaken before sampling. The doubling time of the wild-type and mutant strains was 33 min. During stationary phase, the OD₆₀₀ of both cultures decreased and remained stable after approximately 1 week of incubation (results not shown). The average maximal percent OD₆₀₀ reduction, [(OD_{max} - OD_{7days})/OD_{max}] \times 100%, was 63% for the wild type and 14% for the mutant (mean of results from three independent experiments). Apparently, the major autolysis of *L. lactis* is not only required for cell separation (3) but is also responsible for cell lysis upon prolonged incubation.




FIG. 1. Plasmid maps and nucleotide sequence of the *acmA* promoter region. (A) Map of pGKAL1. The putative ribosome binding site (RBS and lowercase letters), -10 and -35 sequences (underlined), start codon (italic), transcriptional start point (asterisk), and the *SspI* (only half) and *ScaI* restriction enzyme sites are indicated. *EmR* and *CmR*, erythromycin and chloramphenicol resistance genes, respectively; ORI (open square), origin of replication of the lactococcal plasmid pWV01; *repA*, gene encoding the replication initiation protein of pWV01; *acmA*, *N*-acetylmuramidase gene of *L. lactis* MG1363. The fragment from pAL01 (3) containing *acmA* is indicated with a grey bar. The cloning vector (thin line) is pGK13. In pGKAL2, the *ScaI* site was fused to the *EcoRV* site of pGK13. (B) Map of pAL12. The abbreviations are the same as in panel A. P1 and P2 (\blacktriangle), promoters from bacteriophage λ ; T, transcription terminator of the lactococcal proteinase gene *prtP*; *rro*, λ 1 repressor gene; *tec*, topological equivalent of lambda *cro*; *ORF5'*, 5'-end of ORF5 of phage λ 1. Only relevant restriction enzyme sites are shown for both plasmids.

Complementation of *acmA* Δ 1 and localization of the *acmA* promoter. A putative -35 hexanucleotide and a -10 sequence preceded by the sequence TGN, found in more than 40% of the lactococcal promoters analyzed so far (10), is present upstream of the start codon of *acmA* (Fig. 1). The spacing between the two consensus sequences (23 nucleotides) is exceptionally large. To examine whether this sequence is functional, pGKAL1 and pGKAL2 were constructed. pGKAL1 contains a 138-bp *SspI-ScaI* fragment carrying this sequence, whereas pGKAL2 does not (see Fig. 1). *L. lactis* MG1363(pGK13) and *L. lactis* MG1363*acmA* Δ 1 containing either pGK13, pGKAL1, or pGKAL2 were patched onto a GM17 plate containing 0.15% autoclaved *M. lysodeikticus* cells, and the plate was incubated for 36 h at 30°C. The results are presented in the inset in Fig. 2 and show that no halo had formed around the colony of cells containing pGKAL2 but that a large halo was present around the cells containing pGKAL1. The halo was even larger than that formed by *L. lactis* MG1363. Apparently, *L. lactis* can cope with multiple copies of *acmA* and with the increased amount of the deleterious enzyme AcmA. This result also indicates that the 138-bp *SspI-ScaI* fragment is required for *acmA* expression. This fragment, when cloned upstream of the promoterless *E. coli lacZ* gene in plasmid pORI13 (32), drove β -galactosidase expression in *E. coli* but not in *L. lactis* (results not shown). Primer extension analysis performed on RNA isolated from MG1363 cells revealed that the *acmA* mRNA starts at the T residue 6 bases downstream of the -10 hexanucleotide (result not shown). Whereas the same RNA sample gave normal primer extension products of the transcripts of two other genes, an exposure of 1 week was needed to visualize a faint band of the extension product, indicating that the promoter is only very weakly expressed. This is in agreement with the fact that we were unable to identify a protein band in a

200-fold-concentrated sample of culture supernatant of *L. lactis* MG1363 run on a PAA gel and stained with Coomassie brilliant blue which would correspond to the position of AcmA clearing bands in an activity gel.

Increased production of AcmA leads to more lysis. Overnight cultures of MG1363(pGK13) and MG1363*acmA* Δ 1 con-




FIG. 2. GM17 growth curves and halo formation, on a GM17 plate containing 0.15% autoclaved *M. lysodeikticus* cells (inset), of *L. lactis* MG1363(pGK13) (\blacktriangle , A), and *L. lactis* MG1363*acmA* Δ 1 containing pGK13 (\blacksquare , C), pGKAL1 (\bullet , D), or pGKAL2 (B).




FIG. 3. GM17 growth curves of *L. lactis* MG1363 (●), *L. lactis* MG1363*pepT* (▲), *L. lactis* MG1363*acmAΔI* (▼), and a coculture of *L. lactis* MG1363 and *L. lactis* MG1363*acmAΔI* mixed 1:1 at the end of the exponential phase (■). The numbered arrowheads at the top of the figure indicate the points at which samples were taken for further analysis (see Fig. 4).

taining pGK13 or pGKAL1 were diluted 100-fold in fresh medium ($0.5\times$ M17), and the OD_{600} was monitored (Fig. 2). During the exponential growth phase the strains grew equally fast. During the following 70 h of incubation, the reduction in the OD_{600} of MG1363*acmAΔI*(pGKAL1) was much higher than that of MG1363(pGK13). As expected, during the same period, nearly no reduction in OD_{600} was observed with the deletion mutant containing pGK13. Apparently, increased production of AcmA from pGKAL1 (see the inset in Fig. 2) results in a higher reduction of the OD compared to the wild-type situation.

AcmA acts intercellularly. Overnight cultures of MG1363, MG1363*acmAΔI*, and MG1363*pepT* were diluted 100-fold in fresh $0.5\times$ M17 medium, and their growth was monitored (Fig. 3). At the end of the exponential phase of growth, equal amounts of the cultures of the *acmA* and *pepT* deletion mutants were mixed. The presence of AcmA activity (Fig. 4A), the release of proteins into the culture medium (Fig. 4B), and the presence of PepT in the supernatant fractions (Fig. 4C) of all four cultures were monitored during 80 h of incubation at 30°C. The reduction of OD_{600} during the prolonged stationary phase of the mixed culture is nearly equal to that of the cultures of MG1363 and the *pepT* deletion mutant (Fig. 3). The average chain length in the mixed culture was equal to that of the chains of MG1363 and MG1363*pepT*, while the chains were very long in the *acmAΔI* culture (reference 3 and results not shown). As expected, AcmA activity was seen in the supernatants of MG1363 and MG1363*pepT* but was absent in MG1363*acmAΔI*. The supernatant of the mixed culture contains AcmA produced by the *pepT* cells (Fig. 4A). Clearly, the activity in the mixture is lower than that in the *pepT* culture, due to the presence of equal amounts of nonexpressing MG1363*acmAΔI* cells. AcmA produces the typical banding pattern due to proteolytic degradation (3). Autolysis results in the release of proteins into the culture medium of the AcmA-producing strains MG1363 and MG1363*pepT* (Fig. 4B). As the protein banding pattern was the same as that of a cell extract of *L. lactis* (results not shown, but compare with Fig. 7A, lane 5), intracellular proteins are liberated. This was confirmed (Fig. 4C) by the presence of the intracellular peptidase PepT (22) among the proteins released from MG1363. Of course, no PepT antigen was present in MG1363*pepT* in the supernatant fraction of this strain (Fig. 4C) or in the cell extract (results not




FIG. 4. (A) Detection of AcmA activity in renaturing SDS-12.5% PAA gels containing 0.15% autoclaved *M. lysodeikticus* cells. (B) Analysis by SDS-12.5% PAGE of proteins present in culture supernatants. The gels were stained with Coomassie brilliant blue. (C) Detection of PepT among the proteins in the supernatant fractions with PepT-specific antibodies. Only the results for the supernatant fractions of the samples taken as indicated in Fig. 3 are shown. Lanes 1 to 7 correspond to the time points indicated in Fig. 3. The equivalent of 60 μ l (lane 1) or 30 μ l (all other lanes) of supernatant was applied to the gels. Molecular masses (in kilodaltons) of standard proteins are shown in the left margins.




FIG. 5. Effect of mitomycin (1 μ g/ml) addition at time zero on OD₆₀₀ of *L. lactis* MG1363(pGK13) (▲) and *L. lactis* LL302 containing pIR12 (●) or pAL12 (▼). A control culture of *L. lactis* LL302(pAL12) which was not induced by mitomycin is also included (■). The arrowhead (▼) at the top of the figure indicates the time point at which 1-ml samples were taken and processed for the analysis of AcmA activity (Fig. 6) and protein and peptidase antigen (Fig. 7).

shown). MG1363*acmA* Δ I does not autolyze (Fig. 3) and consequently does not release intracellular proteins (Fig. 4B). Although PepT antigen was not found in the supernatant of this culture, it was clearly present in the cell extract of this strain (results not shown). Intracellular proteins, including PepT antigen present in the cells of MG1363*acmA* Δ I, were liberated in the mixed culture (Fig. 4B and C). This must have been caused by AcmA, produced and released from MG1363*pepT*, degrading the cell walls of the MG1363*acmA* Δ I cells. Both the total amount of released proteins and the AcmA activity decreased over time (Fig. 4), probably due to the action of released intracellular proteolytic enzymes.

Induced expression of AcmA. The *acmA* gene lacking its native promoter but retaining its own ribosome binding site was taken from pAL01 and inserted into pIR12 (25). In the resulting plasmid, pAL12 (Fig. 1), expression of *acmA* is repressed by the repressor Rro and is induced by mitomycin (25). Plasmid pAL12 was used to transform *L. lactis* LL302 which contains a copy of the pWV01 *repA* gene on the chromosome (19) to ensure efficient replication of pWV01-derived vectors. All strains used for this experiment grew with the same μ_{\max} and reached similar final OD values in the absence of mitomycin. After 2 to 3 h after mitomycin addition, the OD₆₀₀ of LL302(pAL12) decreased gradually and steadily (Fig. 5). MG1363(pGK13) and LL302(pIR12), the latter of which produces *E. coli* β -galactosidase from the r1t promoter/operator cassette (25), did not show detectable lysis in this assay. Clearly, the addition of 1 μ g of mitomycin per ml resulted in a reduction in the growth rates of all cultures. In MG1363(pGK13) the maximal OD₆₀₀ reached was 1.6, whereas LL302(pIR12) did not reach this OD₆₀₀ and appeared to lyse slightly.

To examine the level to which AcmA was induced, samples were taken 4 h after mitomycin addition and inspected by renaturing SDS-PAGE (Fig. 6). Increased activity of AcmA was present in an *L. lactis* LL302(pAL12) cell extract compared to the amount in cell extracts of *L. lactis* LL302(pIR12). Qualitatively, the same was true for the supernatant fractions of the strains: only *L. lactis* LL302(pAL12) produced enhanced clearing bands at a position corresponding to proteins of approximately 30 kDa, which have previously been shown to be active degradation products of AcmA (3).

Although the level of AcmA production was clearly increased, the OD measurements did not conclusively show that




FIG. 6. Renaturing SDS-PAGE analysis of the autolysin activity of *L. lactis* LL302 containing pIR12 (lanes 1 and 2) and pAL12 (lanes 3 and 4). Samples of 1 ml were taken 4 h after the addition of mitomycin to exponentially growing cells (arrowhead in Fig. 5). Cell extracts (lanes 1 and 3) and supernatant fractions (lanes 2 and 4) were loaded on a 12.5% PAA gel containing 0.15% autoclaved *M. lysodeikticus* cells. Molecular masses of standard proteins (lane M) are shown on the left, and the clearing bands due to mature AcmA (40.3 kDa) activity are indicated on the right, (all in kilodaltons). Clearing bands caused by degradation products of AcmA (3) are indicated by an arrow.

it resulted in cell lysis. To examine this in a more direct way, the supernatant fractions of cultures induced for 4 h were assayed for the presence of intracellular proteins by SDS-PAGE. The results (Fig. 7A) show that only one protein is detectable in uninduced cultures. Most probably, this protein is the previously described major secreted protein, Usp45, of *L. lactis* (40). Upon induction, proteins normally present in the cell extracts only are, to a considerable extent, extruded into the culture medium in the case of *L. lactis*(pAL12) (Fig. 7A, lane 4). To a lesser extent, *L. lactis*(pIR12) released proteins into the supernatant. To ascertain that cytoplasmic proteins were indeed present in the culture medium after mitomycin induction, immunoblots were performed on supernatants of cells carrying the various plasmids. The results in Fig. 7B show that antibodies raised against the cytoplasmic lactococcal peptidase PepO (23) gave a strong signal with the supernatant of *L. lactis*(pAL12) and only a weak one with that fraction of *L. lactis*(pIR12). The reacting band at a position of around 40 kDa is caused by an impurity in the antibody preparation (23).




FIG. 7. (A) Analysis by SDS-12.5% PAGE of proteins present in the culture supernatants of induced (lanes 2 and 4) and noninduced (lanes 1 and 3) *L. lactis* LL302 containing pIR12 (lanes 1 and 2) or pAL12 (lanes 3 and 4). The samples were taken at the time point indicated in Fig. 5. Lane 5 contains a cell extract of noninduced *L. lactis* LL302(pIR12). The gel was stained with Coomassie brilliant blue. Molecular masses (in kilodaltons) of standard proteins (M) are shown on the left. (B) Western blot analysis of the gel shown in panel A with polyclonal antibodies raised against the lactococcal intracellular endopeptidase PepO (23). The arrowhead indicates the position of the endopeptidase.

DISCUSSION

In this work we have clearly shown that AcmA of *L. lactis* is required for autolysis of this organism during stationary phase. Deletion of the *acmA* gene resulted in complete loss of the autolytic behaviour. Autolysis resulted in the release of intracellular proteins, including the intracellular peptidases PepT and PepO. The reduction in OD₆₀₀ of MG1363*acmA*Δ1 was at most 15% during stationary phase. This decrease occurred immediately after the culture had reached its maximum OD₆₀₀. Thereafter, the OD₆₀₀ of the culture remained constant for at least 7 days. The OD reduction was not accompanied by a release of intracellular proteins (Fig. 4B and C), indicating that it is not caused by (auto)lysis. In other words, in *L. lactis* MG1363, AcmA is the only enzyme responsible for autolysis. The initial steep drop in the OD₆₀₀ of approximately 15% after reaching stationary phase was observed in all the strains examined. Since the viable count of MG1363*acmA*Δ1(pGK13) did not change from the point of maximum OD₆₀₀ to 10 h thereafter (unpublished data), the initial OD₆₀₀ reduction has to be explained by general changes in cell morphology and/or intracellular components influencing light scattering and thus reducing OD₆₀₀.

Although Mou et al. (24) and Niskasaari (26) detected only muramidase activity in two strains of *L. lactis*, Østlie et al. (28) have recently shown that three other *L. lactis* strains contained a glucosidase and an *N*-acetylmuramoyl-L-alanine amidase or endopeptidase activity. Also, Crow et al. (6) suggested the presence of more than one autolytic enzyme in lactococci on the basis of activity profiles in renaturing SDS-PAGE activity assays. From the literature, it is clear that autolytic behavior is different among lactococcal strains, and it will be interesting to determine the actual contribution of each of these (putative) enzymatic activities to autolysis. Based on the data presented here and our unpublished results that an active copy of *acmA* is present in more than 15 different (industrial) strains of *L. lactis*, we postulate that AcmA is the only or major enzyme involved in stationary phase autolysis in many, if not all, lactococci.

Loss of autolysis was also seen in other gram-positive bacteria when expression of peptidoglycan hydrolases was prevented. Insertional inactivation of the gene encoding the major autolysin *N*-acetylmuramoyl-L-alanine amidase (*cwlB*) of *B. subtilis* led to loss of approximately 90% of the total cell wall hydrolytic activity of stationary-phase cells. The mutant strain was extremely resistant to cell lysis but did not grow in filaments (15). Interruption of *Streptococcus pneumoniae* *lytA*, the gene encoding *N*-acetylmuramoyl-L-alanine amidase, resulted in loss of autolysis during stationary phase. No significant difference in chain formation was observed between the wild-type and mutant strains (37). Two mutants of *Staphylococcus aureus* showing negligible autolysis during a prolonged stationary phase were created by Tn917-*lacZ* insertion mutagenesis (21). The strains lacked the endo-β-*N*-acetylglucosaminidase (51-kDa) and *N*-acetylmuramoyl-L-alanine amidase (62-kDa) activities, which Sugai et al. (36) later showed were involved in the separation of daughter cells.

Vegarud et al. (44) have shown that changes in the composition of M17 leading to a reduction in maximal OD generally resulted in a reduction of autolysis. This is in agreement with observations made in this study. As detailed in Results, autolysis of MG1363 grown in M17 medium was estimated to be 63%, which is similar to that measured by Østlie et al. for two lactococcal strains grown under the same conditions (29). When MG1363 was grown in 0.5× M17 (Fig. 3), the decrease in OD was only 35%. The difference in autolysis cannot be

explained by a difference in final culture medium pH, an important factor for AcmA activity (reference 24 and unpublished results), because the pHs reached in both media used in this study were 5.2.

AcmA was shown to also act intercellularly, releasing the cellular content of an AcmA-nonproducing strain. Although AcmA is normally attached to the cell wall through its C-terminal repeat domain (reference 3 and unpublished results), the enzyme is apparently not covalently linked. It can be released and can subsequently recognize, bind, and hydrolyze the wall of another cell. This observation opens the possibility of using *L. lactis* for the controlled overexpression of AcmA and adding such a strain to the mixture of strains present in a cheese starter culture. Induction of the *acmA* gene in the adjunct strain could lead to the enhanced lysis of all strains in the starter. Among the proteins released would be flavor-enhancing enzymes. To have such an inducible lysis system at one's disposal could be of great industrial interest. As a first step toward an inducible system for *L. lactis*, *acmA* lacking its own promoter was cloned downstream of the promoter/operator region of the temperate lactococcal bacteriophage r1t. Expression of AcmA from this construct was inducible by the addition of mitomycin. Increased expression of AcmA was observed 4 h after induction of the lactococcal strain containing pAL12. Mitomycin did not induce expression of the chromosomal copy of *acmA*. AcmA induction was much lower than β-galactosidase induction (25) when the same genetic element was used. Among other possibilities, this may be due to factors needed for (extracellular) AcmA activity that become limiting. In this respect, it is interesting that part of an operon which is involved in the secretion of the strongly homologous muramidase-2 of *Enterococcus hirae* was recently cloned and sequenced (9).

A decrease in OD₆₀₀ with the release of intracellular proteins was seen in cultures of the strain overexpressing AcmA, but limited lysis of cells was also observed in the strain overexpressing β-galactosidase. The slow decrease in the OD₆₀₀ of the latter strain may be caused by the production of deleterious quantities of β-galactosidase only or in combination with the presence of mytomycin, a substance which clearly inhibits cell growth.

Although we have successfully overproduced AcmA with concomitant cell lysis, it is clear that the system is not yet optimal and cannot be used for industrial fermentations. Research is currently focused on the isolation of a temperature-sensitive mutant of the repressor (Rro), which would allow us to lyse cells and release important proteins and enzymes in cibo in a food-grade way.

ACKNOWLEDGMENTS

We thank Henk Mulder for preparation of the photographs, Anne de Jong for advice about and support of the computer work, and Jan Willem Sanders for assistance in the RNA work. We thank Aat Ledebor and Wouter Musters for their discussions and suggestions.

This study was supported by Unilever Research Laboratorium, Vlaardingen, The Netherlands. Jan Kok is the recipient of a fellowship of the Royal Netherlands Academy of Arts and Sciences.

REFERENCES

- Béliveau, C., C. Potvin, J. Trudel, A. Asselin, and G. Bellemare. 1991. Cloning, sequencing and expression in *Escherichia coli* of a *Streptococcus faecalis* autolysin. *J. Bacteriol.* **173**:5619–5623.
- Bie, R., and G. Sjöström. 1975. Autolytic properties of some lactic acid bacteria used in cheese production. II. Experiments with fluid substrates and cheese. *Milchwissenschaft* **30**:739–747.
- Buist, G., J. Kok, K. J. Leenhouts, M. Dabrowska, G. Venema, and A. J. Haandrikman. 1995. Molecular cloning and nucleotide sequence of the gene encoding the major peptidoglycan hydrolase of *Lactococcus lactis*, a mur-

- amidase needed for cell separation. *J. Bacteriol.* **177**:1554–1563.
4. Casadaban, M. J., and S. N. Cohen. 1980. Analysis of gene control signals by DNA fusion and cloning in *Escherichia coli*. *J. Mol. Biol.* **138**:179–207.
 5. Chapot-Chartier, M.-P., C. Deniel, M. Rousseau, L. Vassal, and J.-C. Gripon. 1994. Autolysis of two strains of *Lactococcus lactis* during cheese ripening. *Int. Dairy J.* **4**:251–269.
 6. Crow, V. L., T. Coolbear, P. K. Gopal, F. G. Martley, L. L. McKay, and H. Riepe. 1995. The role of autolysis of lactic acid bacteria in the ripening of cheese. *Int. Dairy J.* **5**:855–875.
 7. Crow, V. L., T. Coolbear, R. Holland, G. G. Pritchard, and F. G. Martley. 1993. Starters as finishers: starter properties relevant to cheese ripening. *Int. Dairy J.* **3**:423–460.
 8. Dako, E., M. El Soda, J.-C. Vuilleumard, and R. E. Simard. 1995. Autolytic properties and aminopeptidase activities of lactic acid bacteria. *Food Res. Int.* **28**:503–509.
 9. Del Mar Lleò, M., R. Fontana, and M. Solioz. 1995. Identification of a gene (*arpU*) controlling muramidase-2 export in *Enterococcus hirae*. *J. Bacteriol.* **177**:5912–5917.
 10. de Vos, W. M., and G. Simons. 1994. Gene cloning and expression systems in lactococci, p. 52–105. *In* M. J. Gasson and W. M. de Vos (ed.), Genetics and biotechnology of lactic acid bacteria. Blake Academic and Professional, London, United Kingdom.
 11. Feirtag, J. M., and L. L. McKay. 1987. Isolation of *Streptococcus lactis* C2 mutants selected for temperature sensitivity and potential use in cheese manufacture. *J. Dairy Sci.* **70**:1773–1778.
 12. Gasson, M. J. 1983. Plasmid complements of *Streptococcus lactis* NCDO 712 and other lactic streptococci after protoplast-induced curing. *J. Bacteriol.* **154**:1–9.
 13. Kok, J. 1992. Special-purpose vectors for lactococci, p. 97–102. *In* G. Dunny, P. P. Cleary, and L. L. McKay (ed.), Genetics and molecular biology of streptococci, lactococci, and enterococci. American Society for Microbiology, Washington, D.C.
 14. Kok, J., and W. M. de Vos. 1994. The proteolytic system of lactic acid bacteria, p. 169–210. *In* M. J. Gasson and W. M. de Vos (ed.), Genetics and biotechnology of lactic acid bacteria. Blake Academic and Professional, London, United Kingdom.
 15. Kuroda, A., and J. Sekiguchi. 1991. Molecular cloning and sequencing of a major *Bacillus subtilis* autolysin gene. *J. Bacteriol.* **173**:7304–7312.
 16. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680–685.
 17. Langsrud, T., A. Landaas, and H. B. Castberg. 1987. Autolytic properties of different strains of group N streptococci. *Milchwissenschaft* **42**:556–560.
 18. Law, B. A., M. E. Sharpe, and B. Reiter. 1974. The release of intracellular dipeptidase from starter streptococci during Cheddar cheese ripening. *J. Dairy Sci.* **41**:137–146.
 19. Leenhouts, K., G. Buist, A. Bolhuis, A. ten Berge, J. Kiel, I. Mierau, M. Dabrowska, G. Venema, and J. Kok. 1996. A general system for generating unlabelled gene replacement in the bacterial chromosomes. *Mol. Gen. Genet.* **253**:217–224.
 20. Leenhouts, K. J., and G. Venema. 1993. Lactococcal plasmid vectors, p. 65–94. *In* K. G. Hardy (ed.), Plasmids, a practical approach. Oxford University Press, Oxford, United Kingdom.
 21. Mani, N., P. Tobin, and R. K. Jayaswal. 1993. Isolation and characterisation of autolysis-defective mutants of *Staphylococcus aureus* created by Tn917-*lacZ* mutagenesis. *J. Bacteriol.* **175**:1493–1499.
 22. Mierau, I., A. J. Haandrikman, O. Velterop, P. S. T. Tan, K. J. Leenhouts, W. N. Konings, G. Venema, and J. Kok. 1994. Tripeptidase gene (*pepT*) of *Lactococcus lactis*: molecular cloning and nucleotide sequencing of *pepT* and construction of a chromosomal deletion mutant. *J. Bacteriol.* **176**:2854–2861.
 23. Mierau, I., P. S. T. Tan, A. J. Haandrikman, J. Kok, K. J. Leenhouts, W. N. Konings, and G. Venema. 1993. Cloning and sequencing of the gene for a lactococcal endopeptidase, an enzyme with sequence similarity to mammalian enkephalinase. *J. Bacteriol.* **175**:2087–2096.
 24. Mou, L., J. J. Sullivan, and G. R. Jargo. 1976. Autolysis of *Streptococcus cremoris*. *J. Dairy Res.* **43**:275–282.
 25. Nauta, A., D. van Sinderen, H. A. Karsens, E. Smit, G. Venema, and J. Kok. 1996. Inducible gene expression mediated by a repressor-operator system isolated from *Lactococcus lactis* bacteriophage r1t. *Mol. Microbiol.* **19**:1331–1341.
 26. Niskasaari, K. 1989. Characteristics of the autolysis of variants of *Lactococcus lactis* subsp. *cremoris*. *J. Dairy Res.* **56**:639–649.
 27. Olson, N. F. 1991. The impact of lactic acid bacteria on cheese flavour. *FEMS Microbiol. Rev.* **87**:131–148.
 28. Østlie, H. M., G. Vegarud, and T. Langsrud. 1995. Autolysis of lactococci: detection of lytic enzymes by polyacrylamide gel electrophoresis and characterization in buffer systems. *Appl. Environ. Microbiol.* **61**:3598–3603.
 29. Østlie, H. M. 1995. Autolysis of lactococci and dairy propionibacteria. Ph.D. thesis. Agricultural University of Norway, Ås.
 30. Rogers, H. J., H. R. Perkins, and J. B. Ward. 1980. The bacterial autolysins, p. 437–460. *In* H. J. Rogers, H. R. Perkins, and J. B. Ward (ed.), Microbial cell walls and membranes. Chapman & Hall, Ltd., London, United Kingdom.
 31. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
 32. Sanders, J. W., G. Venema, J. Kok, and K. Leenhouts. Unpublished data.
 33. Sanger, F. S., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
 34. Segers, J. F. M. L., S. Bron, C. M. Franke, G. Venema, and R. Kiewiet. 1994. The majority of lactococcal plasmids carry a highly related replicon. *Microbiology* **140**:1291–1300.
 35. Shearman, C. A., K. Jurry, and M. J. Gasson. 1992. Autolytic *Lactococcus lactis* expressing a lactococcal bacteriophage lysin gene. *Bio/Technology* **10**:196–199.
 36. Sugai, M., H. Komatsuzawa, T. Akiyama, Y.-M. Hong, T. Oshida, Y. Miyake, T. Yamaguchi, and H. Suganaka. 1994. Identification of endo- β -N-acetylglucosaminidase and N-acetylmuramyl-L-alanine amidase as cluster-dispersing enzymes in *Staphylococcus aureus*. *J. Bacteriol.* **177**:1491–1496.
 37. Tomasz, A., P. Moreillon, and G. Pozzi. 1988. Insertional inactivation of the major autolysin gene of *Streptococcus pneumoniae*. *J. Bacteriol.* **170**:5931–5934.
 38. Towbin, H., T. Staehelin, and J. Gordon. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. USA* **76**:4350–4354.
 39. Van Asseldonk, M., A. Simons, H. Visser, W. M. de Vos, and G. Simons. 1993. Cloning, nucleotide sequence, and regulatory analysis of the *Lactococcus lactis dnaJ* gene. *J. Bacteriol.* **175**:1637–1644.
 40. Van Asseldonk, M., G. Rutten, M. Oteman, R. J. Siezen, W. M. de Vos, and G. Simons. 1990. Cloning of *usp45*, a gene encoding a secreted protein from *Lactococcus lactis* subsp. *lactis* MG1363. *Gene* **95**:155–160.
 41. van de Guchte, M., J. Kodde, J. M. B. M. van der Vossen, J. Kok, and G. Venema. 1990. Heterologous gene expression in *Lactococcus lactis* subsp. *lactis*: synthesis, secretion, and processing of the *Bacillus subtilis* neutral protease. *Appl. Environ. Microbiol.* **56**:2606–2611.
 42. Van Sinderen, D., G. Galli, P. Cosmina, F. De Ferra, S. Withoff, G. Venema, and G. Grandi. 1993. Characterization of the *srfA* locus of *Bacillus subtilis*: only the valine-activating domain of *srfA* is involved in the establishment of genetic competence. *Mol. Microbiol.* **8**:833–841.
 43. Van Sinderen, D., H. A. Karsens, J. Kok, P. Terpstra, M. H. J. Ruiters, G. Venema, and A. Nauta. 1996. Sequence analysis and molecular characterization of the temperate lactococcal bacteriophage r1t. *Mol. Microbiol.* **19**:1343–1355.
 44. Vegarud, G., H. B. Castberg, and T. Langsrud. 1983. Autolysis of group N streptococci. Effects of media composition modifications and temperature. *J. Dairy Sci.* **66**:2294–2302.
 45. Visser, S. 1993. Proteolytic enzymes and their relation to cheese ripening and flavour: an overview. *J. Dairy Sci.* **76**:329–350.
 46. Wilkinson, M. G., T. P. Guinee, D. M. O'Callaghan, and P. F. Fox. 1994. Autolysis and proteolysis in different strains of starter bacteria during Cheddar cheese ripening. *J. Dairy Res.* **61**:249–262.
 47. Zabarovsky, E. R., and G. Winberg. 1990. High efficiency electroporation of ligated DNA into bacteria. *Nucleic Acids Res.* **18**:5912.