

University of Groningen

## Star formation and the interstellar medium in galaxy simulations

Gerritsen, Jeroen Peter Erik

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1997

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Gerritsen, J. P. E. (1997). *Star formation and the interstellar medium in galaxy simulations*. s.n.

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# 4

---

## Why LSB galaxies are blue, thin and poor in molecular gas

J.P.E. Gerritsen & W.J.G. de Blok

We present  $N$ -body simulations of Low Surface Brightness (LSB) galaxies and their interstellar medium to investigate the cause for their low star formation rates (SFRs). Due to their massive halos, stellar disks of LSB galaxies are very stable and thin. Lack of dust makes the projected edge-on surface brightness of LSB galaxies comparable to the projected edge-on surface brightness of dust-rich High Surface Brightness (HSB) galaxies of similar size. We show that the low surface densities found in LSB galaxies are by themselves not enough to explain the slow evolution of LSB galaxies. A low metal content of the gas is essential. As a consequence the gas cools inefficiently, resulting in an almost negligible cold gas fraction. We show that LSB galaxies must have molecular gas fractions of less than 5 per cent. Our best model has a SFR which is on average low but fluctuates strongly. This causes the large spread in colors of LSB galaxies. From a distribution of birthrate parameters we conclude that the presently-known and modeled gas-rich blue LSB galaxies constitute the majority of the total population of gas-rich LSB disk galaxies. We deduce the existence of an additional red, quiescent and gas-rich population which constitutes less than 20 per cent of the total population. This does not rule out the existence of a large number of gas-poor LSB galaxies. These must however have had an evolutionary history dramatically different from that of the gas-rich galaxies.

DEEP surveys of the night-sky have uncovered a large population of disk galaxies with properties quite different from those of the extensively studied “normal” high surface brightness (HSB) galaxies. These so-called Low Surface Brightness (LSB) galaxies have, as their name already implies, surface brightnesses much lower than what was until quite recently assumed to be representative for disk galaxies (Freeman 1970). The LSB galaxies we will be discussing are generally dominated by an exponential disk, with scale lengths of a few kpc. Morphologically they form an extension of the Hubble sequence towards very late-type galaxies. Observations suggest that LSB galaxies are unevolved galaxies, as shown by e.g. the sub-

solar metallicities (McGaugh 1994) and the colors (McGaugh & Bothun 1994; De Blok, Van der Hulst & Bothun 1995). The evolutionary rate of a galaxy may in fact be reflected in its surface brightness. For example, the gas fraction ( $M_{\text{gas}}/M_{\text{gas+stars}}$ ) increases systematically with surface brightness, from a few percent for early type spirals to much higher values approaching unity for late-type LSB galaxies (De Blok, McGaugh & Van der Hulst 1996).

In many LSB galaxies the gas mass exceeds the stellar mass (even for such extreme assumptions as maximum disk). It is still unclear what the physical driver is for the difference between HSB and LSB galaxies. Investigations of their dynamics, using HI ob-

servations (De Blok, McGaugh & Van der Hulst 1996), suggest that LSB galaxies are low-density galaxies (De Blok & McGaugh 1996). This is one of the favored explanations for the low evolution rate of LSB galaxies (see e.g. Van der Hulst et al. 1987), as this implies a large dynamical time-scale.

The main purpose of this paper is to investigate whether the low density alone can explain the properties of LSB galaxies. We use 3D numerical simulations to address this problem. The dynamical simulations include both stars and gas and incorporate a parameterized description of star formation, including feedback on the gas.

The prescription of star formation in numerical simulations is not straightforward, given the limited resolution of the models and the fragmentary knowledge about the physics governing star formation on kpc scales. In the literature various different star formation algorithms can be found. For instance, Friedly & Benz (1995) use a criterion based on the Toomre stability parameter  $Q$  to study star formation in barred systems; Mihos & Hernquist (1994a, 1994b, 1996) adopt a Schmidt law based on the gas density (star formation rate (SFR)  $\propto \rho^{1.5}$ ) to study mergers of galaxies; other authors require gas to be in a convergent and Jeans unstable flow to form stars (Katz 1992; Katz, Weinberg & Hernquist 1996; Navarro & White 1993; Steinmetz 1996)

We employ the method of Gerritsen & Icke (1997a and b, Chapters 2 and 3 respectively), which uses a Jeans criterion to define star forming regions, coupled with an estimate of the cloud collapse time. In these simulations gas is treated fully radiative with allowed temperatures between 10 K and  $10^7$  K; cooling is described by standard cooling functions, heating is assumed to be provided by far-ultraviolet (FUV) radiation and mechanical heating from stars. In this way the simulated interstellar medium (ISM) mimics a multi-phase ISM (Field, Goldsmith & Habing 1996; McKee & Ostriker 1997). This is an improvement over other simulations which try to create a multi-phase ISM but do

not allow radiative cooling below  $10^4$  K (e.g. Hernquist 1989, Katz, Weinberg & Hernquist 1996). The multi-phase ISM allows us to restrict ourselves to considering only relatively cold ( $T < 10^3$  K) regions as the sites for star formation.

Here we apply this method to study the ISM and star forming properties of LSB galaxies, and our results will be valid under the assumption that the physical processes governing star formation are the same for HSB and LSB galaxies. We test whether a low mass-density is sufficient to explain the properties of LSB galaxies by discussing two different models of a specific LSB galaxy: the first model has the cooling properties of a solar abundance ISM, while in the second model the cooling efficiency is lowered, thus mimicking a metal-poor ISM. The main result is that low density alone is not enough to correctly model the current properties of LSB galaxies, but that low metallicities in the gas are essential.

The structure of the paper is as follows. In Section 4.1 we describe the numerical techniques. The construction of a model LSB galaxy is presented in Section 4.2. This model galaxy is evolved for a few Gyr, using two different prescriptions for the cooling properties of the gas (Section 4.3). The implications of the simulations are discussed in Section 4.4. We conclude this paper with a summary in Section 4.5.

## 4.1 Numerical technique

We model the evolution of galaxies using a hybrid  $N$ -body/hydrodynamics code (TREESPH; Hernquist & Katz 1989). A brief summary of the techniques pioneered by these authors follows. An extensive description is given in Chapters 2 and 3. A tree algorithm (Barnes & Hut 1996; Hernquist 1987) determines the gravitational forces on the collisionless and gaseous components of the galaxies. The hydrodynamic properties of the gas are modeled using smoothed particle hydrodynamics (SPH; Lucy 1977, Gingold & Monaghan 1977). The gas evolves accord-

ing to hydrodynamic conservation laws, including an artificial viscosity for an accurate treatment of shocks. Each particle is assigned an individual smoothing length,  $h$ , which determines the local resolution and an individual time step. Estimates of the gas properties are found by smoothing over 32 neighbors within  $2h$ . For example, the smoothed density is given by

$$\rho(\vec{r}) = \sum_{j=1}^{32} m_j W(\vec{r} - \vec{r}_j, h), \quad (4.1)$$

where  $W(\vec{r}, h)$  is the smoothing kernel and  $m_j$  is the particle mass.

We adopt the equation of state

$$P = (\gamma - 1)\rho u, \quad (4.2)$$

where  $P$  is the pressure,  $\rho$  the density,  $u$  the thermal energy density, and  $\gamma = 5/3$  for an ideal gas.

In the Lagrangian formulation of SPH energy conservation can be expressed as

$$\frac{du}{dt} = -\frac{P}{\rho} \nabla \vec{v} + (\Gamma - \rho\Lambda), \quad (4.3)$$

where  $\vec{v}$  is the velocity, and  $(\Gamma - \rho\Lambda)$  represents the radiative heating and cooling processes.

We allow radiative cooling of the gas according to the cooling function for a standard hydrogen gas mix with a helium mass fraction of 0.25 (Dalgarno & McCray 1972). Radiative heating is modeled as photo-electric heating of small grains and PAHs by the FUV field (Wolfire et al. 1995), produced by the stellar distribution.

All our simulations are advanced in time steps of  $3 \times 10^6$  yr for star particles, while the time steps for SPH particles can be 8 times shorter ( $3.8 \times 10^5$  yr). The gravitational softening length is 150 pc. A tolerance parameter  $\theta = 0.6$  is used for the force calculation, which includes quadrupole moments.

#### 4.1.1 Star formation and feedback

Chapters 2 and 3 extensively describe the recipe for transforming gas into stars and the

method for supplying feedback onto the gas. The recipe works well for normal HSB galaxies, with the energy budget of the ISM as prime driver for the star formation. The simulations allow for a multi-phase ISM with temperature between  $10 < T < 10^7$  K. This allows us to consider cold  $T < 10^3$  K regions as places for star formation (giant molecular clouds in real life). Below we summarize the important ingredients of the method.

From our SPH particle distribution we select conglomerates where the Jeans mass is below the mass of a typical giant molecular cloud. In the simulations performed here we use  $M_c = 10^6 M_\odot$  for this aggregate mass (the method is not very sensitive to the exact value for this mass). The Jeans mass depends on the local gas properties and is calculated from the SPH estimates of the density  $\rho$  and sound speed  $s$ ,

$$M_J = \frac{1}{6} \pi \rho \left( \frac{\pi s^2}{G\rho} \right)^{\frac{3}{2}}, \quad (4.4)$$

with  $G$  the constant of gravity. During the simulations the maximum number density achieved is of the order  $1 \text{ cm}^{-3}$ . It follows that only regions below  $10^3$  K are unstable and may form stars.

We follow unstable regions during their dynamical and thermal evolution and if an SPH particle resides in such a region longer than the collapse time,

$$t_u > t_c = \frac{1}{\sqrt{4\pi G\rho}}, \quad (4.5)$$

half of its mass is converted into a star particle.

Important for our calculations is that we consider star particles as stellar clusters with an age. Thus for each individual star particle we can attribute quantities like the SN-rate, the mass loss, and the FUV-flux, according to its age. We use the spectral synthesis models of Bruzual & Charlot (1993) to determine these quantities, where we adopt a Salpeter initial mass function (IMF) with slope 1.35 and with lower and upper mass limits of  $0.1 M_\odot$  and  $125 M_\odot$  respectively.

The radiative heating for a gas particle is calculated by adding the FUV-flux contributions from all stars, which is done together with the force calculation. The mechanical luminosity from a star particle is determined by both the SN-rate and the mass loss rate. We assume that each SN injects  $10^{51}$  ergs of energy and that the energy injected by stellar winds is  $E_w = \frac{1}{2}\Delta m_* v_\infty^2$ , with  $v_\infty$  the wind terminal velocity and  $\Delta m_*$  the stellar mass loss. For massive stars  $v_\infty$  depends critically on the stellar mass, luminosity, effective temperature, and metallicity (e.g. Leitherer, Robert & Drissen 1992; Lamers & Leitherer 1993). For simplicity we adopt  $v_\infty = 2000$  km/s. After  $3.3 \times 10^7$  yr the last  $8 M_\odot$  stars explode and no more mechanical energy is supplied to the gas. Thus we return mechanical energy from massive stars into the ISM, and ignore the mechanical energy from low mass stars.

In the simulations the parent SPH particle is the carrier of the mechanical energy from the new star particle. This SPH particle (“SN particle”) mimics a hot bubble interior. Radiative cooling is temporarily switched off (the resolution does not allow the creation of a low-density, hot bubble), the temperature of the SN particle is set to the mechanical energy density (of a few  $10^6$  K), and the particle evolves adiabatically. The first  $10^7$  yr, the position and velocity of the SN particle are equal to the position and velocity of the associated star particle. Afterwards, the SN particle evolves freely. After  $3 \times 10^7$  yr radiative cooling is switched on again, and the particle behaves like an ordinary SPH particle.

## 4.2 Modeling an LSB galaxy

There are many ways to construct model galaxies. For our purpose we prefer to build a model galaxy after an existing galaxy with well-determined properties. Here we choose LSB galaxy F563-1; this galaxy is a late-type LSB galaxy, representative of the field LSB galaxies found in the survey by Schombert et al. (1992). The optical properties of this galaxy are described in De Blok, Van der Hulst & Bothun (1995); measurements of

TABLE 4.1— Parameters for F563-1

$L_B$	$1.35 \times 10^9 L_\odot$	
$h_*$	2.8 kpc	center ( $R < 5$ kpc)
$h_*$	5.0 kpc	outside ( $R > 5$ kpc)
SFR	$0.05 M_\odot/\text{yr}$	
$M_{\text{HI}}$	$2.75 \times 10^9 M_\odot$	
$v_{\text{max}}$	113 km/s	
$\rho_0^{\text{halo}}$	$0.0751 M_\odot \text{pc}^{-3}$	
$R_c^{\text{halo}}$	1.776 kpc	

metallicities in H II regions are described in De Blok & Van der Hulst (1997); a neutral hydrogen map and rotation curve are given in De Blok, McGaugh & Van der Hulst (1996). Parameters such as stellar velocity dispersion, which cannot be measured directly, are set in comparison with values measured locally in the Galaxy. The current SFR as deduced from H $\alpha$  imaging is taken from Van den Hoek et al. (1997). For convenience these data are summarized in Table 4.1.

The most difficult problem we face in constructing a model is converting the measured luminosity to a stellar disk mass. This is one of the most persistent problems in analyzing the dynamics of galaxies, and, unfortunately, the present observations do not provide a unique answer for this stellar disk mass-to-light ratio  $(M/L_B)_*$ . Rather than using the so-called “maximum disk” value  $(M/L_B)_* = 9$ , which is an upper limit to the possible values of  $(M/L_B)_*$ , we adopt a value based on colors and velocity dispersions of  $(M/L_B)_* = 1.75$ . An extensive motivation for this choice is given by De Blok & McGaugh (1997). The implications of choosing a different value of  $(M/L_B)_*$  for the evolution of the stellar disk will be discussed later.

### 4.2.1 The model

The stellar disk particles are distributed radially according to the (measured) surface density profile. We adopt a vertical distribution of the form  $\text{sech}^2(z/z_*)$  (e.g. Van der Kruit & Searle 1982), with a constant vertical scale height  $z_* = 0.5$  kpc. The disk is truncated at 25 kpc. The luminosity yields a total stellar mass of  $M_* = 2.36 \times 10^9 M_\odot$ . The velocity

dispersion of the stars is fixed via the relation

$$\sigma_z = \sqrt{\pi G z_* \Sigma_*}, \quad (4.6)$$

where  $\Sigma_*$  is the stellar surface density and  $G$  the gravitational constant. This implies a stability parameter  $3 < Q < 3.5$  throughout for the disk (dotted line in Fig. 4.2).

Particle velocities are assigned according to the (gas) rotation curve corrected for asymmetric drift. Dispersions in  $R, z, \theta$  directions are drawn from Gaussian distributions with dispersions of  $\sigma_R, \sigma_z, \sigma_\theta$  respectively, using the relations valid for the solar neighborhood

$$\sigma_z = 0.6 \sigma_R, \quad \sigma_\theta = \frac{\kappa}{2\omega} \sigma_R, \quad (4.7)$$

with  $\omega$  and  $\kappa$  the orbital and epicyclic frequencies respectively.

The gas surface density distribution is modeled after the HI distribution. The total gas mass is  $M_g = 3.85 \times 10^9 M_\odot$  (this is the total HI mass multiplied by 1.4 to include He); the surface density decreases almost linearly with radius out to 33 kpc. The vertical distribution is assumed to decline exponentially. The scale height  $z_g$  of the gas layer can be calculated using

$$\sigma_g^2 = 2\pi G(\Sigma_g + \Sigma_*) \frac{z_g^2}{\frac{1}{2}z_* + z_g}, \quad (4.8)$$

(Dopita & Ryder 1994) where  $\sigma_g = 3$  km/s is the (adopted) gas velocity dispersion. The gas particles are assigned velocities according to the rotation curve, with isotropic dispersion  $\sigma_g$ .

The halo is included in the calculations as a rigid potential. This is justified since the model galaxy evolves in isolation. We will thus also ignore any contraction of the halo under the influence of the disk potential. We do not expect this effect to be important anyway as the mass of the disk (assuming  $(M/L_B)_* = 1.75$ ) is only 4 per cent of the measured halo mass (De Blok & McGaugh 1997). As advantages we do not have to make assumptions about halo particle orbits, and we do not have to spend time in calculating

the force of the halo particles on the galaxy. We assign an isothermal density distribution to the halo,

$$\rho_h = \frac{\rho_0}{1 + (r/R_c)^2}, \quad (4.9)$$

with central volume density  $\rho_0 = 0.0751 M_\odot \text{pc}^{-3}$  and core radius  $R_c = 1.776$  kpc. These are the values derived from a rotation curve decomposition assuming  $(M/L_B)_* = 1.75$ . This will correctly put the maximum rotation velocity at 113 km/s.

For the simulations we use 40,000 SPH particles and 80,000 star particles initially. This corresponds to an SPH particle mass of  $9.6 \times 10^4 M_\odot$  and a star particle mass of  $3.0 \times 10^4 M_\odot$ .

#### 4.2.2 Implications of $(M/L_B)_*$ for star formation history

We can show that a maximum disk value for  $(M/L_B)_*$  is not a plausible option for our modeling exercise, and that in fact the “most likely” value we use in constructing models may still overestimate the true value of  $(M/L_B)_*$ , making the stellar disk a really insignificant component of the whole galaxy system.

The star formation history of galaxies is commonly parameterized by an exponential function:

$$\text{SFR} = -\dot{M}_g = M_g/\tau_* \quad (4.10)$$

where  $\tau_*$  is the star formation time scale (e.g. Guiderdoni & Rocca-Volmerange 1987 and Charlot & Bruzual 1991). This ignores any gas locked up in long-lived stars, but this effect will not be important in LSB galaxies due to their large gas-fractions. A small value for  $\tau_*$  means that star formation proceeds rapidly, while a large  $\tau_*$  indicates that star formation proceeds very slowly.

For HSB galaxies the star formation time scale is normally a few Gyr (Kennicutt, Tamblyn & Congdon 1994). For F563-1 the measured current SFR as derived from H $\alpha$  imaging is approximately  $0.05 M_\odot/\text{yr}$ ; this yields a star formation time scale of 77 Gyr.

Thus star formation in this disk proceeds much slower than in HSB galaxies.

Since we know the stellar and gas mass, and hence the total disk mass  $M_t = M_* + M_g$  of the galaxy, we can use the exponential parameterization and the known SFR to compute the “age”  $t_d$  of the disk (i.e. the time elapsed since star formation started):

$$M_* = - \int_0^{t_d} \dot{M}_g dt = M_t \left( 1 - e^{-t_d/\tau_*} \right). \quad (4.11)$$

For the adopted stellar mass of F563-1 Eq. 4.11 yields an age for the galaxy of 37 Gyr, much larger than the age of the universe.

If the exponentially declining SFR describes the star formation history adequately and the galaxy is not in a state of unusually low star formation activity (note that the blue colors for these LSB galaxies suggest a relatively *high* current star formation activity), then the old implied age of the disk could mean that the mass of the disk should in reality be even less.

We show this in Fig. 4.1 where we present a few solutions for Eqs. 4.10 and 4.11 for various values of the SFR. If we adopt  $0.05 M_\odot/\text{yr}$  as the true SFR then  $(M/L_B)_* \approx 0.6$  if the galaxy is 15 Gyr old. Higher values for  $(M/L_B)_*$  would make the galaxy older than the universe. A maximum disk solution for the stellar mass-to-light ratio (Van Albada & Sancisi 1986) yields  $(M/L_B)_* = 9$ , which leads to an age of 110 Gyr. A current SFR of order  $0.4 M_\odot/\text{yr}$  is needed to reconcile the maximum disk with the exponentially declining SFR model, which implies that the measurements underestimate the true SFR by an order of magnitude. Thus seems highly unlikely (it would give LSB galaxies SFRs comparable to those of actively star forming late type galaxies), and therefore we reject the maximum disk as an acceptable solution for the stellar disk mass.

The adopted  $(M/L_B)_* = 1.75$  which we use in the galaxy model is only consistent with the age of the universe if the true SFR is of order  $0.10 M_\odot/\text{yr}$ , otherwise the stellar mass is




FIGURE 4.1— Relation between galaxy age,  $t_d$  (vertical axis), and stellar mass,  $M_*$  (horizontal axis), according to Eqs. 4.10 and 4.11. The different lines are solutions to these equations for various values of the current SFR, where the measured value is  $0.05 M_\odot/\text{yr}$ . The top axis shows the  $(M/L_B)_*$  ratio corresponding to the stellar mass. The input model assumes  $(M/L_B)_* = 1.75$ , the maximum disk model has  $(M/L_B)_* = 9$ , which yields a very high age for the galaxy.

probably overestimated. However if the stellar disk is indeed less massive than our model galaxy, then any conclusions concerning the disk stability and thickness of the disk will be even stronger.

It might be argued that the exponential star formation history is not the right choice for this type of galaxy, and that other functional forms of star formation history, in combination with the measurements, will yield reasonable ages. This is not true, as any other (reasonable) star formation history will result in even larger ages for the disk. Taking for example another extreme functionality – a constant SFR of  $0.05 M_\odot/\text{yr}$  – yields an age  $t_d = 47$  Gyr. The only way to get a reasonable age, given the measured luminosity and assumed  $(M/L_B)_*$ , is to assume that the current SFR in F563-1 (and LSB galaxies in general) has been underestimated, which is unlikely. We refer to McGaugh & De Blok (1997) for an extensive discussion on the various functionalities of the star formation history. We there-

fore conclude that the disk of F563-1 must have a small stellar mass-to-light ratio.

### 4.3 Evolution

In this section we demonstrate that the low density as found in LSB galaxies, by itself is not sufficient to reproduce the low observed SFRs of LSB galaxies. *Low metallicity gas is required to explain the properties of LSB galaxies.*

To show this we construct two model galaxies using the structural parameters of F563-1. Model *H* represents an LSB galaxy with a solar metallicity gas. Although we use the structural parameters relevant for F563-1, the model is in effect a model HSB galaxy, which is “stretched out” to give the low (surface) densities found in LSB galaxies. This model therefore tests the low-density hypothesis.

The other model, *L*, has the same structural parameters as *H*, but in addition we lowered the cooling efficiency of the gas below  $10^4$  K by a factor of seven. Cooling below  $10^4$  K is dominated by metals, so lowering the efficiency is equivalent to lowering the metallicity by an equal amount. Model *L* thus most closely approximates what is currently known observationally about LSB galaxies.

We assume that the physics regulating the star formation is the same for LSB and HSB galaxies, and therefore use the star formation recipe described in Section 4.1 and applied to HSB galaxies in Chapters 2 and 3. After initialization we let the two model galaxies evolve for 2.3 Gyr. The models rapidly settled into equilibrium, any longer time interval would have produced identical results.

In this section we first discuss the evolution of the stellar disk. The stellar disk evolves rather independently from the gas disk and star formation. Hence we effectively explore the consequences of the disk/halo decomposition, notably on the stability and thickness of the stellar disk. Second, we discuss the evolution of the SFR with time for both simulations. For a physical interpretation of the difference in SFRs we present phase diagrams of the ISM in both simulations.




FIGURE 4.2— The stability parameter  $Q$  versus radius. The dotted line shows the  $Q$ -value upon initialization of the model galaxy. The solid line denotes simulation *L*, the dashed line simulation *H*. Both simulation yield approximately the same stability for the stellar disk.

#### 4.3.1 Stellar disk

Figure 4.2 shows the stability parameter  $Q$  for the stellar disk at  $t = 2$  Gyr (that is 2 Gyr after the start of the simulation which started at  $t = 0$ ), where  $Q$  is defined as

$$Q = \frac{\kappa \sigma_R}{3.36 G \Sigma_*}, \quad (4.12)$$

with  $\kappa$  the epicycle frequency (Toomre 1964).  $Q < 1$  denotes a (local) instability, while  $Q > 1$  means stability. For both simulations  $Q \approx 2.5$  in the center and rises to  $Q \approx 5$  in the outer parts of the disk. There is no evolution in the stability. Once the system has settled after the start of the simulation, the radial behavior of  $Q$  as shown is reached. The value of course depends on the input parameters, notably the stellar mass, but it is clear that the stellar disks of LSB galaxies are more stable than the stellar disks of HSB galaxies, where  $Q$  is of order 2 (Bottema 1993, Van der Hulst et al. 1993).

In both simulations the scale height of the stellar disk decreases by approximately a factor of 1.3. The initial scale height corresponded to the scale height of an Sc galaxy (Van der Kruit & Searle 1982). The final axial ratio for the model galaxy is about 15. As particle scattering during the simulation






FIGURE 4.3— The evolution of the SFR versus time. The lower line shows the SFR from simulation *L*, the upper line represents simulation *H*.

tends to increase the thickness of the disk, we conclude that the stellar disks of LSB galaxies are thinner than those of HSB galaxies. We will return to this in more detail in Section 4.4.

#### 4.3.2 Star formation & ISM

Figure 4.3 shows the SFR versus time. In model *H* the SFR settles after a short adjustment at a value of  $0.33 M_{\odot}/\text{yr}$  and declines slowly with time. The SFR in model *L* settles at  $0.08 M_{\odot}/\text{yr}$ .

For both simulations the SFR varies on time scales of a few tens of Myr and the amplitude of these variations can exceed  $0.1 M_{\odot}/\text{yr}$ . The rapid variability in the SFR is due to the discrete nature of star formation in our simulation. New star particles have a mass of approximately  $5 \times 10^4 M_{\odot}$  and thus represent (large) stellar clusters. In this way our simulations incorporate the idea that most stars in real galaxies form in stellar clusters. For a small number of star forming regions (as in LSB galaxies) this will in real life also lead to a rapidly fluctuating, but on average low SFR.

In simulation *L* the strength of the star formation peak is large compared to the average SFR. These fluctuations in star formation activity will actually dominate the color of the galaxy, with large fluctuations giving rise to blue colors. The same star formation

fluctuations will also be present in HSB galaxies, but due to the higher SFR the large number of fluctuations will give the impression of a high, relatively constant SFR.

We thus find that it is not so much the absolute value of the average SFR which determines the colors, but the contrast of any SFR fluctuation with respect to the average SFR. For the LSB galaxies the large contrast leads to blue colors.

In order to understand the behavior of the star formation activity in the simulations we constructed phase diagrams of the ISM. Figure 4.4 shows the temperature versus density for all gas particles in both simulations. The top panel shows simulation *L* while the bottom panel shows the phase diagram for simulation *H*. Most particles have a temperature of  $10^4$  K. In simulation *H* a large fraction of the particles has a temperature of about 100 K. This simulation shows a dominant two-phase structure, and resembles the ISM in simulations of HSB galaxies (Chapter 3). Quantitatively, the cold gas fraction ( $T < 1000$  K) makes up 37 per cent of the total gas mass in simulation *H* and only 4 per cent in simulation *L*. This reflects the cooling properties of the gas: in simulation *L* seven times less heat input is required to keep the gas at  $10^4$  K as in simulation *H*.

Consequently simulation *L* contains virtually only a warm, one-phase ISM. The simulated absence of metals prevents the ISM from cooling efficiently. As molecular gas is only formed in/from the cold component of the ISM, the 4 per cent mentioned above is actually an upper limit to the amount of molecular gas that could form in such a galaxy. The amount of cold molecular gas in sites for star formation would thus be negligible. From these models we expect the disks of LSB galaxies to contain only negligible amounts of molecular gas. This is consistent with observations by Schombert et al. (1990) and De Blok & Van der Hulst (1997), who find upper limits of less than 10 per cent.




FIGURE 4.4— Phase diagrams (temperature versus number density) for the simulations. Top panel shows simulation *L*, bottom panel shows simulation *H*; each dot represents an SPH particle. The individual particles at the top of each diagram are hot SN particles. Simulation *H* clearly shows a two-phase structure, while in simulation *L* almost all gas is in the warm ( $T \approx 10^4$  K) phase (85 per cent).

## 4.4 Discussion

The structural parameters of the modeled disks appear to be independent from the star formation activity. In this section we will discuss the appearance of the stellar disk of LSB galaxies, and the ISM in LSB galaxies.

### 4.4.1 Stellar disk

LSB galaxies are flatter than normal HSB galaxies. From a study of the catalog of edge-on galaxies by Karachentzev, Karechentzeva & Parnovsky (1993), Kudrya et al. (1994) concluded that galaxies become increasingly thinner towards later Hubble types, with the very thinnest galaxies having major over minor axis ratios of 20. Our model galaxies have axial ratios of 15, that is, almost as flat as the flattest galaxies in the Karachentzev, Karechentzeva & Parnovsky catalog. The top panel in Fig. 4.5 shows the edge-on view of mass distribution in LSB galaxy *L*.

Our model LSB galaxies support claims made by Dalcanton & Shectman (1996) that some of the extremely flat galaxies (“chain galaxies”) detected in medium-redshift HST images (Cowie, Hu & Songaila 1995) are simply edge-on LSB galaxies. Cowie, Hu & Songaila present HST wide-*I* band observations of these chain galaxies which they describe as “extremely narrow, linear structures ... with superposed bright ‘knots.’” A few of these galaxies were found to lie at a redshift  $z \sim 0.5$ . As an *I*-band view of that redshift corresponds approximately to a *V*-band view locally, we have attempted to simulate a *V*-band observation of model *L* edge-on, by assigning to each stellar particle the *V*-band flux corresponding to its age (Charlot & Bruzual 1993). This is shown in the bottom panel of Fig. 4.5, where the dynamic range of the picture has been adjusted to correspond approximately with the contrast that the observed chain galaxies have with respect to the sky-background. Comparison with any of the galaxies in Fig. 20 of Cowie, Hu & Songaila will show the resemblance.

How will the edge-on surface brightness distributions of LSB galaxies compare with those of HSB galaxies of similar scale length? Compared to an equally large HSB galaxy, the scale height of an LSB galaxy is smaller. Due to the low dust content in LSB galaxies (McGaugh 1994), effects of edge-brightening will be stronger in the LSB galaxy compared to the dust-rich HSB galaxy. An edge-on LSB




FIGURE 4.5— Edge-on views of the mass and light distribution in LSB galaxy *L*. The top panel shows a linear grey-scale representation of the projected mass surface density. The bottom panel shows a grey-scale representation of the projected *V*-band surface brightness. Here, the star particles formed during the run-time of the simulation are assigned a *V*-band flux according to their age following Charlot & Bruzual (1993). The contrast has been adjusted to a dynamic range of 5 magnitudes. This view resembles the so-called “chain galaxies” described by Cowie, Hu & Songaila (1995).

might have a higher apparent surface brightness than an edge-on HSB galaxy of the same size. For example, for HSB galaxies in the Ursa Major cluster, Tully et al. (1997) find a differential extinction in the *B*-band going from face-on to edge-on of  $1.8 \text{ mag arcsec}^{-2}$ . The LSB galaxies in their sample are found to be almost transparent with a differential extinction of only  $0.1 \text{ mag arcsec}^{-2}$ . The almost two magnitudes extra extinction in the HSB can thus easily compensate the few magnitudes difference in *intrinsic* surface brightness, making edge-on LSB galaxies as bright in surface brightness as edge-on HSB galaxies.

Hence it is much more difficult to distinguish between edge-on LSB and HSB galaxies than between face-on LSB and HSB galaxies. A smaller number of star forming regions, extreme flatness and blue colors will be the only easily visible distinguishing characteristics.

The reason why the stellar disks of LSB galaxies are so thin is that they are very stable to local instabilities (Fig. 4.2). As a con-

sequence, bars and spiral structure, which are the most efficient methods for heating a stellar disk (Sellwood & Carlberg 1984), are unlikely to develop spontaneously in LSB galaxies. Thus there is no natural way to make the disks thick.

This explanation is supported by the rarity of barred LSB galaxies. In the LSB galaxy catalog by Impey et al. (1996), only 4 per cent of the galaxies are barred, while the frequency of barred galaxies in the RC2 is some 30 per cent (Elmegreen, Elmegreen & Bellin 1990). The stability of LSB disks is confirmed in a numerical study of the dynamical stability of these systems by Mihos, McGaugh & De Blok (1997). Whereas we adopted the “most likely” solution for the mass-to-light ratio of stellar disk, these authors considered the “worst case” scenario of maximum disk. However their conclusion is the same as ours: the disks of LSB galaxies are extremely stable.

The lack of truly LSB edge-on galaxies (Schombert et al. 1992) could tell us something about the existence of very LSB galax-

ies. As shown above, the presently known population of LSB galaxies, when turned edge-on, brightens to surface brightnesses comparable to those of similar-sized HSB galaxies. In principle, the edge-on counterparts of the currently known LSB galaxies should thus already be in the conventional galaxy catalogs, showing up as “streaks on the sky”. As truly LSB edge-on galaxies seem to be lacking from the LSB catalogs, this suggests that galaxies with face-on surface brightnesses  $\mu_0^B > 25 \text{ mag arcsec}^{-2}$  are probably very rare. This is consistent with the idea that at low surface densities galaxies become completely ionized by the UV radiation field.

#### 4.4.2 Blue and red LSB galaxies

Fluctuations in the SFR as shown in Fig. 4.3, may very well explain why most of the LSB galaxies detected in surveys are blue. As McGaugh (1996) argues, the selection effects against finding red LSB galaxies on the blue sensitive plates on which surveys have been carried out are quite severe. It will be very hard to find LSB galaxies with colors  $B - V \simeq 1$  or redder. So if the absence of red LSB galaxies from the current catalogs is caused purely by these selection effects, the blue LSB galaxies could simply be the bursting tip of a proverbial iceberg.

In order to explain the colors of the bluest LSB galaxies in the sample of De Blok, McGaugh & Van der Hulst (1996), Van den Hoek et al. (1997) had to invoke bursts with a duration of between 0.5 and 5 Myr and an amplitude between 1 and 5  $M_\odot/\text{yr}$ . The fluctuations found in our simulation  $L$  have an amplitude of 0.15  $M_\odot/\text{yr}$  with a duration of about 20 Myr. This duration is determined by the lifetime for OB stars and the collapse time for molecular clouds, which are both of the order of a few  $10^7$  yr. They are therefore slightly milder than the bursts invoked by Van den Hoek et al. (1997), but keep in mind that the latter bursts were used to explain the *bluest* galaxies. Our models attempt to simulate an average LSB galaxy, and it should not come as a surprise that the fluctuations

we find are milder. Given the many assumptions and uncertainties in both modeling and observations, it is actually quite encouraging that the parameters agree to within an order of magnitude.

Assuming that blue LSB galaxies are currently undergoing a period of enhanced star formation, implies that there exists a population of *red, non-bursting, quiescent* LSB galaxies. These should then also be metal-poor and gas-rich, and share many of the properties of the galaxies we have been modeling.

We can estimate the fraction of red LSB galaxies by calculating the distribution of the birthrate parameter  $b$  for simulation  $L$ . The  $b$  parameter is the ratio of the present SFR over the average past SFR and is a useful tool for studying the star formation history of galaxies. Birthrate parameters have been determined for a large sample of spiral galaxies by Kennicutt, Tamblyn & Congdon (1994). The trend is that early type galaxies have small values for  $b$ , thus most star formation occurred in the past, while late type and irregular galaxies have large values for  $b$ , often exceeding unity, indicating that those galaxy are still actively forming stars, and more so than in the past. Here we apply this analysis to simulation  $L$  in order to estimate the fractions of blue and red LSB galaxies.

In Fig. 4.6a we plot the distribution of  $b$  values for simulation  $L$  (solid line), where we have followed the value of  $b$  over the duration of the simulation in steps of 15 Myr. Thus if the SFR peaks in a particular time interval, the corresponding value of  $b$  will be high. If the SFR is low in this time interval  $b$  is also low. In total we have 200  $b$  values.

Also shown in Fig. 4.6a are the  $b$  distribution for a simulation of an HSB Sc galaxy (Chapter 3, dotted line) and the mean values for different galaxy types (from Kennicutt, Tamblyn & Congdon 1994). Due to the low average SFR the distribution for the LSB simulation is much broader than the distribution for the HSB simulation, and the average  $b$  value is larger.

Also it is clear that the LSB galaxy has  $b$




FIGURE 4.6— (a) Distribution of the birthrate parameter  $b$ , the ratio of the current SFR to the average past SFR. Solid line shows the  $b$  values derived for simulation  $L$ , dotted line shows the  $b$  distribution for a simulation of an Sc galaxy (Chapter 3). On the top are average  $b$  values for different types of galaxies (Kennicutt, Tamblyn & Congdon 1994). The bottom panel shows the cumulative distribution for the two simulations. Less than 20 per cent of the  $b$  values for the LSB galaxy are below  $b = 0.4$ , hence we expect at most 20 per cent of LSB galaxies to be “red”.

values larger than the average for early type galaxies for most of the time. “Classical” LSB galaxies are blue compared to HSB galaxies, they thus have an excess of recent star formation or equivalently a higher  $b$  value. We now define a LSB galaxy to be “blue” if its birthrate parameter  $b$  exceeds the average value of  $b$  for a HSB late-type galaxy (see Kennicutt, Tamblyn & Congdon 1994 for relations between birthrate parameter and color). Fig. 4.6a shows that this requires that  $b_{\text{LSB}} > \langle b_{\text{HSB}} \rangle \approx 0.4$ .

LSB galaxies that do not meet this requirement are “red”: non-bursting, but nevertheless still gas-rich. From Fig. 4.6b (which shows the cumulative  $b$  distribution) we can see that over 80 per cent of the fluctuations result in blue LSB galaxies. Less than 20 per cent of the fluctuations therefore results in red LSB galaxies.

We can estimate colors of the red population using the bursts from van den Hoek et al. (1997). These change the colors and surface brightnesses by  $\Delta(B - V) \simeq -0.5$ ,  $\Delta(R - I) \simeq -0.2$ ,  $\Delta B \simeq -1$  and  $\Delta I \simeq -0.4$  (cf. Table 3.5). Using these values together with the meas-

ured colors of the bluest LSB galaxies in De Blok, Van der Hulst & Bothun (1995), yields  $B - V \sim 1$ ,  $R - I \sim 0.6$  and  $\mu_0^B = 24.5$  for the red population.

A recent CCD survey (O’Neill, Bothun & Cornell 1997) has picked up a class of LSB galaxies which have  $\mu_0^B \simeq 24$  and  $B - V \simeq 0.8$ . As no red-shifts are available yet it is difficult to know at what distance these galaxies are, but if they are indeed the non-bursting counterparts of the blue LSB galaxies, they should also be metal-poor and gas-rich, and share many of the properties of the galaxies we have been modeling.

In summary, if the blue colors found in LSB galaxies are the result of fluctuations in the SFR, then this implies that the red LSB galaxies constitute less than 20 per cent of this particular class of LSB disk galaxies. This does not rule out the existence of a population of red, gas-poor LSB galaxies. These must however have had an evolutionary history quite different from those discussed here and possibly have consumed or expelled all their gas quite early in their life.

### 4.4.3 ISM

The SFRs from both simulations are larger than the observed rate of about  $0.05 M_{\odot}/\text{yr}$ . This latter value is probably correct within a factor of 2. Especially the high SFR of simulation *H* ( $0.33 M_{\odot}/\text{yr}$ ) is almost an order of magnitude larger than what is needed and it seems impossible to reconcile this value with the observations. Instead the global SFR is consistent with the global SFR for equal-luminosity galaxies (e.g. Kennicutt 1983). For instance the Sc galaxy NGC 6503 with a maximum rotation velocity of  $v = 120 \text{ km/s}$  has a total measured SFR of  $0.4 M_{\odot}/\text{yr}$ , and a simulated SFR of  $0.35 M_{\odot}/\text{yr}$  (Chapters 2 and 3). In general the SFRs of equal-luminosity galaxies differ by a factor 10 between LSB and HSB galaxy (Van den Hoek et al. 1997; Kennicutt 1983), despite the copious amounts of gas available in the LSB galaxies (De Blok et al. 1996).

$\text{SFR}_L$  approaches the correct value and the physical reason is shown in Fig. 4.4. The essential information to retain from this phase diagram is that we need a different ISM for LSB galaxies, where the bulk of the gas is not directly available for star formation, as it is too warm (of order  $10^4 \text{ K}$ ). The simulations do not include phase transitions from neutral to molecular gas, but as an estimate for the  $\text{H}_2$  mass we can consider all star forming gas ( $T \lesssim 300 \text{ K}$ ) to be molecular. This gas represents less than 2 per cent of the total gas mass. In this respect it is interesting to note that for a small sample of LSB galaxies CO is not detected, yielding a very hard upper limit of  $M(\text{H}_2)/M(\text{H I}) \ll 30$  per cent (De Blok & Van der Hulst 1997, Schombert et al. 1990). A few galaxies have upper limits less than 10 per cent.

As discussed in the previous section the warm ISM may be caused by a low metallicity for the gas. Below  $10^4 \text{ K}$ , cooling is dominated by heavy elements like  $\text{C}^+$ ,  $\text{Si}^+$ ,  $\text{Fe}^+$ , O. If these elements are rare then it is difficult for the gas to radiate its energy away.

Direct observational support for a low metallicity ISM in LSB galaxies comes from

oxygen abundance measurements of H II regions in LSB galaxies. Those studies yield metallicities of approximately 0.5 times solar metallicity (McGaugh 1994). Measurements of the oxygen abundance in F563-1 (De Blok and Van der Hulst 1997) give an average oxygen abundance of  $0.15 Z_{\odot}$  (compare with the difference in metallicity between models *H* and *L*).

A point of concern is the IMF. Our premise is that the IMF is universal, however it can differ from our adopted Salpeter IMF (see 4.1.1). Especially a low upper-mass cutoff could easily lead to an underestimation of the SFR (e.g. an upper mass cutoff at  $30 M_{\odot}$  reduces the measured SFR by 2-3; Kennicutt 1983). However current determinations of the IMF in external galaxies point towards a universal IMF (Kennicutt 1989; Kennicutt, Tamblyn & Congdon 1994). Furthermore, observations suggest that stars more massive than  $60 M_{\odot}$  are present in the H II regions of LSB galaxies (McGaugh 1994).

## 4.5 Conclusions

What causes the low evolution rate for LSB galaxies? The low density has often been invoked to explain this, since the dynamical time scales with  $1/\sqrt{\rho}$ . This scenario is exactly what is tested in simulation *H*. The only difference with a normal HSB galaxy is the scale length of the galaxy, which is for instance three times larger than the scale length for the equally bright galaxy NGC 6503. The result is striking: adopting “standard” values for the star formation process results in a SFR identical to the rates of HSB galaxies.

Thus the low density in itself seems not capable of doing the job, and we have to rely on a scarcity of heavy elements to reproduce a true LSB galaxy. This fits in logically with the notion that stars are the producers of these elements; the low star formation activity prevents metal enrichment of the ISM. It implies that the SFR has been low throughout the evolution of LSB galaxies, and that these galaxies are “trapped” in their current evolutionary state: low density prevents rapid star

formation, which prevents enrichment of the ISM, which prevents cooling, resulting in a warm one-phase ISM. So although the lack of metals is directly responsible for the low SFR, the low density may ultimately determine the fate of LSB galaxies.

In summary we reach the following conclusions on the physical properties of LSB galaxies.

*Dominant halo:* The dominance of the dark matter halos in LSB galaxies (De Blok & McGaugh 1997) results in very thin and stable disks. The major over minor axis ratios of the stellar disks are larger than 15. Lack of dust in LSB galaxies will result in edge-on LSB galaxies having apparent surface brightnesses equal to those of edge-on HSB galaxies of similar size. The lack of truly LSB edge-on galaxies may indicate that disk galaxies with  $\mu_0^B > 25$  mag arcsec<sup>-2</sup> are rare.

*Low metallicity:* Modeling the low SFR found in LSB galaxies, demands that the metallicity in the ISM must be approximately 0.2 solar, which is consistent with observation by McGaugh (1994) and De Blok & Van der Hulst (1997). These low metallicities prevent effective cooling so that the ISM in LSB galaxies primarily consists of warm ( $\sim 10^4$  K) gas. The amount of cold (molecular) gas is probably less than 5 per cent of the H I mass, supporting claims by Schombert et al. (1990) and De Blok & Van der Hulst (1997) who derive small molecular gas fractions.

*Low average SFR:* Due to the fluctuations in the SFR in LSB galaxies and their large contrast with the average SFR, the spread in colors among LSB galaxies will be larger than among HSB galaxies. From the distribution of birthrate parameters in our simulations we deduce that, if the currently known blue gas-rich LSB galaxies are the most actively star forming LSB galaxies, they constitute over 80 per cent of the total population of field LSB disk galaxies. This implies that there is at most an additional 20 per cent of quiescent,

gas-rich LSB disk galaxies. This does not preclude the existence of an additional red, gas-poor population. However, this population must have an evolutionary history quite different from that described in this work.

## Acknowledgements

We are much indebted to Lars Hernquist for generously providing the TREESPH code. We also thank Frank Briggs and Thijs van der Hulst for their comments on a draft of this paper. The research of JPEG was supported by the Netherlands Foundation for Research in Astronomy (NFRA) with financial aid from the Netherlands Organization for Scientific Research (NWO).

## References

- Barnes J.E., Hut P., 1986, *Nature*, 324, 466  
 Bottema R., 1993, *A&A* 275, 16  
 Bruzual B.A., Charlot S., 1993, *ApJ* 405, 538  
 Charlot S., Bruzual B.A., 1991, *ApJ* 367, 126  
 Cowie L.L., Hu E.M., Songaila A., 1995, *AJ* 110, 1576  
 Dalcanton J.J., Shtetman S.A., 1996, *ApJL* 465, L9  
 Dalgarno A., McCray R., 1972, *ARA&A* 10, 375  
 De Blok W.J.G., 1997, PhD thesis, University of Groningen  
 De Blok W.J.G., McGaugh S.S., 1996, *ApJ* 169, L89  
 De Blok W.J.G., McGaugh S.S., 1997, *MNRAS*, in press  
 De Blok W.J.G., Van der Hulst J.M., 1997, *A&A*, submitted  
 De Blok W.J.G., Van der Hulst J.M., Bothun G.D., 1995, *MNRAS* 274, 235  
 De Blok W.J.G., McGaugh S.S., Van der Hulst J.M., 1996, *MNRAS* 283, 18  
 Dopita M.A., Ryder S.D., 1994, *ApJ* 430, 163  
 Elmegreen D.M., Elmegreen B.G., Bellin A.D., 1990, *ApJ* 364, 415  
 Field G.B., Goldsmith D.W., Habing H.J., 1969, *ApJ* 155, L149  
 Friedli D., Benz W., 1995, *A&A* 301, 649  
 Gerritsen J.P.E., Icke V., 1997a, *A&A* 325, 972  
**chapter 2**  
 Gerritsen J.P.E., Icke V., 1997b, submitted, **chapter 3**  
 Gingold R.A., Monaghan J.J., 1977, *MNRAS* 181, 375  
 Guiderdoni G., Rocca-Volmerange B., 1987, *A&A* 186, 1  
 Hernquist L., 1987, *ApJS* 64, 715

- Hernquist L., 1989, *Nature* 340, 687  
Hernquist L., Katz N., 1989, *ApJS* 70, 419  
Impey C.D., Sprayberry D., Irwin M.J., Bothun G.D., 1996, *ApJS* 105, 209  
Karachentzev I.D., Karechentzeva V.E., Parnovsky S.L., 1993, *Astron. Nach.* 314, 97  
Katz N., 1992, *ApJ* 391, 502  
Katz N., Weinberg D.H., Hernquist L., 1996, *ApJ* 105, 19  
Kennicutt R.C., 1983, *ApJ* 272, 54  
Kennicutt R.C., 1989, *ApJ* 344, 685  
Kennicutt R.C., Tamblyn P., Congdon C.W., 1994, *ApJ* 435, 22  
Kudrya Y.N., Karachentzev J.D., Karachentzeva V.E., Parnovsky S.L., 1994, *Astron. Lett.* 20, 8  
Lamers H.J.G.M.L., Leitherer C., 1993, *ApJ* 412, 771  
Leitherer C., Robert C., Drissen L., 1992, *ApJ* 401, 596  
Lucy L.B., 1977, *AJ* 82, 1013  
McGaugh S.S., 1994, *ApJ* 426, 135  
McGaugh S.S., 1996, *MNRAS* 280, 337  
McGaugh S.S., Bothun G.D., 1994, *AJ* 107, 530  
McGaugh S.S., De Blok W.J.G., 1997, *ApJ* 481, 689  
McKee C.F., Ostriker J.P., 1977, *ApJ* 218, 148  
Mihos J.C., Hernquist L., 1994, *ApJ* 437, 611  
Mihos J.C., Hernquist L., 1994, *ApJ* 431, L9  
Mihos J.C., Hernquist L., 1996, *ApJ* 464, 641  
Mihos J.C., McGaugh S.S., De Blok W.J.G., 1997, *ApJ* 477, 79L  
Navarro J.F., White S.D.M., 1993, *MNRAS* 265, 271  
O'Neil K., Bothun G.D., Cornell M.E., 1997, *AJ* 113, 1212  
Schombert J.S., Bothun G.D., Impey D.D., Mundy L.G., 1990, *AJ* 100, 1523  
Schombert J.S., Bothun G.D., Schneider S.E., McGaugh S.S., 1992, *AJ* 103, 1107  
Sellwood J.A., Carlberg R.G., 1984, *ApJ* 282, 61  
Steinmetz M., 1996, *MNRAS* 278, 1005  
Toomre A., 1964, *ApJ* 139, 1217  
Tully R.B., Verheijen M.A.W., Pierce M.J., Huang J.-S., Wainscoat R.J., 1996, *AJ* 112, 2471  
Van Albada T.S., Sancisi R., 1986, *Phil. Trans. R. Soc. Lond. A* 320, 447  
Van den Hoek L.B., De Blok W.J.G., Van der Hulst J.M., De Jong T., 1997, *A&A*, submitted  
Van der Hulst J.M., Skillman E.D., Kennicutt R.C., Bothun G.D., 1987, *A&A* 117, 63  
Van der Hulst J.M., Skillman E.D., Smith T.R., Bothun G.D., McGaugh S.S., De Blok W.J.G., 1993, *AJ* 106, 548  
Van der Kruit P.C., Searle L., 1982, *A&A* 110, 61  
Wolfire M.G., Hollenbach D., McKee C.F., Tielens



