

University of Groningen

## Ecological notes on *Carex aquatilis* communities

Grootjans, A.P.; van Tooren, B.F.

*Published in:*  
Vegetatio

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1984

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*  
Grootjans, A. P., & van Tooren, B. F. (1984). Ecological notes on *Carex aquatilis* communities. *Vegetatio*, 57(2-3), 79-89.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Ecological notes on *Carex aquatilis* communities\*

A. P. Grootjans\*\* & B. F. van Tooren\*\*.

\*\*\*  
Department of Plant Ecology, University of Groningen, Biological Centre, P.O. Box 14, 9750 AA Haren (Gn), The Netherlands

Keywords: *Carex aquatilis*, Groundwater, NW Europe, Preservation, Seepage areas

### Abstract

*Carex aquatilis* spp. *aquatilis* grows in mesotrophic fens and has a boreal circumpolar distribution. The species is most frequent in areas north of the 66° N line. In Europe the southern limits of its distribution is ca 53°N. Here *Carex aquatilis* is found in eutrophic flood plains. However, mesotrophic communities of *C. aquatilis* have been reported from these areas as well. Nowadays the peripheral populations in NW Europe appear to be relics of former mesotrophic communities which at present are under heavy pressure from human activities such as drainage and fertilizing. In the northern part of the Netherlands *C. aquatilis* stands are restricted to areas with a steady supply of cool groundwater. A decreased groundwater discharge has a negative influence on the development of inflorescences. It is expected that further interference with the hydrology will promote a further decline of the peripheral populations. The species will not be able to survive in drained, eutrophic, warmed up habitats.

\* Nomenclature of phanerogams follows Heukels - van Oostroom (1977), that of Bryophytes Smith (1978). Nomenclature of northern plant communities follows Dierssen (1982), that of Dutch and NW German communities Westhoff & den Held (1969) and Ellenberg (1978).

\*\* *Acknowledgements.* This paper is dedicated to the late Prof. Dr D. Bakker who started the floristic surveys in the Drentse Aa catchment and rose our interest in ecological factors influencing plant species distribution. The authors wish to thank Prof. Dr J. J. Barkman (Wijster), Prof. Dr K. Dierssen (Kiel), Dr Denis Ratcliffe (Leicester), and our colleagues at the Department of Plant Ecology (Haren) for their comments on the manuscript. We also would like to thank Mr F. Adema (Rijksherbarium, Leiden), the Provinciale Planologische Dienst Drenthe and Mr F. Rudophy for providing information on the occurrence of *Carex aquatilis* stands. Mr Leeuwenga prepared the figures and Mrs Biewenga typed various versions of the manuscript. Their help is gratefully acknowledged.

\*\*\* Present address: Vakgroep Vegetatiekunde en Botanische Oecologie, Lange Nieuwstraat 106, 3512 PN Utrecht, The Netherlands.

### Introduction

*Carex aquatilis* Wahlenb. is a swamp species with a boreal circumpolar distribution (Cajander, 1903; Alechin, 1927; Andreas, 1951; Kern & Reichelt, 1954; Churchill, 1955; Bakker & Westhoff, 1957; Neumann, 1957; Hultén, 1962; Komarov, 1964; Tutin *et al.*, 1980; Dierssen, 1982).

Three subspecies of *Carex aquatilis* have been distinguished (Hultén, 1962; *Carex aquatilis* ssp. *stans* (Drejer) Hultén, *Carex aquatilis* ssp. *altior* Rydb., and *Carex aquatilis* ssp. *aquatilis*). Subspecies *stans* has a more northern distribution than ssp. *aquatilis*, while the American ssp. *altior* has a more southern distribution. In Europe ssp. *altior* is absent. This article deals with *C. aquatilis* ssp. *aquatilis* only (see Fig. 1).

In Europe the main distribution of *Carex aquatilis* is north of the 66°N line (Dierssen, 1982) where the species grows in reed swamps, fens, wet meadows and even in relatively saline habitats (Siira,




Fig. 1. Distribution of *Carex aquatilis* ssp. *aquatilis* in the northern hemisphere. In America the distribution of *Carex aquatilis* includes both *C. a. ssp. aquatilis* and *C. a. ssp. altior* (after Hultén, 1962).

1970; Jefferies, 1977). In northern Sweden the species has a remarkable peat-forming ability (Elveland, 1976; Waldemarson-Jensen, 1979).

In the Netherlands and in NW Germany *C. aquatilis* was known only as a sub-fossil remain until 1948 (Andreas, 1951). In the period 1951–1955 many localities of *C. aquatilis* were discovered (Andreas, 1951; Küsel, 1956 pers. comm.; Bakker, 1957; Neumann, 1957; Dierschke, 1968). Before 1951 the species must have been confused with *Carex acuta*. Up to now the most southern locality of *C. aquatilis* in Europe appears to be the Overijsselse Vecht in the Netherlands at 53° N.

In this article we will discuss how *C. aquatilis* is able to survive at the southern border of its distribution, where conditions are quite different from those in the main distribution area. The occurrence of differences in the ecology of central and peripheral populations has been suggested (Bakker, 1979; Schmidt, 1969; see also Hengeveld & Haeck, 1982). Thus the NW European lowland habitats of the species may differ from those of more northern

localities. The synecological conditions of *Carex aquatilis* stands were studied on the Drenthian Plateau, N Netherlands, which is similar to several Pleistocene plateau's in the NW German lowland.

The species composition and performance of *C. aquatilis* stands were studied in relation to hydrological features. The results are compared with published research on *C. aquatilis* communities in NW Germany, Norway, N Sweden and Scotland.

#### Materials and methods

The study area is situated south of Groningen (53°08'N, 6°37'E) in the catchment of the Drentse Aa, which drains 30 000 ha of a pleistocene landscape. Detailed information on hydrological and pedological aspects of this region was presented by Gischler (1967) and de Gans (1982). The vegetation of the Drentse Aa catchment area has been discussed by de Bruijn (1980), Everts *et al.* (1980) and Grootjans (1980).

In 1978 *Carex aquatilis* stands in this area were described according to the Braun-Blanquet approach, using the scale of Londo (1976). Below 5% cover a combination of letter and number is given:  $p$  = less than 20 individuals on 4 m<sup>2</sup>,  $a$  = 20–100,  $m$  = more than 100; 1 = less than 1% cover, 2 = 1–3%, 3 = 3–5%. Above 5% cover only a number is given: 1 = 5–15%, 2 = 15–25%, . . . , 9 = 85–95%, 10 = 95–100%.

The performance of *C. aquatilis* was estimated by assessing the percentage of flowering shoots and the cover of the species. The percentage of flowering shoots is thought to be a measure of the vitality of the plants. For sedge species this may not always be a correct assumption. For instance shoots of *Carex lacustris* remain vegetative in most years (Bernard, 1975). Flowers were formed only during one summer following severe flooding. In *C. rostrata* all vegetative mature shoots will flower; the number of flowering shoots is dependent on shoot mortality (Bernard & Gorham, 1978). *C. aquatilis* has a life cycle similar to that of *C. rostrata* (Gorham & Somers, 1973), so the percentage of flowering shoots appears to be a good measure of the vitality of the plants.

Observations of the water table were made in a number of *C. aquatilis* stands. The measurements were carried out in PVC tubes (40 mm diameter) every two weeks. All the data have been transformed to cumulative frequency histograms. These lines represent the residence time of groundwater in the soil profile (for more detailed information see Niemann, 1973; Grootjans & ten Klooster, 1980).

## Results

### *Carex aquatilis* in Dutch catchment areas

In the Netherlands *Carex aquatilis* is restricted to the northeastern part of the country (Fig. 2). The main distribution area is the fringe of the Drenthian Plateau where deep groundwater, originating from the plateau, emerges. In Figure 3 the distribution of *Carex aquatilis* is shown. Seepage areas and flow patterns of deep groundwater are shown too (see Working Group Geohydrological Research in Drenthe, 1978).

*Carex aquatilis* seems to prefer seepage areas in the lower reaches of the Drenthian streams. This




Fig. 2. Distribution of *Carex aquatilis* in the Netherlands. Each square represents 28 km<sup>2</sup>.

suggests a preference of the species for enriched groundwater.

Although the preference of *Carex aquatilis* for certain seepage areas is clear, the species is absent from areas where agricultural improvement projects have been carried out. Some stream systems have been devastated almost completely, leaving none of the marsh vegetation.

In the eastern part of Drenthe, the absence of *Carex aquatilis* is probably due to the presence of extensive oligotrophic bog complexes preventing the mineral rich deep groundwater from reaching the surface. Most of these mires have been reclaimed for agricultural purposes, and only small remnants of more natural vegetation are still present. Relevés of *Carex aquatilis* from Dutch stream basins are shown in Table 1. Five groups have been distinguished (A–E); they will be discussed only briefly here. Figure 4 presents a sketch of hydrological conditions and occurrence in the landscape.

*Carex aquatilis* is most abundant in the lower basins where the species is able to form extensive mats on soils that are rich in silt and organic material and which are flooded regularly. The relevés of group A must be classified as *Lysimachio-Caricetum aquatilis inops*. The species richness of these *Magnocaricion* stands is very low. The rest of the relevés of the lower basin (group B) and the majority of the relevés of the middle basin (groups D and E) belong to the *Senecioni-Brometum racemosi*




Fig. 3. Distribution of *Carex aquatilis* in the N Netherlands. Each square represents 1 km<sup>2</sup>. Indicated are seepage areas, former bogs and flow direction of the deep groundwater. The former coastline is thought to correspond to the present 2 m above Amsterdam ordnance datum (NAP) line. The occurrence of brackish groundwater is indicated by the 500 mg/l Cl<sup>-</sup> line.

and are rich in species. Relevés from group C belong to the *Caricetum curto-echinatae* and are rich in species as well.

*Carex aquatilis* stands in the lower basins are bordered by *Carex acuta* stands belonging to the *Caricetum gracilis* and by communities of small sedges (*Caricion curto-nigrae*).

In the middle reaches of undisturbed catchments *C. aquatilis* is fairly common. It grows in damp depressions just behind the stream banks. In these stands *Caltha palustris*, *Myosotis scorpioides*, *Lychnis flos-cuculi*, *Filipendula ulmaria* and many more species of the *Molinio-Arrhenatheretea* are frequent.

In the upper reaches *C. aquatilis* stands are rare. They can be classified as *Calthion palustris* as well.

Summarizing it can be said that most of the *C. aquatilis* stands of the middle and upper reaches belong to the *Calthion palustris* alliance, while quite a number of stands in the lower basin can be

assigned to the *Magnocaricion* alliance. This points to prevailing less wet conditions in the middle and upper reaches compared to the lower basin. To verify this the groundwater regime of *Carex aquatilis* stands will be considered.

#### *Carex aquatilis* and groundwater regime

From the catchment area of the Drentse Aa, all available measurements of groundwater tables in *Carex aquatilis* stands were gathered. The data have been transformed into cumulative frequency histograms (Fig. 5). Unfortunately few series of measurements were available, so a direct comparison of *C. aquatilis* stands of lower and middle basins was not possible from a statistical point of view. We can however compare the grouped frequency lines with references from the literature (Grootjans, 1980; Grootjans & ten Klooster, 1980). For that purpose we chose the *Senecioni-Brometum race-*

Table 1. Relevés of *Carex aquatilis* stands in the middle and lower reaches of the catchment of the Drentse Aa (N Netherlands).

Group code	A			B			C			D			E																					
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32		
Relevé number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32		
L = lower basin M = middle basin	L	L	L	L	L	L	L	M	M	M	L	L	L	L	L	L	L	M	M	M	M	M	M	M	M	M	M	M	M	M	M	M		
Surface in m <sup>2</sup>	4	4	4	4	4	4	4	2	2	2	4	4	4	4	4	4	4	4	4	2	4	4	4	4	2	2	4	4	4	4	4	4		
Cover (%) of phanerogams	30	30	50	50	35	80	30	30	40	95	30	35	30	30	35	30	60	40	60	40	60	50	15	40	70	50	40	85	95	70	70	60		
Cover (%) of mosses	-	-	-	-	-	-	-	3	5	<1	10	3	20	50	50	85	25	1	2	2	5	2	5	25	10	35	20	20	<1	6	<1	2		
Number of taxa	13	5	7	6	7	6	9	16	20	15	16	17	21	17	15	14	17	16	23	29	33	29	25	27	28	29	33	36	34	33	23	33		
<i>Carex aquatilis</i>	1	3	3	4	2	7	3	1	2	4	1	1	2	2	2	3	5	2	m <sub>2</sub>	1	1	2	1	2	3	3	2	3	2	1	1	1		
<i>Equisetum fluviatile</i>	a <sub>1</sub>	a <sub>1</sub>	l	p <sub>1</sub>	m <sub>2</sub>	p <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	r	p <sub>1</sub>	p <sub>1</sub>	a <sub>1</sub>	p <sub>1</sub>	p <sub>1</sub>	p <sub>1</sub>	p <sub>1</sub>	p <sub>1</sub>	m <sub>1</sub>	m <sub>4</sub>	m <sub>3</sub>	m <sub>1</sub>	m <sub>1</sub>	p <sub>1</sub>	p <sub>1</sub>			p <sub>1</sub>	m <sub>1</sub>	m <sub>2</sub>	p <sub>1</sub>	a <sub>1</sub>			
<i>Galium palustre</i>	r	p <sub>1</sub>	p <sub>1</sub>			p <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	m <sub>1</sub>	m <sub>1</sub>	m <sub>2</sub>	m <sub>1</sub>	a <sub>1</sub>	m <sub>1</sub>			m <sub>1</sub>	m <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	m <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	m <sub>1</sub>	m <sub>1</sub>	m <sub>1</sub>	m <sub>1</sub>	m <sub>1</sub>	a <sub>1</sub>			
<i>Glyceria maxima</i>	m <sub>2</sub>	m <sub>2</sub>	l	a <sub>2</sub>	p <sub>1</sub>	m <sub>2</sub>	a <sub>1</sub>	p <sub>1</sub>																										
<i>Carex rostrata</i>																																		
<i>Phalaris arundinacea</i>	a <sub>1</sub>				a <sub>4</sub>		a <sub>1</sub>	p <sub>1</sub>	a <sub>2</sub>	p <sub>1</sub>	p <sub>1</sub>	p <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	a <sub>1</sub>	p <sub>1</sub>																		
<i>Lysimachia vulgaris</i>	a <sub>1</sub>			r	r																													
<i>Agrostis stolonifera</i>																																		
<i>Myosotis palustris</i>																																		
<i>Galtha palustris</i>																																		
<i>Ranunculus repens</i>																																		
<i>Poa trivialis</i>																																		
<i>Holcus lanatus</i>																																		
<i>Calliergon cordifolium</i>																																		
<i>Calliergonella cuspidata</i>																																		
<i>Plagiomium affine</i>																																		
<i>Brachythecium rutabulum</i>																																		
<i>Amblystegium riparium</i>																																		
<i>Callamagrostis canescens</i>																																		
<i>Cardamine pratensis</i> ssp. <i>palustris</i>																																		
<i>Ranunculus flammula</i>																																		
<i>Agrostis canina</i>																																		
<i>Viola palustris</i>																																		
<i>Hydrocotyle vulgaris</i>																																		
<i>Filipendula ulmaria</i>																																		
<i>Rumex acetosa</i>																																		
<i>Cardamine pratensis</i> ssp. <i>pratensis</i>																																		
<i>Anthoxanthum odoratum</i>																																		
<i>Lychnis flos-cuculi</i>																																		
<i>Cirsium palustre</i>																																		
<i>Mentha aquatica</i>																																		
<i>Equisetum palustre</i>																																		
<i>Lotus uliginosus</i>																																		
<i>Stellaria palustris</i>																																		
<i>Galium uliginosum</i>																																		
<i>Cynosurus cristatus</i>																																		
<i>Cardamine amara</i>																																		
<i>Menyanthes trifoliata</i>																																		
<i>Carex nigra</i>																																		
<i>Trifolium repens</i>																																		
<i>Cerastium holosteoideum</i>																																		
<i>Festuca pratensis</i>																																		
<i>Ranunculus acris</i>																																		
<i>Festuca rubra</i>																																		
<i>Plantago lanceolata</i>																																		
<i>Trifolium pratense</i>																																		
<i>Rhinanthus serotinus</i>																																		

Addenda: other accompanying species

*Peucedanum palustre* 1/p<sub>1</sub>, 5/r; *Rorippa amphibia* 1/pl, 3/2; *Cicuta virosa* 1/p<sub>1</sub>, 6/r; *Iris pseudacorus* 1/p<sub>1</sub>, 7/r; *Typha latifolia* 2/r, 4/p<sub>1</sub>; *Carex disticha* 5/1, 7/m<sub>2</sub>, 13/a<sub>1</sub>; *Carex hudsonii* 1/1, 5/a<sub>2</sub>, 12/a<sub>1</sub>, 13/1; *Stellaria alsine* 8/p, 9/r, 18/m, 19/m; *Carex acutiformis* 8/2, 10/p<sub>1</sub>, 18/p<sub>1</sub>, 22p<sub>1</sub>; *Juncus effusus* 7/r, 8/r, 9/a<sub>1</sub>, 26/a, 27/a, 28/p<sub>1</sub>, 29/aa, 31/a<sub>1</sub>; *Lythrum salicaria* 21/p<sub>1</sub>, 22/r, 24/r, 29/p<sub>1</sub>; *Taraxacum* sect. *vulgaria* 19/p<sub>1</sub>, 24/p<sub>1</sub>, 25/r, 26/p<sub>1</sub>, 29/r, 32/p<sub>1</sub>; *Achillea ptarmica* 19/a<sub>1</sub>, 24/m<sub>2</sub>, 25/p<sub>1</sub>; *Carex curta* 21/r, 29/r, 30/a<sub>1</sub>; *Climacium dendroideum* 22/p<sub>1</sub>, 26/a<sub>1</sub>, 32/a<sub>1</sub>; *Rhynchospora squarrosa* 23/a<sub>1</sub>, 26/a<sub>1</sub>, 31/m<sub>1</sub>; *Senecio aquaticus* 28/a<sub>1</sub>, 29/r, 32/r.




Fig. 4. Occurrence of *Carex aquatilis* stands in upper, middle and lower reaches of Drenthian streams. Shaded arrows indicate mesotrophic groundwater or eutrophic floodwater. Light arrows indicate oligotrophic subsurface groundwater above boulder clay lenses.




Fig. 5. Cumulative frequency histograms of groundwater tables ((Dauerlinien) of three plant communities during the vegetation period. The lines represent the residence time of groundwater in the soil profile. The results of a statistical analysis (using the Wilcoxon-Mann-Whitney test) are at the base of the figure. L = lower reaches; M = middle reaches.

mosi (a wet *Calthion palustris* community) and the *Caricetum curto-echinatae* (a very wet *Parvocaricetea* community). In our region both communities are found in close contact with *Carex aquatilis* communities. The data from these communities include a wet (1979) and a dry (1976) year.

From the statistical analysis of the data, it can be concluded that the groundwater regime of the *Carex aquatilis* and the *Caricetum curto-echinatae* community are very similar.

Although some of the *Carex aquatilis* stands (especially those in the lower reaches) show more profound flooding than the small sedge community, these differences are not significant. Groundwater regimes of both communities differ, however, from the *Senecioni-Brometum racemosi* association in all levels. The differences in the highest levels are most pronounced. In contrast to what would be expected, the analysis shows that the cumulative frequency lines of the *Caricetum curto-echinatae* are more different from the *Senecioni-Brometum* lines, than from those of the *Carex aquatilis* groups, especially in water levels slightly below the surface. So the groundwater regime of *Carex aquatilis* stands have more in common with that of the drier *Calthion palustris* association than with that of the wetter *Pervocaricetea* community. Figure 5 shows that the lines from the middle basin are responsible for this effect.

As most *Carex aquatilis* stands of the middle reaches are situated just behind the stream banks, drainage is most probably caused by the stream itself. Superficial drainage by ditches, discharging in the stream, may enhance this effect.

Recent hydrological studies (Streefkerk & van Hoorn, 1984) have shown that after radical interference with the hydrology during the last 30 years, the stream levels in the middle course have indeed been lowered, especially in spring and winter. These lower levels may well be responsible for the unexpected results of the analysis of groundwater levels

and may account for the many *Carex aquatilis* stands that must be classified as *Calthion palustris*.

#### *Observations on reproduction in Carex aquatilis stands*

If the supposition presented above is correct, one would expect that the habitats of *Carex aquatilis* stands in the middle reaches has become less favourable. If so, *Carex aquatilis* plants in the lower basin would perform better in some way than the plants in the middle reaches. To test this we estimated the cover percentage and the percentage of flowering shoots in 32 *Carex aquatilis* stands. If the percentage of flowering shoots and the percentage cover would be a measure of the vitality of the plants, one would expect this to be lower in the middle reaches than in the lower reaches.

The results show that in 1978 the percentage of flowering shoots in the middle basin was indeed lower than in the lower basin (Table 2). In the middle reaches only 2% of the shoots had inflorescences, while in the lower reaches this was 12–28%.

In 1980 the same plots were visited again. In the lower basin the percentage of flowering shoots was almost the same as in 1978. The flowers however, withered before maturing as a consequence of a very dry period in spring. In the middle reaches the percentage of flowering shoots had increased considerably (to 13%) and the flowers did not wither. Although the high values of the lower basin were

Table 2. Percentage of flowering shoots and cover of *Carex aquatilis*. The group codes (A–E) correspond to those mentioned in Table 1.

Group code (see Table 1)	A	B	C	D	E
L = Lower basin / M = Middle basin	L	L/M	M	M	M
Average % cover of <i>C. aquatilis</i>	32	23	11	21	18
% of flowering shoots					
1978 mean	28.3	11.6	0.3	2.1	1.5
S.D.	9.8	21.8	0.6	2.3	1.8
1980 mean	20–30	10–15	13.0	6.8	2.8
S.D.			6.9	7.8	4.2






Fig. 6. Synecological conditions of *Carex aquatilis* communities in N and NW Europe. Eutrophic corresponds to substrates rich in silt and groundwater rich in dissolved minerals (high pH and E.C.). Oligotrophic corresponds to nutrient-poor organic substrates influenced by mineral-poor groundwater. The wetness gradient ranges from drained to immersed (flooded). In N Sweden flooding may occur by brackish water (Elveland, 1976). Double arrows indicate predominant vegetation development in the region. Information on ecological conditions in N European communities has been derived from Elveland (1976), Waldemarson-Jensén (1979), and Dierssen (1982). Additional information on NW European *C. aquatilis* communities originate from Neumann (1957), Dierschke (1968), and Tüxen & Dierschke (1975).

not reached, the flowering plants were well developed. The observed phenomena could be related to a gradual increase in precipitation since 1978 after the exceptionally dry year 1976; this led to higher water tables in the Drenthian Plateau. Higher water tables in the plateau were associated with increased water discharge of deep groundwater in the river valleys, especially in the middle reaches of the Drentse Aa, where the seepage is more intense and where fewer drainage channels (adjacent land consolidation) are present. Thus the effect of the dry spring period of 1980 was better compensated by the higher groundwater discharge in the middle reaches than in the lower reaches. In the middle reaches *Carex aquatilis* stands responded with an increased percentage of flowering shoots compared with 1978 and the flowers remained in good condition. Despite this the plants did not subsequently form ripe seed in either area.

#### *Carex aquatilis* in NW Europe

Communities of *Carex aquatilis* have been reported mainly from areas north of the 66°N line (Cajander, 1903; Alechin, 1927; Elveland, 1976; Waldemarson-Jensen, 1979; Dierssen, 1982). More to the south *Carex aquatilis* communities are described from subalpine and alpine areas (McVean & Ratcliffe, 1962; Dierssen, 1982) or from river valleys in lowland areas (Küsel, 1956; Bakker, 1957; Neumann, 1957; Dierschke, 1968; Tüxen & Dierschke, 1975).

Since most authors present rather limited information on habitat characteristics we will compare synecological conditions of the different communities in the following way. The associations with *C. aquatilis* have been ranked along two environmental gradients (see Fig. 6):

- A trophic gradient, from eutrophic to oligotrophic. By eutrophic we mean that the substrate is rich in silt and the groundwater is rich in minerals (E.C. 25 °C > 300 μS/cm). Oligotrophic means: the substrate is poor in silt and the groundwater is poor in minerals (E.C. 25 °C < 150 μS/cm).
- A wetness gradient, from immersed to drained. Immersed means: high groundwater levels and frequent flooding by (stream) overflow (see Kulczynski, 1949). Emerged means: high, but usually constant groundwater levels sustained

by groundwater emerging at the surface. Flooding can occur, but silt precipitation is of minor importance. Drained means that on average the groundwater table is more than 20 cm below the surface throughout the year.

When we compare the northern *Carex aquatilis* communities with the ones more to the south, large differences and only a few similarities can be observed. In Norway and N Sweden eutrophic floodplain communities such as the *Caricetum acuto-vesicariae* (= *Caricetum gracilis*) and the *Lysimachio-Caricetum aquatilis* are not present. In both the north and the south *Carex aquatilis* is found in *Calthion palustris* meadows which emerge from a very wet fen vegetation dominated by *Carex aquatilis*, after drainage and fertilization (Siira, 1970; Tüxen & Dierschke, 1975; Dierssen, 1982).

The northern *Drepanoclado-Caricetum aquatilis* is a distinct mesotrophic *Caricion lasiocarpae* community (Dierssen, 1982) and lacks *Phragmitetea* species like *Glyceria maxima* and *Rumex hydrolapathum*. The southern *Lysimachio-Caricetum aquatilis* belongs to the *Magnocaricion*, while many *Phragmitetea* species are present. Only the mesotrophic subassociation with *Potentilla palustris*, *Menyanthes trifoliata*, *Lysimachia thyrsiflora* and *Carex rostrata* is almost identical to the *Drepanoclado-Caricetum aquatilis* (see also Dierssen, 1982). This subassociation is very rare in NW Germany and has not yet been found in the Netherlands.

The *Caricetum nigrae* and the *Caricetum curtoechinatae* mentioned in Figure 6 are identical as well. *Carex aquatilis* has a reduced performance here.

*Carex aquatilis* thus has its optimum in habitats where enriched groundwater emerges and where frequent flooding occurs. In northern Europe natural and semi-natural *Carex aquatilis* communities with a distinct mesotrophic character are most frequent according to the literature. Eutrophic floodplain communities with *Carex aquatilis* seem to be rare and lack the *Phragmitetea* species which are characteristic for the NW German and Dutch floodplain communities.

At the southern border of the distribution area, mesotrophic *Carex aquatilis* communities have been registered, but are rare. Most frequent here are eutrophic sedge communities with *Carex acuta*.

Despite the pronounced differences in species

composition between northern and southern *Carex aquatilis* communities, there appears to be little evidence that in former days the species occupied an essentially different habitat at the border of its distribution area compared to the center. At present populations of *Carex aquatilis* in NW Germany and in the Netherlands seem to be under heavy pressure of human activities, which are giving rise to more eutrophic and to drier conditions.

This process of eutrophication in the lowlands is stimulated by both drainage and increased flooding. Drainage of mesotrophic *Carex aquatilis* communities has led to species-rich *Calthion palustris* communities, in which *C. aquatilis* has a reduced fertility. Almost all *C. aquatilis* stands in the middle reaches of Dutch catchments are probably degenerate stages of former mesotrophic communities. Furthermore drainage in upstream areas leads to increased rates of water discharge in the wet season and this is followed by increased flooding of the downstream areas. Thus the downstream areas experience more eutrophic conditions mainly because of the increased silt deposition.

Consequently the *Lysimachio-Caricetum aquatilis comaretosum* is replaced by the *L. - C. a. inops*, or even by the *Caricetum gracilis* (Bakker & Boedeltje, 1980; Brouwer & van den Hof, 1981).

The water surcharge promotes further interference with the hydrology. The construction of large canals with a regulated water discharge will eventually diminish both emersion of groundwater and immersion by floodwater. As a result the disappearance of *Carex aquatilis* stands is almost inevitable.

In former days *Carex aquatilis* communities belonging to the *Caricion lasiocarpae* must have been more widespread in the Dutch and German lowlands. Oligo- and mesotrophic conditions probably prevailed in most of the European brooks, rivulets and lowland marshes (cf. Succow, 1971, 1982; Kulczynski, 1949). These conditions were sustained by a very regular supply of cool groundwater. Because of this welling up of groundwater with a temperature of 9–10 °C the habitat of *C. aquatilis* in the Netherlands is much colder than would be expected climatologically. The findings of Conolly & Dahl (1970) that *C. aquatilis* is not present in areas where summer maximum temperatures exceed 28 °C is no doubt correct, but the temperature in the root zone of *C. aquatilis* is, in fact, at least 10 °C

lower. The maximum temperatures do not exceed 15–17 °C and may be only slightly higher than the maximum temperatures in northern stands. Yet the species seems to be unable to form ripe seeds. Up to now utricles collected in the Netherlands have only been found to contain undeveloped fruits. This might be caused by climatic and/or genetic factors. As the species is at the southern limit of its range the sedge may well be outside the area within which the climate permits regular fruit production. A similar case has been presented by Shaver *et al.* (1979), who found a population of *Carex aquatilis* in the Alaskan tundra that persisted only by means of vegetative propagation. Their research showed distinct ecotypic differentiation in nutrient uptake between the different populations. So gene exchange must have taken place in the distant past and natural selection subsequently gave rise to edaphic ecotypes.

It would be very interesting to center on ecotypic differentiation in Europe and find out exactly by what mechanisms the populations of *Carex aquatilis* are able to survive in habitats characterized by rather eutrophic conditions. Experiments with reciprocal transplantation both here, in N Sweden, and in Alaska, would make a good start to solving these problems (see Chapin & Chapin, 1981).

## References

- Alechin, W., 1927. Die Alluvionen der Flusstäler in Russland. Rept. Spec. Nov. Regni Veget. Beih. 47: 1-79.
- Andreas, C. H., 1951. *Carex aquatilis* Wahlenb. A glacial relict in the Netherlands. Ned. Kruidk. Arch. 58: 48-56.
- Bakker, D., 1957. On the distribution of *Carex aquatilis* Wahlenb. Acta Bot. Neerl. 6: 93-95.
- Bakker, D. & Westhoff, V., 1957. Over de verspreiding van *Carex aquatilis* Wahlenb. Correspondentieblad 2: 13. Leiden.
- Bakker, P. A., 1979. Vegetation science and nature conservation. In: M. J. A. Werger (ed.). The Study of Vegetation, pp. 247-288. Junk, The Hague. 316 pp.
- Balátová-Tuláčková, E., 1978. Die Nass- und Feuchtwiesen Nordwestbohmens mit besonderer Berücksichtigung der Magnocaricetalia-Gesellschaften. Roxpr. CSAV, Ser. Mat.-Natur., Praha 88: 1-113.
- Barkman, J. J. & Westhoff, V., 1969. Botanical evaluation of the Drenthian District. Vegetatio 19: 330-388.
- Bernard, J. M., 1975. The life history of shoots of *Carex lacustris*. Can. J. Bot. 53: 256-260.
- Bernard, J. M. & Gorham, E., 1978. Life history aspects of primary production in sedge wetlands. In: E. Good *et al.* (eds). Freshwater Wetlands, pp. 39-51. Acad. Press.

- Boedeltje, G. & Bakker, J. P., 1980. Vegetation, soil, hydrology and management in a Drenthian brookland. *Acta Bot. Neerl.* 29: 509-522.
- Brouwer, C. & Hof, B. van den, 1981. Successie van de vegetatie op de Kappersbult in relatie met beheer en waterhuishouding. Internal report Dept. of Plant Ecology, Haren. 97 pp.
- Brujin, O. de, 1980. The *Carex* flora of varied landscapes in the Netherlands: an example of decreasing ecological diversity. *Acta Bot. Neerl.* 29: 359-376.
- Cajander, A. K., 1903. Beiträge zur Kenntniss der Vegetationen der Alluvionen des Nördlichen Eurasiens. I: die Alluvionen des unteren Lenatales. *Acta Soc. Sci. Fenn.* 32: 1-182.
- Chapin, A. S. & Chapin, M. C., 1981. Ecotypic differentiation of growth processes in *Carex aquatilis* along altitudinal and local gradients. *Ecology* 62: 1000-1009.
- Churchill, E. D., 1955. Phytosociological and environmental characteristics of some plant communities in the Umiat region of Alaska. *Ecology* 36: 606-627.
- Conolly, A. & Dahl, E., 1970. Maximum summer temperature in relation to the modern and Quaternary distribution of certain arctic-montane species in the British Isles. In: D. Walker & P. G. West (eds.). *Studies in the Vegetational History of the British Isles*, p. 1-265. Cambridge.
- Dierschke, H., 1968. Über eine Grosseeggen-Riedgesellschaft mit *Carex aquatilis* im Wümmetal östlich von Bremen. *Mitt. Flor.-soz. Arbeitsgem.* NF 13: 48-58.
- Dierssen, K., 1982. Die wichtigsten Pflanzengesellschaften der Moore NW-Europa. *Conservatoire et Jardin botaniques, Genève*. 382 pp.
- Ellenberg, H., 1978. *Vegetation Mitteleuropas mit den Alpen*. Stuttgart. 981 pp.
- Elveland, J., 1976. Myrar på Storön vid Norbottenkusten. *Wahlenbergia* 3: 1-274.
- Everts, F. H., Grootjans, A. P. & Vries, N. P. J. de, 1980. De vegetatie van de madelanden in het stroomdal van de Drentsche Aa. Internal report Dept. of Plant Ecology 2 vols., Haren (Gn). 288 pp.
- Gans, W. de, 1982. The Drentsche Aa valley system, a study in Quaternary geology. Thesis Vrije Universiteit, Amsterdam. 130 pp.
- Gischler, C. E., 1976. A semi qualitative study of the hydrogeology of the North Netherlands. Leiden.
- Gorham, E. & Somers, M. G., 1973. Seasonal changes in the standing crop of two montane sedges. *Can. J. Bot.* 51: 1097-1108.
- Grootjans, A. P., 1980. Distribution of plant communities along rivulets in relation to the hydrology and management. *Ber. Int. Symp. I.V.V.* 1979, pp. 143-170. Cramer Verlag.
- Grootjans, A. P. & Klooster, W. Ph. ten, 1980. Changes in groundwater regime in wet meadows. *Acta Bot. Neerl.* 29: 541-554.
- Hengeveld, R. & Haec, J., 1982. The distribution of abundance. *J. Biogeogr.* 9: 303-316.
- Heukels, H. & Oostroom, S. J. van, 1975. *Flora van Nederland*. Wolters-Noordhoff, Groningen. 909 pp.
- Hultén, E., 1962. *The Circumpolar Plants*. I. Kon. Sv. Akad. Handl. 8, 5. Stockholm.
- Jefferies, R. L., 1977. The vegetation of salt marshes at some coastal sites in arctic North America. *J. Ecol.* 65: 661-672.
- Jermy, A. C. & Tutin, T. G., 1969. *British Sedges*. London. 199 pp.
- Kern, J. H. & Reichelt, Th. J., 1954. *Cyperaceae: Carex*, *Flora Neerl.* 1-3. Amsterdam.
- Komarov, V. L. (chief ed.), 1964. *Flora of the U.S.S.R.* I. Translated from Russian, published in Jerusalem.
- Kulczynski, S., 1949. Peatbogs of Polesie. *Mem. Ac. Sci. Cracovie B*: 1-356.
- Londo, G., 1976. The decimal scale for relevés of permanent quadrats. *Vegetatio* 33: 61-64.
- McVean, D. & Ratcliffe, D., 1962. *Plant communities of the Scottish Highlands. A study of Scottish Mountain, Moorland and Forest Vegetation*. I. London. 429 pp.
- Neumann, A., 1975. *Carex aquatilis* auch in Deutschland. *Mitt. Flor.-soz. Arbeitsgem.* NF 6: 172-182.
- Niemann, E., 1973. Grundwasser und Vegetationsgefuge. *Nova Acta Leopoldina* 38: 1-172.
- Ploeg, D. T. E. and der, 1977. *Atlas fan de Floara fan Fryslan*. Leeuwarden. 171 pp.
- Shaver, G. R., Chapin, F. S. & Dillings, W. D., 1979. Ecotypic differentiation in *Carex aquatilis* on ice-wedge polygons in the Alaskan coastal tundra. *J. Ecol.* 67: 1025-1046.
- Siira, J., 1970. Studies in the ecology in the sea-shore meadows of the Bothnian Bay with special reference to the Liminka area. *Aquilo, Ser. Bot.* 9: 1-109.
- Schmidt, G., 1969. *Vegetations Geographie auf ökologisch-soziologischer Grundlage*. Leipzig.
- Smith, A. J. E., 1978. *The moss Flora of Britain and Ireland*. Univ. Press, Cambridge.
- Streefkerk, J. & Hoorn, D. van, 1984. *Hydrologisch onderzoek in het stroomdal van de Drentsche Aa*. Report Staatsbosbeheer Utrecht.
- Succow, M., 1971. Die Talmoore des norddeutschen Flachlandes, ein Beitrag zur Charakterisierung des Moortypes 'Niedermoor'. *Arch. Naturschutz Landsch.* 11: 133-168.
- Tutin, T. G., Heywood, V. H., Burges, N. A., Moore, D. M., Valentine, D. H., Walters, S. M. & Webb, D. A., 1980. *Flora Europaea*, Vol. 5. Cambridge.
- Tüxen, R. & Dierschke, H., 1975. Die Vegetation des Langholter- und Rhauder Meeres und seiner Randgebiete. *Mitt. Flor.-Soz. Arbeitsgem.* N.F. 18: 157-202.
- Waldemarson-Jensen, E., 1979. Succession in relation to lagoon development in the Laiture delta, North Sweden. *Acta Phytogeogr. Suec.* 66: 1-120.
- Westhoff, V. & Held, A. J. den, 1969. *Plantengemeenschappen in Nederland*. Zutphen. 423 pp.
- Working Group Geohydrological Research in Drenthe, 1978. *Geohydrologisch onderzoek in de provincie Drenthe*. Rijks Instituut voor Drinkwatervoorziening. Voorburg. 96 pp.