

University of Groningen

Purification and Characterization of an L-Aminopeptidase from *Pseudomonas putida* ATCC 12633

Hermes, H.F.M.; Sonke, T.; Peters, P.J.H.; Balken, J.A.M. van; Kamphuis, J.; Dijkhuizen, L.; Meijer, E.M.

Published in:
Applied and environmental microbiology

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1993

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hermes, H. F. M., Sonke, T., Peters, P. J. H., Balken, J. A. M. V., Kamphuis, J., Dijkhuizen, L., & Meijer, E. M. (1993). Purification and Characterization of an L-Aminopeptidase from *Pseudomonas putida* ATCC 12633. *Applied and environmental microbiology*, 59(12), 4330-4334.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Purification and Characterization of an L-Aminopeptidase from *Pseudomonas putida* ATCC 12633

H. F. M. HERMES,¹ T. SONKE,¹ P. J. H. PETERS,¹ J. A. M. VAN BALKEN,¹
J. KAMPHUIS,¹ L. DIJKHUIZEN,^{2*} AND E. M. MEIJER¹

DSM Research, Bio-organic Chemistry section, P.O. Box 18, 6160 MD Geleen,¹ and Department of Microbiology, University of Groningen, Kerklaan 30, 9571 NN Haren,² The Netherlands

Received 5 April 1993/Accepted 25 August 1993

An L-aminopeptidase of *Pseudomonas putida*, used in an industrial process for the hydrolysis of D,L-amino acid amide racemates, was purified to homogeneity. The highly L-enantioselective enzyme resembled thiol reagent-sensitive alkaline serine proteinases and was strongly activated by divalent cations. It possessed a high substrate specificity for dipeptides and α -H amino acid amides, e.g., L-phenylglycine amide.

Various chemoenzymatic processes involving resolution of racemic mixtures for the production of chiral compounds have found commercial applications (16). Examples are processes for enantiomerically pure amino acids, versatile chiral building blocks for the synthesis of pharmaceuticals, agrochemicals, and food or feed additives (7-9, 12, 13). Knowledge on the substrate specificity and enantioselectivity of the proteinases, amidases, and aminopeptidases involved in these processes is limited (6, 7, 10, 11, 14). Here we report the purification and characterization of an L-aminopeptidase from *Pseudomonas putida* ATCC 12633, used as a whole-cell biocatalyst for the enantioselective hydrolysis of a broad range of D,L-amino acid amide racemates (1, 8).

The racemic mixtures of amino acid amides used were obtained as follows. Starting out from the corresponding aldehydes, aminonitriles were derived via the Strecker reaction (HCN, NH₃) and converted into the amides under alkaline conditions in the presence of a catalytic amount of a ketone. Enantiomerically pure amide and acid derivatives of phenylglycine (PG) and valine were prepared according to the enzymatic resolution process of DSM (1, 8). Other enantiomerically pure amides were from Bachem (Bubendorf, Switzerland).

P. putida ATCC 12633 was grown in 10 liters of mineral medium (1.55 g of K₂HPO₄, 0.79 g of NaH₂PO₄ · H₂O, 0.077 g of MgCl₂ · 6H₂O, 0.2 ml of trace element solution [15] in 1 liter of distilled water, all heat sterilized) with 15 mM sodium citrate (heat sterilized) and 10 mM L-valine amide (L-Val-NH₂; filter sterilized) at 30°C and pH 7 to 7.5 in a 15-liter fermentor. Cells (about 40 g [wet weight]) were harvested from the late stationary phase by centrifugation (15,000 × g at 4°C for 15 min).

The L-aminopeptidase, routinely assayed at 40°C in 200 mM Tris(hydroxymethyl)methyl-3-aminopropanesulfonic acid (TAPS) · KOH, pH 9.0, with L-PG-NH₂ (133 mM) as a substrate, was purified (at 4°C) to homogeneity by a combination of French pressure cell disintegration (at 1,000 MPa; 1.4 × 10⁵ kN · m⁻²), isolation of cell envelopes by centrifugation (at 70,000 × g at 4°C for 45 min), extraction of the enzyme from the cell envelopes with MgSO₄ (100 mM), and Mono S HR 5/5 cation-exchange chromatography (Table 1; Fig. 1). Detergents such as Triton X-100, successfully used to extract an arginine aminopeptidase from membranes of

TABLE 1. Purification of L-aminopeptidase from *P. putida* ATCC 12633

| Purification step | Total activity (μmol · min ⁻¹) | Total protein (mg) | Sp act (μmol · min ⁻¹ · mg ⁻¹) | Recovery (%) | Purification (fold) |
|-----------------------|--|--------------------|---|--------------|---------------------|
| Whole cells | 4,507 | 657 | 7 | 100 | 1 |
| Cell envelopes | 2,342 | 194 | 12 | 52 | 1.7 |
| Cell envelope extract | 1,453 | 24 | 62 | 32 | 9 |
| Mono S | 946 | 3 | 338 | 21 | 49 |

Streptococcus sanguis 903 (4), were ineffective in this case. Further characterization of the protein revealed a native molecular mass of approximately 400 kDa (by gel filtration chromatography on a Hiloal 26/60 Superdex 200 column) and subunits of 53 kDa (by sodium dodecyl sulfate [SDS]-




FIG. 1. SDS-PAGE of the enzyme fractions obtained during purification of the L-aminopeptidase. Electrophoresis was performed with molecular weight standard proteins (lanes a, d, e, and h), 100 ng of cell envelope extract (lanes b and c), and 50 ng of protein after Mono S HR 5/5 column chromatography (lanes f and g). Protein bands were visualized by silver staining. (Coreldraw 3.0 format.)

* Corresponding author.




FIG. 2. Effects of several proteinase inhibitors on the L-aminopeptidase activity. Purified enzyme (1.3 μg) was incubated with compounds to be tested in 12.5 mM Tris \cdot H₂SO₄, pH 8.0, at 40°C for 30 min, and the residual activity was determined in the standard assay. Activity is expressed as a percentage of the activity of the nonpreincubated enzyme, with 100% activity corresponding to 363 $\mu\text{mol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$. Symbols: ○, diisopropylfluorophosphate; ●, iodoacetamide; ▼, phenylmethylsulfonyl fluoride; ◆, *o*-phenanthroline; ▲, *p*-chloromercuribenzoate; ■, EDTA; +, dithiothreitol.

polyacrylamide gel electrophoresis [PAGE] [Fig. 1]), suggesting that it has an octameric structure. The native molecular weight of the *P. putida* enzyme is relatively high but comparable to, for instance, that of a *Sulfolobus* aminopeptidase (5). The isoelectric point of the protein was estimated at pH 10.5 (by using an IEF Phastgel with an expanded pH range), which is extremely high compared with those of other aminopeptidases (11). Highest enzyme activity was observed at pH 9.5 and 40°C.

Divalent cations had a marked effect on enzyme activity: Mg²⁺, Co²⁺ (2- to 3-fold), and especially Mn²⁺ ions (12-fold) stimulated activity (at 0.2 to 2 mM), whereas treatment with Cu²⁺ and Ca²⁺ ions (at 2 mM) caused 70 and 40% inhibition, respectively. Similar observations have been made for various other aminopeptidases (11).





The *P. putida* L-aminopeptidase was sensitive to various proteinase inhibitors (Fig. 2), e.g., the serine protease inhibitors phenylmethylsulfonyl fluoride and diisopropylfluorophosphate. The enzyme thus resembles the alkaline serine proteinases (6). In addition, the enzyme was inhibited by the chelators EDTA and *o*-phenanthroline and the thiol reagents *p*-chloromercuribenzoate and iodoacetamide but was stimulated by dithiothreitol (Fig. 2). These phenomena may be due to the presence of a cysteine residue near the active site and the involvement of divalent metal ions in the catalytic mechanism.

The substrate specificity of the purified enzyme (1.2 to 1.8 μg) was tested with a range of amide substrates (50 to 100 mM) (Table 2) and dipeptides (2 mM). The ammonium ions produced from the amide substrates were determined with

an ammonia electrode system (Orion model 95-12). Amino acids produced from dipeptides were qualitatively analyzed on silica gel 60 F₂₅₄ thin-layer chromatography plates (E. Merck AG, Darmstadt, Germany) by using a chloroform-methanol-ammonia (60:45:20) mixture as a mobile phase, followed by spraying with a 0.3% ninhydrin solution and incubation at 125°C for 5 min. The enzyme was only active with α -amino acid amides with a H atom at the C _{α} position. Highest activities were observed with L-Leu-NH₂ and L-PG-NH₂. The enzyme was inactive with L- α -CH₃-Val-NH₂, the single α -methyl-substituted amino acid amide tested (Table 2). High activities were also observed with the four aliphatic and aromatic dipeptides tested (L-Phe-L-Phe, L-Phe-L-Leu, L-Leu-L-Phe, and L-Leu-L-Leu), designating the purified enzyme as an L-aminopeptidase.

The kinetic properties (K_m and V_{max} values) of the purified enzyme (1.8 μg) were estimated from double-reciprocal and Eadie-Hofstee plots of initial rates of hydrolysis of amino acid amides as a function of substrate concentration (Table 3). The enzyme displayed normal Michaelis-Menten type of kinetics for the six L- α -amino acid amides tested (Table 3), the products of which were determined by high-pressure liquid chromatography (HPLC) analysis (2). The additional methyl groups at the C atoms (C _{β}) adjacent to the C _{α} atoms in L-Val-NH₂ and L-Ile-NH₂ appears to result in relatively high K_m and low V_{max} values and consequently a low catalytic efficiency and specificity (k_{cat}/K_m values) for these substrates. The highest k_{cat}/K_m value was observed for L-Leu-NH₂, where this methyl group at the C _{β} atom is missing (Tables 2 and 3). Similarly, amongst the two aro-

TABLE 2. Substrate specificity of the purified L-aminopeptidase of *P. putida* ATCC 12633

| Substrate | Formula | R ¹ | R ² | R ³ | Relative activity |
|--|---|------------------|---|------------------|-------------------|
| Acetamide ^a | CH ₃ -CONH ₂ | -H | -H | -H | <5 |
| Propionamide ^a | CH ₃ -CH ₂ -CONH ₂ | -H | -CH ₃ | -H | <5 |
| Butyramide ^a | CH ₃ -CH ₂ -CH ₂ -CONH ₂ | -H | -CH ₂ -CH ₃ | -H | <5 |
| Isobutyramide ^a | (CH ₃) ₂ CH-CONH ₂ | -H | -CH ₃ | -CH ₃ | <5 |
| Acrylamide ^a | CH ₂ =CH-CONH ₂ | -H | -CH ₂ | - | <5 |
| Fluoroacetamide ^a | F-CH ₂ -CONH ₂ | -H | -H | -F | <5 |
| D,L-Mandelic acid amide ^a | C ₆ H ₅ -CH(OH)-CONH ₂ | -H | -C ₆ H ₅ | -OH | <5 |
| Pivalamide ^a | (CH ₃) ₃ C-CONH ₂ | -CH ₃ | -CH ₃ | -CH ₃ | <5 |
| Methacrylamide ^a | CH ₂ =C(CH ₃)-CONH ₂ | -CH ₃ | -CH ₂ | - | <5 |
| Nicotinamide ^a |  | - | - | - | <5 |
| Glycine amide ^b | H-CH(NH ₂)-CONH ₂ | -H | -H | -NH ₂ | <5 |
| L-Alanine amide ^b | CH ₃ -CH(NH ₂)-CONH ₂ | -H | -CH ₃ | -NH ₂ | <5 |
| L-α-Aminobutyramide ^b | CH ₃ -CH ₂ -CH(NH ₂)-CONH ₂ | -H | -CH ₂ -CH ₃ | -NH ₂ | 10 |
| L-Valine amide ^b | (CH ₃) ₂ CH-CH(NH ₂)-CONH ₂ | -H | -CH(CH ₃) ₂ | -NH ₂ | 5 |
| L-Leucine amide ^b | (CH ₃) ₂ CH-CH ₂ -CH(NH ₂)-CONH ₂ | -H | -CH ₂ -CH(CH ₃) ₂ | -NH ₂ | 215 |
| L-Isoleucine amide ^b | CH ₃ -CH ₂ -CH(CH ₃)-CH(NH ₂)-CONH ₂ | -H | -CH(CH ₃)-CH ₂ -CH ₃ | -NH ₂ | 15 |
| L-Phenylglycine amide ^{a,b} | C ₆ H ₅ -CH(NH ₂)-CONH ₂ | -H | -C ₆ H ₅ | -NH ₂ | 100 |
| L-Phenylalanine amide ^b | C ₆ H ₅ -CH ₂ -CH(NH ₂)-CONH ₂ | -H | -CH ₂ -C ₆ H ₅ | -NH ₂ | 15 |
| L-Methionine amide ^b | CH ₃ -S-CH ₂ -CH ₂ -CH(NH ₂)-CONH ₂ | -H | -CH ₂ -CH ₂ -S-CH ₃ | -NH ₂ | 60 |
| D,L-Proline amide ^a |  | -H | -CH ₂ -CH ₂ -CH ₂ - | -NH- | <5 |
| L-Tryptophan amide ^b |  | -H | -CH ₂ -  | -NH ₂ | 15 |
| L-Serine amide ^b | HO-CH ₂ -CH(NH ₂)-CONH ₂ | -H | -CH ₂ OH | -NH ₂ | <5 |
| L-Glutamic acid amide ^b | HOOC-CH ₂ -CH ₂ -CH(NH ₂)-CONH ₂ | -H | -CH ₂ -CH ₂ -COOH | -NH ₂ | <5 |
| D,L-α-Methyl valine amide ^a | (CH ₃) ₂ CH-C(CH ₃)(NH ₂)-CONH ₂ | -CH ₃ | -CH(CH ₃) ₂ | -NH ₂ | <5 |

^a Reactions were performed in Tris-H₂SO₄ (pH 8.0); activity towards L-PG-NH₂ (100%) was 180 μmol · min⁻¹ · mg of protein⁻¹.

^b Reactions were performed in TAPS-KOH (pH 9.0); activity towards L-PG-NH₂ was 490 μmol · min⁻¹ · mg of protein⁻¹.

TABLE 3. Kinetic parameters for the hydrolysis of different amino acid amides by purified L-aminopeptidase of *P. putida* ATCC 12633

| Substrate | -R | K_m^a | V_{max}^b | k_{cat}^c | k_{cat}/K_m^d |
|----------------------------|--|---------|-------------|-------------|-----------------|
| L-PG-NH ₂ | -C ₆ H ₅ | 65 | 1,565 | 1,380 | 20,580 |
| L-Phe-NH ₂ | -CH ₂ -C ₆ H ₅ | 15 | 80 | 68 | 4,525 |
| L-Phe-NH ₂ .HCl | -CH ₂ -C ₆ H ₅ | 15 | 70 | 65 | 3,910 |
| L-Val-NH ₂ .HCl | -CH(CH ₃) ₂ | 130 | 110 | 95 | 715 |
| L-Leu-NH ₂ .HCl | -CH ₂ -CH(CH ₃) ₂ | 15 | 1,915 | 1,690 | 99,215 |
| L-Ile-NH ₂ .HCl | -CH(CH ₃)-CH ₂ -CH ₃ | 150 | 270 | 240 | 1,604 |

^a K_m is expressed in millimolar.

^b V_{max} is expressed in micromoles per minute per milligram of protein.

^c k_{cat} is expressed per second.

^d k_{cat}/K_m is expressed per second per molar.

matic amino acid amides tested, the lowest K_m value was observed with L-Phe-NH₂ containing a C_β atom without further substituents. However, in this case the highest catalytic efficiency was observed with L-PG-NH₂, indicating that the presence of an aromatic ring directly adjacent to the C_α atom has a strongly positive effect.

The enantioselectivity of the purified enzyme (2.1 μg) towards racemic mixtures of Leu-NH₂ and PG-NH₂ (133 mM) was studied by chiral HPLC (3) to determine the concentrations of both the amino acid and the amino acid amide enantiomers after overnight incubation. The enzyme system displayed high L-enantioselectivity towards both the aliphatic and the aromatic α-H amino acid amide. The enantiomeric excess of the L-amino acids formed varied from 96% for PG to more than 99% for Leu, corresponding with calculated E values of 79 and more than 1,000, respectively (Table 4).

Comparison of the characteristics of this *P. putida* aminopeptidase with those of other aminopeptidases and (amino) amidases described (e.g., reference 11) leads to the conclusion that this enzyme is unique not only with respect to its physicochemical characteristics but especially also concerning its high enantioselectivity and substrate specificity. Further studies are required to elucidate the physiological role of the enzyme and its reaction mechanism.

Thanks are due to W. Kortenoeven for preparation of the figures and tables, L. Kerkhoffs and B. Dassen for the preparation of the amino acids and derivatives, B. Kaptein for evaluations on the

enantioselectivity and substrate specificity, and W. Harder for stimulating discussions.

REFERENCES

- Boesten, W. H. J. December 1986. European patent 0.232.562.
- Duchateau, A., and M. Crombach. 1987. Determination of α-aminonitriles, α-amino acid amides and α-amino acids by means of HPLC, post-column reaction and fluorescence detection. *Chromatographia* 24:339-344.
- Duchateau, A., M. Crombach, M. Aussems, and J. Bongers. 1989. Determination of the enantiomers of α-amino acids and α-amino acid amides by high-performance liquid chromatography with a chiral mobile phase. *J. Chromatogr.* 461:419-428.
- Floderus, E., L. E. Linder, and M.-L. Sund. 1990. Characterization of membrane-associated arginine aminopeptidase in *Streptococcus sanguis* 903. *Curr. Microbiol.* 21:145-149.
- Hanner, M., B. Redl, and G. Stöffler. 1990. Isolation and characterization of an intracellular aminopeptidase from the extreme thermophilic archaeobacterium *Sulfolobus solfataricus*. *Biochim. Biophys. Acta* 1033:148-153.
- Kalisz, H. M. 1988. Microbial proteinases. *Adv. Biochem. Eng. Biotechnol.* 36:1-65.
- Kamphuis, J., W. H. J. Boesten, Q. B. Broxterman, H. F. M. Hermes, J. A. M. van Balken, E. M. Meijer, and H. E. Schoemaker. 1990. New developments in the chemo-enzymatic production of amino acids. *Adv. Biochem. Eng. Biotechnol.* 42:133-186.
- Kamphuis, J., H. F. M. Hermes, J. A. M. van Balken, H. E. Schoemaker, W. H. J. Boesten, and E. M. Meijer. 1990. Chemo-enzymatic synthesis of enantiomerically pure α-H and α-alkyl α-amino acids and derivatives, p. 119-125. *In* G. Lubec and G. A. Rosenthal (ed.), *Amino acids: chemistry, biology and medicine*. ESCOM Science Publishers B.V., Leiden, The Netherlands.
- Kamphuis, J., M. Kloosterman, H. E. Schoemaker, W. H. J. Boesten, and E. M. Meijer. 1987. Chiral intermediates and applications, p. 331-348. *In* O. M. Neijssel, R. R. van der Meer, and K. C. A. M. Luyben (ed.), *Proceedings of the 4th European Congress on Biotechnology*, vol. 4. Elsevier, Amsterdam.
- Lazdunski, A. M. 1989. Peptidases and proteases of *Escherichia coli* and *Salmonella typhimurium*. *FEMS Microbiol. Rev.* 63: 265-276.
- Maestracci, M., K. Bui, A. Thiéry, A. Arnaud, and P. Galzy. 1988. The amidases from a *Brevibacterium* strain: study and

TABLE 4. Enantioselectivity of the L-aminopeptidase of *P. putida* ATCC 12633

| Substrate | ee _{L-acid} ^a | c ^b | E ^c |
|-------------------------|-----------------------------------|----------------|----------------|
| D,L-PG-NH ₂ | 0.96 | 38 | 79 |
| D,L-Leu-NH ₂ | >0.99 | 46 | >1,000 |

^a ee_{L-acid} = (L-acid - D-acid)/(L-acid + D-acid). ee, enantiomeric excess.

^b c = (L-acid + D-acid)/(L-amide_{t=0} + D-amide_{t=0}) × 100. c, conversion.

^c E = ln[1 - c(1 + ee_{L-acid})]/ln[1 - c(1 - ee_{L-acid})].

- applications. *Adv. Biochem. Eng. Biotechnol.* **36**:66–115.
12. **Meijer, E. M., W. H. J. Boesten, H. E. Schoemaker, and J. A. M. van Balken.** 1985. Use of biocatalysts in the industrial production of specialty chemicals, p. 135–156. *In* J. Tramper, H. C. van der Plas, and P. Linko (ed.), *Biocatalysts in organic syntheses*. Elsevier, Amsterdam.
 13. **Meijer, E. M., J. Kamphuis, J. A. M. van Balken, H. F. M. Hermes, W. J. J. van den Tweel, M. Kloosterman, W. H. J. Boesten, and H. E. Schoemaker.** 1990. Advantages and limitations of applied biocatalysis, p. 363–382. *In* V. Claassen (ed.), *Trends in drug research*. Elsevier, Amsterdam.
 14. **Morihara, K.** 1974. Comparative specificity of microbial proteinases. *Adv. Enzymol.* **41**:179–243.
 15. **Vishniac, W., and M. Santer.** 1975. The thiobacilli. *Bacteriol. Rev.* **21**:195–213.
 16. **Yonaha, K., and K. Soda.** 1986. Applications of stereoselectivity of enzymes: synthesis of optically active amino acids and α -hydroxy acids, and stereospecific isotope-labelling of amino acids, amines and coenzymes. *Adv. Biochem. Eng. Biotechnol.* **33**:95–130.