

University of Groningen

## Gamma - ray emission in yhe CF-252 spontaneous fission process

van der Ploeg, H; Postma, R; Bacelar, JCS; van den Berg, T; Jacob, VE; Jongman, J.R.; van der Woude, A

*Published in:*  
Physical Review Letters

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1995

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

van der Ploeg, H., Postma, R., Bacelar, JCS., van den Berg, T., Jacob, VE., Jongman, J. R., & van der Woude, A. (1995). Gamma - ray emission in yhe CF-252 spontaneous fission process: Reply. *Physical Review Letters*, 74(15), 3082-3082.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Gamma Ray Emission in the $^{252}\text{Cf}$ Spontaneous Fission Process

In a recent Letter, van der Ploeg *et al.* [1] reported the observation of a large anisotropy of photons emitted in the spontaneous fission of  $^{252}\text{Cf}$ . In the energy region between 8 and 12 MeV, gamma-ray emission parallel to the fission direction was found to be strongly enhanced. In the analysis of van der Ploeg *et al.* sum-up effects were neglected. However, sum up of gamma rays and neutrons from the same fission process may be very important, since neutrons are strongly boosted in the fission direction.

We have measured the angular distribution of gamma rays accompanying the spontaneous fission of  $^{252}\text{Cf}$  with the setup shown in Fig. 1. Fission fragments were detected by means of five  $\text{Al}_2\text{O}_3$  foils and microchannel plate (MCP) detectors. For the gamma rays we used six  $\text{BaF}_2 + \text{BGO}$  detectors. Neutrons with an energy below 25 MeV could be discriminated by their time of flight.

Because of the combined symmetry of the fission process and the measuring apparatus, the gamma-ray anisotropy could be determined,

$$\frac{W(0^\circ)}{W(90^\circ)} = \frac{\{N[(\gamma_1 \vee \gamma_2) \wedge (\text{MCP}_1 \vee \text{MCP}_2)] - N[(\gamma_1 \wedge \gamma_2) \wedge (\text{MCP}_1 \vee \text{MCP}_2)]\}N[(\gamma_3 \vee \gamma_4) \wedge \text{MCP}_3]}{\{N[(\gamma_3 \vee \gamma_4) \wedge (\text{MCP}_1 \vee \text{MCP}_2)] - N[(\gamma_3 \wedge \gamma_4) \wedge (\text{MCP}_1 \vee \text{MCP}_2)]\}N[(\gamma_1 \vee \gamma_2) \wedge \text{MCP}_3]}$$

$N$  [coincidence requirement] denotes the number of coincidences indicated. The second term within the curly brackets is a correction for sum-up effects. The  $(\gamma_1 \wedge \gamma_2)$  in this term not only stands for a coincidence requirement but also for a software summation for each registered event of the energies in the two detectors. The anisotropy was determined for different energy intervals both with and without sum-up correction. The result is present in Fig. 2.

The importance of the sum-up correction is clearly demonstrated in our result. A large anisotropy, similar to that reported in Ref. [1], was found for the 8–12 MeV energy region when no sum-up correction was applied. However, after the proper correction no anisotropy remains for high-energy gamma rays.

Chr. Bargholtz and Kj. Fransson  
Fysikum, University of Stockholm, Box 6730  
S-113 85, Stockholm, Sweden

O. Batenkov, M. Majorov, S. Smirnov, and A. Veshikov  
V. G. Khlopin Radium Institute, 28 Murinskij Avenue  
194021 St. Petersburg, Russia

Received 12 December 1994

PACS numbers: 25.85.Ca, 21.10.Re, 27.90.+b

- [1] H. van der Ploeg, R. Postma, J.C. Bacelar, T. van den Berg, V.E. Jacob, J.R. Jongman, and A. van der Woude, *Phys. Rev. Lett.* **68**, 3145 (1992); **69**, 1148(e) (1992).




FIG. 1. Experimental setup. The  $^{252}\text{Cf}$  source is sandwiched between  $\text{Al}_2\text{O}_3$  foils providing the electrons for the start signal from the two ring shaped MCP's.




FIG. 2. Angular correlation ratios (cf. text) as a function of the energy deposited in the gamma-ray detectors. Filled circles and unfilled diamonds show the ratios without and with sum-up corrections, respectively. The inserted table gives the integrated ratios for two energy intervals.