

University of Groningen

## Deep wide field HI imaging of Messier 31

Chemin, Laurent; Carignan, Claude; Foster, Tyler

*Published in:*

Panoramic Radio Astronomy: Wide-field 1-2 GHz research on galaxy evolution

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2009

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Chemin, L., Carignan, C., & Foster, T. (2009). Deep wide field HI imaging of Messier 31. In *Panoramic Radio Astronomy: Wide-field 1-2 GHz research on galaxy evolution* University of Groningen.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Deep wide field HI imaging of M31

---

### Laurent Chemin\*

*GÉPI, Observatoire de Paris, section Meudon, CNRS & Université Paris 7 UMR 8111, 5 Place Jules Janssen, 92195 Meudon, France*  
E-mail: laurent.chemin@obspm.fr

### Claude Carignan

*Laboratoire d'Astrophysique Expérimentale (LAE), Observatoire du mont Mégantic, and Département de physique, Université de Montréal, C.P. 6128, Succ. Centre-Ville, Montréal, QC, Canada H3C 3J7*  
*Observatoire d'Astrophysique de l'Université de Ouagadougou (UFR/SEA), 03 BP 7021 Ouagadougou 03, Burkina Faso*

### Tyler Foster

*Department of Physics & Astronomy, Brandon University, Brandon, MB, Canada R7A 6A9*

We report on preliminary results from a new deep 21-cm survey of the Andromeda galaxy, based on observations performed with the Synthesis Telescope and the 26-m antenna at DRAO. The HI distribution and kinematics of the disc are analyzed and basic dynamical properties are derived. New HI structures are discovered, like thin HI spur-like structures and an external arm in the disc outskirts. The HI spurs are related to perturbed stellar clumps outside the main disc of M31. The external arm lies on the far, receding side of the galaxy and has no obvious counterpart in the opposite side. These HI perturbations probably result from tidal interactions with companions. It is found a dynamical mass of  $(3.9 \pm 0.3) \times 10^{11} M_{\odot}$  enclosed within a radius  $R = 36$  kpc and a total mass of  $\sim 7.9 \times 10^{11} M_{\odot}$  inside the virial radius.

*Panoramic Radio Astronomy: Wide-field 1-2 GHz research on galaxy evolution - PRA2009*  
*June 02 - 05 2009*  
*Groningen, the Netherlands*

---

\*Speaker.

## 1. Introduction

Understanding the formation and evolution of galaxies like the Milky Way is one of the major goal of astrophysics. The Andromeda galaxy (M31) is very well suited to put constraints on the physical processes that control the evolution of spiral galaxies because of its proximity. The important stellar features related to the evolution of M31 are the faint, extended and perturbed structures seen in the disc outskirts, in addition to many of the dwarf companions that have been detected in its close neighborhood ([8]). They are undoubtedly the imprints of the hierarchical growth of the stellar disc and halo of M31, similar to those seen in numerical models of dark matter and galaxy evolution in the framework of the Cold Dark Matter paradigm (e.g. [11]).

In this work we are interested in providing a detailed view of the neutral hydrogen disc of M31 from recent, deep, wide-field and high angular HI imaging. Our direct objectives are (i) to study the most extended HI distribution of M31, (ii) to derive an accurate HI rotation curve for it, and (iii) to derive its basic dynamical parameters in order to put further constraints on the history of its mass assembly. Preliminary results are described hereafter. A complete analysis of the data are presented in Chemin et al. (2009, [5]).

## 2. Observations


The HI observations were performed with the Synthesis Telescope and the 26-m dish at the Dominion Radio Astrophysical Observatory (DRAO) between September and December 2005. Five fields were observed in the direction of M31 for a total exposure time of 144 hours per field. The spectral resolution is  $5.3 \text{ km s}^{-1}$  and the angular size of the synthesized beam is  $\sim 60'' \times 90''$ . It samples a linear scale of  $\sim 230 \text{ pc} \times 340 \text{ pc}$  at the M31 distance (785 kpc, [9]).

## 3. Results

The integrated emission is displayed in Figure 1 (left-hand panel). The high resolution HI map shows a disc with very little gas in its central regions, as usually observed in early-type discs. Faint spiral-like or ring-like structures are observed at  $R \sim 3 \text{ kpc}$  and  $R \sim 8 \text{ kpc}$ . They coincide with dusty ring-like structures observed in NIR images from SPITZER-IRAC data ([1]) as well as with molecular gas ring-like structures ([10]). Other brighter spiral- or ring-like structures are observed at  $R \sim 10 \text{ kpc}$ ,  $13 \text{ kpc}$  and  $16 \text{ kpc}$ . Part of these HI structures have already been presented in previous studies of M31 ([3, 12]).

New faint structures that were not seen in old HI images are discovered. First the two disc extremities exhibit thin spur-like extensions, particularly towards the North-East. Their kinematics are in good continuity to the adjacent inner disc. Velocity gradients are detected in these spurs ( $\sim 20 \text{ km s}^{-1}$  along  $\sim 7 - 10 \text{ kpc}$ ). These spurs appear tightly linked to stellar clumps (the ‘‘G1’’ clump and the NE extension, as identified in [8]).

Then an external spiral arm is discovered on the edge of the receding half of the disc. It is outlined with dashed lines in Figure 1 (right-hand panel). The HI mass of this new arm is  $\sim 10^8 M_{\odot}$ . Its apparent length is  $\sim 32 \text{ kpc}$ . It is connected to another more extended, brighter spiral arm. The arm is clumpy towards its northeastern end. Part of the thin NE HI spurs is a kinematical



**Figure 1: Left-hand panel:** HI integrated emission of Messier 31. **Right-hand panel:** 3D view of the HI datacube of M31. The top panel is the position-velocity diagram of the full datacube projected onto the photometric major axis. The bottom panel is the same map as in the right-hand panel but displayed with the major axis parallel to the horizontal axis. Dashed lines show the location of the newly discovered external arm (see text for details).

extension of that external arm, as seen in the position-velocity plot of Fig. 1. It is striking that the external arm has no evident morphological and kinematical counterpart in the approaching half of the disc with respect to the galactic centre. Moreover, it is obvious that its kinematics is very peculiar compared to the disc velocities. As a consequence it is very likely that this spiral-like structure has been generated by external effects to M31, either by tidal effects after the passage of a galaxy satellite or even by gas accretion from the intergalactic medium or from a gas rich companion. Numerical simulations are necessary to investigate the different possibilities and find the origin of the faint new perturbations. Notice that we are confident about their detection because they are also observed in the recent HI map of M31 by Braun et al. (2009, [2]).

#### 4. Dynamical analysis

Our previous dynamical analysis of M31 allowed us to derive a total mass of  $3.4 \times 10^{11} M_{\odot}$  for  $R < 35$  kpc from single dish data obtained at Effelsberg and GBT ([4]). A new, more extended rotation curve is derived from a tilted-ring analysis of the velocity field of M31. A mass distribution model is fitted to the rotation curve. As provisional results, it is derived a dynamical mass of  $M_{\text{Dyn}} = (3.9 \pm 0.3) \times 10^{11} M_{\odot}$  inside  $R = 36$  kpc and a dark-to-baryonic mass ratio of  $M_{\text{Dark}}/M_{\text{Baryon}} \sim 3.0$  (75% of dark matter, 25% of luminous baryons). Here  $M_{\text{Baryon}}$  represents the sum of the black hole, gaseous and stellar masses,  $M_{\text{Dark}}$  the dark matter mass, all values integrated within  $R = 36$  kpc. The total mass of M31 extrapolated to the virial radius ( $R \sim 190$  kpc, [8]) is  $M_{\text{Vir}} \sim 7.9 \times 10^{11} M_{\odot}$ . All these measurements are in excellent agreement with results found from other dynamical tracers ([6], [7]). A very good consensus seems to have been obtained for the

enclosed mass inside the inner 36 kpc of M31, as well as for its total mass inside 190 kpc. We refer to Chemin et al. (2009, [5]) for a more detailed discussion of these results.

## Acknowledgements

We are very grateful to the staff of the Dominion Radio Astronomy Observatory at Penticton for their support in obtaining the observations, and especially T. Landecker for encouraging us to pursue this project. We thank the scientific and local committees at ASTRON and Groningen for having organized the PRA 2009 conference and for all stimulating discussions during the conference.

## References

- [1] Barmby P., Ashby M. L. N., Bianchi L., Engelbracht C. W., Gehrz R. D., Gordon K. D., Hinz J. L., Huchra J. P., et al. 2006, *ApJ*, 650, L45
- [2] Braun R., Thilker D. A., Walterbos R. A. M., & Corbelli E. 2009, *ApJ*, 695, 937
- [3] Brinks E., & Shane W. W. 1984, *A&ASS*, 55, 179
- [4] Carignan C., Chemin L., Huchtmeier W. K., & Lockman F. J. 2006, *ApJ*, 641, L109
- [5] Chemin L., Carignan C., Foster T., 2009, *ApJ*, submitted
- [6] Evans N. W., Wilkinson M. I., Guhathakurta P., Grebel E. K., & Vogt S. S. 2000, *ApJ*, 540, L9
- [7] Ibata R., Chapman S., Ferguson A. M. N., Irwin M., & Lewis G., 2004, *MNRAS*, 351, 117
- [8] Ibata R., Martin N. F., Irwin M., Chapman S., Ferguson A. M. N., Lewis G., & McConnachie A. W. 2007, *ApJ*, 671, 1591
- [9] McConnachie A. W., Irwin M. J., Ferguson R. A., Ibata R. A., Lewis G. F., & Tanvir N. 2005, *MNRAS*, 356, 979
- [10] Nieten C., Neiningen N., Guélin M., Ungerechts H., Lucas R., Berkhuijsen E. M., Beck R., & Wielebinski R., 2006, *A&A*, 453, 459
- [11] Springel V., White S. D. M., Jenkins A., Frenk C. S., Yoshida N., Gao L., Navarro J. F., Thacker R., et al., 2005, *Nature*, 435, 629
- [12] Unwin S. C. 1983, *MNRAS*, 205, 787