

University of Groningen

Computational Lipidomics and the Lipid Organization of Cell Envelopes

Ingolfsson, Helgi I.; Melo, Manuel N.; Wassenaar, Tsjerk A.; Periole, Xavier; de Vries, Alex H.; Tieleman, D. Peter; Marrink, Siewert J.

Published in:
Biophysical Journal

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Ingolfsson, H. I., Melo, M. N., Wassenaar, T. A., Periole, X., de Vries, A. H., Tieleman, D. P., & Marrink, S. J. (2015). Computational Lipidomics and the Lipid Organization of Cell Envelopes. *Biophysical Journal*, 108(2), 342A-342A.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

1719-Plat**Analysis of PI(4,5)P2 Lateral Organization at the Plasma Membrane of Living Cells Through FRET**

Maria João Sarmiento¹, Ana Coutinho^{1,2}, Manuel Prieto¹, Fábio Fernandes¹.

¹Centro de Química-Física Molecular and IN, Instituto Superior Técnico - Univ. Lisbon, Lisbon, Portugal, ²Dep. Química e Bioquímica, Faculty of Sciences - Univ. Lisbon, Lisbon, Portugal.

Phosphatidylinositol (4,5)-bisphosphate (PI(4,5)P2) is an important component of the inner leaflet of the plasma membrane of eukaryotic cells. Despite the fact that it only comprises approximately 1 mol% of the total membrane phospholipids, this phosphoinositide has been associated with many different cell functions including membrane trafficking, actin cytoskeleton remodeling and cell motility, among others. The physiological functions of this lipid seem to depend on localized concentration fluctuations within the plasma membrane. In fact, the distribution of this lipid in the plasma membrane has been proposed to be heterogeneous, and PI(4,5)P2 clustering is detected on model membranes under specific conditions. Domains highly enriched in PI(4,5)P2 were also reported at the plasma membrane of specific cell types. However, for most cellular models, scarce evidence has been found for PI(4,5)P2 segregation/clustering in the plasma membrane.

In this context, our main goal was to study the distribution of PI(4,5)P2 molecules in cell lines where no heterogeneity in PI(4,5)P2 lateral distribution had been previously observed. The distribution of PI(4,5)P2 was assessed from FRET microscopy measurements with pleckstrin homology (PH) domains tagged with different fluorescent proteins. We applied a FRET methodology capable of discriminating between FRET from aggregates/clusters from non-interacting molecules within the plasma membrane of living cells. Our results clearly show distinct PI(4,5)P2 local densities in different cellular models, suggesting different patterns of PI(4,5)P2 lateral distributions within the plasma membrane. In addition, the effect of cholesterol removal on PI(4,5)P2 lateral organization is significantly different in distinct cell lines, suggesting that the role of cholesterol in the formation of PI(4,5)P2 enriched domains varies considerably.

This work was supported by FCT - Foundation of Science and Technology (PTDC/QUI-BIQ/119494/2010 and RECI/CTM-POL/0342/2012). M.J.S. and F.F. acknowledge research grants (SFRH/BD/80575/2011 and SFRH/BPD/64320/2009) from FCT.

1720-Plat**Impact of PI(3,4,5)P3-Mediated Beta-Arrestin-1 Recruitment on Structure of Asymmetric Lipid Bilayers**

Achebe N.O. Nzulumike¹, Signe Mathiasen¹, Jacob P. Mahoney², Marité Cárdenas Gómez¹, Dimitrios G. Stamou¹, Kell Mortensen¹.

¹Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark,

²Department of Pharmacology, University of Michigan Medical School, Ann Arbor, MI, USA.

The membrane spanning G-protein coupled receptors (GPCRs) facilitate crucial physiological responses to a variety of extracellular ligands, such as hormones, neurotransmitters, ions, photons, and other stimuli [1], thus 50% of drugs available in the market today target GPCRs [2]. The biological mechanisms behind GPCR-signaling involves several other biomolecules, including the key protein group of β -arrestins. These cytosolic adaptor proteins regulate signaling through several different pathways, classically by removing receptors from the plasma membrane via clathrin-mediated endocytosis [3].

In membrane binding experiments using both confocal fluorescence microscopy and quartz crystal microbalance, we found that β -arrestin-1 binds specifically to supported lipid bilayers (SLBs) containing phosphatidylinositol trisphosphate lipids, PI(3,4,5)P3. We have consistently shown, using both methods, that such binding is lipid specific and not driven by membrane charge. We have also shown that the membrane-protein interaction depends on protein concentration and membrane composition, and that β -arrestins can induce membrane curvature, which may play a role in the endocytic pathway. This is for the first time quantified with respect to dynamics and respective contributions of specific vs non-specific binding.

Having established an assay for PI(3,4,5)P3-mediated recruitment of β -arrestin-1, we conducted a series of neutron reflection experiments to study the protein-lipid interaction on a structural level. Both hydrogenated and deuterated protein was used in reflectometry experiments with diverse contrast matching of the bulk media. We present here the detailed structures of asymmetric lipid bilayers containing PI(3,4,5)P3, and demonstrate that β -arrestin-1 not only binds but also reorganizes the membrane structure.

1. Rosenbaum, D. M. *et al. Nature.* **459**(7245), 356-363 (2009).
2. Gudermann, T. *et al. J Mol Med-Jmm.* **73**(2), 51-63 (1995).
3. DeWire, S. M. *et al. Annu Rev Physiol.* **69**, 483-510 (2007).

1721-Plat**Computational Lipidomics and the Lipid Organization of Cell Envelopes**

Helgi I. Ingólfsson¹, Manuel N. Melo¹, Tsjerk A. Wassenaar¹, Xavier Periole¹, Alex H. de Vries¹, D. Peter Tieleman², Siewert J. Marrink¹.

¹University of Groningen, Groningen, Netherlands, ²University of Calgary, Calgary, AB, Canada.

The detailed lipid organization of cellular membranes remains elusive. A typical plasma membrane contains hundreds of different lipid species that are actively regulated by the cell. Currently over 30,000 biologically relevant lipids have been identified and specific organisms often synthesize thousands of different lipid types. This is far greater diversity than is needed to maintain bilayer barrier properties and to solvate membrane proteins. Why organisms go through the costly progress of creating and maintaining such a large diversity of lipids is one of the big open questions in biology. What is the individual role of these lipids, and how do they interact and organize in the membrane plane?

To start to address these questions we optimized and developed the Martini coarse-grained force field lipidome. We systematically explore over a hundred different lipid types using the Martini model. Bulk properties of each individual lipid type (e.g. bilayer thickness, area per lipid, diffusion, order parameter and area compressibility) were analyzed and overall trends compared to experimental values. Using pre-existing Martini lipids and the newly characterized ones, idealized plasma membranes containing dozens of different lipid types asymmetrically distributed between the membrane leaflets were constructed and simulated on the multi microsecond time scale. In terms of lipid composition these are by an order of magnitude the most complex simulations to date. These large-scale simulations provide a high-resolution view on the lipid organization of plasma membranes at an unprecedented level of complexity and allow us to analyze a variety of plasma membrane physicochemical properties, including: lipid-lipid interactions, bilayer bulk material properties, domain formation and coupling between the bilayer leaflets. Overall, the plasma membranes show global non-ideal mixing of different lipid species at different spatiotemporal scales.

1722-Plat**Phase Behavior of Synaptosomal Membranes: The Effect of Lipid Composition and Temperature**

Atsuko Kimura¹, Gulcin Pekkurmaz², Tomohiro Kimura¹,

Sierra C. Germeyan¹, Jessica Zimmerberg-Helms¹, James Loewke¹,

Paul S. Blank¹, Thomas S. Reese¹, Klaus Gawrisch¹, Ludmila Bezrukov¹,

Joshua Zimmerberg¹.

¹National Institutes of Health, Bethesda, MD, USA, ²F.M. Kirby Neurobiology Center, Boston Children's Hospital, Boston, MA, USA.

The phase behavior of synaptosomal membranes isolated from mammalian rat and marine invertebrate squid intact nerve endings was compared. Homogeneous liquid disordered ($L\alpha$) phase was observed at their body temperatures of 37 and $\sim 20^\circ\text{C}$ for the rat and squid, respectively, using fluorescence microscopy and ^1H -MAS NMR. Temperature decrease resulted in co-existence of an ordered phase with onset temperatures of 24°C for rat and 8°C for squid. Comprehensive analyses of the lipid composition by LC/MS explained the significant difference in the onset temperature: $>2x$ higher content of 18:0 and 18:1 lipid chains in rat, which contributes to a higher transition temperature. The amount of ω -3 chains was $\sim 3x$ higher in squid than in rat due to a difference in the amounts of 22:6 and 20:5: squid membranes contained $2x$ more total 22:6 in major phospholipids (PC, PE, and PS) and cardiolipin (17:0,22:6,22:6,22:6-species). While prominent in rat, 20:4 was not found in squid. It is known for $L\alpha$ phase PC that 16:0,22:6 is more ordered and thicker with a smaller area/lipid than 16:0,20:4. The high content of 22:6 in squid may contribute to order in the $L\alpha$ phase at the body temperature ($\sim 20^\circ\text{C}$) in contrast to the high content of 20:4 in rat at a higher body temperature (37°C). Fluorescence microscopy data showing the temperature-dependent phase behavior using giant unilamellar vesicles prepared with lipid compositions based on the synaptosomal measurements above will be discussed. Synaptosomal preparations of squid with body temperatures differing by $\sim 9^\circ\text{C}$ will be compared.

1723-Plat**Morphology Induced Receptor Trapping in Artificial Dendritic Spines**

Wim Pomp, Thomas Schmidt.

Leiden University, Leiden, Netherlands.

Memory and learning are believed to be regulated by the strength of the connections in the synapses of the 100 billion neurons in the human brain. In the synapse the signal is transmitted between the presynaptic axon and the dendritic spine by neurotransmitters. The number of receptors in the membrane of the dendritic spine defines the strength of a synapse. The persistent increased local concentration of receptors however, is contradicted by the finding of high receptor mobility within the synapse, dependent on spine morphology. The