

University of Groningen

Long Term Outcomes of a Geriatric Liaison Intervention in Frail Elderly Cancer Patients

Hempenius, Liesbeth; Slaets, Joris; van Asselt, Dieneke; de Bock, Truuske H; Wiggers, Theo; van Leeuwen, Barbara L

Published in:
 PLoS ONE

DOI:
[10.1371/journal.pone.0143364](https://doi.org/10.1371/journal.pone.0143364)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hempenius, L., Slaets, J., van Asselt, D., de Bock, T. H., Wiggers, T., & van Leeuwen, B. L. (2016). Long Term Outcomes of a Geriatric Liaison Intervention in Frail Elderly Cancer Patients. *PLoS ONE*, 11(2), [e0143364]. <https://doi.org/10.1371/journal.pone.0143364>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

RESEARCH ARTICLE

Long Term Outcomes of a Geriatric Liaison Intervention in Frail Elderly Cancer Patients

Liesbeth Hempenius^{1,2*}, Joris P. J. Slaets¹, Dienneke van Asselt³, Truuske H. de Bock⁴, Theo Wiggers⁵, Barbara L. van Leeuwen⁵

1 University Center for Geriatric Medicine, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands, **2** Geriatric Center, Medical Center Leeuwarden, Leeuwarden, the Netherlands, **3** Department of Geriatrics, Radboud University Medical Center, Nijmegen, the Netherlands, **4** Department of Epidemiology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands, **5** Department of Surgery, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

* l.hempenius@umcg.nl



CrossMark
click for updates

OPEN ACCESS

Citation: Hempenius L, Slaets JPJ, van Asselt D, de Bock TH, Wiggers T, van Leeuwen BL (2016) Long Term Outcomes of a Geriatric Liaison Intervention in Frail Elderly Cancer Patients. PLoS ONE 11(2): e0143364. doi:10.1371/journal.pone.0143364

Editor: Apar Kishor Ganti, University of Nebraska Medical Center, UNITED STATES

Received: July 19, 2014

Accepted: October 31, 2015

Published: February 22, 2016

Copyright: © 2016 Hempenius et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Due to ethical restrictions with respect to participant identifying information, data are available upon request to the Corresponding Author.

Funding: Funding was obtained from the Netherlands Organisation for Health Research and Development, trial number 945-07-516 (<http://www.zonmw.nl>). The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Background

The aim of this study was to evaluate the long term effects after discharge of a hospital-based geriatric liaison intervention to prevent postoperative delirium in frail elderly cancer patients treated with an elective surgical procedure for a solid tumour. In addition, the effect of a postoperative delirium on long term outcomes was examined.

Methods

A three month follow-up was performed in participants of the Liaison Intervention in Frail Elderly study, a multicentre, prospective, randomized, controlled trial. Patients were randomized to standard treatment or a geriatric liaison intervention. The intervention consisted of a preoperative geriatric consultation, an individual treatment plan targeted at risk factors for delirium and daily visits by a geriatric nurse during the hospital stay. The long term outcomes included: mortality, rehospitalisation, Activities of Daily Living (ADL) functioning, return to the independent pre-operative living situation, use of supportive care, cognitive functioning and health related quality of life.

Results

Data of 260 patients (intervention n = 127, Control n = 133) were analysed. There were no differences between the intervention group and usual-care group for any of the outcomes three months after discharge. The presence of postoperative delirium was associated with: an increased risk of decline in ADL functioning (OR: 2.65, 95% CI: 1.02–6.88), an increased use of supportive assistance (OR: 2.45, 95% CI: 1.02–5.87) and a decreased chance to return to the independent preoperative living situation (OR: 0.18, 95% CI: 0.07–0.49).

Conclusions

A hospital-based geriatric liaison intervention for the prevention of postoperative delirium in frail elderly cancer patients undergoing elective surgery for a solid tumour did not improve

outcomes 3 months after discharge from hospital. The negative effect of a postoperative delirium on late outcome was confirmed.

Trial Registration

Nederlands Trial Register, [Trial ID NTR 823](#).

Introduction

Hospitalized elderly are at increased risk for functional decline resulting in adverse health outcomes such as mortality, prolonged hospital stay, nursing home placement and increased dependency at home. It is estimated that approximately 35% of patients aged 75 and older develop a new disability after hospitalization or suffer functional decline [1–3].

To limit functional decline after hospital stay, the prevention of delirium is of great importance. Delirium is a common and serious complication in hospitalized elderly people. It is associated with persistent functional and cognitive decline, increased morbidity and mortality, longer hospital stays, higher rates of nursing home placement and increased health-care costs [4–7]. Mortality rates vary from 4% to 20% in patients who develop delirium during their hospital stay [8,9].

We performed a randomized controlled trial to evaluate the effect of a multicomponent intervention compared to standard care, on the incidence of postoperative delirium in frail elderly cancer patients undergoing surgery for a solid tumour [10]. The intervention was targeted at risk factors for postoperative delirium: cognitive impairment, visual impairment, hearing impairment, malnutrition, pain, sleep disturbance, defecation problems, infection and impaired mobility. Delirium was chosen as the primary outcome measure because it could be determined within the intervention period during hospital stay. The intervention has not shown to be effective for preventing postoperative delirium [10]. Three months after discharge, a follow-up was performed. The follow-up measurements were focused on postoperative functional outcomes such as Activities of Daily Living (ADL) functioning, return to the independent pre-operative living situation, use of supportive care, cognitive functioning and health related quality of life, next to mortality and rehospitalisation. Most previous studies on adverse outcomes after cancer surgery in the elderly were targeted at outcomes such as postoperative complications, mortality, length of hospital stay and readmissions [11–14], while ADL functioning and quality of life (QOL) are at least as important outcomes of surgical treatment for the elderly.

In this manuscript, the long term results, three months after discharge, and the effect of postoperative delirium on long term outcomes are described. The long term outcomes included: mortality, rehospitalisation, ADL functioning, return to the independent pre-operative living situation, use of supportive care, cognitive functioning and health related quality of life.

Methods

Ethics statement

The study was approved by the Medical Ethical Committee of the University Medical Center Groningen, trial ID NTR 823 ([S1](#) and [S2 Texts](#)). Written informed consent was obtained from the participants.

Study design

The study, entitled Liaison Intervention in Frail Elderly (LIFE), was a multicentre, randomized clinical trial [10]. The participating centres were the University Medical Center Groningen

(serving a population of three million people), the Medical Center Leeuwarden (a large teaching hospital) and Diaconessenhuis Leiden (a community hospital). All participating centres are located in the Netherlands.

The primary short term outcome of this study was the incidence of postoperative delirium up to 10 days postoperatively. The reported incidence of postoperative delirium varies widely from less than 10% to 50%. Based on these data and the fact that this study included a high-risk population, a delirium incidence of 30% was expected in the study population. An absolute reduction of 15% was expected in the intervention group based on Inouye's results [15]. To achieve a power of 80% with an α of 5% (one-sided), a β of 95% and an expected drop-out rate of 10%, it was calculated that a total of at least 294 patients would need to be included in this study.

Participants

From June 2007 to June 2010 all consecutive patients over 65 years of age undergoing elective surgery for a solid tumour were screened with the Groningen Frailty Indicator (GFI) [16–18] at the outpatient departments of general surgery, gynaecology, ear, nose and throat medicine and maxillofacial surgery at the participating centres. The GFI is an internally consistent 15-item screening instrument used to determine an individual's level of frailty [16,18]. The GFI is widely used in clinical practice, in outpatient settings, and in clinical studies [19–23]. It was shown that frail older persons (as identified with the GFI) had higher levels of case complexity, disability, and lower quality of life and life satisfaction [15]. Patients with a GFI score greater than 3 were regarded as frail and recruited to the LIFE study. The participants were randomly allocated to either the control group or the geriatric liaison intervention group. The randomization was stratified by tumour type. A distinction was made between tumours in the chest or abdomen and tumours elsewhere. The research nurses used an interactive voice response telephone service provided by the University Medical Center Groningen for the randomization.

Patients were excluded if the research nurse or the responsible physician estimated they were unable to complete the study protocol and follow-up schedule before inclusion (e.g. for logistical reasons or if any extra hospital visits would be too burdensome). Patients unable to fill in the questionnaires used in this study were also excluded.

Intervention

The multicomponent intervention focused on best supportive care and the prevention of delirium. Patients in the intervention group were assessed preoperatively by a geriatric team and monitored during their hospital stay. As the three participating centres are heterogeneous and this could cause variance in how the intervention was conducted, checklists were used to standardize the intervention as much as possible.

The geriatric team was supervised by a geriatrician, and helped devise the individual care plan. The preoperative comprehensive geriatric assessment by a geriatrician consisted of a medical history, physical examination and follow-up examinations on indication resulting in an individual treatment plan, with specific attention to patient-related risk factors for delirium.

During their hospital stay, the patients in the intervention group were assessed daily by a geriatric nurse. If a problem was encountered, the geriatric nurse or geriatrician contacted the treatment team to discuss the proposed intervention and establish a treatment plan, checking daily to determine whether the advice had been followed. For a detailed description of the intervention we refer to [10].

Standard care. Patients in the usual-care group received standard care, meaning that additional geriatric care was only provided at the request of the treating physician.

Surgical procedure. Surgical procedures were divided into three categories: minor, intermediate and major according to the duration of the operation and the localization of the tumour (intracavitary versus superficial (Table 1)).

Long term outcomes

The long term outcomes considered in the here presented analyses were mortality, rehospitalisation, ADL functioning, return to the independent pre-operative living situation, supportive care, cognitive functioning, and health related quality of life. The measurement instruments that were used for these outcomes are described in the assessments section.

Assessments. The baseline assessment was completed by the research nurses at least 24 hours before surgery and was performed prior to randomization. Data on long-term outcomes were collected by the research nurses 3 months following hospital discharge during a telephone interview or a home visit, between August 2007 and November 2010.

At baseline, demographic data were collected. Both the baseline assessment and the follow-up assessment included the measurement of the health related quality of life by the Physical Component Summary measure (PCS) and the Mental Component Summary measure (MCS) of the Short Form-36 (SF-36) score [24–26]; basic ADL functioning by the Care Dependency Scale (CDS) [27] and cognitive functioning by the Mini-Mental State Examination (MMSE) [28]. Data regarding the living situation and supportive care (domestic help, care assistance and informal care) were also collected.

To screen for delirium during hospital stay, the Delirium Observation Scale (DOS) was used in both groups [29]. The DOS was recorded three times a day, up to 10 days postoperatively. In the case of a mean DOS score ≥ 3 (possible delirium) a geriatrician or psychiatrist examined the patient to confirm the diagnosis according to the criteria of the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition* (DSM IV).

A paper-based standardized form was used to collect data. Data were entered into Oracle Clinical Remote Data Capture program by trained research nurses. After entry, the data were checked by an independent individual. The research nurses were not blinded to the group the patients had been assigned to.

Definition of long term outcomes. For the current analysis all long term outcomes were considered as binaries. ADL functioning, was categorized in a lower score at 3-month follow-up compared to the baseline the baseline score (“decreased”) versus a same or higher score (“same/ increased”). Use of supportive care was dichotomized in an increased number of hours supportive care per week at 3-month follow-up compared to baseline (“increased”) versus the same or a decreased number of hours supportive care (“same/ decreased”). Cognitive functioning was categorized as MMSE score decreased ≥ 2 points at 3-month follow-up versus baseline versus MMSE score same or increased. Health related quality of life was dichotomized as a decreased score on the SF-36 physical and mental component scale at 3 month follow up versus baseline (“decreased”) versus a same or increased score (“same/ increased”).

Table 1. Classification of the type of surgery by duration of the procedure and tumour localization.

Surgery load	Tumour localization
Minor	Breast and skin
Intermediate	Vulva, cervix, endometrium, uterus, head/neck and retroperitoneum
Major	Gastrointestinal, liver, pancreas, lung, ovary, oropharynx, larynx and intra-abdominal sarcoma

doi:10.1371/journal.pone.0143364.t001

Statistical analysis

Differences in baseline characteristics between the groups were examined using a Fisher exact test for nominal variables and a two-sample Smirnov test for ordinal or continuous variables.

To examine the effectiveness of the intervention as compared to standard care on the long term outcomes at 3 months follow up, univariate binary logistic regression analysis was used and Odds Ratios (ORs) with a 95% Confidence Intervals (CIs) were estimated, where the intervention was considered as independent and the long-term outcomes were considered as dependents. There was no pre-determined hierarchy between the long term outcomes. The primary outcome of this study, postoperative delirium, was evaluated previously [10]. In case of a p -value < 0.05 , correction for multiple testing was performed.

The effect of postoperative delirium (independent variable) on the outcomes at 3 months follow up (dependent variable) was also calculated using univariate binary logistic regression analysis.

IBM SPSS Statistics Version 20 was used for the statistical analysis.

Results

A description of the flow of participants through each stage of the LIFE study was presented in a previous article [10]. Of the 260 patients who were followed during hospital stay, 33 were lost to follow-up at the time of the 3-month assessment: 14 died during hospital stay, 12 died before follow-up assessment, seven withdrew informed consent post discharge. Therefore the final sample size for this study was 227: 106 intervention group and 121 usual-care group (Fig 1, lower part). There were no significant differences between the groups at discharge (Table 2).

Long term outcomes

The results of the logistic regression analyses for the outcome variables are shown in Table 3. There were no significant differences between the intervention and usual-care group for any of the outcomes.

Influence of postoperative delirium on long term outcomes

In total, 227 patients were analysed for the long term outcomes of delirium. A postoperative delirium occurred in 26 of these patients (11.5%). Delirium increased the risk of a decline in ADL functioning (OR: 2.65, 95% CI: 1.02–6.88) resulting in an increased need for care assistance (OR: 2.45, 95% CI: 1.02–5.87) and a decreased chance to return to the independent pre-operative living situation (OR: 0.18 (0.07–0.49)). These results are presented in Table 4.

Discussion

Three months after discharge from hospital no benefit could be detected from a geriatric liaison intervention targeted at risk factors for postoperative delirium in frail elderly patients undergoing surgery for a solid tumour. Because postoperative delirium is a known risk factor for functional decline after hospital stay [4–7], we, a priori, hypothesized that prevention of postoperative delirium would result in decreased risk for adverse outcomes after hospitalisation. Other studies have shown varying results of multicomponent delirium prevention interventions on long term outcomes [15, 30, 31].

The low delirium incidence rates found in the LIFE study (14.3% in the control group versus 9.4% in the intervention group) may have been of crucial importance for our negative results [10]. This resulted in an underpowered study. The intervention appeared not to be effective in

preventing delirium in the population under study and showed consequent no effect on long term results.

Factors that probably contributed to the low delirium incidence rate in our study are the exclusion of patients with severe cognitive impairment (high risk for postoperative delirium)

CONSORT 2010 Flow Diagram




Fig 1. CONSORT diagram for the study.

doi:10.1371/journal.pone.0143364.g001

Table 2. Characteristics of the patients at discharge according to study group.

Characteristic	Intervention group (n = 127)	Usual-care group (n = 133)	P-value
Age (years), mean (SD)	77.37 (6.88)	77.42 (7.71)	0.49†
Female, n (%)	76 (59.8)	85 (63.9)	0.53‡
Type of surgery^f, n (%)			0.54‡
Minor	32 (25.2)	33 (24.8)	
Intermediate	20 (15.7)	28 (21.1)	
Major	75 (59.1)	72 (54.1)	
Comorbidities*, n (%)			0.47‡
≤ 2	51 (40.2)	55 (41.4)	
>2	76 (59.8)	78 (58.6)	
Living situation, n (%)			0.10‡
independent	113 (89.0)	110 (82.7)	
alone	55 (43.3)	53 (39.8)	
with others	58 (45.7)	57 (42.9)	
dependent	14 (11.0)	23 (17.3)	
protected housing	0 (0.0)	4 (3.0)	
home for the elderly	11 (8.7)	16 (12.0)	
nursing home	3 (2.4)	3 (2.3)	
Supportive care, n (%)			0.46‡
Domestic help			
No	60 (47.2)	61 (46.2)	
Yes	67 (52.8)	71 (53.8)	
Missing	0	1	
Care assistance			0.40‡
No	87 (69.0)	94 (71.2)	
Yes	39 (31.0)	38 (28.8)	
Missing	1	1	
Informal care			0.49‡
No	68 (54.0)	70 (53.0)	
Yes	58 (46)	62 (47.0)	
Missing	1	1	
Care Dependency Score, mean (SD)	72.49 (8.52)	74.23 (6.97)	0.27†
Mini Mental State Examination, mean (SD)	26.97 (2.47)	26.51 (3.74)	0.97†
Missing	19	31	
Short Form-36, mean (SD)			
Physical component summary measure	48.36 (9.07)	49.32 (7.02)	0.17†
Mental component summary measure	44.69 (8.79)	44.38 (8.42)	0.98†

†Kolmogorov-Smirnov test.

‡ Fisher's exact test.

^fSurgery load: Major = gastrointestinal, liver, pancreas, lung, ovary, oropharynx, larynx and intra-abdominal sarcoma. Intermediate = vulva, cervix, endometrium, uterus, head/neck and retroperitoneum. Minor = breast and skin.

*Comorbidities = diabetes, COPD, hypertension, myocardial infarction, other cardiovascular disorders, neurological disorders, cerebrovascular disorders, hearing and vision problems, memory problems in daily life, psychiatric disorders or musculoskeletal disorders.

doi:10.1371/journal.pone.0143364.t002

and the inclusion of patients undergoing superficial surgery (low risk for postoperative delirium). Furthermore the low delirium incidence rate implies a high standard of care for frail elderly patients in the participating hospitals before the start of the study and the introduction

Table 3. Univariate Logistic regression analyses for the effectiveness of the intervention compared to standard care on the long term outcomes (intervention group versus control group).

	Mortality Yes	Mortality No	Missing cases	OR (95% CI)
Control group	9 (6.8)	124 (93.2)		1
<i>During hospital stay</i>	4			
<i>After discharge</i>	5			
Intervention group	17 (13.4)	110 (86.6)		2.13 (0.91–4.97)
<i>During hospital stay</i>	10			
<i>After discharge</i>	7			
	Hospital readmission Yes	Hospital readmission No		
Control group	22 (18.3)	98 (81.7)	1	1
Intervention group	24 (22.9)	81 (77.1)	1	1.32 (0.69–2.53)
	ADL functioning Decreased	ADL functioning Same/ increased		
Control group	68 (56.2)	53 (43.8)		1
Intervention group	64 (60.4)	42 (39.6)		1.19 (0.70–2.02)
	No return to independent preoperative living situation	Return to independent preoperative living situation		
Control group	9 (8.9)	92 (91.1)		1
Intervention group	15 (16.5)	76 (83.5)		2.02 (0.84–4.87)
	Use of supportive care Increased	Use of supportive care Same/ decreased		
Domestic help				
Control group	38 (32.2)	80 (67.8)	4	1
Intervention group	33 (32.4)	69 (67.6)	4	1.01 (0.57–1.78)
Care assistance				
Control group	39 (33.3)	78 (66.7)	4	1
Intervention group	42 (41.2)	60 (58.8)	4	1.40 (0.81–2.43)
Informal care				
Control group	37 (31.6)	80 (68.4)	4	1
Intervention group	39 (38.2)	63 (61.8)	4	1.34 (0.57–1.78)
	Cognitive functioning decreased	Cognitive functioning same/ increased		
Control group	9 (14.1)	55 (85.9)	57	1
Intervention group	15 (23.1)	50 (76.9)	41	1.83 (0.74–4.56)
	Health related quality of life decreased SF-36 Physical component summary measure	Health related quality of life same/ increased SF-36 Physical component summary measure		

(Continued)

Table 3. (Continued)

	Mortality Yes	Mortality No	Missing cases	OR (95% CI)
Control group	80 (66.7)	40 (33.3)	1	1
Intervention group	63 (60)	42 (40)	1	1.33 (0.77–2.30)
	Health related quality of life decreased SF-36 Mental component summary measure	Health related quality of life same/ increased SF-36 Mental component summary measure		
Control group	53 (44.2)	67 (55.8)	1	1
Intervention group	51 (48.6)	54 (51.4)	1	0.84 (0.50–1.42)

doi:10.1371/journal.pone.0143364.t003

Table 4. Univariate logistic regression analyses for the influence of postoperative delirium on 3-month outcomes.

	n (%)	n (%)	Missing cases	OR (95% CI)
	Mortality Yes	Mortality No		
No postoperative delirium	21 (9.5)	201 (90.5)		1
<i>During hospital stay</i>	10			
<i>After discharge</i>	11			
Postoperative delirium	5 (16.1)	26 (83.9)		1.90 (0.66–5.48)
<i>During hospital stay</i>	4			
<i>After discharge</i>	1			
	Hospital readmission Yes	Hospital readmission No		
No postoperative delirium	43 (21.5)	157 (78.5)	1	1
Postoperative delirium	3 (12)	22 (88)	1	0.50 (0.14–1.74)
	ADL functioning Decreased	ADL functioning Same/ increased		
No postoperative delirium	112 (55.7)	89 (44.3)		1
Postoperative delirium	20 (76.9)	6 (23.1)		2.65 (1.02–6.88)*
	No return to independent preoperative living situation	Return to independent preoperative living situation		
No postoperative delirium	17 (9.9)	155 (90.1)		1
Postoperative delirium	8 (38.1)	13 (61.9)		0.18 (0.07–0.49)*
	Use of supportive care Increased	Use of supportive care Same/ decreased		
Domestic help				
No postoperative delirium	65 (33.0)	132 (67)	4	1
Postoperative delirium	6 (26.1)	17 (73.9)	3	0.72 (0.27–1.90)
Care assistance				
No postoperative delirium	68 (34.7)	128 (65.3)	5	1

(Continued)

Table 4. (Continued)

	n (%)	n (%)	Missing cases	OR (95% CI)
	Mortality Yes	Mortality No		
Postoperative delirium	13 (56.5)	10 (43.5)	3	2.45 (1.02–5.87)*
Informal care				
No postoperative delirium	68 (34.7)	128 (65.3)	5	1
Postoperative delirium	8 (34.8)	15 (65.2)	3	1.00 (0.41–2.49)
	Cognitive functioning decreased	Cognitive functioning same/ increased		
No postoperative delirium	21 (18.1)	95 (81.9)	85	1
Postoperative delirium	3 (23.1)	10 (76.9)	3	1.36 (0.34–5.36)
	Health related quality of life decreased SF-36 Physical component summary measure	Health related quality of life same/ increased SF-36 Physical component summary measure		
No postoperative delirium	123 (61.5)	77 (38.5)	1	1
Postoperative delirium	20 (80)	5 (20)	1	2.26 (0.96–5.36)
	Health related quality of life decreased SF-36 Mental component summary measure	Health related quality of life same/ increased SF-36 Mental component summary measure		
No postoperative delirium	88 (44)	112 (56)	1	1
Postoperative delirium	16 (64)	9 (36)	1	(0.90–6.95)

*significant difference

doi:10.1371/journal.pone.0143364.t004

of the Delirium Observation Scale (DOS) [29] on the wards to screen for delirium may have ensured increased alertness among medical staff for the prevention of postoperative delirium, in both the intervention and control group.

The long term results of this type of studies may be influenced by a wash-out effect due to interventions performed after discharge and outside the study protocol. Probably, continuation of in hospital interventions after discharge might overcome this, although, little is known about the effect of prolonged interventions in elderly patients who were hospitalized. One study showed a significantly decreased mortality in older cancer patients after a 4 weeks lasting intervention post discharge [32].

Up to 50% of elderly patients suffer functional decline after hospitalization resulting in a decline in health-related quality of life and loss of independence in (I)ADL functioning [1,33]. In our study, also a considerable part of patients suffered a postoperative decline in ADL functioning (60.4% in the intervention group versus 56.2% in the control group) and health related QOL (physical component: 60% in the intervention group versus 66.7% in the control group; mental component: 48.6% in the intervention group versus 44.2% in the control group) (See Table 2).

For the frail elderly surgical oncology patients participating in the LIFE study, postoperative delirium was a risk factor for functional decline after discharge. Delirium was associated with: an increased risk of a postoperative decline in ADL functioning, an increased use of care assistance and a decreased chance to return to the independent preoperative living situation. Only

26 patients that had developed postoperative delirium versus 201 nondelirious patients were tested in this analysis. Our data confirm that a postoperative delirium is a sign of increased (brain) vulnerability associated with poorer prognosis [34]. Therefore, targeting preventive interventions at those elderly at risk for (postoperative) delirium remains a major concern in minimizing functional decline after hospitalization.

The results shown in this manuscript concern a post hoc analysis. The effectiveness of a geriatric liaison intervention as well as the effect of a postoperative delirium on the outcomes 3 months postoperative (dependent variables) were explored using binary logistic regression analysis. In view of the nature of the analysis (post hoc) and the number of dependent variables tested (11 in total), the result should be interpreted with some caution.

In conclusion, the lower than expected delirium incidence rate and the high standard of basic care may have influenced the long term results. The association between postoperative delirium and functional decline after hospitalization was confirmed in the population under study. Therefore prevention of postoperative delirium seems one of the ways to limit functional decline after surgery in this patient group.

Supporting Information

S1 Text. Study Protocol.

(DOC)

S2 Text. CONSORT checklist.

(DOC)

Author Contributions

Conceived and designed the experiments: BL TW TB JS. Performed the experiments: LH BL. Analyzed the data: LH TB. Wrote the paper: LH BL TW DA JS TB.

References

1. Covinsky KE, Palmer RM, Fortinsky RH, Counsell SR, Stewart AL, Kresevic D, et al. Loss of independence in activities of daily living in older adults hospitalized with medical illnesses: Increased vulnerability with age. *J Am Geriatr Soc.* 2003; 51: 451–458. PMID: [12657063](#)
2. Mahoney JE, Sager MA, Jalaluddin M. Use of an ambulation assistive device predicts functional decline associated with hospitalization. *J Gerontol A Biol Sci Med Sci.* 1999; 54: M83–8. PMID: [10051860](#)
3. Sager MA, Rudberg MA. Functional decline associated with hospitalization for acute illness. *Clin Geriatr Med.* 1998; 14: 669–679. PMID: [9799473](#)
4. McCusker J, Cole M, Dendukuri N, Belzile E, Primeau F. Delirium in older medical inpatients and subsequent cognitive and functional status: A prospective study. *CMAJ.* 2001; 165: 575–583. PMID: [11563209](#)
5. Inouye SK, Rushing JT, Foreman MD, Palmer RM, Pompei P. Does delirium contribute to poor hospital outcomes? A three-site epidemiologic study. *J Gen Intern Med.* 1998; 13: 234–242. PMID: [9565386](#)
6. O'Keeffe S, Lavan J. The prognostic significance of delirium in older hospital patients. *J Am Geriatr Soc.* 1997; 45: 174–178. PMID: [9033515](#)
7. Witlox J, Eurelings LS, de Jonghe JF, Kalisvaart KJ, Eikelenboom P, van Gool WA. Delirium in elderly patients and the risk of postdischarge mortality, institutionalization, and dementia: A meta-analysis. *JAMA.* 2010; 304: 443–451. doi: [10.1001/jama.2010.1013](#) PMID: [20664045](#)
8. Marcantonio ER, Goldman L, Mangione CM, Ludwig LE, Muraca B, Haslauer CM, et al. A clinical prediction rule for delirium after elective noncardiac surgery. *JAMA.* 1994; 271: 134–139. PMID: [8264068](#)
9. Robinson TN, Raeburn CD, Tran ZV, Angles EM, Brenner LA, Moss M. Postoperative delirium in the elderly: Risk factors and outcomes. *Ann Surg.* 2009; 249: 173–178. doi: [10.1097/SLA.0b013e31818e4776](#) PMID: [19106695](#)

10. Hempenius L, Slaets JP, van Asselt D, de Bock GH, Wiggers T, van Leeuwen BL. Outcomes of a geriatric liaison intervention to prevent the development of postoperative delirium in frail elderly cancer patients: Report on a multicentre, randomized, controlled trial. *PLoS One*. 2013; 8: e64834. PMID: [23840308](#)
11. Dekker JW, Gooiker GA, van der Geest LG, Kolfschoten NE, Struikmans H, Putter H, et al. Use of different comorbidity scores for risk-adjustment in the evaluation of quality of colorectal cancer surgery: Does it matter? *Eur J Surg Oncol*. 2012; 38: 1071–1078. doi: [10.1016/j.ejso.2012.04.017](#) PMID: [22703758](#)
12. Huisman MG, van Leeuwen BL, Ugolini G, Montroni I, Spiliotis J, Stabilini C, et al. "Timed up & go": A screening tool for predicting 30-day morbidity in onco-geriatric surgical patients? A multicenter cohort study. *PLoS One*. 2014; 9: e86863. doi: [10.1371/journal.pone.0086863](#) PMID: [24475186](#)
13. Kristjansson SR, Farinella E, Gaskell S, Audisio RA. Surgical risk and post-operative complications in older unfit cancer patients. *Cancer Treat Rev*. 2009; 35: 499–502. doi: [10.1016/j.ctrv.2009.04.004](#) PMID: [19435652](#)
14. Robinson TN, Wu DS, Sauaia A, Dunn CL, Stevens-Lapsley JE, Moss M, et al. Slower walking speed forecasts increased postoperative morbidity and 1-year mortality across surgical specialties. *Ann Surg*. 2013; 258: 582–590. doi: [10.1097/SLA.0b013e3182a4e96c](#) PMID: [23979272](#)
15. Inouye SK, Bogardus ST Jr., Charpentier PA, Leo-Summers L, Acampora D, Holford TR, et al. A multi-component intervention to prevent delirium in hospitalized older patients. *N Engl J Med*. 1999; 340: 669–676. PMID: [10053175](#)
16. Peters LL, Boter H, Buskens E, Slaets JP. Measurement properties of the groningen frailty indicator in home-dwelling and institutionalized elderly people. *J Am Med Dir Assoc*. 2012; 13: 546–551. doi: [10.1016/j.jamda.2012.04.007](#) PMID: [22579590](#)
17. Schuurmans H, Steverink N, Lindenberg S, Frieswijk N, Slaets JP. Old or frail: What tells us more? *J Gerontol A Biol Sci Med Sci*. 2004; 59: M962–M965. PMID: [15472162](#)
18. Steverink N, Slaets JP, Schuurmans H, van Lis M. Measuring frailty: Development and testing of the groningen frailty indicator (GFI). *Gerontologist*. 2009; 41: 236–237.
19. Aaldriks AA, van der Geest LG, Giltary EJ, le Cessie S, Portielje JE, Tanis BC, et al. Frailty and malnutrition predictive of mortality risk in older patients with advanced colorectal cancer receiving chemotherapy. *J Geriatr Oncol*. 2013; 4: 218–26. doi: [10.1016/j.jgo.2013.04.001](#) PMID: [24070460](#)
20. Aaldriks AA, Giltary EJ, Nortier JW, van der Geest LG, Tanis BC, Ypma P, et al. Prognostic significance of geriatric assessment in combination with laboratory parameters in elderly patients with aggressive non-Hodgkin lymphoma. *Leuk Lymphoma*. 2015; 19: 1–9.
21. Andela RM, Dijkstra A, Slaets JP, Sanderma R. Prevalence of frailty on clinical wards: description and implications. *Int J Nurs Pract*. 2010; 16 (1): 14–9.
22. Biesma B, Wijmenga AW, Vincent A, Dalesio O, Smit HJ, Stigt JA, et al. Quality of life, geriatric assessment and survival in elderly patients with non-small-cell lung cancer treated with carboplatin-gemcitabine or carboplatin-paclitaxel: NVALT-3 a phase III study. *Ann Oncol*. 2011; 22 (7):1520–7. doi: [10.1093/annonc/mdq637](#) PMID: [21252061](#)
23. Pol RA, van Leeuwen BL, Visser L, Izaks GJ, van den Dungen JJ, Tielliu IF, et al. Standardised frailty indicator as predictor for postoperative delirium after vascular surgery: a prospective cohort study. *Eur J of Vasc Endovasc*. 2011; 42 (6): 824–30.
24. McHorney CA, Ware JE Jr., Lu JF, Sherbourne CD. The MOS 36-item short-form health survey (SF-36): III. tests of data quality, scaling assumptions, and reliability across diverse patient groups. *Med Care*. 1994; 32: 40–66. PMID: [8277801](#)
25. Ware JE, Snow KK, Kosinski M, Gandek B. SF-36 health survey: Manual and interpretation guide. Boston: The Health Institute: New England Medical Center; 1993.
26. Aaronson NK, Muller M, Cohen PD, Essink-Bot ML, Fekkes M, Sanderma R, et al. Translation, validation, and norming of the dutch language version of the SF-36 health survey in community and chronic disease populations. *J Clin Epidemiol*. 1998; 51: 1055–1068. PMID: [9817123](#)
27. Dijkstra A, Buist G, Dassen T. Nursing-care dependency. development of an assessment scale for demented and mentally handicapped patients. *Scand J Caring Sci*. 1996; 10: 137–143. PMID: [9060783](#)
28. Tombaugh TN, McIntyre NJ. The mini-mental state examination: A comprehensive review. *J Am Geriatr Soc*. 1992; 40: 922–935. PMID: [1512391](#)
29. Schuurmans MJ, Shortridge-Baggett LM, Duursma SA. The delirium observation screening scale: A screening instrument for delirium. *Res Theory Nurs Pract*. 2003; 17: 31–50. PMID: [12751884](#)
30. Counsell SR, Holder CM, Liebenauer LL, Palmer RM, Fortinsky RH, Kresevic DM, et al. Effects of a multicomponent intervention on functional outcomes and process of care in hospitalized older patients:

- A randomized controlled trial of acute care for elders (ACE) in a community hospital. *J Am Geriatr Soc.* 2000; 48: 1572–1581. PMID: [11129745](#)
31. Slaets JP, Kauffmann RH, Duivenvoorden HJ, Pelemans W, Schudel WJ. A randomized trial of geriatric liaison intervention in elderly medical inpatients. *Psychosom Med.* 1997; 59: 585–591. PMID: [9407576](#)
 32. Bogardus ST Jr, Desai MM, Williams CS, Leo-Summers L, Acampora D, Inouye SK. The effects of a targeted multicomponent delirium intervention on postdischarge outcomes for hospitalized older adults. *Am J Med.* 2003; 114: 383–390. PMID: [12714128](#)
 33. Lawrence VA, Hazuda HP, Cornell JE, Pederson T, Bradshaw PT, Mulrow CD, et al. Functional independence after major abdominal surgery in the elderly. *J Am Coll Surg.* 2004; 199: 762–772. PMID: [15501119](#)
 34. Inouye SK, Westendorp RG, Saczynski JS. Delirium in elderly people. *Lancet.* 2014; 383 (9920); 911–22. doi: [10.1016/S0140-6736\(13\)60688-1](#) PMID: [23992774](#)