

University of Groningen

N-acetylglucosamine-Mediated Expression of nagA and nagB in *Streptococcus pneumoniae*

Afzal, Muhammad; Shafeeq, Sulman; Manzoor, Irfan; Henriques-Normark, Birgitta; Kuipers, Oscar P.

Published in:
Frontiers in Cellular and Infection Microbiology

DOI:
[10.3389/fcimb.2016.00158](https://doi.org/10.3389/fcimb.2016.00158)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Afzal, M., Shafeeq, S., Manzoor, I., Henriques-Normark, B., & Kuipers, O. P. (2016). N-acetylglucosamine-Mediated Expression of nagA and nagB in *Streptococcus pneumoniae*. *Frontiers in Cellular and Infection Microbiology*, 6, 1-12. [158]. <https://doi.org/10.3389/fcimb.2016.00158>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



N-acetylglucosamine-Mediated Expression of *nagA* and *nagB* in *Streptococcus pneumoniae*

Muhammad Afzal^{1,2}, Sulman Shafeeq³, Irfan Manzoor^{1,2}, Birgitta Henriques-Normark³ and Oscar P. Kuipers^{1*}

¹ Department of Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Groningen, Netherlands, ² Department of Bioinformatics and Biotechnology, Government College University, Faisalabad, Pakistan, ³ Department of Microbiology, Tumor and Cell Biology, Karolinska Institutet, Stockholm, Sweden

In this study, we have explored the transcriptomic response of *Streptococcus pneumoniae* D39 to N-acetylglucosamine (NAG). Transcriptome comparison of *S. pneumoniae* D39 wild-type grown in chemically defined medium (CDM) in the presence of 0.5% NAG to that grown in the presence of 0.5% glucose revealed elevated expression of many genes/operons, including *nagA*, *nagB*, *manLMN*, and *nanP*. We have further confirmed the NAG-dependent expression of *nagA*, *nagB*, *manLMN*, and *nanP* by β -galactosidase assays. *nagA*, *nagB* and *glmS* are putatively regulated by a transcriptional regulator NagR. We predicted the operator site of NagR (*dre* site) in *PnagA*, *PnagB*, and *PglmS*, which was further confirmed by mutating the predicted *dre* site in the respective promoters (*nagA*, *nagB*, and *glmS*). Growth comparison of Δ *nagA*, Δ *nagB*, and Δ *glmS* with the D39 wild-type demonstrates that *nagA* and *nagB* are essential for *S. pneumoniae* D39 to grow in the presence of NAG as a sole carbon source. Furthermore, deletion of *ccpA* shows that CcpA has no effect on the expression of *nagA*, *nagB*, and *glmS* in the presence of NAG in *S. pneumoniae*.

Keywords: N-acetylglucosamine, NAG, NagA, NagB, GlmS, pneumococcus, CcpA

OPEN ACCESS

Edited by:

Jorge Eugenio Vidal,
Emory University, USA

Reviewed by:

Samantha J. King,
Ohio State University, USA
Marco Rinaldo Oggioni,
University of Leicester, UK
Zehava Eichenbaum,
Georgia State University, USA

*Correspondence:

Oscar P. Kuipers
o.p.kuipers@rug.nl

Received: 17 January 2016

Accepted: 02 November 2016

Published: 16 November 2016

Citation:

Afzal M, Shafeeq S, Manzoor I,
Henriques-Normark B and Kuipers OP
(2016) N-acetylglucosamine-Mediated
Expression of *nagA* and *nagB* in
Streptococcus pneumoniae.
Front. Cell. Infect. Microbiol. 6:158.
doi: 10.3389/fcimb.2016.00158

INTRODUCTION

Pneumonia, sepsis, meningitis, otitis media, and conjunctivitis are a few of the diseases caused by the major human pathogen *Streptococcus pneumoniae* that results in over a million deaths each year worldwide (Ispahani et al., 2004; O'Brien et al., 2009). *S. pneumoniae* relies on several nutrient sources and virulence factors to colonize in the human nasopharynx (Phillips et al., 1990; Titgemeyer and Hillen, 2002; Carvalho et al., 2011). Regulatory mechanisms of a number of carbon and nitrogen sources important for the lifestyle and virulence of *S. pneumoniae* have already been studied (Kloosterman et al., 2006b; Carvalho et al., 2011; Kloosterman and Kuipers, 2011; Afzal et al., 2015c,d). *S. pneumoniae* has been shown to metabolize 32 carbohydrates including the three-carbon molecule glycerol, nine hexoses or hexose derivatives (ascorbate, fructose, galactose, glucosamine, glucose, mannose, N-acetylglucosamine, N-acetyl-mannosamine, and N-acetylneuraminic acid), three α -galactosides (melibiose, raffinose, and stachyose), two β -galactosides (lactose, and lactulose), four α -glucosides (maltose, maltotriose, sucrose, and trehalose), seven β -glucosides (amygdalin, arbutin, 1-O-methyl- β -glucoside, cellobiose, gentiobiose, aesculin, and salicin) and six polysaccharides (glycogen, hyaluronate, inulin, maltodextrin, pectin, and pullulan) (Bidossi et al., 2012).

The importance of carbon sources in the life-style of *S. pneumoniae* can be judged from the fact that over 30% of all the transporters in the genome are presumably involved in sugar uptake (Tettelin et al., 2001; Bidossi et al., 2012), a considerably larger number than that present in the other microorganisms inhabiting the same niche (Paulsen et al., 2000; Tettelin et al., 2001). The glycoproteins lining the epithelial surfaces in the human nasopharynx might be good carbon and energy sources for pneumococcal growth. Notably, *S. pneumoniae* has the ability to grow on mucin as a sole carbon source (Yesilkaya et al., 2008). Mucins are constituents of the mucus that span the epithelial surfaces (Rose and Voynow, 2006). These structures are largely O-glycosylated glycoproteins and are usually composed of N-acetylglucosamine (NAG), N-acetylgalactosamine, N-acetylneuraminic acid, galactose, fucose, and sulphated sugars connected to the protein core, mostly via an N-acetylgalactosamine moiety (Rose and Voynow, 2006; Terra et al., 2010). *S. pneumoniae* has several extracellular glycosidases with an extensive variety of substrates specificities and can make use of the other host glycans, such as N-glycans and glycosaminoglycans (Burnaugh et al., 2008; King, 2010; Marion et al., 2012). These enzymes produce a number of free sugars that potentially can be used by the pneumococcus. The deglycosylation activity of both exo- and endoglycosidases has previously been demonstrated in *S. pneumoniae* (King et al., 2006; Burnaugh et al., 2008; Jeong et al., 2009; Marion et al., 2009). The ability to utilize complex glycans present at the site of colonization contributes to the successful survival and virulence of *S. pneumoniae* in the host (Buckwalter and King, 2012; Linke et al., 2013). Besides, the role of these enzymes in *in vivo* fitness is demonstrated by the findings that glycosidase mutants show attenuated capacity to colonize and to cause disease in mouse models (Tong et al., 2000; Jeong et al., 2009; Marion et al., 2009; Terra et al., 2010).

NAG is an important amino-carbon source for several bacteria due to its role as an energy resource and in peptidoglycan synthesis (Dobrogosz, 1968; Mobley et al., 1982). Several studies highlighted the importance of NAG as a preferred carbon source in bacteria (Dobrogosz, 1968; Mobley et al., 1982). The involvement of NAG for both catabolic and anabolic purposes requires proper regulatory mechanisms for its utilization, as shown in model microorganisms, such as *Bacillus subtilis*, *Escherichia coli*, *Streptococcus mutans*, and *Streptomyces coelicolor* (Plumbridge, 2001; Nothaft et al., 2010; Bertram et al., 2011; Zeng and Burne, 2015). The NAG regulon consists of *nagA*, *nagB*, and *glmS* in *S. mutans*, and is regulated by a GntR-family transcriptional regulator NagR (Moye et al., 2014). NagA is an NAG-6-phosphate deacetylase, whereas NagB is a GlcN-6-P deaminase, and GlmS is a Fru-6-P amidotransferase. NagB was upregulated in the presence of NAG while GlmS expression decreased, signifying that the regulatory mode of these enzymes depends on the concentration of environmental NAG. A *glmS*-inactivated mutant could not grow in the absence of NAG, whereas the growth of *nagB*-inactivated mutant was decreased in the presence of NAG (Kawada-Matsuo et al., 2012). *nagB* inactivation led to a decrease in the expression of virulence factors, including cell-surface protein antigen and

glucosyltransferase, and also impeded biofilm formation and saliva-induced aggregation in *S. mutans* (Kawada-Matsuo et al., 2012). NagA has been shown to be important for the growth of pneumococcus in the presence of NAG as a sole carbon source (Paixão et al., 2015).

Diverse bacterial groups including streptomycetes, firmicutes, and enterobacteriaceae universally use phosphotransferase systems (PTSs) for uptake and phosphorylation of NAG (Simoni et al., 1976; Mobley et al., 1982; Alice et al., 2003; Nothaft et al., 2003, 2010). ManLMN PTS has been shown to be a glucose and mannose PTS in *Streptococcus salivarius* (Vadeboncoeur, 1984) and was also responsible for uptake of fructose, and NAG (Lortie et al., 2000). ManLMN transports glucose and mannose, and also shows specificity for galactose, NAG, and glucosamine in *S. pneumoniae* (Bidossi et al., 2012). NagP (PTS EIIBC component) is the main transporter of NAG in *B. subtilis* and *S. mutans* (Reizer et al., 1999; Saier et al., 2002; Moye et al., 2014).

Here, we demonstrate the effect of NAG on the global gene expression of *S. pneumoniae* and NAG-dependent expression of *nagA*, *nagB*, *manLMN*, and *nanP*. We further hypothesize that a GntR-family transcriptional regulator, NagR, might be involved in the regulation of *nagA*, *nagB*, and *glmS* and predict a putative operator site for NagR (*dre* site). We also explored the global impact of *ccpA* deletion on the transcriptome of *S. pneumoniae* in the presence of NAG, showing that *ccpA* has no effect on the expression of *nagA*, *nagB*, and *glmS*. Furthermore, we show that *nagA* and *nagB* are essential for *S. pneumoniae* to grow on NAG validating the previous study, where essentiality of *nagA* in the growth of *S. pneumoniae* was demonstrated (Paixão et al., 2015).

MATERIALS AND METHODS

Bacterial Strains, Growth Conditions, and DNA Isolation and Manipulation

Bacterial strains and plasmids used in this study are listed in **Table 1**. *S. pneumoniae* D39 was grown as described previously (Kloosterman et al., 2006a; Afzal et al., 2014). For β -galactosidase assays, derivatives of *S. pneumoniae* D39 were grown in chemically defined medium (CDM) (Neves et al., 2002) supplemented either with 0.5% glucose or with 0.5% NAG. For selection on antibiotics, medium was supplemented with the following concentrations of antibiotics: 150 μ g/ml spectinomycin, 15 μ g/ml trimethoprim and 2.5 μ g/ml tetracycline for *S. pneumoniae*; and 100 μ g/ml ampicillin for *E. coli*. All bacterial strains used in this study were stored in 10% (v/v) glycerol at -80°C . For PCR amplification, chromosomal DNA of *S. pneumoniae* D39 (Lanie et al., 2007) was used. Primers used in this study are based on the sequence of the *S. pneumoniae* D39 genome and listed in **Table 2**.

Construction of *nagA*, *nagB*, and *glmS* Mutants

nagA and *nagB* deletion mutants were made by allelic replacement with trimethoprim- and spectinomycin-resistance cassettes, respectively. Briefly, primers *nagA*-1/*nagA*-2 and

TABLE 1 | List of strains and plasmids used in this study.

Strain/plasmid	Description*	Source
S. PNEUMONIAE		
D39	Serotype 2 strain	Laboratory of P. Hermans
MA700	D39 $\Delta nagA$; Trm ^R	This study
MA701	D39 $\Delta nagB$; Spec ^R	This study
MA702	D39 $\Delta glmS$	This study
MA703	D39 $\Delta bgaA::PnagA-lacZ$; Tet ^R	This study
MA704	D39 $\Delta bgaA::PnagB-lacZ$; Tet ^R	This study
MA705	D39 $\Delta bgaA::PglmS-lacZ$; Tet ^R	This study
MA706	D39 $\Delta bgaA::PmanL-lacZ$; Tet ^R	This study
MA707	D39 $\Delta bgaA::PnagA-M-lacZ$; Tet ^R	This study
MA708	D39 $\Delta bgaA::PnagB-M-lacZ$; Tet ^R	This study
MA709	D39 $\Delta bgaA::PglmS1-M-lacZ$; Tet ^R	This study
MA710	D39 $\Delta bgaA::PglmS3-M-lacZ$; Tet ^R	This study
MA203	D39 $\Delta bgaA::PnanE-lacZ$; Tet ^R	Afzal et al., 2015b
E. COLI		
EC1000		Laboratory collection
PLASMIDS		
pPP2	Amp ^R Tet ^R ; promoter-less <i>lacZ</i> . For replacement of <i>bgaA</i> with promoter <i>lacZ</i> fusion. Derivative of pPP1	Halfmann et al., 2007
pORI280	Erm ^R ; <i>ori⁺ repA⁻</i> ; deletion derivative of pWV01; constitutive <i>lacZ</i> expression from P32 promoter	Leenhouts et al., 1998
pMA700	pORI280 carrying <i>glmS</i> deletion	This study
pMA701	pPP2 <i>PnagA-lacZ</i>	This study
pMA702	pPP2 <i>PnagB-lacZ</i>	This study
pMA703	pPP2 <i>PglmS-lacZ</i>	This study
pMA704	pPP2 <i>PmanL-lacZ</i>	This study
pMA705	pPP2 <i>PnagA-M-lacZ</i>	This study
pMA706	pPP2 <i>PnagB-M-lacZ</i>	This study
pMA707	pPP2 <i>PglmS1-M-lacZ</i>	This study
pMA708	pPP2 <i>PglmS3-M-lacZ</i>	This study

*Amp^R, Spec^R, Tet^R, and Trm^R confer ampicillin, spectinomycin, tetracycline and trimethoprim resistance gene, respectively.

nagA-3/*nagA*-4 were used to generate PCR fragments of the left and right flanking regions of *nagA*. Similarly, primers *nagB*-1/*nagB*-2 and *nagB*3/*nagB*-4 were used to generate PCR fragments of the left and right flanking regions of *nagB*. PCR products of left and right flanking regions of *nagA* and *nagB* contain *AscI* and *NotI* restriction enzyme sites, respectively. The trimethoprim- and spectinomycin-resistance cassettes that are amplified by primers Spec-F/Spec-R and Trmp-F/Trmp-R, respectively from pORI38 and pKOT, also contain *AscI* and *NotI* restriction enzyme sites on their ends. Then, by restriction and ligation, the left and right flanking regions of *nagA* and *nagB* were fused to the trimethoprim- and spectinomycin-resistance genes, respectively. The resulting ligation products were transformed to *S. pneumoniae* D39 wild-type and selection of the mutant strains was done on appropriate concentrations of trimethoprim and spectinomycin.

A markerless *glmS* mutant ($\Delta glmS$) was constructed in the *S. pneumoniae* D39 wild-type using the pORI280 plasmid, as described before (Kloosterman et al., 2006a). Primer pairs, *glmS*-1/*glmS*-2 and *glmS*-3/*glmS*-4, were used to generate PCR fragments of the left and right flanking regions of *glmS*, respectively. These PCR fragments were inserted into pORI280 using *XbaI* and *BglII* restriction sites, resulting in pMA700. All mutants were further confirmed by PCR and DNA sequencing.

Growth Analysis

For growth analysis of $\Delta nagA$, $\Delta nagB$, and $\Delta glmS$, *S. pneumoniae* D39 wild-type and its isogenic mutants ($\Delta nagA$, $\Delta nagB$, and $\Delta glmS$) were grown microaerobically at 37°C in 5 ml tubes containing 3 ml CDM supplemented either with 0.5% NAG or with 0.5% Glucose. Cultures were maintained at 37°C for 11 h and optical density at 600 nm was recorded with 1 h interval. CDM without inoculum was taken as blank. The growth of each strain was monitored from six biological replicates from at least two different days.

Construction of Promoter *lacZ*-fusions and β -Galactosidase Assays

Chromosomal transcriptional *lacZ*-fusions to the *nagA*, *nagB*, *glmS*, and *manL* promoters were constructed in the integration plasmid pPP2 (Halfmann et al., 2007) with primer pairs mentioned in Table 2, resulting in pMA701-04 respectively. Briefly, PCR products of *nagA*, *nagB*, *glmS*, and *manL* promoters were obtained using primer pairs mentioned in Table 2. These PCR fragments contain *EcoRI* and *BamHI* restriction sites at their ends. pPP2 also has *EcoRI* and *BamHI* restriction sites in its multiple cloning site (MCS). Then, by restriction and ligation, these PCR fragments were cloned into pPP2. pMA701-04 were further introduced into the *S. pneumoniae* D39 wild-type resulting in strains MA703-06, respectively. All plasmid constructs were checked by PCR and DNA sequencing. β -galactosidase assays were performed as described before (Israelsen et al., 1995; Kloosterman et al., 2006a) using cells that were grown in CDM with appropriate sugar mentioned in Results section and harvested in the mid-exponential phase of growth.

To study the functionality of *dre* site, the following promoter *lacZ*-fusions of *nagA*, *nagB*, and *glmS* with mutated *dre* sites were made in pPP2 (Halfmann et al., 2007) using the primer pairs mentioned in Table 2: *PnagA-M* (mutation in the *dre* site), *PnagB-M* (mutation in the *dre* site), *PglmS1-M* (mutation in the 1st *dre* site), and *PglmS3-M* (mutation in the 3rd *dre* site), resulting in plasmid pMA705-08, respectively. The mutations were incorporated into the primers used to amplify the target promoter regions containing the *dre* sites. These constructs were introduced into the *S. pneumoniae* D39 wild-type, resulting in strains MA707-10.

Microarray Analysis

For DNA microarray analysis in the presence of NAG, the transcriptome of *S. pneumoniae* D39 wild-type, grown in biological replicates in CDM with 0.5% NAG was compared to D39 wild-type grown in CDM with 0.5% glucose. Similarly, for DNA microarray analysis of $\Delta ccpA$, the transcriptome of

TABLE 2 | List of primers used in this study.

Name	Nucleotide sequence (5'→3')*	Restriction site
nagA-R	CATGGGATCCGTCCACAAGTTCCAAGTAACC	<i>Bam</i> HI
nagA-F	CATGGAATTCCGACAGACAGCTCAAGACAAGC	<i>Eco</i> RI
nagA-F-M	CATGGAATTCTATCTCCAAAAATAGGTGCTGTCATTTACAAAT	<i>Eco</i> RI
nagB-R	CATGGGATCCGCAACTTTCCACCTTGAACCTGG	<i>Bam</i> HI
nagB-F	CATGGAATTCCGGCAATCAATTCCTCTGGC	<i>Eco</i> RI
nagB-F-M	CATGGAATTCCGTTTTCACTTGACAAAAATTGGTCGCTGCATATAATAA	<i>Eco</i> RI
glmS-R	CATGGGATCCCAACACCAACAATCCACAC	<i>Bam</i> HI
glmS-F	CATGGAATTCCGTCGCTGAAGAAATCAGG	<i>Eco</i> RI
manL-F	CATGGAATTCCAGTAGAAGATGCTGTTG	<i>Eco</i> RI
manL-R	CATGGGATCCCTGACTGATGAATACCC	<i>Bam</i> HI
glmS1-F-M	CATGGAATTCCACAGGAGCTTAATTTGAACGCTGTCAATTTTACTC	<i>Eco</i> RI
glmS3-R-M	CATGGGATCCACATAGTATATACGACACAGCAAGCTGTGCTTTCTCCTT AAAATTGGGCGCGCTCAATTC	<i>Bam</i> HI
nagA-1	GACGGTGGTCATTGCGACTG	–
nagA-2	GCATAGGCGCGCCCTCGACGAACTCCGTGTG	<i>Asc</i> I
nagA-3	CGATTGCGGCGCGGCTAGCAACCTACCTAGATGG	<i>Not</i> I
nagA-4	CGTAGATATTCAGCCTGCATACC	–
nagB-1	GGGTGTCGTTTCATGACAAGGG	–
nagB-2	GCATAGGCGCGCGCTACTTCCCTGTGCGCAAGTCC	<i>Asc</i> I
nagB-3	CGATTGCGGCGCGCGCAGATGCTGAAGCGCTTAGC	<i>Not</i> I
nagB-4	CCATAGACAATGTCTAGTCTAAGC	–
glmS-1	TGCCTAGAGGTCATCTTCGTGAACCTTCACCG	<i>Xba</i> I
glmS-2	CCGCAGAATCATAGCCACGG	–
glmS-3	GCTATGATTCTGCGGCGACTGTACACCCTTACCTCTC	–
glmS-4	GAAGATCTCCAGGACAATCTCTGGGGC	<i>Bgl</i> II
Spec-R	GCTAAGCGGCGCGCCTAAACGAAATAAACGC	<i>Not</i> I
Spec-F	GCTATGGCGCGCGCCTAATCAAAAATAGTGAGGAGG	<i>Asc</i> I
Trmp-R	GCATGCGGCGCGGTTACGACGCGCATAGACGG	<i>Asc</i> I
Trmp-F	GCATGCGGCGCGGATTTTGTGAGCTTGA	<i>Not</i> I

*The underlined sequences represent the respective restriction sites.

S. pneumoniae D39 wild-type was compared to D39 Δ ccpA, grown in biological replicates in CDM with 0.5% NAG. The cells were harvested at their respective mid-exponential growth phases. All other procedures regarding the DNA microarray experiment and data analysis were performed as previously described (Afzal et al., 2015a; Shafeeq et al., 2015). For the identification of differentially expressed genes a Bayesian $p < 0.001$ and a fold change cut-off ≥ 2 was applied. Microarray data have been submitted to GEO (Gene Expression Omnibus) database under the accession number GSE89589 and GSE89590.

RESULTS

The Putative NAG Regulon in *S. pneumoniae*

The NAG regulon consists of *nagA*, *nagB*, and *glmS* in *S. mutans* and is regulated by a GntR-family transcriptional regulator NagR (Moye et al., 2014). NagA is an NAG-6-phosphate deacetylase, whereas NagB is a GlcN-6-P deaminase, and GlmS is a Fru-6-P amidotransferase. *S. pneumoniae* D39 also possess the genes that encode proteins putatively involved in the transport and

utilization of NAG. These genes are *nagA*, *nagB*, *manLMN*, *nanP*, and *glmS*. In *S. pneumoniae*, it appears that NAG enters the cell through NanP PTS (SPD-1496) and/or ManLMN (SPD-0262-64) and is subsequently phosphorylated (Kanehisa et al., 2014). NanP (PTS) is encoded by the gene that is part of *nan* operon-I (*spd_1488-97*) of the *nan* gene cluster and is proposed to transport amino sugars (Bidossi et al., 2012; Afzal et al., 2015b). *nanP* codes for EIIBC components of the PTS and therefore, needs EIIA component of another PTS to phosphorylate the incoming NAG. The phosphorylated NAG is deacetylated to glucosamine-6-P by NagA (Kanehisa et al., 2014). NagB converts glucosamine-6-P to fructose-6-P, whereas GlmS converts fructose-6-P to glucosamine-6-P. The role of NAG on the gene expression of *S. pneumoniae* has not been investigated before. Therefore, we decided to explore the effect of NAG on the whole transcriptome of *S. pneumoniae*.

NAG-Dependent Gene Expression in *S. pneumoniae*

To study the transcriptomic response of *S. pneumoniae* D39 to NAG, microarray comparison of *S. pneumoniae* D39 wild-type grown in CDM with 0.5% NAG to that grown in CDM with

0.5% glucose was performed. Presence of NAG in the medium resulted in altered expression of a number of genes/operons after applying the criteria of ≥ 2.0 -fold and $p < 0.001$ (Table S1). **Table 3** summarizes the transcriptome changes incurred in *S. pneumoniae* in the presence of NAG and lists the fold-change in the expression of the putative NAG transport and utilization genes.

The glutamine regulon was downregulated around 4-fold in the presence of NAG. This regulon consists of genes involved in glutamine synthesis and uptake (*glnA* and *glnPQ*), glutamate synthesis (*gdhA*), and the gene coding for the pentose phosphate pathway enzyme Zwf, which forms an operon with *glnPQ* (Kloosterman et al., 2006b). The glutamine regulon is shown to be repressed in the presence of a nitrogen source (Kloosterman et al., 2006b). The presence of nitrogen in NAG might be the reason of down-regulation of the glutamine regulon. A putative operon *spd_1969-72* was highly upregulated in the presence of NAG. This operon encodes proteins that are putatively involved in the utilization of carbohydrates. *spd_1970* codes for a ROK-family protein (RokD), but it lacks an HTH (helix-turn-helix) domain making it a less probable candidate as a transcriptional regulator of this operon (Shafeeq et al., 2012). ROK-family proteins are a class of transcriptional regulators involved in carbohydrate-dependent transcriptional control (repressor, ORF and kinase). They also contain sugar kinases and many functionally uncharacterized proteins (Titgemeyer et al., 1994). This operon was also upregulated in the presence of cellobiose (Shafeeq et al., 2013) and some other sugars making it a candidate for the utilization of multiple sugars. *strH* is another gene that was highly upregulated in the presence of NAG. StrH is a β -N-acetylhexosaminidase and is an important virulence factor of *S. pneumoniae*. StrH is a cell-surface attached β -N-acetylglucosaminidase that is used by *S. pneumoniae* to process the termini of host complex N-linked glycans (Pluvinage et al., 2013). StrH and SPD-1969 are also possibly being involved in the conversion of chitobiose into NAG based presumably on bioinformatics (Kanehisa et al., 2014). Similarly, *spd_1973-74* was also upregulated in our microarray analysis. Both these genes are annotated to be involved in carbohydrate metabolism, where SPD-1973 is a putative α -1,2-mannosidase and SPD-1974 is a hypothetical protein.

cel gene cluster (*spd-0277-0283*) putatively involved in the utilization of cellobiose is upregulated in the presence of NAG. This gene cluster is shown to be activated by transcriptional regulator CelR in the presence of cellobiose (Shafeeq et al., 2011). Tagatose pathways (*lac* gene cluster) and Leloir pathway genes (*galK* and *galT*) involved in the utilization of lactose and galactose (Afzal et al., 2014, 2015e) are significantly upregulated in the presence of NAG. *lac* gene cluster comprises of two operons in *S. pneumoniae*, i.e., *lac* operon-I (*lacABCD*) and *lac* operon-II (*lacFEG*) (Afzal et al., 2014). LacR, a DeoR-type regulator acts as a transcriptional repressor of *lac* operon-I in the absence of lactose/galactose (Afzal et al., 2014). Whereas, BglG-family transcriptional antiterminator LacT acts as a transcriptional activator of the *lac* operon-II in the presence of lactose (Afzal et al., 2014). Putative Raffinose uptake genes

TABLE 3 | Summary of data from Table S1 showing transcriptome comparison of *S. pneumoniae* D39 wild-type grown in CDM with 0.5% NAG to that grown in CDM with 0.5% glucose.

D39 tag ^a	Function ^b	Ratio ^c
<i>spd_1971</i>	Glycosyl hydrolase-related protein	22.8
<i>spd_0063</i>	β -N-acetylhexosaminidase, StrH	14.2
<i>spd_1970</i>	ROK family protein	13.9
<i>spd_1050</i>	Tagatose 1,6-diphosphate aldolase, LacD	13.7
<i>spd_1972</i>	hypothetical protein	13.6
<i>spd_0277</i>	6-phospho- β -glucosidase	13.1
<i>spd_1969</i>	Glycosyl hydrolase-related protein	11.9
<i>spd_1051</i>	Tagatose-6-phosphate kinase, LacC	11.0
<i>spd_1052</i>	Galactose-6-phosphate isomerase, LacB	10.7
<i>spd_1053</i>	Galactose-6-phosphate isomerase, LacA	9.9
<i>spd_0280</i>	Transcriptional regulator, CelR	7.7
<i>spd_1677</i>	Sugar ABC transporter, RafE	7.4
<i>spd_1676</i>	Sugar ABC transporter, RafF	6.6
<i>spd_1634</i>	Galactokinase, GalK	6.5
<i>spd_1633</i>	Galactose-1-phosphate uridylyltransferase, GalT	6.3
<i>spd_1675</i>	Sugar ABC transporter, RafG	5.6
<i>spd_1047</i>	PTS system, lactose-specific IIBC components, LacE	5.4
<i>spd_0283</i>	PTS system, IIC component	5.2
<i>spd_1974</i>	Hypothetical protein	4.8
<i>spd_1046</i>	6-phospho- β -galactosidase, LacG	4.4
<i>spd_1049</i>	Transcription antiterminator, LacT	4.4
<i>spd_0282</i>	Hypothetical protein	4.4
<i>spd_0281</i>	PTS system, IIA component	4.2
<i>spd_1664</i>	PTS system, trehalose-specific IABC components	4.2
<i>spd_1663</i>	α -phosphotrehalase, TreC	4.0
<i>spd_0279</i>	PTS system, IIB component	3.9
<i>spd_1495</i>	Sugar ABC transporter, sugar-binding protein	3.7
<i>spd_1973</i>	α -1,2-mannosidase, putative	3.6
<i>spd_1496</i>	PTS system, IIBC components	3.4
<i>spd_0263</i>	PTS system, mannose-specific IIC component, ManM	3.2
<i>spd_0262</i>	PTS system, mannose/fructose/sorbose family protein, IID component	3.0
<i>spd_1494</i>	Sugar ABC transporter, permease protein	2.7
<i>spd_1493</i>	Sugar ABC transporter, permease protein	2.4
<i>spd_1866</i>	N-acetylglucosamine-6-phosphate deacetylase, NagA	2.4
<i>spd_1846</i>	PTS system, IIB component	2.3
<i>spd_1246</i>	glucosamine-6-phosphate isomerase, NagB	2.3
<i>spd_0264</i>	PTS system, mannose-specific IAB components, ManL	2.1
<i>spd_1492</i>	Hypothetical protein	2.0
<i>spd_1491</i>	Hypothetical protein	2.0
<i>spd_1100</i>	Glucose-6-phosphate 1-dehydrogenase, Zwf	-2.2
<i>spd_0448</i>	Glutamine synthetase, GlnA	-3.3
<i>spd_1099</i>	Amino acid ABC transporter, ATP-binding protein	-3.3
<i>spd_0447</i>	Transcriptional regulator, GlnR	-3.7
<i>spd_1098</i>	Amino acid ABC transporter, amino acid-binding protein	-4.4

^aGene numbers refer to D39 locus tags.

^bD39 annotation (Lanie et al., 2007).

^cRatio represents the fold increase/decrease in the expression of genes in the presence of 0.5% NAG compared to 0.5% glucose.

rafEFG (*spd-1675-77*) are highly upregulated in the presence of NAG. Glucose and sucrose are shown to inhibit raffinose uptake (Tyx et al., 2011). A putative trehalose system (*spd-1663-64*) is highly expressed under our tested conditions. Upregulation of these different sugar systems under our tested conditions might be due to absence of CCR in the presence of NAG as a sole carbon source and further experiments are required to explore the role of these genes in the utilization of NAG.

Expression of genes that are putatively part of NAG utilization and transport pathway was also altered in our transcriptome analysis (Table 3). Expression of *manLMN* is upregulated around 3-fold in the presence of NAG. We observed upregulation of the *nan* operon-I, which is involved in the transport and utilization of sialic acid (an amino carbon source) (Marion et al., 2011). Moreover, expression of *nagA* and *nagB* was upregulated more than two folds in the presence of NAG (Table 3). No change in the expression of *glmS* is observed in our transcriptome in the presence of NAG. Upregulation of *nanP*, *manLMN*, *nagA*, and *nagB* in our transcriptome supports that these genes are important in NAG utilization in *S. pneumoniae* and strengthens the notion that NanP and ManLMN might be very important for NAG transport. Therefore, we decided to further explore the regulation of these genes in the presence of NAG.

NAG Induces the Expression of the Genes Involved in the Putative Transport and Utilization of Amino Sugars

In order to investigate in more detail the transcriptional regulation of *nanP*, *manLMN*, *nagA*, and *nagB* in the presence of NAG and to confirm our microarray results, we made ectopic transcriptional *lacZ*-fusions of *nanE*, *manL*, *nagA*, and *nagB* promoters (*PnagA-lacZ*, *PnagB-lacZ*, *PnanE-lacZ*, and *PmanL-lacZ*) and performed β -galactosidase assays (Figure 1). Our β -galactosidase assays data revealed that the expression of *PnagA-lacZ*, *PnagB-lacZ*, *PnanE-lacZ*, and *PmanL-lacZ* was strikingly higher in the presence of NAG compared to glucose in CDM (Figure 1). These data further confirm our microarray data mentioned above. We did not observe any change in the expression of *glmS* in our microarray analysis in the presence of NAG. To further study the expression of *glmS* in the presence of NAG and confirm our microarray analysis, we constructed ectopic transcriptional *lacZ*-fusion of *glmS* promoter (*PglmS-lacZ*) and performed β -galactosidase assays. We could not see any significant change in the expression of *PglmS-lacZ* under our tested conditions.

Predication and Confirmation of the *dre* Sites in the Promoter Regions of *glmS*, *nagA*, and *nagB*

Recently, NagR was characterized as a transcriptional repressor and shown to bind with specific DNA sequences named as *dre* sites, present in the promoter regions of the *nagAB* and *glmS* genes in *S. mutans* (Zeng and Burne, 2015). Blast search in *S. pneumoniae* D39 for NagR revealed the presence of a putative GntR-family transcriptional regulator NagR (SPD-1275)




FIGURE 1 | Expression levels (in Miller units) of *PnagA-lacZ*, *PnagB-lacZ*, *PglmS-lacZ*, *PnanE-lacZ*, and *PmanL-lacZ* in *S. pneumoniae* D39 wild-type grown in CDM either with 0.5% glucose or with 0.5% NAG. Standard deviations of three independent experiments are indicated in bars. Statistical significance of the differences in the expression levels was determined by one-way ANOVA (NS, not significant and * $P < 0.0001$).**

in *S. pneumoniae* D39. Presence of a NagR ortholog in *S. pneumoniae* might suggest its role in the regulation of the *glmS*, *nagA*, and *nagB*. We decided to delete *nagR* in *S. pneumoniae* D39 and study its role. However, we could not delete *nagR* in *S. pneumoniae* D39, suggesting the essentiality of NagR in the lifestyle of *S. pneumoniae*.

To explore the NagR regulated genes in *S. pneumoniae* D39, we decided to explore the genome of *S. pneumoniae* D39 for *dre* sites by using the Genome2D tool (Baerends et al., 2004) and a MEME motif sampler search (Bailey and Elkan, 1994). A 20-bp consensus sequence was found upstream of *nagA* (5'-AAATAGGTCTATACCATTTA-3') and *nagB* (5'-AAATTGGTCTATACCATATA-3') in *S. pneumoniae* D39 (Figure S1). We also found three *dre* sites in the promoter region of *glmS* (5'-AATTTGAACTATAACCAATTT-3', 5'-AAACAAGTATATACTGTTT-3' and 5'-GAATTAGACTATACCAATTT-3'). These DNA stretches may serve as *dre* sites in *S. pneumoniae*. We further checked the conservation of this *dre* site in other streptococcal species (*Streptococcus mitis*, *Streptococcus agalactiae*, *Streptococcus dysgalactiae*, *Streptococcus equi*, *S. mutans*, *Streptococcus pyogenes*, *Streptococcus sanguinis*, *Streptococcus gallolyticus*, *Streptococcus suis*, and *Streptococcus uberis*) and constructed weight matrix of the putative *dre* sites presents in different streptococci (Figure S2). We found that the *dre* sequence is highly conserved in the promoter regions of *nagA*, *nagB* and *glmS* in these streptococci as well (Figure S1).


To determine if the located stretch of DNA mediates the NagR-dependent transcriptional control of the *glmS*, *nagA*, and *nagB*, we made a number of transcriptional *lacZ*-fusions, where conserved bases in the putative *dre* sites were mutated in *PnagA* (5'-AAATAGGTCTATACCATTTA-3' to 5'-AAATAGGTCGCTGTCATTTA-3'), *PnagB* (5'-AAATTGGTCTATACCATATA-3' to 5'-AAATTGGTCGCTGTCATATA-3') and *PglmS* (first site: 5'-AATTTGAACTATAACCAATTT-3' to 5'-AATTTGAACTGTCATTT-3' and third site: 5'-GAATTAGACTATACCAATTT-3' to 5'-GAATTAGAC

GCGCCAATTT-3'). We could not mutate the second *dre* site in *PglmS* as it overlaps with core promoter sequence. The expression of *PnagA-M-lacZ* and *PnagB-M-lacZ* (few conserved bases of the *dre* sites were mutated) compared to that of the promoters with the intact *dre* sites (*PnagA-lacZ* and *PnagB-lacZ*) was considerably higher in the presence of glucose and NAG (Figures 2A,B). A derepression of the expression of *PglmS* was observed when either of the putative *dre* sites in *PglmS* (*dre* site 1 and 3) was mutated. These results suggest that *dre* sites present in *PglmS*, *PnagA* and *PnagB* are active and may act as an operator site for NagR in *S. pneumoniae*.

nagA and *nagB* Are Essential for *S. pneumoniae* D39 to Grow in the Presence of NAG as a Sole Carbon Source

nagA, *nagB*, and *glmS* encode important enzymes for the metabolism of NAG in bacteria. To elucidate the significance

of these genes on the growth of *S. pneumoniae*, we made knockout mutants of these genes ($\Delta nagA$, $\Delta nagB$, and $\Delta glmS$), and explored the impact of mutation of these genes on the growth of *S. pneumoniae* D39 in the presence of 0.5% NAG or glucose in CDM. The genetic organization and PCR confirmation of *nagA*, *nagB*, and *glmS* mutants are given in the Figure 3 and Figure S3, respectively. All three mutants ($\Delta nagA$, $\Delta nagB$, and $\Delta glmS$) had approximately the same growth as D39 wild-type in the presence of glucose in the medium (Figure 4). $\Delta glmS$ also showed the same growth pattern as the D39 wild-type in the presence of NAG. However, in contrast to D39 wild-type, $\Delta nagA$ and $\Delta nagB$ were not able to grow in the presence of NAG (Figure 4). These results suggest that *nagA* and *nagB* are necessary for *S. pneumoniae* to grow in the presence of NAG. These results are also in accordance with recently published data, where they showed that a mutant of *nagA* did not grow in the presence of NAG as a sole carbon source (a phenotype that could be complemented) (Paixão et al., 2015).





Role of CcpA in Regulation of *nagA*, *nagB*, *nanP*, *glmS*, and *manL*

CcpA is the master transcriptional regulator in *S. pneumoniae* that represses the expression of genes involved in the utilization of non-preferred sugars in the presence of a preferred one. To explore the effect of *ccpA* deletion on the transcriptome of *S. pneumoniae* and in NAG-dependent regulation of NAG utilization and transport genes, we performed transcriptome comparison of D39 $\Delta ccpA$ to D39 wild-type in CDM with 0.5% NAG. Expression of a number of genes was altered significantly (Table S1). These genes have been categorized according to their protein function in COG categories (Table 4). We did not observe any significant change in the expression of *nagA*, *nagB*, or *glmS*, suggesting CcpA independent expression of these genes. However, expression of *manLMN* and *nan* operon-I was upregulated in $\Delta ccpA$, which might suggest a putative role of CcpA in regulation of the *manLMN* and *nan* operon-I. *nan* operon-I was already shown to be regulated by CcpA and to have a *cre* box (Afzal et al., 2015b). Therefore, upregulation of *nan* operon-I in the absence of *ccpA* strengthens the previous observation (Afzal et al., 2015b).

To further confirm the role of CcpA in the regulation of *nagA*, *nagB*, *glmS*, and *manL*, we analyzed the promoter regions of these genes for the presence of *cre* boxes. We could not find a *cre* box in the promoter regions of *nagA*, *nagB*, and *glmS*, which might confirm the CcpA-independent regulation of the *nagA*, *nagB*, and *glmS* by transcriptional regulator NagR. However, we found a putative *cre* box (5'-ATGAAAACGGTTTATA-3') in the promoter regions of *manL*, further confirming the role of CcpA in the regulation of *manLMN*.

DISCUSSION AND CONCLUSIONS

The existence of well-developed sugar transport mechanisms in the opportunistic respiratory human pathogen, *S. pneumoniae*, emphasizes the importance of carbohydrates in the lifestyle of pneumococcus and confers an extra advantage to survive in a



changing nutritional environment (Tettelin et al., 2001). Glucose is the most preferred carbon source for *S. pneumoniae* but the presence of several other sugar-specific systems in *S. pneumoniae* indicates its ability to use other available sugars (Hoskins et al., 2001; Lanie et al., 2007; Bidossi et al., 2012). Extensive studies have been performed regarding regulatory mechanisms of different dedicated systems for sugars, including maltose, raffinose, cellobiose, sialic acid, and others in *S. pneumoniae* (Tyx et al., 2011; Shafeeq et al., 2013; Afzal et al., 2015b,f). Lack of free carbohydrates in the human airway makes modification and import of complex glycans much more critical for pneumococci to obtain the necessary carbon (Buckwalter and King, 2012). At least nine surface-associated glycosidases have been shown to modify host glycans in pneumococci, which makes bacterial survival better in the host (King et al., 2006; Burnaugh et al., 2008; Dalia et al., 2010). Data suggests that NAG may be an important carbohydrate for pneumococci (Bidossi et al., 2012). The regulatory mechanisms of genes putatively involved in NAG utilization have not been explored in *S. pneumoniae*. The current study sheds light on the regulatory mechanism of the *nagA*, *nagB*, and *glmS* in *S. pneumoniae*.

nagA, *nagB*, and *glmS* are annotated as a part of the amino sugar metabolism pathways in *S. pneumoniae* (Kanehisa et al., 2014). In our transcriptome comparison of *S. pneumoniae* D39 grown in CDM with 0.5% NAG to that grown in CDM with 0.5% glucose revealed increased expression of *nagA*, *nagB*, *manLMN*, and *nanP*. In *S. mutans*, expression of *glmS* is repressed in the presence of NAG compared to glucose (Zeng and Burne, 2015). This repression of *glmS* in the presence of NAG was relieved

TABLE 4 | Number of genes significantly affected in *S. pneumoniae* D39 $\Delta ccpA$ compared to the D39 wild-type grown in CDM with 0.5% NAG.

Functional categories	Total	Up	Down
C: Energy production and conversion	10	4	6
D: Cell cycle control, cell division, chromosome partitioning	2	0	2
E: Amino acid transport and metabolism	4	3	1
F: Nucleotide transport and metabolism	1	0	1
G: Carbohydrate transport and metabolism	19	17	2
H: Coenzyme transport and metabolism	1	0	1
I: Lipid transport and metabolism	3	0	3
J: Translation, ribosomal structure and biogenesis	15	3	12
K: Transcription	3	2	1
L: Replication, recombination and repair	3	2	1
M: Cell wall/membrane/envelope biogenesis	6	2	4
O: Posttranslational modification, protein turnover, chaperones	3	3	0
P: Inorganic ion transport and metabolism	3	1	2
Q: Secondary metabolites biosynthesis, transport and catabolism	2	1	1
R: General function prediction only	5	3	2
S: Function unknown	42	22	20
T: Signal transduction mechanisms	3	3	0
U: Intracellular trafficking, secretion, and vesicular transport	2	2	0
V: Defense mechanisms	8	7	1
Others	34	19	15
Total number of genes	169	94	75

Genes affected with more than 2-fold in D39 $\Delta ccpA$ compared to the D39 wild-type are shown in COG functional categories.

in *nagR* mutant (Zeng and Burne, 2015). However, no change in the expression of *glmS* is observed in our NAG-dependent transcriptome and no effect of *ccpA* deletion on the expression of *glmS* is observed. Mutating *dre* site 1 or 3 in the *PglmS* led to increase in expression of *PglmS* in the presence of glucose and NAG. This might indicate that NagR represses the expression of *glmS* in the presence of glucose and NAG.

The transport of amino-sugars has been attributed to a PTS (NanP) and *manLMN* in *S. mutans* (Moye et al., 2014). Similarly, a NAG-specific PTS (NagE) and a mannose-specific PTS ManXYZ have been shown to be involved in the NAG transport in *E. coli* (White, 1970; Alvarez-Añorve et al., 2009). *manLMN* has also been proposed to be involved in NAG transport in *S. pneumoniae* (Bidossi et al., 2012). Similarly, a PTS present in *nan* operon-I (putatively called *nanP*) has been suggested to play a part in the transport of glucosamine in *S. pneumoniae* (Kanehisa et al., 2014). *manLMN* and *nanP* are upregulated in our NAG-dependent transcriptome analysis, which is further confirmed by β -galactosidase assays. These observations confirm the findings of the previous studies and strengthen the involvement of *nanP* and *manLMN* in the transport of NAG.

NagA, NagB, and GlmS are very important for the metabolism of NAG and these three factors are associated with the synthesis of GlcN-6-P, a precursor for cell wall peptidoglycan synthesis in *E. coli* (Plumbridge et al., 1993; Plumbridge and Vimr, 1999). Here, we have studied the impact of *nagA*, *nagB*, and *glmS* deletions on the growth of *S. pneumoniae* in the presence of

NAG. Our studies suggest that *nagA* and *nagB* are important for pneumococcus to grow on NAG as their deletion mutants failed to grow in the presence of NAG in the medium as a sole carbon source. NagA has also been shown to be essential for growth in the presence of NAG as a sole carbon source (Paixão et al., 2015). However, no impact of *glmS* deletion on the growth of *S. pneumoniae* was observed. In *B. subtilis*, *nagB* has been shown to be essential for growth in the presence of NAG (Gaugué et al., 2013). NagB and GlmS have been shown to be involved in virulence in *S. mutans* (Kawada-Matsuo et al., 2012). Inactivation of *nagB* led to a decrease in the expression of virulence factors, including cell-surface protein antigen and glucosyltransferase, and also impeded biofilm formation and saliva-induced aggregation in *S. mutans* (Kawada-Matsuo et al., 2012). Pneumococcal *nagA* mutant was tested in mouse model of colonization and of model of bronchopneumonia with bacteremia, and no difference in virulence was observed (Paixão et al., 2015). It might be still interesting to further explore the role of the *nagB* and *glmS* in virulence of *S. pneumoniae*.

In *E. coli*, a ROK-family protein (NagC) acts as a transcriptional repressor of the NAG regulon (*nagE* and *nagBACD*), which encodes genes that are involved in the uptake and metabolism of NAG. Furthermore, it has been shown that NAG binds to NagC to relieve the repression caused by NagC (Plumbridge, 1991; Titgemeyer et al., 1994). Similarly, a GntR family transcriptional regulator NagR has been shown to act as a transcriptional regulator of the genes involved in NAG utilization in *B. subtilis*, *S. mutans* and in some other bacteria

(Bertram et al., 2011; Moye et al., 2014). In *S. mutans*, NagR has been shown to regulate the expression of *glmS* and *nagAB* by binding to the NagR operator sites called *dre* sites (Zeng and Burne, 2015). Our study suggests that NagR is present in *S. pneumoniae* and might regulate the expression of the *nagA*, *nagB*, and *glmS* by binding to the *dre* sites present in the promoter regions of these genes. We could not delete *nagR*, which might indicate about its essentiality or its involvement in some important cell process directly or indirectly. However, we mutated the conserved bases in the *dre* sites present in the promoter regions of *nagA*, *nagB*, and *glmS* which might suggest the importance of these bases in the regulation of these genes. To explore more putative *dre* sites in the D39 genome, we conducted a genome-wide search for putative pneumococcal *dre* sites. A *dre* site was only found in the promoter regions of *nagA* and *nagB*, and three *dre* sites were found in the promoter region of *glmS*. This predicted *dre* site was also found to be highly conserved in other streptococcal species as well (Novichkov et al., 2010), suggesting a similar function of NagR in these organisms.

The master transcriptional regulator, CcpA (Carbon catabolite protein A), was shown to be involved in the repression of non-preferred sugar metabolism genes in the presence of a preferred carbon source, and has a role in pneumococcal pathogenesis (Lulko et al., 2007; Zomer et al., 2007; Carvalho et al., 2011). A number of non-preferred sugar systems have also been shown to be regulated independently of CcpA by other transcriptional regulators, like the *cel* gene cluster activated by CelR in *S. pneumoniae* (Shafeeq et al., 2011). In this study, we elucidated the role of CcpA in the regulation of *nagA*, *nagB*, *glmS*, *manLMN*, and the *nan* operon-I by elaborating the impact of a *ccpA* deletion on the whole transcriptome of *S. pneumoniae* in the presence of NAG as a sole carbon source in CDM. Our transcriptome data demonstrated the CcpA-independent

expression of *nagA*, *nagB*, and *glmS*, and CcpA-dependent expression of *manLMN* and the *nan* operon-I. We further analyzed the promoter regions of these genes for the presence of a *cre* box and found *cre* boxes only in the promoter regions of *manLMN* and the *nan* operon-I. The absence of *cre* boxes in the promoter regions of *nagA*, *nagB*, and *glmS* confirms that CcpA may not have a role in the regulation of *nagA*, *nagB*, and *glmS*. However, the presence of a *cre* box in the promoter regions of *manLMN* and the *nan* operon-I further supports the role of CcpA in the regulation of *manLMN* and the *nan* operon-I.

AUTHOR CONTRIBUTIONS

Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work: MA, SS, IM, BHN, and OPK. Drafting the work or revising it critically for important intellectual content: MA, SS, IM, BHN, and OPK. Final approval of the version to be published: MA, SS, IM, BHN, and OPK. Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved: MA, SS, IM, BHN, and OPK.

ACKNOWLEDGMENTS

MA and IM are supported by Government College University, Faisalabad, Pakistan under faculty development program of HEC Pakistan.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fcimb.2016.00158/full#supplementary-material>

REFERENCES

- Afzal, M., Manzoor, I., and Kuipers, O. P. (2015a). A fast and reliable pipeline for bacterial transcriptome analysis case study: serine-dependent gene regulation in *Streptococcus pneumoniae*. *J. Vis. Exp. JoVE*. e52649. doi: 10.3791/52649
- Afzal, M., Shafeeq, S., Ahmed, H., and Kuipers, O. P. (2015b). Sialic acid-mediated gene expression in *Streptococcus pneumoniae* and role of NanR as a transcriptional activator of the *nan* gene cluster. *Appl. Environ. Microbiol.* 81, 3121–3131. doi: 10.1128/AEM.00499-15
- Afzal, M., Shafeeq, S., Henriques-Normark, B., and Kuipers, O. P. (2015c). UlaR activates expression of the *ula* operon in *Streptococcus pneumoniae* in the presence of ascorbic acid. *Microbiol. Read. Engl.* 161, 41–49. doi: 10.1099/mic.0.083899-0
- Afzal, M., Shafeeq, S., and Kuipers, O. P. (2014). LacR is a repressor of *lacABCD* and LacT is an activator of *lacTFEG*, constituting the *lac*-gene cluster in *Streptococcus pneumoniae*. *Appl. Environ. Microbiol.* 80, 5349–5358. doi: 10.1128/AEM.01370-14
- Afzal, M., Shafeeq, S., and Kuipers, O. P. (2015d). Ascorbic acid-dependent gene expression in *Streptococcus pneumoniae* and the activator function of the transcriptional regulator UlaR2. *Front. Microbiol.* 6:72. doi: 10.3389/fmicb.2015.00072
- Afzal, M., Shafeeq, S., Manzoor, I., and Kuipers, O. P. (2015e). GalR acts as a transcriptional activator of *galKT* in the presence of galactose in *Streptococcus pneumoniae*. *J. Mol. Microbiol. Biotechnol.* 25, 363–371. doi: 10.1159/000439429
- Afzal, M., Shafeeq, S., Manzoor, I., and Kuipers, O. P. (2015f). Maltose-dependent transcriptional regulation of the *mal* regulon by MalR in *Streptococcus pneumoniae*. *PLoS ONE* 10:e0127579. doi: 10.1371/journal.pone.0127579
- Alice, A. F., Pérez-Martínez, G., and Sánchez-Rivas, C. (2003). Phosphoenolpyruvate phosphotransferase system and N-acetylglucosamine metabolism in *Bacillus sphaericus*. *Microbiol. Read. Engl.* 149, 1687–1698. doi: 10.1099/mic.0.26231-0
- Alvarez-Añorve, L. I., Bustos-Jaimes, I., Calcagno, M. L., and Plumbridge, J. (2009). Allosteric regulation of glucosamine-6-phosphate deaminase (NagB) and growth of *Escherichia coli* on glucosamine. *J. Bacteriol.* 191, 6401–6407. doi: 10.1128/JB.00633-09
- Baerends, R. J. S., Smits, W. K., de Jong, A., Hamoen, L. W., Kok, J., and Kuipers, O. P. (2004). Genome2D: a visualization tool for the rapid analysis of bacterial transcriptome data. *Genome Biol.* 5:R37. doi: 10.1186/gb-2004-5-5-r37
- Bailey, T. L., and Elkan, C. (1994). Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proc. Int. Conf. Intell. Syst. Mol. Biol. ISMB Int. Conf. Intell. Syst. Mol. Biol.* 2, 28–36.
- Bertram, R., Rigali, S., Wood, N., Lulko, A. T., Kuipers, O. P., and Titgemeyer, F. (2011). Regulon of the N-acetylglucosamine utilization regulator NagR in *Bacillus subtilis*. *J. Bacteriol.* 193, 3525–3536. doi: 10.1128/JB.00264-11
- Bidossi, A., Mulas, L., Decorosi, F., Colomba, L., Ricci, S., Pozzi, G., et al. (2012). A functional genomics approach to establish the complement of carbohydrate transporters in *Streptococcus pneumoniae*. *PLoS ONE* 7:e33320. doi: 10.1371/journal.pone.0033320

- Buckwalter, C. M., and King, S. J. (2012). Pneumococcal carbohydrate transport: food for thought. *Trends Microbiol.* 20, 517–522. doi: 10.1016/j.tim.2012.08.008
- Burnaugh, A. M., Frantz, L. J., and King, S. J. (2008). Growth of *Streptococcus pneumoniae* on human glycoconjugates is dependent upon the sequential activity of bacterial exoglycosidases. *J. Bacteriol.* 190, 221–230. doi: 10.1128/JB.01251-07
- Carvalho, S. M., Kloosterman, T. G., Kuipers, O. P., and Neves, A. R. (2011). CcpA ensures optimal metabolic fitness of *Streptococcus pneumoniae*. *PLoS ONE* 6:e26707. doi: 10.1371/journal.pone.0026707
- Dalia, A. B., Standish, A. J., and Weiser, J. N. (2010). Three surface exoglycosidases from *Streptococcus pneumoniae*, NanA, BgaA, and StrH, promote resistance to opsonophagocytic killing by human neutrophils. *Infect. Immun.* 78, 2108–2116. doi: 10.1128/IAI.01125-09
- Dobrogosz, W. J. (1968). Effect of amino sugars on catabolite repression in *Escherichia coli*. *J. Bacteriol.* 95, 578–584.
- Gaugué, I., Oberto, J., Putzer, H., and Plumbridge, J. (2013). The use of amino sugars by *Bacillus subtilis*: presence of a unique operon for the catabolism of glucosamine. *PLoS ONE* 8:e63025. doi: 10.1371/journal.pone.0063025
- Halfmann, A., Hakenbeck, R., and Bruckner, R. (2007). A new integrative reporter plasmid for *Streptococcus pneumoniae*. *FEMS Microbiol. Lett.* 268, 217–224. doi: 10.1111/j.1574-6968.2006.00584.x
- Hoskins, J., Alborn, W. E. Jr., Arnold, J., Blaszcak, L. C., Burgett, S., DeHoff, B. S., et al. (2001). Genome of the bacterium *Streptococcus pneumoniae* strain R6. *J. Bacteriol.* 183, 5709–5717. doi: 10.1128/JB.183.19.5709-5717.2001
- Ispahani, P., Slack, R. C. B., Donald, F. E., Weston, V. C., and Rutter, N. (2004). Twenty year surveillance of invasive pneumococcal disease in Nottingham: serogroups responsible and implications for immunisation. *Arch. Dis. Child.* 89, 757–762. doi: 10.1136/adc.2003.036921
- Israelsen, H., Madsen, S. M., Vrang, A., Hansen, E. B., and Johansen, E. (1995). Cloning and partial characterization of regulated promoters from *Lactococcus lactis* Tn917-lacZ integrants with the new promoter probe vector, pAK80. *Appl. Environ. Microbiol.* 61, 2540–2547.
- Jeong, J. K., Kwon, O., Lee, Y. M., Oh, D. B., Lee, J. M., Kim, S., et al. (2009). Characterization of the *Streptococcus pneumoniae* BgaC protein as a novel surface beta-galactosidase with specific hydrolysis activity for the Galbeta1-3GlcNAc moiety of oligosaccharides. *J. Bacteriol.* 191, 3011–3023. doi: 10.1128/JB.01601-08
- Kanehisa, M., Goto, S., Sato, Y., Kawashima, M., Furumichi, M., and Tanabe, M. (2014). Data, information, knowledge and principle: back to metabolism in KEGG. *Nucleic Acids Res.* 42, D199–D205. doi: 10.1093/nar/gkt1076
- Kawada-Matsuo, M., Mazda, Y., Oogai, Y., Kajiyama, M., Kawai, T., Yamada, S., et al. (2012). GlnS and NagB regulate amino sugar metabolism in opposing directions and affect *Streptococcus mutans* virulence. *PLoS ONE* 7:e33382. doi: 10.1371/journal.pone.0033382
- King, S. J. (2010). Pneumococcal modification of host sugars: a major contributor to colonization of the human airway? *Mol. Oral Microbiol.* 25, 15–24. doi: 10.1111/j.2041-1014.2009.00564.x
- King, S. J., Hippe, K. R., and Weiser, J. N. (2006). Deglycosylation of human glycoconjugates by the sequential activities of exoglycosidases expressed by *Streptococcus pneumoniae*. *Mol. Microbiol.* 59, 961–974. doi: 10.1111/j.1365-2958.2005.04984.x
- Kloosterman, T. G., Bijlsma, J. J. E., Kok, J., and Kuipers, O. P. (2006a). To have neighbour's fare: extending the molecular toolbox for *Streptococcus pneumoniae*. *Microbiol. Read. Engl.* 152, 351–359. doi: 10.1099/mic.0.28521-0
- Kloosterman, T. G., Hendriksen, W. T., Bijlsma, J. J., Bootsma, H. J., van Hijum, S. A., Kok, J., et al. (2006b). Regulation of glutamine and glutamate metabolism by GlnR and GlnA in *Streptococcus pneumoniae*. *J. Biol. Chem.* 281, 25097–25109. doi: 10.1074/jbc.M601661200
- Kloosterman, T. G., and Kuipers, O. P. (2011). Regulation of arginine acquisition and virulence gene expression in the human pathogen *Streptococcus pneumoniae* by transcription regulators ArgR1 and AhrC. *J. Biol. Chem.* 286, 44594–44605. doi: 10.1074/jbc.M111.295832
- Lanie, J. A., Ng, W. L., Kazmierczak, K. M., Andrzejewski, T. M., Davidsen, T. M., Wayne, K. J., et al. (2007). Genome sequence of Avery's virulent serotype 2 strain D39 of *Streptococcus pneumoniae* and comparison with that of unencapsulated laboratory strain R6. *J. Bacteriol.* 189, 38–51. doi: 10.1128/JB.01148-06
- Leenhouts, K., Venema, G., and Kok, J. (1998). A lactococcal pWV01 based integration toolbox for bacteria. *Methods Cell Sci.* 20, 35–50. doi: 10.1023/A:1009862119114
- Linke, C. M., Woodiga, S. A., Meyers, D. J., Buckwalter, C. M., Salhi, H. E., and King, S. J. (2013). The ABC transporter encoded at the pneumococcal fructooligosaccharide utilization locus determines the ability to utilize long- and short-chain fructooligosaccharides. *J. Bacteriol.* 195, 1031–1041. doi: 10.1128/JB.01560-12
- Lortie, L. A., Pelletier, M., Vadeboncoeur, C., and Frenette, M. (2000). The gene encoding IIAB(Man)L in *Streptococcus salivarius* is part of a tetracistronic operon encoding a phosphoenolpyruvate: mannose/glucose phosphotransferase system. *Microbiol. Read. Engl.* 146(Pt 3), 677–685. doi: 10.1099/00221287-146-3-677
- Lulko, A. T., Buist, G., Kok, J., and Kuipers, O. P. (2007). Transcriptome analysis of temporal regulation of carbon metabolism by CcpA in *Bacillus subtilis* reveals additional target genes. *J. Mol. Microbiol. Biotechnol.* 12, 82–95. doi: 10.1159/000096463
- Marion, C., Burnaugh, A. M., Woodiga, S. A., and King, S. J. (2011). Sialic acid transport contributes to pneumococcal colonization. *Infect. Immun.* 79, 1262–1269. doi: 10.1128/IAI.00832-10
- Marion, C., Limoli, D. H., Bobulsky, G. S., Abraham, J. L., Burnaugh, A. M., and King, S. J. (2009). Identification of a pneumococcal glycosidase that modifies O-linked glycans. *Infect. Immun.* 77, 1389–1396. doi: 10.1128/IAI.01215-08
- Marion, C., Stewart, J. M., Tazi, M. F., Burnaugh, A. M., Linke, C. M., Woodiga, S. A., et al. (2012). *Streptococcus pneumoniae* can utilize multiple sources of hyaluronic acid for growth. *Infect. Immun.* 80, 1390–1398. doi: 10.1128/IAI.05756-11
- Mobley, H. L., Doyle, R. J., Streips, U. N., and Langemeier, S. O. (1982). Transport and incorporation of N-acetyl-D-glucosamine in *Bacillus subtilis*. *J. Bacteriol.* 150, 8–15.
- Moye, Z. D., Burne, R. A., and Zeng, L. (2014). Uptake and metabolism of N-acetylglucosamine and glucosamine by *Streptococcus mutans*. *Appl. Environ. Microbiol.* 80, 5053–5067. doi: 10.1128/AEM.00820-14
- Neves, A. R., Ventura, R., Mansour, N., Shearman, C., Gasson, M. J., Maycock, C., et al. (2002). Is the glycolytic flux in *Lactococcus lactis* primarily controlled by the redox charge? Kinetics of NAD(+) and NADH pools determined *in vivo* by 13C NMR. *J. Biol. Chem.* 277, 28088–28098. doi: 10.1074/jbc.M202573200
- Nothaft, H., Dresel, D., Willimek, A., Mahr, K., Niederweis, M., and Titgemeyer, F. (2003). The phosphotransferase system of *Streptomyces coelicolor* is biased for N-acetylglucosamine metabolism. *J. Bacteriol.* 185, 7019–7023. doi: 10.1128/JB.185.23.7019-7023.2003
- Nothaft, H., Rigali, S., Boomsma, B., Swiatek, M., McDowall, K. J., van Wezel, G. P., et al. (2010). The permease gene nagE2 is the key to N-acetylglucosamine sensing and utilization in *Streptomyces coelicolor* and is subject to multi-level control. *Mol. Microbiol.* 75, 1133–1144. doi: 10.1111/j.1365-2958.2009.07020.x
- Novichkov, P. S., Laikova, O. N., Novichkova, E. S., Gelfand, M. S., Arkin, A. P., Dubchak, I., et al. (2010). RegPrecise: a database of curated genomic inferences of transcriptional regulatory interactions in prokaryotes. *Nucleic Acids Res.* 38, D111–D118. doi: 10.1093/nar/gkp894
- O'Brien, K. L., Wolfson, L. J., Watt, J. P., Henkle, E., Deloria-Knoll, M., McCall, N., et al. (2009). Burden of disease caused by *Streptococcus pneumoniae* in children younger than 5 years: global estimates. *Lancet* 374, 893–902. doi: 10.1016/S0140-6736(09)61204-6
- Paixão, L., Oliveira, J., Verissimo, A., Vinga, S., Lourenço, E. C., Ventura, M. R., et al. (2015). Host glycan sugar-specific pathways in *Streptococcus pneumoniae*: galactose as a key sugar in colonisation and infection [corrected]. *PLoS ONE* 10:e0121042. doi: 10.1371/journal.pone.0121042
- Paulsen, I. T., Nguyen, L., Sliwinski, M. K., Rabus, R., and Saier, M. H. (2000). Microbial genome analyses: comparative transport capabilities in eighteen prokaryotes. *J. Mol. Biol.* 301, 75–100. doi: 10.1006/jmbi.2000.3961
- Phillips, N. J., John, C. M., Reinders, L. G., Gibson, B. W., Apicella, M. A., and Griffiss, J. M. (1990). Structural models for the cell surface lipooligosaccharides of *Neisseria gonorrhoeae* and *Haemophilus influenzae*. *Biomed. Environ. Mass Spectrom.* 19, 731–745. doi: 10.1002/bms.1200191112
- Plumbridge, J. (2001). Regulation of PTS gene expression by the homologous transcriptional regulators, Mlc and NagC, in *Escherichia coli* (or how two similar repressors can behave differently). *J. Mol. Microbiol. Biotechnol.* 3, 371–380.

- Plumbridge, J. A. (1991). Repression and induction of the nag regulon of *Escherichia coli* K-12: the roles of nagC and nagA in maintenance of the uninduced state. *Mol. Microbiol.* 5, 2053–2062. doi: 10.1111/j.1365-2958.1991.tb00828.x
- Plumbridge, J. A., Cochet, O., Souza, J. M., Altamirano, M. M., Calcagno, M. L., and Badet, B. (1993). Coordinated regulation of amino sugar-synthesizing and -degrading enzymes in *Escherichia coli* K-12. *J. Bacteriol.* 175, 4951–4956.
- Plumbridge, J., and Vimr, E. (1999). Convergent pathways for utilization of the amino sugars N-acetylglucosamine, N-acetylmannosamine, and N-acetylneuraminic acid by *Escherichia coli*. *J. Bacteriol.* 181, 47–54.
- Pluvinae, B., Stubbs, K. A., Hattie, M., Vocadlo, D. J., and Boraston, A. B. (2013). Inhibition of the family 20 glycoside hydrolase catalytic modules in the *Streptococcus pneumoniae* exo- β -D-N-acetylglucosaminidase, StrH. *Org. Biomol. Chem.* 11, 7907–7915. doi: 10.1039/c3ob41579a
- Reizer, J., Bachem, S., Reizer, A., Arnaud, M., Saier, M. H. Jr., and Stulke, J. (1999). Novel phosphotransferase system genes revealed by genome analysis - the complete complement of PTS proteins encoded within the genome of *Bacillus subtilis*. *Microbiol. Read. Engl.* 145(Pt 12), 3419–3429. doi: 10.1099/00221287-145-12-3419
- Rose, M. C., and Voynow, J. A. (2006). Respiratory tract mucin genes and mucin glycoproteins in health and disease. *Physiol. Rev.* 86, 245–278. doi: 10.1152/physrev.00010.2005
- Saier, M. H., Goldman, S. R., Maile, R. R., Moreno, M. S., Weyler, W., Yang, N., et al. (2002). Transport capabilities encoded within the *Bacillus subtilis* genome. *J. Mol. Microbiol. Biotechnol.* 4, 37–67.
- Shafeeq, S., Afzal, M., Henriques-Normark, B., and Kuipers, O. P. (2015). Transcriptional profiling of UlaR-regulated genes in *Streptococcus pneumoniae*. *Genomics Data* 4, 57–59. doi: 10.1016/j.gdata.2015.02.004
- Shafeeq, S., Kloosterman, T. G., and Kuipers, O. P. (2011). CelR-mediated activation of the cellobiose-utilization gene cluster in *Streptococcus pneumoniae*. *Microbiol. Read. Engl.* 157, 2854–2861. doi: 10.1099/mic.0.051359-0
- Shafeeq, S., Kloosterman, T. G., Rajendran, V., and Kuipers, O. P. (2012). Characterization of the ROK-family transcriptional regulator RokA of *Streptococcus pneumoniae* D39. *Microbiol. Read. Engl.* 158, 2917–2926. doi: 10.1099/mic.0.062919-0
- Shafeeq, S., Kuipers, O. P., and Kloosterman, T. G. (2013). Cellobiose-mediated gene expression in *Streptococcus pneumoniae*: a repressor function of the novel GntR-type regulator BguR. *PLoS ONE* 8:e57586. doi: 10.1371/journal.pone.0057586
- Simoni, R. D., Roseman, S., and Saier, M. H. (1976). Sugar transport. Properties of mutant bacteria defective in proteins of the phosphoenolpyruvate: sugar phosphotransferase system. *J. Biol. Chem.* 251, 6584–6597.
- Terra, V. S., Homer, K. A., Rao, S. G., Andrew, P. W., and Yesilkaya, H. (2010). Characterization of novel beta-galactosidase activity that contributes to glycoprotein degradation and virulence in *Streptococcus pneumoniae*. *Infect. Immun.* 78, 348–357. doi: 10.1128/IAI.00721-09
- Tettelin, H., Nelson, K. E., Paulsen, I. T., Eisen, J. A., Read, T. D., Peterson, S., et al. (2001). Complete genome sequence of a virulent isolate of *Streptococcus pneumoniae*. *Science* 293, 498–506. doi: 10.1126/science.1061217
- Titgemeyer, F., and Hillen, W. (2002). Global control of sugar metabolism: a gram-positive solution. *Antonie Van Leeuwenhoek* 82, 59–71. doi: 10.1023/A:1020628909429
- Titgemeyer, F., Reizer, J., Reizer, A., and Saier, M. H. Jr. (1994). Evolutionary relationships between sugar kinases and transcriptional repressors in bacteria. *Microbiol. Read. Engl.* 140(Pt 9), 2349–2354. doi: 10.1099/13500872-140-9-2349
- Tong, H. H., Blue, L. E., James, M. A., and DeMaria, T. F. (2000). Evaluation of the virulence of a *Streptococcus pneumoniae* neuraminidase-deficient mutant in nasopharyngeal colonization and development of otitis media in the chinchilla model. *Infect. Immun.* 68, 921–924. doi: 10.1128/IAI.68.2.921-924.2000
- Tyx, R. E., Roche-Hakansson, H., and Hakansson, A. P. (2011). Role of dihydrolipoamide dehydrogenase in regulation of raffinose transport in *Streptococcus pneumoniae*. *J. Bacteriol.* 193, 3512–3524. doi: 10.1128/JB.01410-10
- Vadeboncoeur, C. (1984). Structure and properties of the phosphoenolpyruvate: glucose phosphotransferase system of oral streptococci. *Can. J. Microbiol.* 30, 495–502. doi: 10.1139/m84-073
- White, R. J. (1970). The role of the phosphoenolpyruvate phosphotransferase system in the transport of N-acetyl-D-glucosamine by *Escherichia coli*. *Biochem. J.* 118, 89–92. doi: 10.1042/bj1180089
- Yesilkaya, H., Manco, S., Kadioglu, A., Terra, V. S., and Andrew, P. W. (2008). The ability to utilize mucin affects the regulation of virulence gene expression in *Streptococcus pneumoniae*. *FEMS Microbiol. Lett.* 278, 231–235. doi: 10.1111/j.1574-6968.2007.01003.x
- Zeng, L., and Burne, R. A. (2015). NagR differentially regulates the expression of the glmS and nagAB genes required for amino sugar metabolism by *Streptococcus mutans*. *J. Bacteriol.* 197, 3533–3544. doi: 10.1128/JB.00606-15
- Zomer, A. L., Buist, G., Larsen, R., Kok, J., and Kuipers, O. P. (2007). Time-resolved determination of the CcpA regulon of *Lactococcus lactis* subsp. cremoris MG1363. *J. Bacteriol.* 189, 1366–1381. doi: 10.1128/JB.01013-06

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2016 Afzal, Shafeeq, Manzoor, Henriques-Normark and Kuipers. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.