

University of Groningen

The genetics of spinocerebellar ataxia and dystonia

Nibbeling, Esther

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Nibbeling, E. (2017). *The genetics of spinocerebellar ataxia and dystonia*. Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Using the shared genetics of dystonia and ataxia to unravel their pathogenesis

Esther AR Nibbeling¹, Cathérine CS Delnooz², Tom J de Koning^{1,2},
Richard J Sinke¹, Hyder A Jinnah³, Marina AJ Tijssen² and Dineke S Verbeek¹

1. *University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, the Netherlands*
2. *University of Groningen, University Medical Center Groningen, Department of Neurology, Groningen, the Netherlands*
3. *Departments of Neurology, Human Genetics and Pediatrics, Emory Clinic, Atlanta, USA*

Neurosci Biobehav Rev. 2017;75:22-39; doi: 10.1016/j.neubiorev.2017.01.033



Abstract

In this review we explore the similarities between spinocerebellar ataxias and dystonias, and suggest potentially shared molecular pathways using a gene co-expression network approach. The spinocerebellar ataxias are a group of neurodegenerative disorders characterized by coordination problems caused mainly by atrophy of the cerebellum. The dystonias are another group of neurological movement disorders linked to basal ganglia dysfunction, although evidence is now pointing to cerebellar involvement as well. Our gene co-expression network approach identified 99 shared genes and showed the involvement of two major pathways: synaptic transmission and neurodevelopment. These pathways overlapped in the two disorders, with a large role for GABAergic signaling in both. The overlapping pathways may provide novel targets for disease therapies. We need to prioritize variants obtained by whole exome sequencing in the genes associated with these pathways in the search for new pathogenic variants, which can then be used to help in the genetic counseling of patients and their families.

Keywords

Spinocerebellar ataxia; dystonia; gene network; synaptic transmission; neurodevelopment; neurodegeneration; molecular pathways; pathophysiology

Introduction

The cerebellar ataxias are a heterogeneous group of movement disorders characterized by degeneration of Purkinje cells (PCs) and atrophy of the cerebellum. Motor symptoms include loss of balance and coordination, unstable gait, dysarthria and abnormal eye movements. Cerebellar ataxias can be primary (genetic), congenital (brain malformations) or acquired (e.g. after stroke). The spinocerebellar ataxias (SCAs), the genetically dominant forms of cerebellar ataxia, have an estimated prevalence of 1-3 per 100,000 in Europe with onset usually occurring in adulthood.¹ Dystonia is a neurological movement disorder characterized by involuntary muscle contractions that cause abnormal twisting movements and postures. It has many clinical manifestations, ranging from isolated and focal to generalized dystonia, or dystonia in combination with other neurological symptoms such as myoclonus or ataxia. The list of diseases that can cause or present with dystonia is extensive.²

Many patients show a combination of cerebellar ataxia and dystonia. Dystonia is frequently seen in SCA2 (14%), SCA3 (24%) and SCA17 (53%), and regularly seen in SCA types 1, 6, 12, 14, 15/16 and 20, in ataxia telangiectasia, in Friedreich's ataxia, and in ataxia with oculomotor apraxia.³⁻⁵ Kuoppamaki and Van de Warrenburg reported eleven patients in total who showed early onset, primarily cervical, dystonia in combination with slowly progressive cerebellar ataxia. All had tested negative for the most common SCA types, although some patients had a positive family history, and all the patients showed cerebellar atrophy.^{6,7}

Taken into account the clinical and etiological heterogeneity, the exact pathophysiological mechanisms of SCA and dystonia are not exactly clear. For SCA, several etiological roles have been identified that lead to neurotransmission deficits and result in PC death, including transcriptional dysregulation, autophagy, mitochondrial defects and alterations in calcium homeostasis.⁸ In dystonia, the basal ganglia have classically been attributed a key role. However, recent theories support a pathophysiological model in which dystonia is seen as a network disorder involving several brain regions, including the sensorimotor cortex, brainstem, thalamus and cerebellum.^{4,5} Nevertheless, it remains uncertain whether dysfunction of a single brain area, combined dysfunction of multiple areas, or abnormal communication between several brain areas leads to dystonia. Dystonia is regarded as a disorder of motor control⁹ involving the cerebellum¹⁰ and the basal ganglia, which are interconnected (Figure 1). The cerebellum also plays a role in cerebellar ataxias. In this review, we therefore focus on the potentially shared pathophysiology of the cerebellum in SCA and dystonia.

Evidence for overlap in pathology between ataxia and dystonia

Evidence from clinical studies

Evidence of cerebellar involvement in dystonia comes from several lines of research. By the beginning of the 20th century it had already been recognized that tumors in the posterior fossa could result in the abnormal postures of the head that we would now classify as dystonia.^{11–14} These clinical findings were replicated in a larger cohort of 25 cervical dystonia patients, in which almost half of the patients had a lesion in the brainstem or cerebellum, whereas lesions in the basal ganglia were seen in only a quarter of them.¹⁵ Batla et al. showed cerebellar abnormalities in 26 out of 188 (14%) cervical dystonia patients.¹⁶ For secondary blepharospasm, a focal form of dystonia, lesions were found mostly in the thalamus, with the remainder equally split between basal ganglia and cerebellum.¹⁷ Other cases reported oromandibular dystonia and blepharospasm after cerebellar infarction,^{18–20} hemidystonia caused by vertebral artery occlusion,²¹ and focal limb dystonia after isolated cerebellar tuberculoma.²²

Only small case series have been reported for neuropathological changes in dystonia, specifically in isolated cervical dystonia. A recent review showed that no pathological abnormalities were found in almost all reported cases that had a high probability of suffering from cervical dystonia.²³ These case studies have, however, several shortcomings. The major limitation is that most were focused on specific brain regions that did not include the cerebellum or the brainstem. The authors also found that the cerebellums of the cervical dystonia patients had significantly lower PC density compared to healthy controls.^{23,24} This implies a role for the cerebellum in dystonia, and it is further worth noting that loss of PCs is also associated with other neurodegenerative disorders, such as SCA.

Evidence from imaging studies

In addition to alterations in the sensorimotor cortices and the basal ganglia, structural abnormalities in the cerebellum or cerebellar projections have been found in several types of dystonia. Diffusion tensor imaging was used to assess microstructural white matter integrity in different non-hereditary isolated dystonias and demonstrated alterations varying from the white matter tracts underlying several cortical areas^{25,26} to connections to the cerebellar lobules and peduncles.^{27–32} See reviews by Neychev et al. and Zoons et al.^{4,33} for a complete overview of the evidence for the role of the cerebellum based on imaging studies. More recent findings come from studies of several groups of hereditary dystonia. In DYT1, DYT6 and DYT11 patients, microstructural abnormalities were found close to the superior cerebellar peduncle.^{34–36} A reduction in structural connectivity of the cerebellothalamic pathway in DYT1 and DYT6 patients was also seen by tractography.³⁷

Metabolic imaging using [18F]-fluorodeoxyglucose-PET also shows involvement of the cerebellum in dystonia. The most reported pattern of altered metabolic activity involves the basal ganglia, pre-motor and motor areas, and the cerebellum, and is present in both hereditary as well as non-hereditary isolated dystonia.³⁸ More recently, Niethammer et al. described a motor-related activation pattern characterized by cerebello-thalamo-cortical motor circuits, that is increased in both hereditary and non-hereditary dystonia and even in non-manifesting carriers of DYT1 and DYT6.²⁸ Furthermore, the results of the eight voxel-based morphometry studies published to date have been inconsistent. An *increase* in gray matter volume of several parts of the cerebellum was seen in various types of isolated focal dystonia,^{29,30,39–41} while a *decrease* in gray matter volume was shown in other studies.^{30,42,43}

Lastly, cerebellar abnormalities have been frequently reported in a number of studies using fMRI. In addition to alterations in sensorimotor cortical areas and the basal ganglia, altered task-related activity (e.g. finger tapping, writing or speaking) was found in several cerebellar areas, including the cerebellar nuclei, posterior vermis, and paramedian cerebellar hemisphere. Again, however, the effects were inconsistent across and between different types of isolated dystonia.^{44–51}

Evidence from electrophysiological studies

In the past few years, electrophysiological studies on dystonia have identified three common themes that explain its pathophysiology: loss of inhibition, maladaptive plasticity and defective sensorimotor integration.⁹ Classically in isolated dystonia, there is a reduction in cortical inhibition (i.e. an increase in motor-evoked potentials and a reduction of the short intracortical inhibition using transcranial magnetic stimulation). However, in some types of dystonia (e.g. myoclonus dystonia), a contrasting pattern is seen that resembles the excitability profile of cerebellar pathology.^{52,53} Maladaptive plasticity, a second common theme, has been demonstrated in several types of dystonia.⁵⁴ It is known, for instance, that beyond the basal ganglia, the cerebellum also plays a role in plasticity and motor learning.⁵⁵ Eye blink classical conditioning (EBCC) is a paradigm for associative motor learning that has been shown to be highly dependent on cerebellar functioning without basal ganglia involvement.^{56,57} Teo and co-authors demonstrated significant abnormal EBCC in isolated dystonia patients,⁵⁸ which could be normalized by continuous theta burst stimulation; this suggests a secondary role for the cerebellum.⁵⁹ A study on sensorimotor adaptation using a motor learning paradigm that relies on both the cerebellum and the sensorimotor network (split-belt paradigm) demonstrated abnormalities in patients with blepharospasm and writer's cramp.⁶⁰ Defective sensorimotor adaptation was also seen in writer's cramp patients performing a visuomotor adaptation task, suggesting that the cerebellum had lost its ability to modulate sensorimotor plasticity of the motor cortex.⁶¹ In contrast to these findings,

normal sensorimotor adaptation was seen in cervical dystonia patients.^{60,62} Experiments with a therapeutic focus also demonstrated cerebellar involvement in dystonia. In focal hand dystonia patients, cerebellar transcranial direct current stimulation improved handwriting and cyclic drawing kinematics, most likely by reducing cerebellar-brain inhibition.⁶³ In cervical dystonia patients, two weeks of cerebellar stimulation resulted in a small but significant, clinical improvement of approximately 15% measured by the Toronto Western Spasmodic Torticollis Rating Scale.⁶⁴

While these effects of non-invasive stimulation have been modest or transient so far, they show that manipulating cerebellar physiology can influence the severity of dystonia. More direct manipulations of the cerebellum, analogous to pallidotomy or deep brain stimulation of the basal ganglia, may be required to produce robust and lasting effects. Cerebellar dentatectomy⁶⁵ and direct electrical stimulation of the cerebellum⁶⁶ were once routinely applied as treatments for dystonia, but these procedures were abandoned because the benefits were unpredictable. However, the indication for these procedures was “hypertonia” or “cerebral palsy”, which could reflect any combination of dystonia, spasticity and/or rigidity, and this grouping together of many disorders with potentially different causes may have been responsible for the unpredictable outcomes. Now that the various types of hypertonia are better distinguished, more recent studies have begun to discriminate between the beneficial effects of dentatectomy⁶⁷ and cerebellar stimulation,⁶⁸ although further studies are still needed.

Evidence from animal models

In line with the evidence from patients who show both dystonia and cerebellar ataxia, several animal models confirm the role of the cerebellum in the etiology of dystonia. Ataxia and dystonia were identified in a mouse model (leaner) with a spontaneous missense mutation in the splice donor consensus sequence, at the 5' end of the affected intron in *Cacna1a*, which encodes the voltage-gated calcium channel Ca_v2.1.^{69–71} This phenotype is associated with abnormal physiological activity and slow degeneration of cerebellar Purkinje neurons.^{70,72–77} Mutations in *CACNA1A* have been linked to a range of movement disorders including SCA6 and benign paroxysmal torticollis of infancy.^{78,79} Notably, in the leaner mouse model, the dystonia abated while the ataxia worsened as PCs were lost over time.⁸⁰ This finding is evidence for the hypothesis that PC-dysfunction leads to dystonia, whereas PC-loss causes ataxia. Additionally, *Cacna1a* null mice also exhibited dystonia and highly selective cerebellar degeneration, further confirming a key role for PC functioning in a shared pathology of dystonia and ataxia.⁶⁹ Additionally, a mouse model for rapid-onset dystonia-Parkinsonism (RDP), which mimicked the effect of mutations in the $\alpha 3$ isoform of the Na(+)/K(+)-ATPase (sodium pump), exhibited ataxia quickly followed by a dystonic phenotype upon blockage of $\alpha 3$ -sodium pumps with ouabain.⁸¹ Fremont et al. showed that restricted cerebellar perfusion with ouabain

is sufficient to induce dystonia, and *in vivo* recordings from these dystonic mice showed persistent high-frequency-burst firing of PCs.⁸² Moreover, selective knock down of the $\alpha 3$ -sodium pump in the substantia nigra resulted in a Parkinsonism phenotype, while knockdown in other basal ganglia regions had no apparent effect. Cerebellum-specific knockdown of the $\alpha 3$ -sodium pump recapitulated the phenotype of the RDP mouse, which was again associated with altered intrinsic pacemaking of PCs, but it had no effect on the firing rate of neurons from the deep cerebellar nuclei (DCN) (see Figure 1 for projections between brain regions).⁸³ However, DCN neurons fired irregularly in dystonic animals compared to controls, a difference most likely caused by aberrant PC input to the DCN.⁸³ Furthermore, a disynaptic connection between the cerebellum and the basal ganglia via the thalamic intralaminar nuclei, was shown to underlie the dystonic phenotype of the RDP mouse, as lesions of the centrolateral nucleus alleviated the dystonia.⁸⁴




Figure 1. Schematic representation of connections between basal ganglia and cerebellum. Blue arrows represent excitatory glutamatergic projections. Green arrows represent inhibitory GABAergic projections. Yellow arrow represents dopaminergic projections. CB = cerebellum; GC = granule cells; PC = Purkinje cells; DCN = deep cerebellar nuclei; STN = subthalamic nucleus; SNr = substantia nigra pars reticulata; SNc = substantia nigra pars compacta; GPe = globus pallidus externus; GPi = globus pallidus internus.

In another model, *Atcay*^{ii-hes} mice, which show very low levels of the caytaxin protein, action-induced stiff dystonic legs can be turned into broad-based ataxic gait by partial cerebellectomy or lesions of the DCN.⁸⁵ In contrast, homozygous missense mutations in *ATCAY*, the human orthologue of *Atcay*, cause autosomal recessive Cayman ataxia but not dystonia.⁸⁶ The dystonic phenotype in mice is most likely the consequence of increased repetitive firing of DCN neurons caused by absence of inhibitory repetitive firing of PCs. Thus, it is the combination of aberrant firing patterns of PCs and DCN neu-

rons that likely exaggerates the hyperexcitability of the DCN neurons above a threshold sufficient to directly activate muscle groups resulting in dystonia and ataxia.

Finally, knockout of *fgf14*, the human disease gene underlying SCA27, resulted in ataxia and a paroxysmal hyperkinetic movement disorder mimicking a form of dystonia.⁸⁷ With the notion that *fgf14* was most abundantly expressed in cerebellar granule cells of the cerebellum, PC dysfunction rather than a direct PC deficit seems to cause the ataxia and dystonia-like phenotype.

Overall, these studies show a clear role for the cerebellum in the etiology of dystonia. We therefore hypothesize that shared genetic pathways might underlie the pathogenesis of dystonia and SCA, and that the cerebellum plays a major role in both disorders. To further elucidate this hypothesis, we first explore the current knowledge about the genetic background of both groups of disorders.

Genetics of dystonia and SCA

The introduction of high-throughput next generation sequencing (NGS), which rapidly detects all protein-coding variants, has led to the discovery of multiple disease genes for SCA (Table 1) and dystonia (Table 2). Exome sequencing has become common practice for gene identification of Mendelian forms of dystonia and SCA. To date, 16 genes have been identified using exome sequencing, including the recently discovered mutations in *COL6A3*⁸⁸ and *TRPC3*.⁸⁹ However, linkage analysis is still commonly used to pinpoint the region of interest and to filter exome sequencing results. Exome sequencing has its limitations, and an alternative method would be genome sequencing of the complete genome (coding and non-coding parts). However, genome sequencing remains relatively expensive and produces a very long list of putative candidates with undefined pathogenicity, many of which will be located in non-coding regions, making pathogenicity very difficult to establish. Furthermore, many diagnostic requests come from independently referred patients (singletons) who lack the additional affected family members necessary for co-segregation analysis. In reality, conventional genetic testing is usually only performed for the most common SCA and dystonia types, leaving a large group of patients genetically undiagnosed.

Despite the many disease genes that have been identified, our knowledge of the underlying biological pathways and pathogenesis of dystonia and ataxia is still limited. The main biological pathway linked to dystonia before the introduction of NGS was dopamine synthesis, which was implicated via mutations in *GCH1* and *TH*, which encode GTP cyclohydrolase 1 and tyrosine hydroxylase, respectively.^{90,91} This finding was reinforced by the observation that these patients are levodopa (L-DOPA) responsive.⁹²

Table 1. Known SCA genes

SCA type	OMIM Number	Locus	Gene Name	Mutation type	References
SCA1	164400	6p22.3	<i>ATXN1</i>	CAG repeat	240
SCA2	183090	12q24.13	<i>ATXN2</i>	CAG repeat	241,242
SCA3	109150	14p32.12	<i>ATXN3</i>	CAG repeat	243,244
SCA5	600224	11q13.2	<i>SPTBN2</i>	Deletion, MM	218,245
SCA6	183086	19p13.13	<i>CACNA1A</i>	CAG repeat	78
SCA7	164500	3p14.1	<i>ATXN7</i>	CAG repeat	246
SCA8	608768	13q21	<i>ATXN8OS</i>	CTG repeat (non-coding)	247
SCA10	603516	22q13.31	<i>ATXN10</i>	ATTCT repeat	248–250
SCA11	604432	15q15.2	<i>TTBK2</i>	Deletion	251
SCA12	604326	5q32	<i>PPP2R2B</i>	CAG repeat (non-coding)	252
SCA13	605259	19q13.33	<i>KCNC3</i>	MM	102,253
SCA14	605361	19q13.42	<i>PRKCG</i>	MM	254
SCA15	606658	3p26.1	<i>ITPR1</i>	Deletion	255,256
SCA16	606658	3p26.1	<i>ITPR1</i>	Deletion	255,256
SCA17	607136	6q27	<i>TBP</i>	CAG repeat	257
SCA19/22	607346	1p13.2	<i>KCND3</i>	MM	103,104,258,259
SCA20	608687	11q12	<i>DAGLA</i>	12 genes duplication	260,261
SCA21	607454	1p36.33	<i>TMEM240</i>	MM	262
SCA23	610245	20p13	<i>PDYN</i>	MM, frameshift	116,263
SCA26	609306	19p13.3	<i>EEF2</i>	MM	264,265
SCA27	609307	13q13.1	<i>FGF14</i>	MM	266
SCA28	610246	18p11.21	<i>AFG3L2</i>	MM	267,268
SCA29	117360	3p26.1	<i>ITPR1</i>	MM	106,269
SCA31	117210	16q22	<i>BEAN</i>	TGGAA repeat (non-coding)	270,271
SCA34	133190	6q14	<i>ELOVL4</i>	MM	272
SCA35	613908	20p13	<i>TGM6</i>	MM	273
SCA36	614153	20p13	<i>NOP56</i>	non-coding repeat	274
SCA38	615957	6p12	<i>ELOVL5</i>	MM	275
SCA40	616053	14q32.12	<i>CCDC88C</i>	MM	276
SCA41	616410	4q27	<i>TRPC3</i>	MM	89
SCA42	616795	17q21.33	<i>CACNA1G</i>	MM	100

MM= missense mutation

However, the majority of dystonia patients do not respond to L-DOPA treatment, indicating that there are other biological pathways involved. At first, no main biological pathway was recognized for SCA, as the functions of *ATXN1* and *ATXN3* had not then been determined. Now, however, the main pathway involved has been identified as altered synaptic transmission due to mutations in *KCNC3*, *KCND3*, *CACNA1A*, *ITPR1*, *TRPC3* and *PDYN*. These encode various ion channels, receptors and a neuropeptide precursor,

affecting potassium (*KCNC3* and *KCND3*), calcium (*CACNA1A*, *ITPR1* and *TRPC3*) and opioid receptor signaling (*PDYN*). See reviews of the genetics of dystonia and SCA by Matilla-Dueñas, Storey and Phil, Balint and Bhatia, and Xiao et al.^{93–96} for further details.

Table 2. Known dystonia genes

Dystonia type	OMIM Number	Locus	Gene Name	Inheritance pattern	References
DYT1	128100	9q34.11	<i>TOR1A</i>	AD	277,278
DYT2	224500	1p35.1	<i>HPCA</i>	AR	279
DYT3	314250	Xq13.1	<i>TAF1</i>	XLR	280,281
DYT4	128101	19p13.3	<i>TUBB4A</i>	AD	282
DYT5	128230	14q22.2	<i>GCH1</i>	AD	90,283
DYT6	602629	8p11.21	<i>THAP1</i>	AD	284–286
DYT8	118800	2q35	<i>PNKD</i>	AD	287,288
DYT9	601042	1p34.2	<i>SLC2A1</i>	AD	289,290
DYT10	128200	16p11.2	<i>PRRT2</i>	AD	291–293
DYT11	159900	7q21.3	<i>SGCE</i>	AD	294,295
DYT12	128235	19q13.2	<i>ATP1A3</i>	AD	296,297
DYT16	612067	2q31.2	<i>PRKRA</i>	AR	298
DYT18	61216	1p34.2	<i>SLC2A1</i>	AR	299
DYT23	614860	9q34.3	<i>CACNA1B</i>	AD	99
DYT24	615034	11p14.2	<i>ANO3</i>	AD	233
DYT25	615073	18p11	<i>GNAL</i>	AD	178
DYT26	616398	22q12.3	<i>KCTD17</i>	AD	300
DYT27	616411	2q37.3	<i>COL6A3</i>	AR	88
	612716	2q13.2	<i>SPR</i>	AR	301
	605407	11p.15.5	<i>TH</i>	AR	91
		9q34.11	<i>CIZ1</i>	AD	302

AD = Autosomal dominant; AR = Autosomal recessive; XLR = X-linked recessive.

Investigating the shared genetic background

For this review we brought together dystonia and SCA genes to identify the shared genetic components and expose possible common underlying biological pathways. Our focus lies on dominantly inherited SCAs and dystonias and we have therefore used 16 autosomal dominant dystonia genes and 28 SCA genes, including polyQ-SCA genes, as input seeds for further analysis (Table 1 and Table 2). To gain insight into the common biological pathways we used GeneNetwork (<http://129.125.135.180:8080/GeneNetwork/cytoscape.html>), a co-expression tool based on approximately 80,000 microarrays from Gene Expression Omnibus,⁹⁷ and PANTHER software to analyze gene ontologies.⁹⁸ Based on the assumption that co-expressed genes are more likely to be involved in

similar biological pathways, this guilt-by-association approach enabled us to assess which genes generally tend to be activated simultaneously, and are thus under similar transcriptional regulation with known autosomal dominant dystonia and ataxia genes. By applying independent GeneNetwork analyses for both known autosomal dominant dystonia and ataxia genes (Supp. Figures 1 and 2) and with manual assessment of the overlap in genes between these gene networks, we identified 99 genes shared by the two disorders (Figure 2). No data was filtered out of the analysis. In contrast, when we




Figure 2. Shared genes between SCA and dystonia

We found 99 genes are shared between the SCA and dystonia gene co-expression networks. Two main pathways were identified that involved synaptic transmission (39 genes) and nervous system development (38 genes).

compared either the dystonia gene network or the ataxia gene network with a “control” gene network based on 26 known autosomal dominant Charcot-Marie-Tooth (CMT) genes, we did not find any significant overlap ($p = 0.2153$ for CMT/SCA and $p = 0.1222$ for CMT/Dystonia), thereby validating our approach. Additionally, six of the 99 genes shared between dystonia and SCA were known dystonia or ataxia genes (*ATP1A3*, *ANO3*, *GNAL*, *SPTBN2*, *PRKCG*, and *PDYN*) and their presence further validates our hypothesis that shared co-expression networks likely point towards shared pathogenesis. Using PANTHER, we were able to categorize most of the 99 shared genes into two major neurological pathways: synaptic transmission and nervous system development (Figure 2). We observed a 14-fold enrichment ($P = 8.39E-31$) for genes playing a role in synaptic transmission (39/99 genes; see Figure 2) and a 3.8-fold enrichment ($P = 8.96E-10$) for genes involved in nervous system development (38/99 genes). However, not all genes fit into these two categories (see Figure 2), indicating that there are other biological pathways/mechanisms playing a role in the molecular pathology of ataxia and dystonia. In the next section, we focus on these two major neurological pathways and highlight their role in disease pathogenesis.

Genes involved in synaptic transmission

Ion channels

We observed that 40 of the shared genes identified by our network approach were involved in synaptic transmission, a finding that was validated by recent studies showing that both dystonia and ataxia are characterized by altered excitability of neurons.^{99–101} One-third of the shared genes identified as involved in synaptic transmission encode ion channel subunits that regulate neuronal potassium, calcium or chloride signaling. Notably, 13.1% of the shared genes encode ion channels compared to 2% of all genes in the complete exome ($P < 0.0001$), further emphasizing the importance of ion channels in the etiology ataxia and dystonia. In contrast, no ion channels were present among the genes that were shared between CMT and SCA and between CMT and dystonia (data not shown).

The role of ion channels in the pathophysiology of both dystonia and SCA is underscored by the identification of human mutations leading to these diseases in *CACNA1B* (DYT23; voltage-gated calcium channel Cav2.2), *KCNC3* (SCA13; voltage-gated potassium channel Kv3.3), *KCND3* (SCA19; voltage-gated potassium channel Kv4.3), *CACNA1A* (SCA6; voltage-gated calcium channel Cav2.1), *CACNA1G* (SCA42; voltage-gated calcium channel Cav3.1), *ITPR1* (SCA15/16; Type 1 Inositol 1,4,5-Trisphosphate Receptor) and *TRPC3* (SCA41; Transient Receptor Potential Cation Channel, Subfamily C, Member 3).^{78,89,99,102–106} Additionally, various ion channel mutations in mouse models lead

to dystonia and/or ataxia including *cacna1a* knockout and *cacna1a* mutants, such as leaner mouse with generalized dystonia/ataxia, tottering mouse displaying ataxia plus paroxysmal generalized dystonia and rocker mouse exhibiting ataxia with paroxysmal focal dystonia. In contrast, lethargic *cacna1b* mutant mouse models appear to have ataxia with paroxysmal exertional dystonia.^{107,108} A putative role for calcium channels in human dystonia was proposed based on the fact that mice treated with an L-type calcium channel agonist developed a phenotype resembling generalized dystonia.¹⁰⁹ Furthermore, in addition to ataxia, posture dystonia was observed in the opisthotonos mouse, which carries a spontaneous mutation in *itpr1* (*ITPR1* is the human SCA15/16 gene) that codes for the type 1 inositol 1,4,5-trisphosphate receptor, which is involved in calcium release from the endoplasmic reticulum.¹¹⁰ While many ion channels are associated with various forms of epilepsy, this topic lies outside the scope of this review and will not be described in further detail here. In the following paragraphs, we discuss the evidence for a role of the shared ion channel genes in the pathogenesis of dystonia and ataxia based on both human and mouse data and according to the pathway in which these genes function.

The shared genes shown to be involved in potassium signaling are *KCHIP2* (Kv channel interacting protein 2), *KCNAB1* (regulatory beta subunit Kv β 1), and *KCNJ4* and *KCNJ6* (inwardly rectifying potassium channels Kir2.3 and Kir3.2/GIRK2, respectively). *KCHIP2* has been implicated in the pathogenesis of SCA19¹¹¹ and mutations in *KCNA1* encoding Kv1.1, the alpha subunit of Kv β 1, cause episodic ataxia (EA1) in humans.¹¹² In mouse striatum, Kir2.3 is specifically located in matrix compartments, which may primarily influence motor circuits within basal ganglia, suggesting it has a role in dystonia.¹¹³ A homozygous point mutation in the pore region of *Girk2* (Gly156Ser) causes the weaver phenotype in mice, which is characterized by cerebellar granule cell death, dopaminergic neuronal cell death in substantia nigra, severe ataxia and spontaneous seizures. In contrast, heterozygous mutant mice showed a thinner granule cell layer and a disorganized PC layer, but no motor phenotype.¹¹⁴ Furthermore, GIRK2 modulates the degree of opioid inhibition upon opioid receptor activation,¹¹⁵ a pathway known to be affected in SCA23.¹¹⁶ GIRK2 has also been shown to be a key determinant of the sensitivity of dopaminergic neurons of the ventral tegmental area to the motor-stimulatory effects of opioids.¹¹⁵ However, its effect in dopaminergic neurons of the substantia nigra is not yet known.

There is increasing evidence that GABAergic signaling is altered in various movement disorders, including dystonia, ataxia, epilepsy and tremor.¹¹⁷ The shared GABA type A receptor subunits *GABRA1*, *GABRA4*, *GABRA5* and *GABRD* identified in our network are ligand-gated chloride channels responsible for the inhibitory effects mediated by the neurotransmitter GABA (γ -aminobutyric acid) in the brain. Multiple cerebellar neurons

use GABA as their output neurotransmitter, including GABAergic inhibitory projection neurons and small GABAergic interneurons in the cerebellar nuclei and Purkinje, Golgi, Lugaro, stellate, basket and candelabrum cells in the cerebellar cortex.¹¹⁸ Additionally, GABAergic projections are present in basal ganglia arising from striatum, globus pallidus and substantia nigra reticula (Figure 1).¹¹⁹ Loss of inhibition is thought to be a crucial causal component in the pathogenesis of dystonia,¹²⁰ and alterations in GABA_A receptors may affect inhibition of action potential firing, thus contributing to disease.¹²¹ Differential expression of GABA_A receptors was observed in a SCA6 mouse model caused by mutations in *Cacna1a*.¹²² In contrast, mice with PC-specific knockdown of *Vgat*, which encodes a vesicular GABA transporter, showed an ataxic phenotype that did not coincide with visible brain malformations at 40 weeks of age. This suggests that GABA release is important for PC functioning, and alterations in PC-specific GABA release cause ataxia.¹²³

Shared genes *CACNG2* and *CACNG3*, which encode voltage-dependent calcium channels γ -subunits 2 and 3, play a role in regulation of glutamate signaling by their function as transmembrane AMPA receptor (AMPA) regulatory proteins (TARPs). The AMPAR-TARP complexes are involved in long-term potentiation and long-term depression, and therefore play a role in memory and motor learning, which has been implicated in various mouse ataxia models.^{124–126} Stargazer mice show distinctive head-tossing and ataxic gait caused by mutant *CACNG2* (TARP- γ 2, stargazin).¹²⁷ In zebrafish, *Cacng2a* was shown to be required for trafficking of the AMPAR to the membrane that mediates normal AMPAR functioning.¹²⁸ The role of glutamate in the pathogenesis of both dystonia and ataxia is further highlighted by a shared gene, *GRIN1*, which encodes the *N*-methyl-D-aspartate (NMDA)-type glutamate receptor subunit, GluN1. GluN1 can interact with the dopamine receptor D1R¹²⁹, and mutations in this gene are likely to affect both NMDA and dopamine signaling, causing non-syndromic intellectual disability and epileptic encephalopathy, encephalopathy with infantile-onset epilepsy, and hyperkinetic and stereotyped movement disorders.^{130,131} Glutamate excitotoxicity has been postulated as a possible disease mechanism for various neurodegenerative disorders, including SCA.¹³² The GRIN1 interaction with DR1 suggests that putative mutations in *GRIN1* or alterations in *GRIN1* expression may lead to aberrant dopamine signaling and subsequently to dystonia.¹³³

Sodium signaling, amongst others mediated by the shared gene *SCN2B* that encodes the auxiliary β 2-subunit of voltage-gated sodium channels, has only recently been implicated in dystonia. A spontaneous mouse mutant carrying a mutation in *scna8* developed a chronic movement disorder with early onset tremor and adult onset dystonia.¹³⁴ Given that Nav1.6 (encoded by *scna8*) is important for the initiation of action potentials, loss of Nav1.6 function might affect the repetitive firing of PCs. Thus, genes

encoding components of sodium signaling could be considered as candidate genes for human movement disorders, including dystonia.

Regulation of neurotransmitters

Our search also identified a group of shared genes involved in the regulation of neurotransmitters, and a clear role for neurotransmitter regulation can be seen in the pathogenesis of dopamine-responsive dystonias. Most of the shared genes are associated with neurotransmitter-containing vesicles and thus play a role in the release of neurotransmitters and control synaptic transmission. Interestingly, the majority of these genes are involved in glutamatergic and GABAergic signaling. Shared genes *SYT5* (synaptotagmin V) and *CADPS2* (Ca²⁺-dependent activator protein for secretion) are both linked to dense-core vesicles. *Syt5* is involved in Ca²⁺-dependent exocytosis in mouse brain,¹³⁵ and *cadps2* is involved in secretion of vesicles containing brain-derived neurotrophic factor (BDNF) and neurotrophin-3 (NT-3) that play a pivotal role in neuronal differentiation and survival.¹³⁶ BDNF goes on to induce presynaptic glutamate release, increase NMDA-receptor density and increase the dendritic complexity of the GABAergic neurons.^{137,138} Furthermore, *cadps2* knockout mice showed deficits in cerebellar development accompanied by aberrant motor coordination and eye movement.¹³⁶ Solute carrier family 17, member 7, encoded by shared gene *SLC17A7* (*VGLUT1*), is a vesicular glutamate transporter necessary for filling glutamate vesicles that thereby influences synaptic transmission efficiency.¹³⁹ *VGLUT1*-deficient mice appeared normal until postnatal week two, after which they started to lag in development and differ in movement and behavior, with death occurring between postnatal week P18 and P21.¹⁴⁰

Among the shared genes identified in our search, there are also components of the SNARE-complex involved in exocytosis of neurotransmitter vesicles. We identified *STX1B* (syntaxin-1B) and vesicle-associated membrane proteins (*VAMP1*/synaptobrevin1 and *VAMP2*/synaptobrevin2) in our network. Decreased spontaneous GABAergic transmission frequencies were detected in *STX1B* knockout cerebellar cultures,¹⁴¹ whereas mutations in *VAMP1* have been identified as causing dominant hereditary spastic ataxia (SPAX1).¹⁴² Moreover, *VAMP1*-deficient mice (lethal-wasting (*lew*)) showed premature death at P15 and were found to have neurological deficits, including profound motor impairments that were most likely caused by reduced functioning of neuromuscular junctions.^{143,144} *VAMP2*-deficient mice demonstrated a critical role for *VAMP2* in Ca²⁺-dependent vesicle exocytosis¹⁴⁵ and in postsynaptic insertion of GluA1-containing AMPA receptors into the synaptic plasma membrane.¹⁴⁶ However, an association with a human disorder has yet to be found. Another shared gene is *SYN1* (synapsin I), which encodes a synaptic-vesicle-associated phosphoprotein involved in organization of the vesicles at the presynaptic terminal and in axonal development.^{147,148}

The only gene in this shared group not specifically linked to vesicles is histamine receptor H3 (*H3R*), which is selectively expressed in brain¹⁴⁹ and present on presynaptic terminals, where it acts as an autoreceptor inhibiting histamine production and release. H3R is also present on postsynaptic nerve terminals where it functions as a heteroreceptor that inhibits the release of other neurotransmitters such as norepinephrine, serotonin, GABA, acetylcholine, glutamate and dopamine.¹⁵⁰ While no mutations in *H3R* have been reported, *hrh3*-knockout mice showed reduced locomotor activity, indicating that *H3R* is a plausible candidate gene for movement disorders.^{151,152}

Synapse organization

The third group of shared genes that emerged from our network analysis plays a role in the organization of synapses and encodes scaffolding proteins necessary for proper synapse formation and vesicular structures. On the postsynaptic site, we identified *DLG4* (Disc large 4, or postsynaptic density protein 95; *PSD95*) and *CLSTN3* (calsynenin-3/alcadein- β) as shared genes between ataxia and dystonia. *DLG4* is involved in clustering of NMDA-receptors, potassium channels and interacting proteins,¹⁵³ some of which some, e.g. *KCNC3*, *KCND3*, and *KCTD17*, are already linked to ataxia or dystonia. Additionally, a *DLG4* knockout mouse model showed enhanced long-term potentiation accompanied by impairments in spatial learning, but no motor phenotype.¹⁵⁴ *CLSTN3* is a cadherin protein that forms a functional complex with α -neurexin and is mainly involved in regulating inhibitory synaptic functions including GABAergic signaling.^{155–157} We also identified dopamine receptor D1 (*DRD1*) in the network, which is a G-protein-coupled receptor for the neurotransmitter dopamine that also regulates NMDA receptor functions.¹⁵⁸ Variants in D1R have been associated to tardive-like dystonia¹⁵⁹ and D1R-knockout mice showed reduced dynorphin expression in the striatum and increased locomotor activity.¹⁶⁰ However, loss of dopaminergic neurons does not necessarily lead to Parkinsonism in SCA2 and SCA3 patients.¹⁶¹ Moreover, *SEZ6* (seizure-related 6 homolog) and *SNCB* (β -synuclein) were identified in our search, and the exact functions of these genes are still unclear. However, *sez-6* knockout mice do display altered dendritic branching, a smaller number of excitatory synapses and defects in excitatory synaptic transmission leading to reduced motor coordination.^{162,163} Bassoon, a presynaptic scaffold protein encoded by *BSN*, is primarily involved in ribbon synapses in retina and cochlear hair cells, and thus shows no direct link to movement disorders.¹⁶⁴

Other shared genes involved in synaptic transmission

We identified 13 other shared genes involved in synaptic transmission that did not match the GO-terms *ion channel*, *neurotransmitter regulation* or *synapse organization*. A number of these genes are involved in intracellular signal transduction. *PRKCG*, for example, encodes the neuron-specific protein kinase C gamma that mediates neu-

ronal signal transduction. *PRKCG* is the causative gene for SCA14, in which dystonic phenotypes, including writer's cramp, myoclonus and focal dystonia are regularly reported.^{165–168} We also identified *MAPK8IP2* (mitogen-activated protein kinase 8-interacting protein 2/JNK-interacting protein 2 (JIP2)), a synaptic scaffolding protein involved in the JNK-signaling cascade.¹⁶⁹ Jip2 knockout mice showed sensorimotor deficits and impaired social interaction due to changes in the morphology of dendritic arbors of PCs associating with aberrant NMDA- and AMPA-receptor-mediated glutamatergic signaling.¹⁷⁰ Notably, Jip1/Jip2 double knockouts were severely ataxic,¹⁷¹ demonstrating a role for JIP2 in the underlying pathology of cerebellar ataxias.

Two other shared genes are directly implicated in dystonia: *ADCY5* encoding adenylyl cyclase 5¹⁷² and *GNAL* encoding stimulatory G-protein α -subunit Gaorf.¹⁷³ Mutations in *ADCY5* have been found to cause familial dyskinesia with facial myokymia (FDFM)^{174,175} and (benign) chorea with dystonia,^{176,177} while mutations in *GNAL* underlie adult-onset cervical and segmental dystonia (DYT25).¹⁷⁸ Additionally, *RASD2* (RASD family, member 2/tumor endothelial marker 2 (TEM2)/Ras homolog enriched in striatum (RHES)) encodes a small GTPase that plays a role in dopaminergic signaling^{179,180} and enhances L-dopa induced dyskinesia via activation of mTOR,¹⁸¹ giving it a putative role in dystonia.

Three other shared genes, *SLC12A5*, *HTR2C* and *KIF5A*, are involved in regulation of receptor functioning, thereby mediating post-synaptic neurotransmission. *SLC12A5* (solute carrier family 12, member 5/potassium-chloride cotransporter 2 (KCC2)) encodes a neuron-specific ion transporter that is the main extruder of the intracellular chloride used to create the chloride gradient necessary for GABA- and glycine-receptor functioning, and which ultimately promotes post-synaptic inhibition.^{182,183} *KIF5A* encodes a neuron-specific kinesin heavy chain protein involved in microtubule-based cargo transport of, among others, the GABA_A-receptor.^{184,185}

For the remainder of the genes in this group, brain functions are not yet completely established, but some, including *GAD2*, *PDYN*, *MINK1* and *CPNE6*, seem to link to GABA neurotransmitter synthesis and synaptic plasticity.^{186–189} Of this list, only *PDYN*, the precursor for the neuropeptides α -neoendorphin, dynorphin A (Dyn A) and dynorphin B (Dyn B), is directly linked to ataxia. Notably, dynorphins are involved in pain sensing, addiction and depression via opioid signaling,¹⁹⁰ while mutations in *PDYN* have been shown to cause SCA23 and suggestively affect glutamate signaling.¹¹⁶ Lastly, *CDH8* mediates Ca²⁺-dependent cell-cell adhesion involved in presynaptic organization and synaptic remodeling.¹⁹¹ *Cdh8* knockdown in rat primary neurons led to aberrant dendritic arborization and decreased self-avoidance of dendritic branches.¹⁹²

Genes involved in neurodevelopment

Beyond the large group of genes involved in synaptic transmission, we identified many shared genes involved in development of the nervous system. Indeed, many disorders that combine dystonia with other neurological features are developmental.² Neurodevelopmental abnormalities have already been linked to DYT1 dystonia and include alterations in the cerebello-thalamo-cortical pathways.¹⁹³ Recently, cerebellar neurogenesis was found to be compromised in mouse models of DYT1 dystonia,¹⁹⁴ demonstrating that loss of Tor1A induces important developmental alterations in the cerebellum, and thereby contributing to the development of dystonia in *DYT1* mutation carriers. To date, no clear evidence has been reported for alterations in brain development in the pathology of dominant hereditary ataxias. However, developmental delay has been reported for SCA27, SCA2, SCA5 and SCA13.^{102,195–197} In SCA13, differences in axonal pathfinding were observed in zebrafish motor neurons expressing mutant F448L-KCNC3,¹⁹⁸ which is also indicative of aberrant neurogenesis. More evidence that abnormalities in cerebellum development and deficits in axon elongation may give rise to ataxia came from a mice lacking neuron navigator 2 (Nav2).¹⁹⁹ These mice exhibited a small cerebellum at E17.5 and ataxia at age 5 months due to aberrant migration of post-mitotic granule cells. Given that all these examples showed a young age of onset or juvenile ataxia, in contrast to the late age of onset found in the majority of SCAs, it seems difficult to reconcile them with playing a role in neurodevelopment in disorders with a late age of onset. However, some preliminary evidence has been reported for a role of neurodevelopment in Alzheimer's disease and Parkinson's disease.^{200–202} Therefore, we will now address the overlapping ataxia-dystonia network genes in the context of their putative role in the central nervous system (CNS) development, including neurogenesis and brain/cerebellum development.

Neurogenesis

The shared genes involved in this group are implicated in axon path finding, regulation of dendrite/axon extension, or dendrite morphogenesis. Many of these genes (*GABRA5*, *GRIN1*, *SEZ6*, *ATP2B2*, *BCL11B*, *CDK5R2*, *CEND1* and *SPTBN2*) were also listed in the group for brain/cerebellum development and these will be discussed in the next section. Additionally, some genes that are involved in neurogenesis and/or neurodevelopment (*GABRA5*, *GRIN1*, *SEZ6*, *KCNIP2*, *DLG4 (PSD-95)*, *CPNE6*, *KIF5A*, *MAPK8IP2* and *MINK1*) are also involved in synaptic transmission and have already been discussed in the *Synaptic transmission* section.

Of the genes involved in neurogenesis, Copine1, which is encoded by *CPNE9*, regulates neurite outgrowth and differentiation of hippocampal progenitor cells.²⁰³ In contrast, *LCAM1* and *NCDN*, which encode cell adhesion molecule L1 and neurochondrin,

respectively, modify neurite growth either in cerebellar neurons²⁰⁴ or PC12 cells.²⁰⁵ The paired-like homeobox domain protein Phox2b is a key determinant of neuronal identity, as it is required for the initial phase in neuronal differentiation, giving rise to noradrenergic neurons that may be involved in the pathology of dystonia.^{206,207} For *PSD*, *RSG14* and *SCRT1*, not much is known about their role in neurogenesis. However, the relation of *PSD* to *DLG4* (see section on *Synaptic transmission*), and the fact that *RSG14* and *SCRT1* are highly expressed during early postnatal development,^{208,209} suggest a role for these genes in brain development.

Brain/cerebellum development

Data highlighting GABAergic signaling as a shared pathway in movement disorders including dystonia, ataxia, epilepsy and tremor, was mentioned in the *Synaptic transmission* section, but GABAergic signaling also plays a role in brain development. Cerebellar maturation was reported to depend on modulation of the pool of GABAergic interneurons via exogenous Sonic hedgehog.²¹⁰ For example, direct neuronal migrations of cortical interneurons expressing GABA_B receptors are regulated by Gai-Go-coupled receptor signaling in which the G-protein-regulator of neurite outgrowth 1, encoded by shared gene *GRIN1*, seemingly participates.^{211,212} *In situ* data showed that GRIN1 was expressed on a migrating subpopulation of neurons from the caudal rhombic lip to the precerebellar nuclei of the brainstem, and moderate GRIN1 expression was seen in the cerebellum, suggesting it has a role in cerebellum development.^{212,213} *NEUROD6*, a member of the basic helix-loop-helix transcription factors is involved in neuronal differentiation of GABAergic and glycinergic interneurons.²¹⁴

The role of the plasma membrane Ca²⁺-Pump 2 encoded by shared gene *ATP2B2* in cerebellar development was established by studies of cerebellar organotypic slice cultures. Here, inhibition of the Ca²⁺-pump resulted in a reduction of the PC dendritic tree, confirming the importance of calcium signaling in controlling dendritic growth.²¹⁵ Mutations in *ATP2B3*, which is also expressed in the cerebellum, led to X-linked congenital cerebellar ataxia,²¹⁶ further emphasizing the importance of these pumps in the pathology of cerebellar ataxias. Additionally, cell-type-restricted and time-dependent expression of the neuron-specific calcium-binding protein hippocalcin, which is encoded by shared gene *HPCA* and underlies *DYT2*, was observed in the developing brain; this suggests that hippocalcin plays a role in neuronal differentiation in the early stages of development.²¹⁷ Furthermore, mutations in overlapping gene *SPTBN2* cause SCA5,²¹⁸ and mutant B-III-spectrin led to mislocalization and dysfunction of mGluR1 α at dendritic spines.¹²⁶ These findings further connect glutamatergic dysfunction and aberrant calcium homeostasis to brain/cerebellar development. Potassium signaling may also play a role in the proper development of the CNS, as differential regulation of

distinct potassium channel beta subunits, including Kvbeta 1 (*KCNAB1*), was observed to mediate neuronal survival and maturation.²¹⁹

Neuronal-specific gene Cell Cycle Exit And Neuronal Differentiation 1 (CEND1) is required for normal cerebellar development. Mice lacking Cend1 showed impaired cerebellar development and motor dysfunction due to an increased proliferation of granule cell precursors, delayed granule cell migration, and alterations in PC differentiation.²²⁰ In contrast, *CDK5R2*, which encodes the neuronal-specific activator of CDK5 kinase, indirectly regulates dendrite development and axon guidance by regulating gene expression and phosphorylation of key targets such as Connexin 43 and PIPKI gamma 90 during brain development.^{221–223} However, no direct role for Tau Tubulin Kinase 1, TTBK1, in brain development has yet been reported.

Other genes not tagged as involved in neurogenesis or brain cerebellum development

In this last group, ten genes were identified that did not fit the GO terms *neurogenesis* and/or *brain-cerebellum development*. We have already discussed *GABRA4*, *SCN2B*, *BSN* and *PRKCG*, which have been previously linked to synaptic transmission, but apparently they also play a role in CNS development. Of the remaining genes, not much is known about their respective functions or role in brain development. However, calcium signaling again plays a role, as *CLSTN3*, *HPCAL4*, *PCP4* and *Ly6h* are all seemingly involved in mediating intracellular calcium homeostasis.^{156,224,225}

Shared gene *ACTL6B* is part of a neuronal-specific chromatin-remodeling complex that mediates long-term memory formation via the coordinated regulation of gene expression.²²⁶ This complex also contains the rBAF factor that regulates the expression of genes involved in the dendritic development required for synaptic plasticity and memory. Deficits in plasticity were also already reported in *DYT1* dystonia mice and various ataxia mouse models.^{125,227–229}

Shared gene *RGS9*, which encodes a regulator of G-protein signaling, was found to have a restricted expression in the basal ganglia and suggestively plays a role in local signaling pathways that include mu-opioid receptor signaling.^{230,231} Aberrant opioid signaling was reported in a hamster model of paroxysmal dystonia²³², and mutations in the precursor protein of the opioid dynorphin peptides, prodynorphin, cause SCA23.¹¹⁶ However, no clear evidence has been published with regard to its postulated role in CNS development. Further studies are required to establish a role for these shared genes in neurodevelopment.

Discussion

By combining gene co-expression networks for dystonia and SCA, we have identified a set of 99 shared genes that may play a role in the pathogenesis of both disorders. These results further validate the clinical observations of co-morbidity of ataxia and dystonia and show a role for the cerebellum in dystonia. Additionally, a “control” network based on known CMT genes showed no significant overlap with dystonia or SCA gene network, confirming that dystonia and SCA are biologically closer related. The key shared molecular pathways between ataxia and dystonia were synaptic transmission and neurodevelopment, with the involvement of synaptic transmission in both disorders well recognized, as mutations in many ion channels have already been shown to cause these diseases.^{78,99,100,102–104,233} On the other hand, molecular pathways related to neurodevelopment had not been as clearly linked to ataxia and dystonia, although a recent review mentioned neurodevelopment as a common theme in dystonia,²³⁴ and developmental defects have been described in DYT1¹⁹³ and developmental delay is known in SCA27, SCA13, SCA5 and SCA2.^{102,195–197} Moreover, genes involved in neurodevelopment may not exert their function solely during development of the CNS because they also appear to play a role in Alzheimer’s disease and Parkinson’s disease,^{200–202} thus providing a link between development and degeneration. Identifying common pathways may aid the development of therapies because restoring a pathway will help patients with different genetic backgrounds.

Within both major pathways, the importance of glutamatergic-, and especially GABAergic signaling was strikingly demonstrated by the identification of four GABA receptor genes in the overlapping gene set. While mutations in these genes are not currently known to cause SCA or dystonia, they have been linked to epilepsy,^{121,235,236} making them potential candidate genes for movement disorders. Furthermore, we also identified a number of genes that indirectly affect GABAergic signaling, including *CADPS2*, *STX1B*, *H3R*, *SLC12A5*, *KIF5A*, *GAD2* and *GRIN1*. We therefore suggest prioritizing genes involved in GABAergic signaling in genetic studies of SCA and dystonia patients, as these genes are likely candidate genes in genetically unexplained cases. This is particularly important because GABAergic signaling can be targeted by drugs; this is a treatment approach used in multiple neuropsychiatric disorders²³⁷ that has also been proven to relieve rotenone-induced Parkinsonism-like symptoms.²³⁸

There were 37 genes that did not fit within the GO-terms *synaptic transmission* or *nervous system development*. This does not, however, mean that these genes are not involved in these pathways, but rather that their functions may be unknown or that they might have more than one function, in both cases this would contribute to their lack of representation in the GO annotations. Alternately, the existence of this group of genes may indicate the involvement of other molecular pathways, in addition to the

two major ones we have described here. Other pathways have certainly been described as triggering SCA and/or dystonia pathology and include transcriptional dysregulation, mitochondrial defects, autophagy, dopaminergic signaling, ER stress and alterations in calcium homeostasis.^{8,239}

Further unraveling of the common pathways in SCA and dystonia will improve our understanding of disease pathologies and may open the way to novel therapies by identifying new drug targets. Furthermore, our sophisticated network approach strongly suggests that the molecular pathways of ataxia and dystonia are indeed closely related. Both should be tested for simultaneously in genetic diagnostics work on ataxia and dystonia patients. As up to 100 genes need to be screened, this will be easier to achieve with the implementation of disease-focused gene panels or dedicated exome strategies. Genes that have been identified in the gene set that overlaps between ataxia and dystonia should be prioritized when searching for disease gene mutations in patients without a genetic diagnosis, because variants in these genes are more likely to be involved in their disease pathology. Overall, this gene co-expression network provides useful insights into the possible shared molecular mechanisms of ataxia and dystonia and it will lead to a better understanding of their pathogenesis.

Acknowledgements and funding

We thank Kate Mc Intyre and Jackie Senior for improving the manuscript. This work was funded by a Rosalind Franklin Fellowship from by the University of Groningen awarded to DSV and by grants to MAJT from the Fonds Nuts-Ohra, Prinses Beatrix Fonds, Gossweiler Foundation, Stichting wetenschapsfonds dystonie vereniging, Fonds Psychische Gezondheid, Phelps Stichting, and the Beatrix Kinderziekenhuis Fonds. Further funding came from unrestricted grants from Actelion, Merz, Ipsen, Allergan Farmaceutics and Medtronic and MAJT received an honorarium from the Merz expert meeting in Paris, January 2016 . HAJ is supported in part by grants to the Dystonia Coalition from the Office of Rare Diseases Research in the USA National Center for Advancing Translational Studies (TR001456) and the NIH National Institute for Neurological Disorders and Stroke (NS067501). TjDK received grants from the Metabolic Power Fundation and Metakids foundation and Ride4Kids foundation (all non-profit) for studying movement disorders in metabolic diseases. He received research grants from Actelion pharmaceuticals (profit) for studying movement disorders in NP-C disease and received an honorarium for presenting at a sponsored meeting on NP-C. None of the funding obtained relates to the work presented here.

References

1. Durr, A. (2010). Autosomal dominant cerebellar ataxias: polyglutamine expansions and beyond. *Lancet. Neurol.* *9*, 885–894.
2. Fung, V.S.C., Jinnah, H.A., Bhatia, K., and Vidailhet, M. (2013). Assessment of patients with isolated or combined dystonia: An update on dystonia syndromes. *Mov. Disord.* *28*, 889–898.
3. van Gaalen, J., Giunti, P., and van de Warrenburg, B.P. (2011). Movement disorders in spinocerebellar ataxias. *Mov. Disord.* *26*, 792–800.
4. Neychev, V.K., Gross, R.E., Lehéricy, S., Hess, E.J., and Jinnah, H.A. (2011). The functional neuroanatomy of dystonia. *Neurobiol. Dis.* *42*, 185–201.
5. Prudente, C.N., Hess, E.J., and Jinnah, H.A. (2014). Dystonia as a network disorder: what is the role of the cerebellum? *Neuroscience* *260*, 23–35.
6. Kuoppamaki, M., Giunti, P., Quinn, N., Wood, N., and Bhatia, K. (2003). Slowly progressive cerebellar ataxia and cervical dystonia: Clinical presentation of a new form of spinocerebellar ataxia? *18*, 200–206.
7. van de Warrenburg, B.P.C., Giunti, P., Schneider, S.A., Quinn, N.P., Wood, N.W., and Bhatia, K.P. (2007). The syndrome of (predominantly cervical) dystonia and cerebellar ataxia: new cases indicate a distinct but heterogeneous entity. *J. Neurol. Neurosurg. Psychiatry* *78*, 774–775.
8. Matilla-Dueñas, A., Ashizawa, T., Brice, A., Magri, S., McFarland, K.N., Pandolfo, M., Pulst, S.M., Riess, O., Rubinsztein, D.C., Schmidt, J., et al. (2014). Consensus Paper: Pathological Mechanisms Underlying Neurodegeneration in Spinocerebellar Ataxias. *The Cerebellum* *13*, 269–302.
9. Hallett, M. (2011). Neurophysiology of dystonia: The role of inhibition. *Neurobiol. Dis.* *42*, 177–184.
10. Shadmehr, R., and Krakauer, J.W. (2008). A computational neuroanatomy for motor control. *Exp. Brain Res.* *185*, 359–381.
11. Batten, F.E. (1903). On the diagnostic value of the position of the head in cases of cerebellar disease. *Brain* *26*, 71–80.
12. Grey, E.G. (1916). Studies on the Localization of Cerebellar Tumors: the Position of the Head and Suboccipital Discomforts. *Ann. Surg.* *63*, 129–139.
13. Krauss, J., Seeger, W., and Jankovic, J. (1997). Cervical dystonia associated with tumors of the posterior fossa. *Mov. Disord.* *12*, 443–447.
14. Extremera, V.C., Alvarez-Coca, J., Rodríguez, G.A., Pérez, J.M., de Villanueva, J.L.R., and Díaz, C.P. (2008). Torticollis is a usual symptom in posterior fossa tumors. *Eur. J. Pediatr.* *167*, 249–250.
15. LeDoux, M.S., and Brady, K.A. (2003). Secondary cervical dystonia associated with structural lesions of the central nervous system. *Mov. Disord.* *18*, 60–69.
16. Batla, A., Sánchez, M.C., Erro, R., Ganos, C., Stamelou, M., Balint, B., Brugger, F., Antelmi, E., and Bhatia, K.P. (2015). The role of cerebellum in patients with late onset cervical/segmental dystonia?—Evidence from the clinic. *Parkinsonism Relat. Disord.*
17. Khooshnoodi, M.A., Factor, S.A., and Jinnah, H.A. (2013). Secondary blepharospasm associated with structural lesions of the brain. *J. Neurol. Sci.* *331*, 98–101.
18. Rumbach, L., Barth, P., Costaz, A., and Mas, J. (1995). Hemidystonia consequent upon ipsilateral vertebral artery occlusion and cerebellar infarction. *Mov. Disord.* *10*, 522–525.

19. O'Rourke, K., O'Riordan, S., Gallagher, J., and Hutchinson, M. (2006). Paroxysmal torticollis and blepharospasm following bilateral cerebellar infarction. *J. Neurol.* 253, 1644–1645.
20. Akin, A., Yilmaz, R., Selcuk, F., and Akbostanci, M.C. (2014). Sudden onset of oromandibular dystonia after cerebellar stroke. *Tremor Other Hyperkinet. Mov. (N.Y.)* 4, 262.
21. Waln, O., and LeDoux, M.S. (2010). Delayed-onset oromandibular dystonia after a cerebellar hemorrhagic stroke. *Parkinsonism Relat. Disord.* 16, 623–625.
22. Alarcón, F., Tolosa, E., and Muñoz, E. (2001). Focal limb dystonia in a patient with a cerebellar mass. *Arch. Neurol.* 58, 1125–1127.
23. Prudente, C.N., Pardo, C.A., Xiao, J., Hanfelt, J., Hess, E.J., LeDoux, M.S., and Jinnah, H.A. (2013). Neuropathology of cervical dystonia. *Exp. Neurol.* 247, 95–104.
24. Zoons, E., and Tijssen, M. a J. (2013). Pathologic changes in the brain in cervical dystonia pre- and post-mortem - a commentary with a special focus on the cerebellum. *Exp. Neurol.* 247, 130–133.
25. Fabbrini, G., Pantano, P., Totaro, P., Calistri, V., Colosimo, C., Carmellini, M., Defazio, G., and Berardelli, A. (2008). Diffusion tensor imaging in patients with primary cervical dystonia and in patients with blepharospasm. *Eur. J. Neurol.* 15, 185–189.
26. Delmaire, C., Vidailhet, M., Wassermann, D., Descoteaux, M., Valabregue, R., Bourdain, F., Lenglet, C., Sangla, S., Terrier, A., Deriche, R., et al. (2009). Diffusion abnormalities in the primary sensorimotor pathways in writer's cramp. *Arch. Neurol.* 66, 502–508.
27. Simonyan, K., Tovar-Moll, F., Ostuni, J., Hallett, M., Kalasinsky, V.F., Lewin-Smith, M.R., Rushing, E.J., Vortmeyer, A.O., and Ludlow, C.L. (2008). Focal white matter changes in spasmodic dysphonia: a combined diffusion tensor imaging and neuropathological study. *Brain* 131, 447–459.
28. Niethammer, M., Carbon, M., Argyelan, M., and Eidelberg, D. (2011). Hereditary dystonia as a neurodevelopmental circuit disorder: Evidence from neuroimaging. *Neurobiol. Dis.* 42, 202–209.
29. Prell, T., Peschel, T., Köhler, B., Bokemeyer, M.H., Dengler, R., Günther, A., and Grosskreutz, J. (2013). Structural brain abnormalities in cervical dystonia. *BMC Neurosci.* 14, 123.
30. Ramdhani, R.A., Kumar, V., Velickovic, M., Frucht, S.J., Tagliati, M., and Simonyan, K. (2014). What's special about task in dystonia? A voxel-based morphometry and diffusion weighted imaging study. *Mov. Disord.* 29, 1141–1150.
31. Yang, J., Luo, C., Song, W., Guo, X., Zhao, B., Chen, X., Huang, X., Gong, Q., and Shang, H.-F. (2014). Diffusion tensor imaging in blepharospasm and blepharospasm-oromandibular dystonia. *J. Neurol.* 261, 1413–1424.
32. Sako, W., Fujita, K., Vo, A., Rucker, J.C., Rizzo, J.-R., Niethammer, M., Carbon, M., Bressman, S.B., Uluğ, A.M., and Eidelberg, D. (2015). The visual perception of natural motion: abnormal task-related neural activity in DYT1 dystonia. *Brain* 138, 3598–3609.
33. Zoons, E., Booij, J., Nederveen, A.J., Dijk, J.M., and Tijssen, M.A.J. (2011). Structural, functional and molecular imaging of the brain in primary focal dystonia--a review. *Neuroimage* 56, 1011–1020.
34. Carbon, M., Kingsley, P.B., Tang, C., Bressman, S., and Eidelberg, D. (2008). Microstructural white matter changes in primary torsion dystonia. *Mov. Disord.* 23, 234–239.
35. Carbon, M., Kingsley, P.B., Su, S., Smith, G.S., Spetsieris, P., Bressman, S., and Eidelberg, D. (2004). Microstructural white matter changes in carriers of the DYT1 gene mutation. *Ann. Neurol.* 56, 283–286.
36. van der Meer, J.N., Beukers, R.J., van der Salm, S.M.A., Caan, M.W.A., Tijssen, M.A.J., and Nederveen, A.J. (2012). White matter abnormalities in gene-positive myoclonus-dystonia. *Mov. Disord.* 27, 1666–1672.

37. Argyelan, M., Carbon, M., Niethammer, M., Ulug, A.M., Voss, H.U., Bressman, S.B., Dhawan, V., and Eidelberg, D. (2009). Cerebellothalamocortical connectivity regulates penetrance in dystonia. *J. Neurosci.* 29, 9740–9747.
38. Eidelberg, D., Moeller, J.R., Antonini, A., Kazumata, K., Nakamura, T., Dhawan, V., Spetsieris, P., deLeon, D., Bressman, S.B., and Fahn, S. (1998). Functional brain networks in DYT1 dystonia. *Ann. Neurol.* 44, 303–312.
39. Obermann, M., Yaldizli, O., De Greiff, A., Lachenmayer, M.L., Buhl, A.R., Tumczak, F., Gizewski, E.R., Diener, H.-C., and Maschke, M. (2007). Morphometric changes of sensorimotor structures in focal dystonia. *Mov. Disord.* 22, 1117–1123.
40. Draganski, B., Thun-Hohenstein, C., Bogdahn, U., Winkler, J., and May, A. (2003). “Motor circuit” gray matter changes in idiopathic cervical dystonia. *Neurology* 61, 1228–1231.
41. Simonyan, K., and Ludlow, C.L. (2012). Abnormal structure-function relationship in spasmodic dysphonia. *Cereb. Cortex* 22, 417–425.
42. Delmaire, C., Vidailhet, M., Elbaz, A., Bourdain, F., Bleton, J.P., Sangla, S., Meunier, S., Terrier, A., and Lehéricy, S. (2007). Structural abnormalities in the cerebellum and sensorimotor circuit in writer’s cramp. *Neurology* 69, 376–380.
43. Piccinin, C.C., Piovesana, L.G., Santos, M.C.A., Guimarães, R.P., De Campos, B.M., Rezende, T.J.R., Campos, L.S., Torres, F.R., Amato-Filho, A.C., França, M.C., et al. (2015). Diffuse decreased gray matter in patients with idiopathic craniocervical dystonia: a voxel-based morphometry study. *Front. Neurol.* 5, 1–11.
44. Wu, C.C., Fairhall, S.L., McNair, N.A., Hamm, J.P., Kirk, I.J., Cunnington, R., Anderson, T., and Lim, V.K. (2010). Impaired sensorimotor integration in focal hand dystonia patients in the absence of symptoms. *J. Neurol. Neurosurg. Psychiatry* 81, 659–665.
45. Hu, X., Wang, L., Liu, H., and Zhang, S. (2006). Functional magnetic resonance imaging study of writer’s cramp. *Chin. Med. J. (Engl.)* 119, 1263–1271.
46. Kadota, H., Nakajima, Y., Miyazaki, M., Sekiguchi, H., Kohno, Y., Amako, M., Arino, H., Nemoto, K., and Sakai, N. (2010). An fMRI study of musicians with focal dystonia during tapping tasks. *J. Neurol.* 257, 1092–1098.
47. Beukers, R.J., Foncke, E.M.J., van der Meer, J.N., Nederveen, A.J., de Ruyter, M.B., Bour, L.J., Veltman, D.J., and Tijssen, M.A.J. (2010). Disorganized sensorimotor integration in mutation-positive myoclonus-dystonia: a functional magnetic resonance imaging study. *Arch. Neurol.* 67, 469–474.
48. Haslinger, B., Erhard, P., Dresel, C., Castrop, F., Roettinger, M., and Ceballos-Baumann, A.O. (2005). “Silent event-related” fMRI reveals reduced sensorimotor activation in laryngeal dystonia. *Neurology* 65, 1562–1569.
49. Simonyan, K., and Ludlow, C.L. (2010). Abnormal activation of the primary somatosensory cortex in spasmodic dysphonia: an fMRI study. *Cereb. Cortex* 20, 2749–2759.
50. Preibisch, C., Berg, D., Hofmann, E., Solymosi, L., and Naumann, M. (2001). Cerebral activation patterns in patients with writer’s cramp: a functional magnetic resonance imaging study. *J. Neurol.* 248, 10–17.
51. Obermann, M., Vollrath, C., de Greiff, A., Gizewski, E.R., Diener, H.-C., Hallett, M., and Maschke, M. (2010). Sensory disinhibition on passive movement in cervical dystonia. *Mov. Disord.* 25, 2627–2633.
52. Talelli, P., Hoffland, B.S., Schneider, S.A., Edwards, M.J., Bhatia, K.P., van de Warrenburg, B.P.C., and Rothwell, J.C. (2011). A distinctive pattern of cortical excitability in patients with the syndrome of dystonia and cerebellar ataxia. *Clin. Neurophysiol.* 122, 1816–1819.

53. Meunier, S., Lourenco, G., Roze, E., Apartis, E., Trocello, J.M., and Vidailhet, M. (2008). Cortical excitability in DYT-11 positive myoclonus dystonia. *Mov. Disord.* 23, 761–764.
54. Quartarone, A., and Pisani, A. (2011). Abnormal plasticity in dystonia: Disruption of synaptic homeostasis. *Neurobiol. Dis.* 42, 162–170.
55. Thompson, R.F., and Steinmetz, J.E. (2009). The role of the cerebellum in classical conditioning of discrete behavioral responses. *Neuroscience* 162, 732–755.
56. Gerwig, M., Kolb, F.P., and Timmann, D. (2007). The involvement of the human cerebellum in eyeblink conditioning. *Cerebellum* 6, 38–57.
57. Sommer, M., Grafman, J., Clark, K., and Hallett, M. (1999). Learning in Parkinson's disease: eyeblink conditioning, declarative learning, and procedural learning. *J. Neurol. Neurosurg. Psychiatry* 67, 27–34.
58. Teo, J.T.H., van de Warrenburg, B.P.C., Schneider, S. a, Rothwell, J.C., and Bhatia, K.P. (2009). Neurophysiological evidence for cerebellar dysfunction in primary focal dystonia. *J. Neurol. Neurosurg. Psychiatry* 80, 80–83.
59. Hoffland, B.S., Kassavetis, P., Bologna, M., Teo, J.T.H., Bhatia, K.P., Rothwell, J.C., Edwards, M.J., and van de Warrenburg, B.P. (2013). Cerebellum-dependent associative learning deficits in primary dystonia are normalized by rTMS and practice. *Eur. J. Neurosci.* 38, 2166–2171.
60. Hoffland, B.S., Veugen, L.C., Janssen, M.M.H.P., Pasman, J.W., Weerdesteyn, V., and van de Warrenburg, B.P. (2014). A gait paradigm reveals different patterns of abnormal cerebellar motor learning in primary focal dystonias. *Cerebellum* 13, 760–766.
61. Hubsch, C., Roze, E., Popa, T., Russo, M., Balachandran, A., Pradeep, S., Mueller, F., Brochard, V., Quartarone, A., Degos, B., et al. (2013). Defective cerebellar control of cortical plasticity in writer's cramp. *Brain* 136, 2050–2062.
62. Sadnicka, A., Teo, J.T., Kojovic, M., Pareés, I., Saifee, T.A., Kassavetis, P., Schwingenschuh, P., Katschnig-Winter, P., Stamelou, M., Mencacci, N.E., et al. (2015). All in the blink of an eye: new insight into cerebellar and brainstem function in DYT1 and DYT6 dystonia. *Eur. J. Neurol.* 22, 762–767.
63. Bradnam, L. V, Graetz, L.J., McDonnell, M.N., and Ridding, M.C. (2015). Anodal transcranial direct current stimulation to the cerebellum improves handwriting and cyclic drawing kinematics in focal hand dystonia. *Front. Hum. Neurosci.* 9, 286.
64. Koch, G., Porcacchia, P., Ponzio, V., Carrillo, F., Cáceres-Redondo, M.T., Brusa, L., Desiato, M.T., Arciprete, F., Di Lorenzo, F., Pisani, A., et al. (2014). Effects of two weeks of cerebellar theta burst stimulation in cervical dystonia patients. *Brain Stimul.* 7, 564–572.
65. Zervas, N.T. (1977). Long-term review of dentatectomy in dystonia musculorum deformans and cerebral palsy. *Acta Neurochir. (Wien)*. 49–51.
66. Davis, R. (2000). Cerebellar stimulation for cerebral palsy spasticity, function, and seizures. *Arch. Med. Res.* 31, 290–299.
67. Teixeira, M.J., Schroeder, H.K., and Lepski, G. (2015). Evaluating cerebellar dentatotomy for the treatment of spasticity with or without dystonia. *Br. J. Neurosurg.* 1–6.
68. Sokal, P., Rudaś, M., Harat, M., Szyłberg, Ł., and Zieliński, P. (2015). Deep anterior cerebellar stimulation reduces symptoms of secondary dystonia in patients with cerebral palsy treated due to spasticity. *Clin. Neurol. Neurosurg.* 135, 62–68.

69. Fletcher, C.F., Lutz, C.M., O'Sullivan, T.N., Shaughnessy, J.D., Hawkes, R., Frankel, W.N., Copeland, N.G., and Jenkins, N.A. (1996). Absence Epilepsy in Tottering Mutant Mice Is Associated with Calcium Channel Defects. *Cell* 87, 607–617.
70. Meier, H., and MacPike, A.D. (1971). Three Syndromes Produced by Two Mutant Genes. *J. Hered.* 62, 297–302.
71. Doyle, J., Ren, X., Lennon, G., and Stubbs, L. (1997). Mutations in the Cacn1a4 calcium channel gene are associated with seizures, cerebellar degeneration, and ataxia in tottering and leaner mutant mice. *Mamm. Genome* 8, 113–120.
72. Herrup, K., and Wilczynski, S.L.L. (1982). Cerebellar cell degeneration in the leaner mutant mouse. *Neuroscience* 7, 2185–2196.
73. Heckroth, J.A., and Abbott, L.C. (1994). Purkinje cell loss from alternating sagittal zones in the cerebellum of leaner mutant mice. *Brain Res.* 658, 93–104.
74. Lau, F.C., Frank, T.C., Nahm, S.-S., Stoica, G., and Abbott, L.C. (2004). Postnatal apoptosis in cerebellar granule cells of homozygous leaner (tg1a/tg1a) mice. *Neurotox. Res.* 6, 267–280.
75. Walter, J.T., Alviña, K., Womack, M.D., Chevez, C., and Khodakhah, K. (2006). Decreases in the precision of Purkinje cell pacemaking cause cerebellar dysfunction and ataxia. *Nat. Neurosci.* 9, 389–397.
76. Ovsepian, S. V, and Friel, D.D. (2012). Enhanced synaptic inhibition disrupts the efferent code of cerebellar Purkinje neurons in leaner Cav2.1 Ca²⁺ channel mutant mice. *Cerebellum* 11, 666–680.
77. Ovsepian, S. V, and Friel, D.D. (2008). The leaner P/Q-type calcium channel mutation renders cerebellar Purkinje neurons hyper-excitabile and eliminates Ca²⁺-Na⁺ spike bursts. *Eur. J. Neurosci.* 27, 93–103.
78. Zhuchenko, O., Bailey, J., Bonnen, P., Ashizawa, T., Stockton, D.W., Amos, C., Dobyns, W.B., Subramony, S.H., Zoghbi, H.Y., and Lee, C.C. (1997). Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansion in the α 1A-voltage-dependent calcium channel. *Nat. Genet.* 15, 62–69.
79. Giffin, N.J., Benton, S., and Goadsby, P.J. (2002). Benign paroxysmal torticollis of infancy: four new cases and linkage to CACNA1A mutation. *Dev. Med. Child Neurol.* 44, 490–493.
80. Raïke, R.S., Hess, E.J., and Jinnah, H.A. (2015). Dystonia and cerebellar degeneration in the leaner mouse mutant. *Brain Res.* 1611, 56–64.
81. Calderon, D.P., Fremont, R., Kraenzlin, F., and Khodakhah, K. (2011). The neural substrates of rapid-onset Dystonia-Parkinsonism. *Nat. Neurosci.* 14, 357–365.
82. Fremont, R., Calderon, D.P., Maleki, S., and Khodakhah, K. (2014). Abnormal high-frequency burst firing of cerebellar neurons in rapid-onset dystonia-parkinsonism. *J. Neurosci.* 34, 11723–11732.
83. Fremont, R., Tewari, A., and Khodakhah, K. (2015). Aberrant Purkinje cell activity is the cause of dystonia in a shRNA-based mouse model of Rapid Onset Dystonia-Parkinsonism. *Neurobiol. Dis.* 82, 200–212.
84. Chen, C.H., Fremont, R., Arteaga-Bracho, E.E., and Khodakhah, K. (2014). Short latency cerebellar modulation of the basal ganglia. *Nat. Neurosci.* 17, 1767–1775.
85. Luna-Cancelon, K., Sikora, K.M., Pappas, S.S., Singh, V., Wulff, H., Paulson, H.L., Burmeister, M., and Shakkottai, V.G. (2014). Alterations in cerebellar physiology are associated with a stiff-legged gait in Atcayji-hes mice. *Neurobiol. Dis.* 67, 140–148.
86. Bomar, J.M., Benke, P.J., Slattery, E.L., Puttagunta, R., Taylor, L.P., Seong, E., Nystuen, A., Chen, W., Albin, R.L., Patel, P.D., et al. (2003). Mutations in a novel gene encoding a CRAL-TRIO domain cause human Cayman ataxia and ataxia/dystonia in the jittery mouse. *Nat. Genet.* 35, 264–269.

87. Wang, Q., Bardgett, M.E., Wong, M., Wozniak, D.F., Lou, J., McNeil, B.D., Chen, C., Nardi, A., Reid, D.C., Yamada, K., et al. (2002). Ataxia and paroxysmal dyskinesia in mice lacking axonally transported FGF14. *Neuron* 35, 25–38.
88. Zech, M., Lam, D.D., Francescato, L., Schormair, B., Salminen, A. V., Jochim, A., Wieland, T., Lichtner, P., Peters, A., Gieger, C., et al. (2015). Recessive Mutations in the $\alpha 3$ (VI) Collagen Gene COL6A3 Cause Early-Onset Isolated Dystonia. *Am. J. Hum. Genet.* 96, 883–893.
89. Fogel, B.L., Hanson, S.M., and Becker, E.B.E. (2015). Do mutations in the murine ataxia gene TRPC3 cause cerebellar ataxia in humans? *Mov. Disord.* 30, 284–286.
90. Ichinose, H., Ohye, T., Takahashi, E., Seki, N., Hori, T., Segawa, M., Nomura, Y., Endo, K., Tanaka, H., Tsuji, S., et al. (1994). Hereditary progressive dystonia with marked diurnal fluctuation caused by mutations in the GTP cyclohydrolase I gene. *Nat. Genet.* 8, 236–242.
91. Lüdecke, B., Dworniczak, B., and Bartholomé, K. (1995). A point mutation in the tyrosine hydroxylase gene associated with Segawa's syndrome. *Hum. Genet.* 95, 123–125.
92. Nygaard, T.G., Marsden, C.D., and Duvoisin, R.C. (1988). Dopa-responsive dystonia. *Adv. Neurol.* 50, 377–384.
93. Matilla-Dueñas, A. (2012). The Ever Expanding Spinocerebellar Ataxias. Editorial. *The Cerebellum* 11, 821–827.
94. Storey, E., and Phil, D. (2014). Genetic Cerebellar Ataxias. 280–292.
95. Balint, B., and Bhatia, K.P. (2015). Isolated and combined dystonia syndromes - an update on new genes and their phenotypes. *Eur. J. Neurol.* 22, 610–617.
96. Xiao, J., Vemula, S.R., and LeDoux, M.S. (2014). Recent advances in the genetics of dystonia. *Curr. Neurol. Neurosci. Rep.* 14, 462.
97. Pers, T.H., Karjalainen, J.M., Chan, Y., Westra, H.-J., Wood, A.R., Yang, J., Lui, J.C., Vedantam, S., Gustafsson, S., Esko, T., et al. (2015). Biological interpretation of genome-wide association studies using predicted gene functions. *Nat. Commun.* 6, 5890.
98. Mi, H., Muruganujan, A., Casagrande, J.T., and Thomas, P.D. (2013). Large-scale gene function analysis with the PANTHER classification system. *Nat. Protoc.* 8, 1551–1566.
99. Groen, J.L., Andrade, A., Ritz, K., Jalalzadeh, H., Haagmans, M., Bradley, T.E.J., Jongejan, A., Verbeek, D.S., Nürnberg, P., Denome, S., et al. (2014). CACNA1B mutation is linked to unique myoclonus-dystonia syndrome. *Hum. Mol. Genet.* 24, 987–993.
100. Coutelier, M., Blesneac, I., Monteil, A., Monin, M., Ando, K., Mundwiller, E., Brusco, A., Le Ber, I., Anheim, M., Castrioto, A., et al. (2015). A Recurrent Mutation in CACNA1G Alters Cav3.1 T-Type Calcium-Channel Conduction and Causes Autosomal-Dominant Cerebellar Ataxia. *Am. J. Hum. Genet.* 97, 726–737.
101. Duarri, A., Nibbeling, E.A.R., Fokkens, M.R., Meijer, M., Boerrigter, M., Verschuuren-Bemelmans, C.C., Kremer, B.P.H., van de Warrenburg, B.P., Dooijes, D., Boddeke, E., et al. (2015). Functional Analysis Helps to Define KCNC3 Mutational Spectrum in Dutch Ataxia Cases. *PLoS One* 10, e0116599.
102. Waters, M.F., Minassian, N.A., Stevanin, G., Figueroa, K.P., Bannister, J.P.A., Nolte, D., Mock, A.F., Evidente, V.G.H., Fee, D.B., Müller, U., et al. (2006). Mutations in voltage-gated potassium channel KCNC3 cause degenerative and developmental central nervous system phenotypes. *Nat. Genet.* 38, 447–451.
103. Duarri, A., Jezierska, J., Fokkens, M., Meijer, M., Schelhaas, H.J., den Dunnen, W.F.A., van Dijk, F., Verschuuren-Bemelmans, C., Hageman, G., van de Vlies, P., et al. (2012). Mutations in potassium channel *kcnd3* cause spinocerebellar ataxia type 19. *Ann. Neurol.* 72, 870–880.

104. Lee, Y.-C., Durr, A., Majczenko, K., Huang, Y.-H., Liu, Y.-C., Lien, C.-C., Tsai, P.-C., Ichikawa, Y., Goto, J., Monin, M.-L., et al. (2012). Mutations in KCND3 cause spinocerebellar ataxia type 22. *Ann. Neurol.* *72*, 859–869.
105. Coutelier, M., Stevanin, G., and Brice, A. (2015). Genetic landscape remodelling in spinocerebellar ataxias: the influence of next-generation sequencing. *J. Neurol.*
106. Huang, L., Warman, J., Carter, M.T., Friend, K.L., Dudding, T.E., Schwartzentruber, J., Zou, R., Schofield, P.W., Douglas, S., Bulman, D.E., et al. (2012). Missense mutations in ITPR1 cause autosomal dominant congenital nonprogressive spinocerebellar ataxia. *Orphanet J. Rare Dis.* *7*, 67.
107. Meisler, M.H., Sprunger, L.K., Plummer, N.W., Escayg, A., and Jones, J.M. (1997). Ion channel mutations in mouse models of inherited neurological disease. *Ann. Med.* *29*, 569–574.
108. Shirley, T.L., Rao, L.M., Hess, E.J., and Jinnah, H.A. (2008). Paroxysmal dyskinesias in mice. *Mov. Disord.* *23*, 259–264.
109. Jinnah, H.A., Sepkuty, J.P., Ho, T., Yitta, S., Drew, T., Rothstein, J.D., and Hess, E.J. (2000). Calcium channel agonists and dystonia in the mouse. *Mov. Disord.* *15*, 542–551.
110. Street, V.A., Bosma, M.M., Demas, V.P., Regan, M.R., Lin, D.D., Robinson, L.C., Agnew, W.S., and Tempel, B.L. (1997). The type 1 inositol 1,4,5-trisphosphate receptor gene is altered in the opisthotonos mouse. *J. Neurosci.* *17*, 635–645.
111. Duarri, A., Lin, M.A., Fokkens, M.R., Meijer, M., Smeets, C.J.L.M., Nibbeling, E.A.R., Boddeke, E., Sinke, R.J., Kampinga, H.H., Papazian, D.M., et al. (2015). Spinocerebellar ataxia type 19 / 22 mutations alter heterocomplex Kv4 . 3 channel function and gating in a dominant manner. *Cell. Mol. Life Sci.* *72*, 3387–3399.
112. Browne, D.L., Gancher, S.T., Nutt, J.G., Brunt, E.R.P., Smith, E.A., Kramer, P., and Litt, M. (1994). Episodic ataxia/myokymia syndrome is associated with point mutations in the human potassium channel gene, KCNA1. *Nat. Genet.* *8*, 136–140.
113. Prüss, H., Wenzel, M., Eulitz, D., Thomzig, A., Karschin, A., and Veh, R.W. (2003). Kir2 potassium channels in rat striatum are strategically localized to control basal ganglia function. *Mol. Brain Res.* *110*, 203–219.
114. Signorini, S., Liao, Y.J., Duncan, S.A., Jan, L.Y., and Stoffel, M. (1997). Normal cerebellar development but susceptibility to seizures in mice lacking G protein-coupled, inwardly rectifying K⁺ channel GIRK2. *Proc. Natl. Acad. Sci. U. S. A.* *94*, 923–927.
115. Kotecki, L., Hearing, M., McCall, N.M., Marron Fernandez de Velasco, E., Pravetoni, M., Arora, D., Victoria, N.C., Munoz, M.B., Xia, Z., Slesinger, P. a., et al. (2015). GIRK Channels Modulate Opioid-Induced Motor Activity in a Cell Type- and Subunit-Dependent Manner. *J. Neurosci.* *35*, 7131–7142.
116. Bakalkin, G., Watanabe, H., Jezierska, J., Depoorter, C., Verschuuren-Bemelmans, C., Bazov, I., Artemenko, K.A., Yakovleva, T., Dooijes, D., Van de Warrenburg, B.P.C., et al. (2010). Prodynorphin mutations cause the neurodegenerative disorder spinocerebellar ataxia type 23. *Am. J. Hum. Genet.* *87*, 593–603.
117. Boecker, H. (2013). Imaging the Role of GABA in Movement Disorders. *Curr. Neurol. Neurosci. Rep.* *13*, 385.
118. Hori, K., and Hoshino, M. (2012). GABAergic Neuron Specification in the Spinal Cord, the Cerebellum, and the Cochlear Nucleus. *Neural Plast.* *2012*, 1–11.
119. Segawa, M., and Nomura, Y. (2014). Genetics and Pathophysiology of Primary Dystonia with Special Emphasis on DYT1 and DYT5. *Semin. Neurol.* *34*, 306–311.

120. Quartarone, A., and Hallett, M. (2013). Emerging concepts in the physiological basis of dystonia. *Mov. Disord.* 28, 958–967.
121. Hirose, S. (2014). Mutant GABA(A) receptor subunits in genetic (idiopathic) epilepsy. *Prog. Brain Res.* 213, 55–85.
122. Kaja, S., Payne, A.J., Nielsen, E.Ø., Thompson, C.L., van den Maagdenberg, A.M.J.M., Koulen, P., and Snutch, T.P. (2015). Differential cerebellar GABAA receptor expression in mice with mutations in CaV2.1 (P/Q-type) calcium channels. *Neuroscience* 304, 198–208.
123. Kayakabe, M., Kakizaki, T., Kaneko, R., Sasaki, A., Nakazato, Y., Shibasaki, K., Ishizaki, Y., Saito, H., Suzuki, N., Furuya, N., et al. (2014). Motor dysfunction in cerebellar Purkinje cell-specific vesicular GABA transporter knockout mice. *Front. Cell. Neurosci.* 7, 286.
124. Nomura, T., Kakegawa, W., Matsuda, S., Kohda, K., Nishiyama, J., Takahashi, T., and Yuzaki, M. (2012). Cerebellar long-term depression requires dephosphorylation of TARP in Purkinje cells. *Eur. J. Neurosci.* 35, 402–410.
125. Wozniak, D.F., Xiao, M., Xu, L., Yamada, K.A., and Ornitz, D.M. (2007). Impaired spatial learning and defective theta burst induced LTP in mice lacking fibroblast growth factor 14. *Neurobiol. Dis.* 26, 14–26.
126. Armbrust, K.R., Wang, X., Hathorn, T.J., Cramer, S.W., Chen, G., Zu, T., Kangas, T., Zink, a N., Oz, G., Ebner, T.J., et al. (2014). Mutant beta-III spectrin causes mGluR1alpha mislocalization and functional deficits in a mouse model of spinocerebellar ataxia type 5. *J Neurosci* 34, 9891–9904.
127. Letts, V.A., Felix, R., Biddlecome, G.H., Arikath, J., Mahaffey, C.L., Valenzuela, A., Bartlett, F.S., Mori, Y., Campbell, K.P., and Frankel, W.N. (1998). The mouse stargazer gene encodes a neuronal Ca²⁺-channel gamma subunit. *Nat. Genet.* 19, 340–347.
128. Roy, B., Ahmed, K.T., Cunningham, M.E., Ferdous, J., Mukherjee, R., Zheng, W., Chen, X.-Z., and Ali, D.W. (2015). Zebrafish TARP Cacng2 is required for the expression and normal development of AMPA receptors at excitatory synapses. *Dev. Neurobiol.* 2, n/a-n/a.
129. de Bartolomeis, A., Buonaguro, E.F., Iasevoli, F., and Tomasetti, C. (2014). The emerging role of dopamine-glutamate interaction and of the postsynaptic density in bipolar disorder pathophysiology: Implications for treatment. *J. Psychopharmacol.* 28, 505–526.
130. Ohba, C., Shiina, M., Tohyama, J., Haginoya, K., Lerman-Sagie, T., Okamoto, N., Blumkin, L., Lev, D., Mukaida, S., Nozaki, F., et al. (2015). *GRIN1* mutations cause encephalopathy with infantile-onset epilepsy, and hyperkinetic and stereotyped movement disorders. *Epilepsia* 56, 841–848.
131. Hamdan, F.F., Gauthier, J., Araki, Y., Lin, D.-T., Yoshizawa, Y., Higashi, K., Park, A.-R., Spiegelman, D., Dobrzaniecka, S., Piton, A., et al. (2011). Excess of de novo deleterious mutations in genes associated with glutamatergic systems in nonsyndromic intellectual disability. *Am. J. Hum. Genet.* 88, 306–316.
132. Miladinovic, T., Nashed, M., and Singh, G. (2015). Overview of Glutamatergic Dysregulation in Central Pathologies. *Biomolecules* 5, 3112–3141.
133. Charlesworth, G., Bhatia, K.P., and Wood, N.W. (2013). The genetics of dystonia: New twists in an old tale. *Brain* 136, 2017–2037.
134. Jones, J.M., Dionne, L., Dell'Orco, J., Parent, R., Krueger, J.N., Cheng, X., Dib-Hajj, S.D., Bunton-Stasyshyn, R.K., Sharkey, L.M., Dowling, J.J., et al. (2016). Single amino acid deletion in transmembrane segment D4S6 of sodium channel Scn8a (Nav1.6) in a mouse mutant with a chronic movement disorder. *Neurobiol. Dis.* 89, 36–45.

135. Saegusa, C., Fukuda, M., and Mikoshiba, K. (2002). Synaptotagmin V is targeted to dense-core vesicles that undergo calcium-dependent exocytosis in PC12 cells. *J. Biol. Chem.* *277*, 24499–24505.
136. Sadakata, T., Kakegawa, W., Mizoguchi, A., Washida, M., Katoh-Semba, R., Shutoh, F., Okamoto, T., Nakashima, H., Kimura, K., Tanaka, M., et al. (2007). Impaired cerebellar development and function in mice lacking CAPS2, a protein involved in neurotrophin release. *J. Neurosci.* *27*, 2472–2482.
137. Caldeira, M. V., Melo, C. V., Pereira, D.B., Carvalho, R.F., Carvalho, A.L., and Duarte, C.B. (2007). BDNF regulates the expression and traffic of NMDA receptors in cultured hippocampal neurons. *Mol. Cell. Neurosci.* *35*, 208–219.
138. Vicario-Abejon, C., Collin, C., McKay, R.D.G., and Segal, M. (1998). Neurotrophins Induce Formation of Functional Excitatory and Inhibitory Synapses between Cultured Hippocampal Neurons. *J. Neurosci.* *18*, 7256–7271.
139. Bellocchio, E.E. (2000). Uptake of Glutamate into Synaptic Vesicles by an Inorganic Phosphate Transporter. *Science (80-)*. *289*, 957–960.
140. Wojcik, S.M., Rhee, J.-S., Herzog, E., Sigler, A., Jahn, R., Takamori, S., Brose, N., and Rosenmund, C. (2004). An essential role for vesicular glutamate transporter 1 (VGLUT1) in postnatal development and control of quantal size. *Proc. Natl. Acad. Sci. U. S. A.* *101*, 7158–7163.
141. Wu, Y.-J., Tejero, R., Arancillo, M., Vardar, G., Korotkova, T., Kintscher, M., Schmitz, D., Ponomarenko, A., Tabares, L., and Rosenmund, C. (2015). Syntaxin 1B is important for mouse postnatal survival and proper synaptic function at the mouse neuromuscular junctions. *J. Neurophysiol.* *114*, 2404–2417.
142. Bourassa, C. V., Meijer, I.A., Merner, N.D., Grewal, K.K., Stefanelli, M.G., Hodgkinson, K., Ives, E.J., Pryse-Phillips, W., Jog, M., Boycott, K., et al. (2012). VAMP1 mutation causes dominant hereditary spastic ataxia in Newfoundland families. *Am. J. Hum. Genet.* *91*, 548–552.
143. Nystuen, A.M., Schwendinger, J.K., Sachs, A.J., Yang, A.W., and Haider, N.B. (2007). A null mutation in VAMP1/synaptobrevin is associated with neurological defects and prewean mortality in the lethal-wasting mouse mutant. *Neurogenetics* *8*, 1–10.
144. Liu, Y., Sugiura, Y., and Lin, W. (2011). The role of synaptobrevin1/VAMP1 in Ca²⁺-triggered neurotransmitter release at the mouse neuromuscular junction. *J. Physiol.* *589*, 1603–1618.
145. Schoch, S. (2001). SNARE Function Analyzed in Synaptobrevin/VAMP Knockout Mice. *Science (80-)*. *294*, 1117–1122.
146. Hussain, S., and Davanger, S. (2015). Postsynaptic VAMP/Synaptobrevin Facilitates Differential Vesicle Trafficking of GluA1 and GluA2 AMPA Receptor Subunits. *PLoS One* *10*, e0140868.
147. Chin, L.S., Li, L., Ferreira, a, Kosik, K.S., and Greengard, P. (1995). Impairment of axonal development and of synaptogenesis in hippocampal neurons of synapsin I-deficient mice. *Proc. Natl. Acad. Sci. U. S. A.* *92*, 9230–9234.
148. Li, L., Chin, L.S., Shupliakov, O., Brodin, L., Sihra, T.S., Hvalby, O., Jensen, V., Zheng, D., McNamara, J.O., and Greengard, P. (1995). Impairment of synaptic vesicle clustering and of synaptic transmission, and increased seizure propensity, in synapsin I-deficient mice. *Proc. Natl. Acad. Sci. U. S. A.* *92*, 9235–9239.
149. Lovenberg, T.W., Roland, B.L., Wilson, S.J., Jiang, X., Pyati, J., Huvar, A., Jackson, M.R., and Erlander, M.G. (1999). Cloning and Functional Expression of the Human Histamine H3 Receptor. *Mol. Pharmacol.* *55*, 1101–1107.
150. Schneider, E.H., Neumann, D., and Seifert, R. (2014). Modulation of behavior by the histaminergic system: Lessons from HDC-, H3R- and H4R-deficient mice. *Neurosci. Biobehav. Rev.* *47*, 101–121.

151. Toyota, H., Dugovic, C., Koehl, M., Laposky, A.D., Weber, C., Ngo, K., Wu, Y., Lee, D.H., Yanai, K., Sakurai, E., et al. (2002). Behavioral characterization of mice lacking histamine H(3) receptors. *Mol. Pharmacol.* *62*, 389–397.
152. Takahashi, K., Suwa, H., Ishikawa, T., and Kotani, H. (2002). Targeted disruption of H3 receptors results in changes in brain histamine tone leading to an obese phenotype. *J. Clin. Invest.* *110*, 1791–1799.
153. Kim, E., Cho, K.-O., Rothschild, A., and Sheng, M. (1996). Heteromultimerization and NMDA Receptor-Clustering Activity of Chapsyn-110, a Member of the PSD-95 Family of Proteins. *Neuron* *17*, 103–113.
154. Migaud, M., Charlesworth, P., Dempster, M., Webster, L.C., Watabe, A.M., Makhinson, M., He, Y., Ramsay, M.F., Morris, R.G.M., Morrison, J.H., et al. (1998). Enhanced long-term potentiation and impaired learning in mice with mutant postsynaptic density-95 protein. *Nature* *396*, 433–439.
155. Um, J.W., Pramanik, G., Ko, J.S., Song, M.-Y., Lee, D., Kim, H., Park, K.-S., Südhof, T.C., Tabuchi, K., and Ko, J. (2014). Calsyntenins function as synaptogenic adhesion molecules in concert with neuexins. *Cell Rep.* *6*, 1096–1109.
156. Pettem, K.L., Yokomaku, D., Luo, L., Linhoff, M.W., Prasad, T., Connor, S.A., Siddiqui, T.J., Kawabe, H., Chen, F., Zhang, L., et al. (2013). The specific α -neuexin interactor calsyntenin-3 promotes excitatory and inhibitory synapse development. *Neuron* *80*, 113–128.
157. Lu, Z., Wang, Y., Chen, F., Tong, H., Reddy, M.V.V.S., Luo, L., Seshadrinathan, S., Zhang, L., Holthausen, L.M.F., Craig, A.M., et al. (2014). Calsyntenin-3 molecular architecture and interaction with neuexin 1 α . *J. Biol. Chem.* *289*, 34530–34542.
158. Lee, F.J.S., Xue, S., Pei, L., Vukusic, B., Chéry, N., Wang, Y., Wang, Y.T., Niznik, H.B., Yu, X., and Liu, F. (2002). Dual Regulation of NMDA Receptor Functions by Direct Protein-Protein Interactions with the Dopamine D1 Receptor. *Cell* *111*, 219–230.
159. Groen, J.L., Ritz, K., Warner, T.T., Baas, F., and Tijssen, M.A.J. (2014). DRD1 rare variants associated with tardive-like dystonia: A pilot pathway sequencing study in dystonia. *Park. Relat. Disord.* *20*, 782–785.
160. Xu, M., Moratalla, R., Gold, L.H., Hiroi, N., Koob, G.F., Graybiel, A.M., and Tonegawa, S. (1994). Dopamine D1 receptor mutant mice are deficient in striatal expression of dynorphin and in dopamine-mediated behavioral responses. *Cell* *79*, 729–742.
161. Schöls, L., Reimold, M., Seidel, K., Globas, C., Brockmann, K., Karsten Hauser, T., Auburger, G., Bürk, K., Den Dunnen, W., Reischl, G., et al. (2015). No parkinsonism in SCA2 and SCA3 despite severe neurodegeneration of the dopaminergic substantia nigra. *Brain* *138*, 3316–3326.
162. Gunnarsen, J.M., Kim, M.H., Fuller, S.J., De Silva, M., Britto, J.M., Hammond, V.E., Davies, P.J., Petrou, S., Faber, E.S.L., Sah, P., et al. (2007). Sez-6 proteins affect dendritic arborization patterns and excitability of cortical pyramidal neurons. *Neuron* *56*, 621–639.
163. Miyazaki, T., Hashimoto, K., Uda, A., Sakagami, H., Nakamura, Y., Saito, S., Nishi, M., Kume, H., Tohgo, A., Kaneko, I., et al. (2006). Disturbance of cerebellar synaptic maturation in mutant mice lacking BSRPs, a novel brain-specific receptor-like protein family. *FEBS Lett.* *580*, 4057–4064.
164. Frank, T., Rutherford, M.A., Strenzke, N., Neef, A., Pangršič, T., Khimich, D., Fejtova, A., Fejtova, A., Gundelfinger, E.D., Liberman, M.C., et al. (2010). Bassoon and the synaptic ribbon organize Ca²⁺ channels and vesicles to add release sites and promote refilling. *Neuron* *68*, 724–738.
165. Miura, S., Nakagawara, H., Kaida, H., Sugita, M., Noda, K., Motomura, K., Ohyagi, Y., Ayabe, M., Aizawa, H., Ishibashi, M., et al. (2009). Expansion of the phenotypic spectrum of SCA14 caused by the Gly128Asp mutation in PRKCG. *Clin. Neurol. Neurosurg.* *111*, 211–215.

166. Ganos, C., Zittel, S., Minnerop, M., Schunke, O., Heinbokel, C., Gerloff, C., Zühlke, C., Bauer, P., Klockgether, T., Münchau, A., et al. (2014). Clinical and neurophysiological profile of four German families with spinocerebellar ataxia type 14. *Cerebellum* 13, 89–96.
167. Visser, J.E., Bloem, B.R., and van de Warrenburg, B.P.C. (2007). PRKCG mutation (SCA-14) causing a Ramsay Hunt phenotype. *Mov. Disord.* 22, 1024–1026.
168. van de Warrenburg, B.P.C., Verbeek, D.S., Piersma, S.J., Hennekam, F.A.M., Pearson, P.L., Knoers, N.V.A.M., Kremer, H.P.H., and Sinke, R.J. (2003). Identification of a novel SCA14 mutation in a Dutch autosomal dominant cerebellar ataxia family. *Neurology* 61, 1760–1765.
169. Yasuda, J., Whitmarsh, J., Cavanagh, J., Sharma, M., and Davis, R.J. (1999). The JIP group of mitogen-activated protein kinase scaffold proteins. *Mol. Cell. Biol.* 19, 7245–7254.
170. Giza, J., Urbanski, M.J., Prestori, F., Bandyopadhyay, B., Yam, A., Friedrich, V., Kelley, K., D'Angelo, E., and Goldfarb, M. (2010). Behavioral and Cerebellar Transmission Deficits in Mice Lacking the Autism-Linked Gene *Islet Brain-2*. *J. Neurosci.* 30, 14805–14816.
171. Kennedy, N.J., Martin, G., Ehrhardt, A.G., Cavanagh-Kyros, J., Kuan, C.Y., Rakic, P., Flavell, R. a., Treisman, S.N., and Davis, R.J. (2007). Requirement of JIP scaffold proteins for NMDA-mediated signal transduction. *Genes Dev.* 21, 2336–2346.
172. Ludwig, M.-G., and Seuwen, K. (2002). CHARACTERIZATION OF THE HUMAN ADENYLYL CYCLASE GENE FAMILY: cDNA, GENE STRUCTURE, AND TISSUE DISTRIBUTION OF THE NINE ISOFORMS. *J. Recept. Signal Transduct.* 22, 79–110.
173. Hervé, D. (2011). Identification of a Specific Assembly of the G Protein Golf as a Critical and Regulated Module of Dopamine and Adenosine-Activated cAMP Pathways in the Striatum. *Front. Neuroanat.* 5, 1–9.
174. Chen, Y.-Z., Matsushita, M.M., Robertson, P., Rieder, M., Girirajan, S., Antonacci, F., Lipe, H., Eichler, E.E., Nickerson, D.A., Bird, T.D., et al. (2012). Autosomal dominant familial dyskinesia and facial myokymia: single exome sequencing identifies a mutation in adenylyl cyclase 5. *Arch. Neurol.* 69, 630–635.
175. Chen, Y.-Z., Friedman, J.R., Chen, D.-H., Chan, G.C.-K., Bloss, C.S., Hisama, F.M., Topol, S.E., Carson, A.R., Pham, P.H., Bonkowski, E.S., et al. (2014). Gain-of-function ADCY5 mutations in familial dyskinesia with facial myokymia. *Ann. Neurol.* 75, 542–549.
176. Mencacci, N.E., Erro, R., Wiethoff, S., Hersheson, J., Ryten, M., Balint, B., Ganos, C., Stamelou, M., Quinn, N., Houlden, H., et al. (2015). ADCY5 mutations are another cause of benign hereditary chorea. *Neurology*.
177. Carapito, R., Paul, N., Untrau, M., Le Gentil, M., Ott, L., Alsaleh, G., Jochem, P., Radosavljevic, M., Le Caignec, C., David, A., et al. (2015). A *de novo* ADCY5 mutation causes early-onset autosomal dominant chorea and dystonia. *Mov. Disord.* 30, 423–427.
178. Fuchs, T., Saunders-Pullman, R., Masuho, I., Luciano, M.S., Raymond, D., Factor, S., Lang, A.E., Liang, T.-W., Trosch, R.M., White, S., et al. (2013). Mutations in GNAL cause primary torsion dystonia. *Nat. Genet.* 45, 88–92.
179. Harrison, L.M., and He, Y. (2011). Rhes and AGS1/Dexas1 affect signaling by dopamine D1 receptors through adenylyl cyclase. *J. Neurosci. Res.* 89, 874–882.
180. Quintero, G.C., Spano, D., Lahoste, G.J., and Harrison, L.M. (2008). The Ras homolog Rhes affects dopamine D1 and D2 receptor-mediated behavior in mice. *Neuropharmacol. Neurotoxicology* 19, 1563–1566.

181. Subramaniam, S., Napolitano, F., Mealer, R.G., Kim, S., Errico, F., Barrow, R., Shahani, N., Tyagi, R., Snyder, S.H., and Usiello, A. (2011). Rhes, a striatal-enriched small G protein, mediates mTOR signaling and L-DOPA-induced dyskinesia. *Nat. Neurosci.* *15*, 191–193.
182. Rivera, C., Voipio, J., Payne, J.A., Ruusuvuori, E., Lahtinen, H., Lamsa, K., Pirvola, U., Saarma, M., and Kaila, K. (1999). The K⁺/Cl⁻ co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* *397*, 251–255.
183. Hübner, C.A., Stein, V., Hermans-Borgmeyer, I., Meyer, T., Ballanyi, K., and Jentsch, T.J. (2001). Disruption of KCC2 Reveals an Essential Role of K-Cl Cotransport Already in Early Synaptic Inhibition. *Neuron* *30*, 515–524.
184. Campbell, P.D., Shen, K., Sapio, M.R., Glenn, T.D., Talbot, W.S., and Marlow, F.L. (2014). Unique function of Kinesin Kif5A in localization of mitochondria in axons. *J. Neurosci.* *34*, 14717–14732.
185. Nakajima, K., Yin, X., Takei, Y., Seog, D.-H., Homma, N., and Hirokawa, N. (2012). Molecular motor KIF5A is essential for GABA(A) receptor transport, and KIF5A deletion causes epilepsy. *Neuron* *76*, 945–961.
186. Nakayama, T., Yaoi, T., Yasui, M., and Kuwajima, G. (1998). N-copine: a novel two C2-domain-containing protein with neuronal activity-regulated expression. *FEBS Lett.* *428*, 80–84.
187. Pan, Z.Z. (2012). Transcriptional control of Gad2. *Transcription* *3*, 68–72.
188. Almuedo-Castillo, M., Saló, E., and Adell, T. (2011). Dishevelled is essential for neural connectivity and planar cell polarity in planarians. *Proc. Natl. Acad. Sci. U. S. A.* *108*, 2813–2818.
189. Cerpa, W., Toledo, E.M., Varela-Nallar, L., and Inestrosa, N.C. (2009). The role of Wnt signaling in neuroprotection. *Drug News Perspect.* *22*, 579–591.
190. Hauser, K.F., Aldrich, J. V., Anderson, K.J., Bakalkin, G., Christie, M.J., Hall, E.D., Knapp, P.E., Scheff, S.W., Singh, I.N., Vissel, B., et al. (2005). Pathobiology of dynorphins in trauma and disease. *Front. Biosci.* *10*, 216–235.
191. Togashi, H., Abe, K., Mizoguchi, A., Takaoka, K., Chisaka, O., and Takeichi, M. (2002). Cadherin Regulates Dendritic Spine Morphogenesis. *Neuron* *35*, 77–89.
192. Friedman, L.G., Riemsdijk, F.W., Sullivan, J.M., Mesias, R., Williams, F.M., Huntley, G.W., and Benson, D.L. (2015). Cadherin-8 expression, synaptic localization, and molecular control of neuronal form in prefrontal corticostriatal circuits. *J. Comp. Neurol.* *523*, 75–92.
193. Carbon, M., and Eidelberg, D. (2009). Abnormal structure-function relationships in hereditary dystonia. *Neuroscience* *164*, 220–229.
194. Vanni, V., Puglisi, F., Bonsi, P., Ponterio, G., Maltese, M., Pisani, A., and Mandolesi, G. (2015). Cerebellar synaptogenesis is compromised in mouse models of DYT1 dystonia. *Exp. Neurol.* *271*, 457–467.
195. Planes, M., Rooryck, C., Vuillaume, M.-L., Besnard, L., Bouron, J., Lacombe, D., Arveiler, B., and Goizet, C. (2015). SCA27 is a cause of early-onset ataxia and developmental delay. *Eur. J. Paediatr. Neurol.* *19*, 271–273.
196. Di Fabio, R., Santorelli, F., Bertini, E., Balestri, M., Cursi, L., Tessa, A., Pierelli, F., and Casali, C. (2012). Infantile childhood onset of spinocerebellar ataxia type 2. *Cerebellum* *11*, 526–530.
197. Jacob, F.-D., Ho, E.S., Martinez-Ojeda, M., Darras, B.T., and Khwaja, O.S. (2013). Case of infantile onset spinocerebellar ataxia type 5. *J. Child Neurol.* *28*, 1292–1295.
198. Issa, F.A., Mock, A.F., Sagasti, A., and Papazian, D.M. (2012). Spinocerebellar ataxia type 13 mutation that is associated with disease onset in infancy disrupts axonal pathfinding during neuronal development. *Dis. Model. Mech.* *5*, 921–929.

199. McNeill, E.M., Klöckner-Bormann, M., Roesler, E.C., Talton, L.E., Moechars, D., and Clagett-Dame, M. (2011). Nav2 hypomorphic mutant mice are ataxic and exhibit abnormalities in cerebellar development. *Dev. Biol.* *353*, 331–343.
200. Grilli, M., Toninelli, G.F., Uberti, D., Spano, P.F., and Memo, M. (2003). Alzheimer's disease linking neurodegeneration with neurodevelopment. *Funct. Neurol.* *18*, 145–148.
201. Wilkaniec, A., Czapski, G.A., and Adamczyk, A. (2016). Cdk5 at crossroads of protein oligomerization in neurodegenerative diseases: facts and hypotheses. *J. Neurochem.* *136*, 222–233.
202. Doehner, J., and Knuesel, I. (2010). Reelin-mediated Signaling during Normal and Pathological Forms of Aging. *Aging Dis.* *1*, 12–29.
203. Park, N., Yoo, J.C., Ryu, J., Hong, S.-G., Hwang, E.M., and Park, J.-Y. (2012). Copine1 enhances neuronal differentiation of the hippocampal progenitor HiB5 cells. *Mol. Cells* *34*, 549–554.
204. Huang, X., Hu, J., Li, Y., Zhuyun Yang, Z., Zhu, H., Zhou, L., Ma, K., Schachner, M., Xiao, Z., and Li, Y. (2013). The cell adhesion molecule L1 regulates the expression of FGF21 and enhances neurite outgrowth. *Brain Res.* *1530*, 13–21.
205. Wang, H., Duan, X., Ren, Y., Liu, Y., Huang, M., Liu, P., Wang, R., Gao, G., Zhou, L., Feng, Z., et al. (2013). FoxO3a negatively regulates nerve growth factor-induced neuronal differentiation through inhibiting the expression of neurochondrin in PC12 cells. *Mol. Neurobiol.* *47*, 24–36.
206. Brunet, J.-F., and Pattyn, A. (2002). Phox2 genes - from patterning to connectivity. *Curr. Opin. Genet. Dev.* *12*, 435–440.
207. Adams, L.M., and Foote, S.L. (1988). Possible involvement of brain noradrenergic neurons in dystonia. *Adv. Neurol.* *50*, 313–333.
208. Evans, P.R., Lee, S.E., Smith, Y., and Hepler, J.R. (2014). Postnatal developmental expression of regulator of G protein signaling 14 (RGS14) in the mouse brain. *J. Comp. Neurol.* *522*, 186–203.
209. Marín, F., and Nieto, M.A. (2006). The expression of Scratch genes in the developing and adult brain. *Dev. Dyn.* *235*, 2586–2591.
210. De Luca, A., Parmigiani, E., Tosatto, G., Martire, S., Hoshino, M., Buffo, A., Leto, K., and Rossi, F. (2015). Exogenous Sonic hedgehog modulates the pool of GABAergic interneurons during cerebellar development. *Cerebellum* *14*, 72–85.
211. López-Bendito, G., Luján, R., Shigemoto, R., Ganter, P., Paulsen, O., and Molnár, Z. (2003). Blockade of GABA(B) receptors alters the tangential migration of cortical neurons. *Cereb. Cortex* *13*, 932–942.
212. Masuho, I., Mototani, Y., Sahara, Y., Asami, J., Nakamura, S., Kozasa, T., and Inoue, T. (2008). Dynamic expression patterns of G protein-regulated inducer of neurite outgrowth 1 (GRIN1) and its colocalization with Galphao implicate significant roles of Galphao-GRIN1 signaling in nervous system. *Dev. Dyn.* *237*, 2415–2429.
213. Wang, V.Y., Rose, M.F., and Zoghbi, H.Y. (2005). Math1 expression redefines the rhombic lip derivatives and reveals novel lineages within the brainstem and cerebellum. *Neuron* *48*, 31–43.
214. Kay, J.N., Voinescu, P.E., Chu, M.W., and Sanes, J.R. (2011). Neurod6 expression defines new retinal amacrine cell subtypes and regulates their fate. *Nat. Neurosci.* *14*, 965–972.
215. Sherkhane, P., and Kapfhammer, J.P. (2013). The plasma membrane Ca²⁺-ATPase2 (PMCA2) is involved in the regulation of Purkinje cell dendritic growth in cerebellar organotypic slice cultures. *Neural Plast.* *2013*, 321685.

216. Zanni, G., Cali, T., Kalscheuer, V.M., Ottolini, D., Barresi, S., Lebrun, N., Montecchi-Palazzi, L., Hu, H., Chelly, J., Bertini, E., et al. (2012). Mutation of plasma membrane Ca²⁺ ATPase isoform 3 in a family with X-linked congenital cerebellar ataxia impairs Ca²⁺ homeostasis. *Proc. Natl. Acad. Sci. U. S. A.* *109*, 14514–14519.
217. Saitoh, S., Takamatsu, K., Kobayashi, M., and Noguchi, T. (1994). Expression of hippocalcin in the developing rat brain. *Brain Res. Dev. Brain Res.* *80*, 199–208.
218. Ikeda, Y., Dick, K.A., Weatherspoon, M.R., Gincel, D., Armbrust, K.R., Dalton, J.C., Stevanin, G., Dürr, A., Zühlke, C., Bürk, K., et al. (2006). Spectrin mutations cause spinocerebellar ataxia type 5. *Nat. Genet.* *38*, 184–190.
219. Downen, M., Belkowsky, S., Knowles, H., Cardillo, M., and Prystowsky, M.B. (1999). Developmental expression of voltage-gated potassium channel β subunits. *Dev. Brain Res.* *117*, 71–80.
220. Sergaki, M.C., Guillemot, F., and Matsas, R. (2010). Impaired cerebellar development and deficits in motor coordination in mice lacking the neuronal protein BM88/Cend1. *Mol. Cell. Neurosci.* *44*, 15–29.
221. Liang, Z., Ye, T., Zhou, X., Lai, K.-O., Fu, A.K.Y., and Ip, N.Y. (2015). Cdk5 Regulates Activity-Dependent Gene Expression and Dendrite Development. *J. Neurosci.* *35*, 15127–15134.
222. Qi, G.-J., Chen, Q., Chen, L.-J., Shu, Y., Bu, L.-L., Shao, X.-Y., Zhang, P., Jiao, F.-J., Shi, J., and Tian, B. (2015). Phosphorylation of Connexin 43 by Cdk5 Modulates Neuronal Migration During Embryonic Brain Development. *Mol. Neurobiol.*
223. Tojima, T., Itofusa, R., and Kamiguchi, H. (2014). Steering neuronal growth cones by shifting the imbalance between exocytosis and endocytosis. *J. Neurosci.* *34*, 7165–7178.
224. Puddifoot, C.A., Wu, M., Sung, R.-J., and Joiner, W.J. (2015). Ly6h regulates trafficking of $\alpha 7$ nicotinic acetylcholine receptors and nicotine-induced potentiation of glutamatergic signaling. *J. Neurosci.* *35*, 3420–3430.
225. Mouton-Liger, F., Sahún, I., Collin, T., Lopes Pereira, P., Masini, D., Thomas, S., Paly, E., Lullier, S., Mème, S., Jouhault, Q., et al. (2014). Developmental molecular and functional cerebellar alterations induced by PCP4/PEP19 overexpression: implications for Down syndrome. *Neurobiol. Dis.* *63*, 92–106.
226. Vogel-Ciernia, A., and Wood, M.A. (2014). Neuron-specific chromatin remodeling: a missing link in epigenetic mechanisms underlying synaptic plasticity, memory, and intellectual disability disorders. *Neuropharmacology* *80*, 18–27.
227. Mark, M.D., Krause, M., Boele, H.-J., Kruse, W., Pollok, S., Kuner, T., Dalkara, D., Koekkoek, S., De Zeeuw, C.I., and Herlitze, S. (2015). Spinocerebellar ataxia type 6 protein aggregates cause deficits in motor learning and cerebellar plasticity. *J. Neurosci.* *35*, 8882–8895.
228. Huynh, D.P., Maalouf, M., Silva, A.J., Schweizer, F.E., and Pulst, S.M. (2009). Dissociated Fear and Spatial Learning in Mice with Deficiency of Ataxin-2. *PLoS One* *4*, e6235.
229. Martella, G., Maltese, M., Nisticò, R., Schirinzi, T., Madeo, G., Sciamanna, G., Ponterio, G., Tassone, A., Mandolesi, G., Vanni, V., et al. (2014). Regional specificity of synaptic plasticity deficits in a knock-in mouse model of DYT1 dystonia. *Neurobiol. Dis.* *65*, 124–132.
230. Thomas, E.A., Danielson, P.E., and Sutcliffe, J.G. (1998). RGS9: a regulator of G-protein signalling with specific expression in rat and mouse striatum. *J. Neurosci. Res.* *52*, 118–124.
231. Psifogeorgou, K., Papakosta, P., Russo, S.J., Neve, R.L., Kardassis, D., Gold, S.J., and Zachariou, V. (2007). RGS9-2 is a negative modulator of mu-opioid receptor function. *J. Neurochem.* *103*, 617–625.


232. Nobrega, J., Parkes, J., Wong, P., Raymond, R., and Richter, A. (2004). Altered expression of preproenkephalin and prodynorphin mRNA in a genetic model of paroxysmal dystonia. *Brain Res.* 1015, 87–95.
233. Charlesworth, G., Plagnol, V., Holmström, K.M., Bras, J., Sheerin, U.M., Preza, E., Rubio-Agusti, I., Ryten, M., Schneider, S.A., Stamelou, M., et al. (2012). Mutations in ANO3 cause dominant craniocervical dystonia: Ion channel implicated in pathogenesis. *Am. J. Hum. Genet.* 91, 1041–1050.
234. Domingo, A., Erro, R., and Lohmann, K. (2016). Novel Dystonia Genes: Clues on Disease Mechanisms and the Complexities of High-Throughput Sequencing. *Mov. Disord.* 31, 471–477.
235. Cossette, P., Lachance-touchette, P., and Rouleau, G. a (2012). Mutated GABA A receptor subunits in idiopathic generalized epilepsy. In *Jasper's Basic Mechanisms of the Epilepsies 4th Edn.*, pp. 714–730.
236. Kim, J.-B. (2014). Channelopathies. *Korean J. Pediatr.* 57, 1–18.
237. Goddard, A.W. (2016). Cortical and subcortical gamma amino acid butyric acid deficits in anxiety and stress disorders: Clinical implications. *World J. Psychiatry* 6, 43–53.
238. Sharma, N., Jamwal, S., and Kumar, P. (2016). Beneficial effect of antidepressants against rotenone induced Parkinsonism like symptoms in rats. *Pathophysiology* 23, 123–134.
239. Bragg, D.C., Armata, I. a, Nery, F.C., Breakefield, X.O., and Sharma, N. (2011). Molecular pathways in dystonia. *Neurobiol. Dis.* 42, 136–147.
240. Orr, H.T., Chung, M., Banfi, S., Kwiatkowski, T.J., Servadio, A., Beaudet, A.L., McCall, A.E., Duwick, L.A., Ranum, L.P.W., and Zoghbi, H.Y. (1993). Expansion of an unstable trinucleotide CAG repeat in spinocerebellar ataxia type 1. *Nat. Genet.* 4, 221–226.
241. Pulst, S.M., Nechiporuk, A., Nechiporuk, T., Gispert, S., Chen, X.N., Lopes-Cendes, I., Pearlman, S., Starkman, S., Orozco-Díaz, G., Lunkes, A., et al. (1996). Moderate expansion of a normally biallelic trinucleotide repeat in spinocerebellar ataxia type 2. *Nat. Genet.* 14, 269–276.
242. Sanpei, K., Takano, H., Igarashi, S., Sato, T., Oyake, M., Sasaki, H., Wakisaka, A., Tashiro, K., Ishida, Y., Ikeuchi, T., et al. (1996). Identification of the spinocerebellar ataxia type 2 gene using a direct identification of repeat expansion and cloning technique, DIRECT. *Nat. Genet.* 14, 277–284.
243. Kawaguchi, Y., Okamoto, T., Taniwaki, M., Aizawa, M., Inoue, M., Katayama, S., Kawakami, H., Nakamura, S., Nishimura, M., Akiguchi, I., et al. (1994). CAG expansions in a novel gene for Machado-Joseph Disease at chromosome 14q32.1. *Nat. Genet.* 8, 221–228.
244. Takiyama, Y., Oyanagi, S., Kawashima, S., Sakamoto, H., Saito, K., Yoshida, M., Tsuji, S., Mizuno, Y., and Nishizawa, M. (1994). A clinical and pathologic study of a large Japanese family with Machado-Joseph disease tightly linked to the DNA markers on chromosome 14q. *Neurology* 44, 1302–1308.
245. Bürk, K., Zühlke, C., König, I.R., Ziegler, A., Schwinger, E., Globas, C., Dichgans, J., and Hellenbroich, Y. (2004). Spinocerebellar ataxia type 5: clinical and molecular genetic features of a German kindred. *Neurology* 62, 327–329.
246. David, G., Abbas, N., Stevanin, G., Dürr, A., Yvert, G., Cancel, G., Weber, C., Imbert, G., Saudou, F., Antoniou, E., et al. (1997). Cloning of the SCA7 gene reveals a highly unstable CAG repeat expansion. *Nat. Genet.* 17, 65–70.
247. Koob, M.D., Moseley, M.L., Schut, L.J., Benzow, K. a, Bird, T.D., Day, J.W., and Ranum, L.P. (1999). An untranslated CTG expansion causes a novel form of spinocerebellar ataxia (SCA8). *Nat. Genet.* 21, 379–384.

248. Matsuura, T., Yamagata, T., Burgess, D.L., Rasmussen, A., Grewal, R.P., Watase, K., Khajavi, M., McCall, A.E., Davis, C.F., Zu, L., et al. (2000). Large expansion of the ATTCT pentanucleotide repeat in spinocerebellar ataxia type 10. *Nat. Genet.* *26*, 191–194.
249. Zu, L., Figueroa, K.P., Grewal, R., and Pulst, S.M. (1999). Mapping of a new autosomal dominant spinocerebellar ataxia to chromosome 22. *Am. J. Hum. Genet.* *64*, 594–599.
250. Grewal, R.P., Tayag, E., Figueroa, K.P., Zu, L., Durazo, A., Nunez, C., and Pulst, S.M. (1998). Clinical and genetic analysis of a distinct autosomal dominant spinocerebellar ataxia. *Neurology* *51*, 1423–1426.
251. Houlden, H., Johnson, J., Gardner-Thorpe, C., Lashley, T., Hernandez, D., Worth, P., Singleton, A.B., Hilton, D. a, Holton, J., Revesz, T., et al. (2007). Mutations in TTBK2, encoding a kinase implicated in tau phosphorylation, segregate with spinocerebellar ataxia type 11. *Nat. Genet.* *39*, 1434–1436.
252. Holmes, S.E., O’Hearn, E.E., McInnis, M.G., Gorelick-Feldman, D.A., Kleiderlein, J.J., Callahan, C., Kwak, N.G., Ingersoll-Ashworth, R.G., Sherr, M., Sumner, A.J., et al. (1999). Expansion of a novel CAG trinucleotide repeat in the 5’ region of PPP2R2B is associated with SCA12. *Nat. Genet.* *23*, 391–392.
253. Waters, M.F., Fee, D., Figueroa, K.P., Nolte, D., Müller, U., Advincula, J., Coon, H., Evidente, V.G., and Pulst, S.M. (2005). An autosomal dominant ataxia maps to 19q13: Allelic heterogeneity of SCA13 or novel locus? *Neurology* *65*, 1111–1113.
254. Chen, D.-H., Brkanac, Z., Christophe Verlinde, L.M.J., Tan, X.-J., Bylenok, L., Nochlin, D., Matsushita, M., Lipe, H., Wolff, J., Fernandez, M., et al. (2003). Missense Mutations in the Regulatory Domain of PKC γ : A New Mechanism for Dominant Nonepisodic Cerebellar Ataxia. *Am. J. Hum. Genet.* *72*, 839–849.
255. Storey, E., Gardner, R.J., Knight, M.A., Kennerson, M.L., Tuck, R.R., Forrest, S.M., and Nicholson, G.A. (2001). A new autosomal dominant pure cerebellar ataxia. *Neurology* *57*, 1913–1915.
256. van de Leemput, J., Chandran, J., Knight, M.A., Holtzclaw, L.A., Scholz, S., Cookson, M.R., Houlden, H., Gwinn-Hardy, K., Fung, H.-C., Lin, X., et al. (2007). Deletion at ITPR1 Underlies Ataxia in Mice and Spinocerebellar Ataxia 15 in Humans. *PLoS Genet.* *3*, e108.
257. Nakamura, K. (2001). SCA17, a novel autosomal dominant cerebellar ataxia caused by an expanded polyglutamine in TATA-binding protein. *Hum. Mol. Genet.* *10*, 1441–1448.
258. Verbeek, D.S., Schelhaas, J.H., Ippel, E.F., Beemer, F. a, Pearson, P.L., and Sinke, R.J. (2002). Identification of a novel SCA locus (SCA19) in a Dutch autosomal dominant cerebellar ataxia family on chromosome region 1p21-q21. *Hum. Genet.* *111*, 388–393.
259. Chung, M.Y., Lu, Y.C., Cheng, N.C., and Soong, B.W. (2003). A novel autosomal dominant spinocerebellar ataxia (SCA22) linked to chromosome 1p21-q23. *Brain* *126*, 1293–1299.
260. Knight, M.A., Hernandez, D., Diede, S.J., Dauwerse, H.G., Rafferty, I., van de Leemput, J., Forrest, S.M., Gardner, R.J.M., Storey, E., van Ommen, G.-J.B., et al. (2008). A duplication at chromosome 11q12.2-11q12.3 is associated with spinocerebellar ataxia type 20. *Hum. Mol. Genet.* *17*, 3847–3853.
261. Knight, M.A., Gardner, R.J.M., Bahlo, M., Matsuura, T., Dixon, J.A., Forrest, S.M., and Storey, E. (2004). Dominantly inherited ataxia and dysphonia with dentate calcification: spinocerebellar ataxia type 20. *Brain* *127*, 1172–1181.
262. Delplanque, J., Devos, D., Huin, V., Genet, A., Sand, O., Moreau, C., Goizet, C., Charles, P., Anheim, M., Monin, M.L., et al. (2014). TMEM240 mutations cause spinocerebellar ataxia 21 with mental retardation and severe cognitive impairment. *Brain* *137*, 2657–2663.


263. Verbeek, D.S., van de Warrenburg, B.P., Wesseling, P., Pearson, P.L., Kremer, H.P., and Sinke, R.J. (2004). Mapping of the SCA23 locus involved in autosomal dominant cerebellar ataxia to chromosome region 20p13-12.3. *Brain* 127, 2551–2557.
264. Yu, G.-Y., Howell, M.J., Roller, M.J., Xie, T.-D., and Gomez, C.M. (2005). Spinocerebellar ataxia type 26 maps to chromosome 19p13.3 adjacent to SCA6. *Ann. Neurol.* 57, 349–354.
265. Hekman, K.E., Yu, G.-Y., Brown, C.D., Zhu, H., Du, X., Gervin, K., Undlien, D.E., Peterson, A., Stevanin, G., Clark, H.B., et al. (2012). A conserved eEF2 coding variant in SCA26 leads to loss of translational fidelity and increased susceptibility to proteostatic insult. *Hum. Mol. Genet.* 21, 5472–5483.
266. van Swieten, J.C., Brusse, E., de Graaf, B.M., Krieger, E., van de Graaf, R., de Koning, I., Maat-Kievit, A., Leegwater, P., Dooijes, D., Oostra, B.A., et al. (2003). A mutation in the fibroblast growth factor 14 gene is associated with autosomal dominant cerebellar ataxia [corrected]. *Am. J. Hum. Genet.* 72, 191–199.
267. Cagnoli, C., Mariotti, C., Taroni, F., Seri, M., Brussino, A., Michielotto, C., Grisoli, M., Di Bella, D., Migone, N., Gellera, C., et al. (2006). SCA28, a novel form of autosomal dominant cerebellar ataxia on chromosome 18p11.22-q11.2. *Brain* 129, 235–242.
268. Di Bella, D., Lazzaro, F., Brusco, A., Plumari, M., Battaglia, G., Pastore, A., Finardi, A., Cagnoli, C., Tempia, F., Frontali, M., et al. (2010). Mutations in the mitochondrial protease gene AFG3L2 cause dominant hereditary ataxia SCA28. *Nat. Genet.* 42, 313–321.
269. Dudding, T.E., Friend, K., Schofield, P.W., Lee, S., Wilkinson, I.A., and Richards, R.I. (2004). Autosomal dominant congenital non-progressive ataxia overlaps with the SCA15 locus. *Neurology* 63, 2288–2292.
270. Hirano, R., Takashima, H., Okubo, R., Tajima, K., Okamoto, Y., Ishida, S., Tsuruta, K., Arisato, T., Arata, H., Nakagawa, M., et al. (2004). Fine mapping of 16q-linked autosomal dominant cerebellar ataxia type III in Japanese families. *Neurogenetics* 5, 215–221.
271. Sato, N., Amino, T., Kobayashi, K., Asakawa, S., Ishiguro, T., Tsunemi, T., Takahashi, M., Matsuura, T., Flanigan, K.M., Iwasaki, S., et al. (2009). Spinocerebellar ataxia type 31 is associated with “inserted” penta-nucleotide repeats containing (TGGAA)*n*. *Am. J. Hum. Genet.* 85, 544–557.
272. Cadieux-Dion, M., Turcotte-Gauthier, M., Noreau, A., Martin, C., Meloche, C., Gravel, M., Drouin, C.A., Rouleau, G. a, Nguyen, D.K., and Cossette, P. (2014). Expanding the Clinical Phenotype Associated With ELOVL4 Mutation: Study of a Large French-Canadian Family With Autosomal Dominant Spinocerebellar Ataxia and Erythrokeratoderma. *JAMA Neurol.* 4–9.
273. Wang, J.L.J., Yang, X., Xia, K., Hu, Z.M., Weng, L., Jin, X., Jiang, H., Zhang, P., Shen, L., Guo, J.F., et al. (2010). TGM6 identified as a novel causative gene of spinocerebellar ataxias using exome sequencing. *Brain* 133, 3510–3518.
274. Kobayashi, H., Abe, K., Matsuura, T., Ikeda, Y., Hitomi, T., Akechi, Y., Habu, T., Liu, W., Okuda, H., and Koizumi, A. (2011). Expansion of intronic GGCCTG hexanucleotide repeat in NOP56 causes SCA36, a type of spinocerebellar ataxia accompanied by motor neuron involvement. *Am. J. Hum. Genet.* 89, 121–130.
275. Di Gregorio, E., Borroni, B., Giorgio, E., Lacerenza, D., Ferrero, M., Lo Buono, N., Ragusa, N., Mancini, C., Gausson, M., Calcia, A., et al. (2014). ELOVL5 Mutations Cause Spinocerebellar Ataxia 38. *Am. J. Hum. Genet.* 95, 209–217.
276. Tsoi, H., Yu, A.C.S., Chen, Z.S., Ng, N.K.N., Chan, A.Y.Y., Yuen, L.Y.P., Abrigo, J.M., Tsang, S.Y., Tsui, S.K.W., Tong, T.M.F., et al. (2014). A novel missense mutation in CCDC88C activates the JNK pathway and causes a dominant form of spinocerebellar ataxia. *J. Med. Genet.* 1–6.

277. Ozelius, L.J., Hewett, J.W., Page, C.E., Bressman, S.B., Kramer, P.L., Shalish, C., de Leon, D., Brin, M.F., Raymond, D., Corey, D.P., et al. (1997). The early-onset torsion dystonia gene (DYT1) encodes an ATP-binding protein. *Nat. Genet.* 17, 40–48.
278. Ozelius, L.J., Kramer, P.L., de Leon, D., Risch, N., Bressman, S.B., Schuback, D.E., Brin, M.F., Kwiatkowski, D.J., Burke, R.E., and Gusella, J.F. (1992). Strong allelic association between the torsion dystonia gene (DYT1) and loci on chromosome 9q34 in Ashkenazi Jews. *Am. J. Hum. Genet.* 50, 619–628.
279. Charlesworth, G., Angelova, P.R., Bartolomé-Robledo, F., Ryten, M., Trabzuni, D., Stamelou, M., Abramov, A.Y., Bhatia, K.P., and Wood, N.W. (2015). Mutations in HPCA Cause Autosomal-Recessive Primary Isolated Dystonia. *Am. J. Hum. Genet.* 96, 657–665.
280. Haberman, G., Schmitt, I., Köhler, A., Peters, U., Rider, S., Chelly, J., Terwilliger, J.D., Monaco, A.P., and Müller, U. (1995). Assignment of the dystonia-parkinsonism syndrome locus, DYT3, to a small region within a 1.8-Mb YAC contig of Xq13.1. *Am. J. Hum. Genet.* 57, 644–650.
281. Makino, S., Kaji, R., Ando, S., Tomizawa, M., Yasuno, K., Goto, S., Matsumoto, S., Tabuena, M.D., Maranon, E., Dantes, M., et al. (2007). Reduced neuron-specific expression of the TAF1 gene is associated with X-linked dystonia-parkinsonism. *Am. J. Hum. Genet.* 80, 393–406.
282. Hersheson, J., Mencacci, N.E., Davis, M., MacDonald, N., Trabzuni, D., Ryten, M., Pittman, A., Paudel, R., Kara, E., Fawcett, K., et al. (2013). Mutations in the autoregulatory domain of β -tubulin 4a cause hereditary dystonia. *Ann. Neurol.* 73, 546–553.
283. Nygaard, T.G., Wilhelmsen, K.C., Risch, N.J., Brown, D.L., Trugman, J.M., Gilliam, T.C., Fahn, S., and Weeks, D.E. (1993). Linkage mapping of dopa-responsive dystonia (DRD) to chromosome 14q. *Nat. Genet.* 5, 386–391.
284. Almasy, L., Bressman, S.B., Raymond, D., Kramer, P.L., Greene, P.E., Heiman, G.A., Ford, B., Yount, J., de Leon, D., Chouinard, S., et al. (1997). Idiopathic torsion dystonia linked to chromosome 8 in two Mennonite families. *Ann. Neurol.* 42, 670–673.
285. Saunders-Pullman, R., Raymond, D., Senthil, G., Kramer, P., Ohmann, E., Deligtisch, A., Shanker, V., Greene, P., Tabamo, R., Huang, N., et al. (2007). Narrowing the DYT6 dystonia region and evidence for locus heterogeneity in the Amish-Mennonites. *Am. J. Med. Genet. A* 143A, 2098–2105.
286. Fuchs, T., Gavarini, S., Saunders-Pullman, R., Raymond, D., Ehrlich, M.E., Bressman, S.B., and Ozelius, L.J. (2009). Mutations in the THAP1 gene are responsible for DYT6 primary torsion dystonia. *Nat. Genet.* 41, 286–288.
287. Fink, J.K., Rainer, S., Wilkowsky, J., Jones, S.M., Kume, A., Hedera, P., Albin, R., Mathay, J., Girbach, L., Varvil, T., et al. (1996). Paroxysmal dystonic choreoathetosis: tight linkage to chromosome 2q. *Am. J. Hum. Genet.* 59, 140–145.
288. Rainier, S., Thomas, D., Tokarz, D., Ming, L., Bui, M., Plein, E., Zhao, X., Lemons, R., Albin, R., Delaney, C., et al. (2004). Myofibrillogenesis regulator 1 gene mutations cause paroxysmal dystonic choreoathetosis. *Arch. Neurol.* 61, 1025–1029.
289. Auburger, G., Ratzlaff, T., Lunke, A., Nelles, H.W., Leube, B., Binkofski, F., Kugel, H., Heindel, W., Seitz, R., Benecke, R., et al. (1996). A gene for autosomal dominant paroxysmal choreoathetosis/spasticity (CSE) maps to the vicinity of a potassium channel gene cluster on chromosome 1p, probably within 2 cM between D1S443 and D1S197. *Genomics* 31, 90–94.

290. Weber, Y.G., Kamm, C., Suls, A., Kempfle, J., Kotschet, K., Schüle, R., Wuttke, T.V., Maljevic, S., Liebrich, J., Gasser, T., et al. (2011). Paroxysmal choreoathetosis / spasticity (DYT9) is caused by a GLUT1 defect. *Neurology* 959–964.
291. Tomita, H., Nagamitsu, S., Wakui, K., Fukushima, Y., Yamada, K., Sadamatsu, M., Masui, A., Konishi, T., Matsuishi, T., Aihara, M., et al. (1999). Paroxysmal Kinesigenic Choreoathetosis Locus Maps to Chromosome 16p11.2-q12.1. *Am. J. Hum. Genet.* 65, 1688–1697.
292. Chen, W.-J., Lin, Y., Xiong, Z.-Q., Wei, W., Ni, W., Tan, G.-H., Guo, S.-L., He, J., Chen, Y.-F., Zhang, Q.-J., et al. (2011). Exome sequencing identifies truncating mutations in PRRT2 that cause paroxysmal kinesigenic dyskinesia. *Nat. Genet.* 43, 1252–1255.
293. Wang, J.-L., Cao, L., Li, X.-H., Hu, Z.-M., Li, J.-D., Zhang, J.-G., Liang, Y., San-A, Li, N., Chen, S.-Q., et al. (2011). Identification of PRRT2 as the causative gene of paroxysmal kinesigenic dyskinesias. *Brain* 134, 3493–3501.
294. Nygaard, T.G., Raymond, D., Chen, C., Nishino, I., Greene, P.E., Jennings, D., Heiman, G.A., Klein, C., Saunders-Pullman, R.J., Kramer, P., et al. (1999). Localization of a gene for myoclonus-dystonia to chromosome 7q21-q31. *Ann. Neurol.* 46, 794–798.
295. Zimprich, A., Grabowski, M., Asmus, F., Naumann, M., Berg, D., Bertram, M., Scheidtmann, K., Kern, P., Winkelmann, J., Müller-Myhsok, B., et al. (2001). Mutations in the gene encoding epsilon-sarcoglycan cause myoclonus-dystonia syndrome. *Nat. Genet.* 29, 66–69.
296. Kramer, P.L., Mineta, M., Klein, C., Schilling, K., de Leon, D., Farlow, M.R., Breakefield, X.O., Bressman, S.B., Dobyns, W.B., Ozelius, L.J., et al. (1999). Rapid-onset dystonia-parkinsonism: linkage to chromosome 19q13. *Ann. Neurol.* 46, 176–182.
297. De Carvalho Aguiar, P., Sweadner, K.J., Penniston, J.T., Zaremba, J., Liu, L., Caton, M., Linazasoro, G., Borg, M., Tijssen, M. a J., Bressman, S.B., et al. (2004). Mutations in the Na⁺/K⁺-ATPase α 3 gene ATP1A3 are associated with rapid-onset dystonia parkinsonism. *Neuron* 43, 169–175.
298. Camargos, S., Scholz, S., Simón-Sánchez, J., Paisán-Ruiz, C., Lewis, P., Hernandez, D., Ding, J., Gibbs, J.R., Cookson, M.R., Bras, J., et al. (2008). DYT16, a novel young-onset dystonia-parkinsonism disorder: identification of a segregating mutation in the stress-response protein PRKRA. *Lancet. Neurol.* 7, 207–215.
299. Weber, Y.G., Storch, A., Wuttke, T.V., Brockmann, K., Kempfle, J., Maljevic, S., Margari, L., Kamm, C., Schneider, S.A., Huber, S.M., et al. (2008). GLUT1 mutations are a cause of paroxysmal exertion-induced dyskinesias and induce hemolytic anemia by a cation leak. *J. Clin. Invest.* 118, 2157–2168.
300. Mencacci, N.E., Rubio-Agusti, I., Zdebik, A., Asmus, F., Ludtmann, M.H.R., Ryten, M., Plagnol, V., Hauser, A.-K., Bandres-Ciga, S., Bettencourt, C., et al. (2015). A Missense Mutation in KCTD17 Causes Autosomal Dominant Myoclonus-Dystonia. *Am. J. Hum. Genet.* 96, 938–947.
301. Bonafé, L., Thöny, B., Penzien, J.M., Czarnecki, B., and Blau, N. (2001). Mutations in the sepiapterin reductase gene cause a novel tetrahydrobiopterin-dependent monoamine-neurotransmitter deficiency without hyperphenylalaninemia. *Am. J. Hum. Genet.* 69, 269–277.
302. Xiao, J., Uitti, R.J., Zhao, Y., Vemula, S.R., Perlmutter, J.S., Wszolek, Z.K., Maraganore, D.M., Auburger, G., Leube, B., Lehnhoff, K., et al. (2012). Mutations in CIZ1 cause adult onset primary cervical dystonia. *Ann. Neurol.* 71, 458–469.



Supplementary Figure 1. Gene co-expression network based on known SCA genes
Red dots are the input genes (known SCA genes). Yellow to green dots represent co-expressed genes, darker colored dots have more connections. The thickness of the line indicates the strength of co-expression.



Supplementary Figure 2. Gene co-expression network based on known dystonia genes
Red dots are the input genes (known dystonia genes). Yellow to green dots represent coexpressed genes, darker colored dots have more connections. The thickness of the line indicates the strength of co-expression.

