

University of Groningen

Red knot habits

Oudman, Thomas

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Oudman, T. (2017). *Red knot habits: An optimal foraging perspective on tidal life at Banc d'Arguin*.
Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

References



A

- Abrams, P.A. & Matsuda, H. (2003) Population dynamical consequences of reduced predator switching at low total prey densities. *Population Ecology*, **45**, 175–185.
- Ahmedou Salem, M.V., van der Geest, M., Piersma, T., Saoud, Y. & van Gils, J.A. (2014) Seasonal changes in mollusc abundance in a tropical intertidal ecosystem, Banc d'Arguin (Mauritania): a test of the 'shorebird depletion' hypothesis. *Estuarine, Coastal and Shelf Science*, **136**, 26–34.
- Alatalo, R.V. & Mappes, J. (1996) Tracking the evolution of warning signals. *Nature*, **382**, 708–710.
- Altenburg, W., Engelmoer, M., Mes, R. & Piersma, T. (1982) *Wintering Waders on the Banc d'Arguin, Mauritania. Report of the Netherlands Ornithological Expedition 1980*. Stichting Veth tot steun aan Waddenonderzoek, Leiden, The Netherlands.
- Anderson, A.E. (1995) Metabolic responses to sulfur in Lucinid bivalves. *American Zoologist*, **35**, 121–131.
- Aplin, L.M., Sheldon, B.C. & Morand-Ferron, J. (2013) Milk bottles revisited: social learning and individual variation in the blue tit, *Cyanistes caeruleus*. *Animal Behaviour*, **85**, 1225–1232.
- Araújo, M.S., Bolnick, D.I. & Layman, C.A. (2011) The ecological causes of individual specialisation. *Ecology Letters*, **14**, 948–958.
- Arnold, T.W. (2010) Uninformative parameters and model selection using Akaike's information criterion. *Journal of Wildlife Management*, **74**, 1175–1178.
- Avgar, T., Mosser, A., Brown, G.S. & Fryxell, J.M. (2013) Environmental and individual drivers of animal movement patterns across a wide geographical gradient. *The Journal of Animal Ecology*, **82**, 96–106.
- Avital, E. & Jablonka, E. (2000) *Animal traditions: Behavioural Inheritance in Evolution*. Cambridge University Press, Cambridge.

B

- Baldwin, J.M. (1896) A new factor in evolution. *American Naturalist*, **30**, 441–451.
- Bagarinao, T. (1992) Sulfide as an environmental factor and toxicant - tolerance and adaptations in aquatic organisms. *Aquatic Toxicology*, **24**, 21–62.
- Baker, A.J., Gonzalez, P.M., Piersma, T., Niles, L.J., de Lima S. do Nascimento, I., Atkinson, P.W., Clark, N.A., Minton, C.D.T., Peck, M.K. & Aarts, G. (2004) Rapid population decline in red knots: fitness consequences of decreased refuelling rates and late arrival in Delaware Bay. *Proceedings of the Royal Society of London B*, **271**, 875–882.
- Barnett, C.A., Skelhorn, J., Bateson, M. & Rowe, C. (2012) Educated predators make strategic decisions to eat defended prey according to their toxin content. *Behavioral Ecology*, **23**, 418–424.
- Barraquand, F. & Benhamou, S. (2008) Animals movements in heterogeneous landscapes: identifying profitable places and homogeneous movement bouts. *Ecology*, **89**, 3336–3348.
- Barnett, C.A., Bateson, M. & Rowe, C. (2007) State-dependent decision making: educated predators strategically trade off the costs and benefits of consuming aposematic prey. *Behavioral Ecology*, **18**, 645–651.
- Barnett, C.A., Skelhorn, J., Bateson, M. & Rowe, C. (2012) Educated predators make strategic decisions to eat defended prey according to their toxin content. *Behavioral Ecology*, **23**, 418–424.
- Bartumeus, F., Campos, D., Ryu, W.S., Lloret-Cabot, R., Méndez, V. & Catalan, J. (2016) Foraging success under uncertainty: search tradeoffs and optimal space use. *Ecology Letters*, **19**, 1299–1313.
- Bastille-Rousseau, G., Gibbs, J.P., Yackulic, C.B., Frair, J.L., Cabrera, F., Rousseau, L.-P., Wikelski, M., Kümmeth, F. & Blake, S. (2017) Animal movement in the absence of predation: environmental drivers of movement strategies in a partial migration system. *Oikos*, doi:10.1111/oik.03928
- Bates, D., Maechler, M., Bolker, B. & Walker, S. (2015) Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, **67**, 1–48.
- Bateson, P. (1979) How do sensitive periods arise and what are they for? *Animal Behaviour*, **27**, 470–486.
- Bateson, P. (2005) The return of the whole organism. *Journal of biosciences*, **30**, 31–39.
- Battley, P.F., Rogers, D.I., van Gils, J.A., Piersma, T., Hassell, C.J., Boyle, A. & Hong-Yan, Y. (2005) How do red knots *Calidris canutus* leave Northwest Australia in May and reach the breeding grounds in June? Predictions of stopover times, fuelling rates and prey quality in the Yellow Sea. *Journal of Avian Biology*, **36**, 494–500.

- Beckmann, C. & Shine, R. (2011) Toad's tongue for breakfast: exploitation of a novel prey type, the invasive cane toad, by scavenging raptors in tropical Australia. *Biological Invasions*, **13**, 1447–1455.
- Belovsky, G.E. (1978) Diet optimization in a generalist herbivore: the moose. *Theoretical Population Biology*, **14**, 105–134.
- Belovsky, G.E. (1984) Herbivore optimal foraging - a comparative test of 3 models. *American Naturalist*, **124**, 97–115.
- Belovsky, G.E. & Schmitz, O.J. (1994) Plant defenses and optimal foraging by mammalian herbivores. *Journal of Mammalogy*, **75**, 816–832.
- Bergquist, D.C., Ward, T., Cordes, E.E., McNelis, T., Howlett, S., Kosoff, R., Hourdez, S., Carney, R. & Fisher, C.R. (2003) Community structure of vestimentiferan-generated habitat islands from Gulf of Mexico cold seeps. *Journal of Experimental Marine Biology and Ecology*, **289**, 197–222.
- Beukema, J.J. (1976) Biomass and species richness of the macro-benthic animals living on the tidal flats of the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, **10**, 236–261.
- Bijleveld, A.I. (2015) *Untying the Knot: Mechanistically Understanding the Interactions between Social Foragers and their Prey*. PhD thesis, University of Groningen.
- Bijleveld, A.I., Egas, M., van Gils, J.A. & Piersma, T. (2010) Beyond the information centre hypothesis: communal roosting for information on food, predators, travel companions and mates? *Oikos*, **119**, 277–285.
- Bijleveld, A.I., Folmer, E.O. & Piersma, T. (2012) Experimental evidence for cryptic interference among socially foraging shorebirds. *Behavioral Ecology*, **23**, 806–814.
- Bijleveld, A.I., MacCurdy, R.B., Chan, Y.-C., Penning, E., Gabrielson, R.M., Cluderay, J., Spaulding, E., Dekkinga, A., Holthuijsen, S., ten Horn, J., Brugge, M., van Gils, J.A., Winkler, D. & Piersma, T. (2016) Understanding spatial distributions: negative density-dependence in prey causes predators to trade-off prey quantity with quality. *Proceedings of the Royal Society B: Biological Sciences*, **283**, 20151557.
- Bijleveld, A.I., Massourakis, G., van der Marel, A., Dekkinga, A., Spaans, B., van Gils, J.A. & Piersma, T. (2014) Personality drives physiological adjustments and is not related to survival. *Proceedings of the Royal Society B: Biological Sciences*, **281**, 20133135.
- Bijleveld, A.I., Twietmeyer, S., Piechocki, J., van Gils, J.A. & Piersma, T. (2015a) Natural selection by pulsed predation: survival of the thickest. *Ecology*, **96**, 1943–1956.
- Bijleveld, A.I., van Gils, J.A., Jouta, J. & Piersma, T. (2015b) Benefits of foraging in small groups: an experimental study on public information use in red knots *Calidris canutus*. *Behavioural Processes*, **117**, 74–81.
- Bijleveld, A.I., van Gils, J.A., van der Meer, J., Dekkinga, A., Kraan, C., van der Veer, H.W. & Piersma, T. (2012) Designing a benthic monitoring programme with multiple conflicting objectives. *Methods in Ecology and Evolution*, **3**, 526–536.
- Bijlsma, R.G. (1990) Predation by large falcons on wintering waders on the Banc d'Arguin, Mauritania. *Ardea*, **78**, 82.
- Bolker, B., Holyoak, M., Křivan, V., Rowe, L. & Schmitz, O. (2003) Connecting theoretical and empirical studies of trait-mediated interactions. *Ecology*, **84**, 1101–1114.
- Bolnick, D.I., Amarasekare, P., Araújo, M.S., Bürger, R., Levine, J.M., Novak, M., Rudolf, V.H.W., Schreiber, S.J., Urban, M.C. & Vasseur, D.A. (2011) Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, **26**, 183–192.
- Bolnick, D.I., Svanbäck, R., Fordyce, J.A., Yang, L.H., Davis, J.M., Darrin Hulsey, C. & Forister, M.L. (2003) The ecology of individuals: incidence and implications of individual specialization. *American Naturalist*, **161**, 1–28.
- Bonsall, M.B. & Hassell, M.P. (1997) Apparent competition structures ecological assemblages. *Nature*, **388**, 371–373.
- Börger, L., Dalziel, B.D. & Fryxell, J.M. (2008) Are there general mechanisms of animal home range behaviour? A review and prospects for future research. *Ecology Letters*, **11**, 637–650.
- Buehler, D.M. & Baker, A.J. (2005) Population divergence times and historical demography in red knots and dunlins. *Condor*, **107**, 497–513.
- Buehler, D.M., Baker, A.J. & Piersma, T. (2006) Reconstructing palaeoflyways of the late Pleistocene and early Holocene red knot *Calidris canutus*. *Ardea*, **94**, 485.

- Buehler, D.M. & Piersma, T. (2008) Travelling on a budget: predictions and ecological evidence for bottlenecks in the annual cycle of long-distance migrants. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **363**, 247–266.
- Bulla, M., Oudman, T., Bijleveld, A.I., Piersma, T. & Kyriacou, C.P. (in review) Living with the moon and the tides. Burghardt, G.M. & Hess, E.H. (1966) Food imprinting in the snapping turtle, *Chelydra serpentina*. *Science*, **151**, 108–109.
- Burnham, K.P. & Anderson, D.R. (2002) *Model Selection and Multimodel Inference: a Practical Information-theoretic Approach*. Springer, Heidelberg.
- Butchart, S.H.M., Walpole, M., Collen, B., van Strien, A., Scharlemann, J.P.W., Almond, R.E.A. et al. (2010) Global biodiversity: indicators of recent declines. *Science*, **328**, 1164–1168.

C

- Calleja, M.L., Marbà, N. & Duarte, C.M. (2007) The relationship between seagrass (*Posidonia oceanica*) decline and sulfide porewater concentration in carbonate sediments. *Estuarine, Coastal and Shelf Science*, **73**, 583–588.
- Carney, R.S. (1994) Consideration of the oasis analogy for chemosynthetic communities at Gulf of Mexico hydrocarbon vents. *Geo-Marine Letters*, **14**, 149–159.
- Caro, T. (2005) *Antipredator Defenses in Birds and Mammals*. University of Chicago Press, Chicago.
- Cary, S.C., Vetter, R.D. & Felbeck, H. (1989) Habitat characterization and nutritional strategies of the endosymbiont-bearing bivalve *Lucinoma aequizonata*. *Marine Ecology Progress Series*, **55**, 31–45.
- Charnov, E.L. (1976) Optimal foraging, the marginal value theorem. *Theoretical Population Biology*, **9**, 129–136.
- Colwell, M.A. (2000) A review of territoriality in non-breeding shorebirds (Charadrii). *Wader Study Group Bulletin*, **93**, 58–66.
- Compton, T.J., Holthuijsen, S., Koolhaas, A., Dekkinga, A., ten Horn, J., Smith, J., Galama, Y., Brugge, M., van der Wal, D., van der Meer, J., van der Veer, H.W. & Piersma, T. (2013) Distinctly variable mudscapes: distribution gradients of intertidal macrofauna across the Dutch Wadden Sea. *Journal of Sea Research*, **82**, 103–116.
- Conklin, J.R. & Colwell, M.A. (2007) Diurnal and nocturnal roost site fidelity of dunlin (*Calidris alpina pacifica*) at Humboldt Bay, California. *Auk*, **124**, 677–689.
- Cordes, E.E., Becker, E.L. & Fisher, C.R. (2010) Temporal shift in nutrient input to cold-seep food webs revealed by stable-isotope signatures of associated communities. *Limnology and Oceanography*, **55**, 2537–2548.
- Cox, Laura M., Yamanishi, S., Sohn, J., Alekseyenko, Alexander V., Leung, Jacqueline M., Cho, I., Kim, Sungheon G., Li, H., Gao, Z., Mahana, D., Zárate Rodriguez, Jorge G., Rogers, Arlin B., Robine, N., Loke, P.n. & Blaser, Martin J. (2014) Altering the intestinal microbiota during a critical developmental window has lasting metabolic consequences. *Cell*, **158**, 705–721.
- Curley, E.A.M., Rowley, H.E. & Speed, M.P. (2015) A field demonstration of the costs and benefits of group living to edible and defended prey. *Biology Letters*, **11**, 20150152.

D

- Dall, S.R.X., Bell, A.M., Bolnick, D.I. & Ratnieks, F.L.W. (2012) An evolutionary ecology of individual differences. *Ecology Letters*, **15**, 1189–1198.
- Danchin, É., Giraldeau, L.-A., Valone, T.J. & Wagner, R.H. (2004) Public information: from nosy neighbors to cultural evolution. *Science*, **305**, 487–491.
- Darmaillacq, A.-S., Chichery, R. & Dickel, L. (2006) Food imprinting, new evidence from the cuttlefish *Sepia officinalis*. *Biology Letters*, **2**, 345–347.
- Darwin, C. (1859) *On the Origin of Species by Means of Natural Selection*. Murray, London.
- Darwin, C. (1881) *The Formation of Vegetable Mould through the Action of Worms with Observation of their Habits*. Murray, London.
- Davidson, N.C. & Piersma, T. (2009) Red knots *Calidris canutus*. In: *An Atlas of Wader Populations in Africa and Western Eurasia* (eds S. Delany, D. Scott, T. Dodman & D. Stroud), pp. 362–368. Wetlands International, Wageningen.

- Davidson, N.C. & Wilson, J.R. (1992) The migration system of European-wintering knots *Calidris canutus islandica*. *Wader Study Group Bulletin*, **64**, 39–51.
- Davis, J.M. & Stamps, J.A. (2004) The effect of natal experience on habitat preferences. *Trends in Ecology & Evolution*, **19**, 411–416.
- Dearing, M.D., Foley, W.J. & McLean, S. (2005) The influence of plant secondary metabolites on the nutritional ecology of herbivorous terrestrial vertebrates. *Annual Review of Ecology, Evolution, and Systematics*, **36**, 169–189.
- de Fouw, J. (2016) *Bottom-up and Top-down Forces in a Tropical Intertidal Ecosystem. The Interplay between Seagrasses, Bivalves and Birds*. PhD thesis, University of Groningen.
- de Fouw, J., Govers, L.L., van de Koppel, J., van Belzen, J., Dorigo, W., Cheikh, M.A.S., Christianen, M.J., van der Reijden, K.J., van der Geest, M. & Piersma, T. (2016) Drought, mutualism breakdown, and landscape-scale degradation of seagrass beds. *Current Biology*, **26**, 1051–1056.
- de Fouw, J., van der Heide, T., Oudman, T., Maas, L.R.M., Piersma, T. & van Gils, J.A. (2016) Structurally complex seagrass obstructs the sixth sense of a specialized avian molluscivore. *Animal Behaviour*, **115**, 55–67.
- DeGabriel, J.L., Moore, B.D., Foley, W.J. & Johnson, C.N. (2009) The effects of plant defensive chemistry on nutrient availability predict reproductive success in a mammal. *Ecology*, **90**, 711–719.
- Dekkinga, A., Dietz, M.W., Koolhaas, A. & Piersma, T. (2001) Time course and reversibility of changes in the gizzards of red knots alternately eating hard and soft food. *Journal of Experimental Biology*, **204**, 2167–2173.
- Dekkinga, A. & Piersma, T. (1993) Reconstructing diet composition on the basis of faeces in a mollusc-eating wader, the knot *Calidris canutus*. *Bird Study*, **40**, 144–156.
- Dekker, R. (1979) *Number, Growth, Biomass and Production of Organic and Calcareous Matter of Hydrobia ulvae (Gastropoda: Prosobranchia) in the Western Dutch Wadden Sea*. NIOZ-report 1979–15, Netherlands Institute for Sea Research, Texel, The Netherlands.
- Dennett, D.C. (2001) The evolution of culture. *The Monist*, **84**, 305–324.
- Desai, M. & Hales, C.N. (1997) Role of fetal and infant growth in programming metabolism in later life. *Biological reviews of the Cambridge Philosophical Society*, **72**, 329–348.
- Dhooge, A., Govaerts, W. & Yu, A.K. (2003) MATCONT: a MATLAB package for numerical bifurcation analysis of ODEs. *ACM Transactions on Mathematical Software (TOMS)*, **29**, 141–164.
- Dick, W.J., Piersma, T. & Prokosch, P. (1987) Spring migration of the Siberian Knots *Calidris canutus canutus*: results of a co-operative Wader Study Group project. *Ornis Scandinavica*, **18**, 5–16.
- Dietz, M.W., Dekkinga, A., Piersma, T. & Verhulst, S. (1999) Estimating organ size in small migrating shorebirds with ultrasonography: an intercalibration exercise. *Physiological and Biochemical Zoology*, **72**, 28–37.
- Dingemanse, N.J. & Dochtermann, N.A. (2013) Quantifying individual variation in behaviour: mixed-effect modelling approaches. *Journal of Animal Ecology*, **82**, 39–54.
- Dingemanse, N.J., Kazem, A.J.N., Réale, D. & Wright, J. (2010) Behavioural reaction norms: animal personality meets individual plasticity. *Trends in Ecology & Evolution*, **25**, 81–89.
- Distel, R.A. & Provenza, F.D. (1991) Experience early in life affects voluntary intake of blackbrush by goats. *Journal of Chemical Ecology*, **17**, 431–450.
- Druce, D.J., Brown, J.S., Kerley, G.I.H., Kotler, B.P., Mackey, R.L. & Slotow, R.O.B. (2009) Spatial and temporal scaling in habitat utilization by klipspringers (*Oreotragus oreotragus*) determined using giving-up densities. *Austral Ecology*, **34**, 577–587.
- Dubilier, N., Bergin, C. & Lott, C. (2008) Symbiotic diversity in marine animals: the art of harnessing chemosynthesis. *Nature Reviews Microbiology*, **6**, 725–740.
- Duffy, J.E. (2003) Biodiversity loss, trophic skew and ecosystem functioning. *Ecology Letters*, **6**, 680–687.
- Duijns, S., Knot, I.E., Piersma, T. & Gils, J.A. (2015) Field measurements give biased estimates of functional response parameters, but help explain foraging distributions. *Journal of Animal Ecology*, **84**, 565–575.
- E**
- Endler, J.A. & Mappes, J. (2004) Predator mixes and the conspicuousness of aposematic signals. *American Naturalist*, **163**, 532–547.

- Ens, B.J. (1983) Feeding territories in curlews (20Y revisited). *Wader Study Group Bulletin*, **39**, 49–50.
- Estes, J.A., Riedman, M.L., Staedler, M.M., Tinker, M.T. & Lyon, B.E. (2003) Individual variation in prey selection by sea otters: patterns, causes and implications. *Journal of Animal Ecology*, **72**, 144–155.
- Eklöv, P. & Svanbäck, R. (2006) Predation risk influences adaptive morphological variation in fish populations. *American Naturalist*, **167**, 440–452.

F

- Finke, D.L. & Denno, R.F. (2004) Predator diversity dampens trophic cascades. *Nature*, **429**, 407–410.
- Fisher, J. & Hinde, R.A. (1949) The opening of milk bottles by birds. *British Birds*, **42**, 57.
- Folmer, E.O. (2012) *Self-organization on Mudflats*. PhD thesis, University of Groningen.
- Folmer, E.O., van der Geest, M., Jansen, E., Olff, H., Anderson, T.M., Piersma, T. & van Gils, J.A. (2012) Seagrass-sediment feedback: an exploration using a non-recursive structural equation model. *Ecosystems*, **15**, 1380–1393.
- Fretwell, S.D. (1972) *Populations in a Seasonal Environment*. Princeton University Press, Princeton, NJ.
- Fretwell, S.D. & Lucas, H.L. (1970) On territorial behaviour and other factors influencing habitat distribution in birds. *Acta Biotheoretica*, **19**, 16–36.
- Fryxell, J.M. (1995) Aggregation and migration by grazing ungulates in relation to resources and predators. In: *Serengeti II: Dynamics, Management, and Conservation of an Ecosystem* (eds. Sinclair, A.R.E. & Arcese, P.). University of Chicago Press Chicago, 257–273.
- Fryxell, J.M., Hazell, M., Börger, L., Dalziel, B.D., Haydon, D.T., Morales, J.M., McIntosh, T. & Rosatte, R.C. (2008) Multiple movement modes by large herbivores at multiple spatiotemporal scales. *Proceedings of the National Academy of Sciences USA*, **105**, 19114–19119.
- Fryxell, J.M. & Lundberg, P. (1994) Diet choice and predator-prey dynamics. *Evolutionary Ecology*, **8**, 407–421.

G

- Gabriel, W., Luttbeg, B., Sih, A. & Tollrian, R. (2005) Environmental tolerance, heterogeneity, and the evolution of reversible plastic responses. *American Naturalist*, **166**, 339–353.
- Garcia, R., Sulkin, S. & Lopez, M. (2011) Effects on larval crabs of exposure to algal toxin via ingestion of heterotrophic prey. *Marine Biology*, **158**, 451–460.
- Gilbert, S.F. (2001) Ecological developmental biology: developmental biology meets the real world. *Developmental Biology*, **233**, 1–12.
- Gillingham, M.P. & Bunnell, F.L. (1989) Effects of learning on food selection and searching behaviour of deer. *Canadian Journal of Zoology*, **67**, 24–32.
- Gillings, S., Fuller, R.J., Sutherland, W.J. & Bollinger, E. (2005) Diurnal studies do not predict nocturnal habitat choice and site selection of European golden-plovers (*Pluvialis apricaria*) and northern lapwings (*Vanellus vanellus*). *Auk*, **122**, 1249–1260.
- Giraldeau, L.-A. & Caraco, T. (2000) *Social Foraging Theory*. Princeton University Press, Princeton, NJ.
- Gittleman, J.L. & Harvey, P.H. (1980) Why are distasteful prey not cryptic? *Nature*, **286**, 149–150.
- Goss-Custard, J.D. (1970) Feeding dispersion in some overwintering wading birds. *Social Behaviour in Birds and Mammals: Essays on the Social Ethology of Animals and Man* (ed. J.H. Crook). Academic Press, London.
- Gould, S.J. & Lewontin, R.C. (1979) The spandrels of San Marco and the Panglossian paradigm: a critique of the adaptationist programme. *Proceedings of the Royal Society of London B: Biological Sciences*, **205**, 581–598.
- Gray, H.M., Kaiser, H. & Green, D.M. (2010) Does alkaloid sequestration protect the green poison frog, *Dendrobates auratus*, from predator attacks? *Salamandra*, **46**, 235–238.
- Green, R.F. (1984) Stopping rules for optimal foragers. *American Naturalist*, **123**, 30–43.
- Greenberg, R. & Marra, P.P.e. (2005) *Birds of Two Worlds: Ecology and Evolution of Migration*. Johns Hopkins University Press, Baltimore, MD.
- Greenlees, M.J., Phillips, B.L. & Shine, R. (2010) Adjusting to a toxic invader: native Australian frogs learn not to prey on cane toads. *Behavioral Ecology*, **21**, 966–971.

- Grieshaber, M.K. & Völkel, S. (1998) Animal adaptations for tolerance and exploitation of poisonous sulfide. *Annual review of physiology*, **60**, 33–53.
- Guilford, T., Freeman, R., Boyle, D., Dean, B., Kirk, H., Phillips, R. & Perrins, C. (2011) A dispersive migration in the Atlantic puffin and its implications for migratory navigation. *PLoS ONE*, **6**, e21336.
- Gutiérrez, J.S., Dietz, M.W., Masero, J.A., Gill Jr, R.E., Dekkinga, A., Battley, P.F., Sánchez-Guzmán, J.M. & Piersma, T. (2011a) Functional ecology of saltglands in shorebirds: flexible responses to variable environmental conditions. *Functional Ecology*, **11**, 236–244.
- Gutiérrez, J.S., Masero, J.A., Abad-Gómez, J.M., Villegas, A. & Sánchez-Guzmán, J.M. (2011b) Understanding the energetic costs of living in saline environments: effects of salinity on basal metabolic rate, body mass and daily energy consumption of a long-distance migratory shorebird. *Journal of Experimental Biology*, **214**, 829–835.

H

- Hagemeijer, W., Smit, C., de Boer, P., van Dijk, A., Ravenscroft, N., van Roomen, M. & Wright, M. (2004) Wader and Waterbird at the Banc d'Arguin, Mauritania, January 2000. *WIWO Report 81*. Foundation Working Group International Waterbird and Wetland Research (WIWO), Beek-Ubbergen, The Netherlands.
- Halpin, C.G. & Rowe, C. (2010) Taste-rejection behaviour by predators can promote variability in prey defences. *Biology Letters*, **6**, 617–619.
- Hall, J. O. 2007. Sulfur. Pages 465–469 in *Veterinary Toxicology* (ed. C.G. Ramesh), Academic Press, Oxford.
- Hammond, K.A., Konarzewski, M., Torres, R.M. & Diamond, J. (1994) Metabolic ceilings under a combination of peak energy demands. *Physiological Zoology*, **67**, 1479–1506.
- Healy, S.D. & Rowe, C. (2014) Animal cognition in the wild. *Behavioural Processes*, **109, Part B**, 101–102.
- Hendriks, I.E., Basso, L., Deudero, S., Cabanellas-Reboredo, M. & Alvarez, E. (2012) Relative growth rates of the noble pen shell *Pinna nobilis* throughout ontogeny around the Balearic islands (western Mediterranean, Spain). *Journal of Shellfish Research*, **31**, 749–756.
- Hirakawa, H. (1995) Diet optimization with a nutrient or toxin constraint. *Theoretical Population Biology*, **47**, 331–346.
- Hobbs, N.T. (1990) Diet selection by generalist herbivores: a test of the linear programming model. *Behavioral Mechanisms of Food Selection* (ed. R.N. Hughes). Springer, Berlin Heidelberg New York.
- Hochman, V. & Kotler, B.P. (2006) Effects of food quality, diet preference and water on patch use by Nubian ibex. *Oikos*, **112**, 547–554.
- Holling, C.S. (1959) Some characteristics of simple types of predation and parasitism. *Canadian Entomologist*, **91**, 385–398.
- Holt, R.D. (1977) Predation, apparent competition, and the structure of prey communities. *Theoretical Population Biology*, **12**, 197–229.
- Holt, R.D. (1984) Spatial heterogeneity, indirect interactions, and the coexistence of prey species. *American Naturalist*, **124**, 377–406.
- Holt, R.D. & Kotler, B.P. (1987) Short-term apparent competition. *American Naturalist*, **130**, 412–430.
- Honkoop, P.J.C., Berghuis, E.M., Holthuijsen, S., Lavaleye, M.S.S. & Piersma, T. (2008) Molluscan assemblages of seagrass-covered and bare intertidal flats on the Banc d'Arguin, Mauritania, in relation to characteristics of sediment and organic matter. *Journal of Sea Research*, **60**, 255–263.
- Hopcraft, J.G.C., Morales, J.M., Beyer, H.L., Borner, M., Mwangomo, E., Sinclair, A.R.E., Olff, H. & Haydon, D.T. (2014) Competition, predation, and migration: individual choice patterns of Serengeti migrants captured by hierarchical models. *Ecological Monographs*, **84**, 355–372.
- Hoppitt, W. & Laland, K.N. (2013) *Social Learning: an Introduction to Mechanisms, Methods and Models*. Princeton University Press, Princeton, NJ.
- Houston, A.I. & McNamara, J.M. (1999) *Models of Adaptive Behavior: an Approach Based on State*. Cambridge University Press, Cambridge, UK.
- Hutchinson, G.E. (1959) Homage to Santa Rosalia or why are there so many kinds of animals? *American Naturalist*, **93**, 145–159.
- Huxley, J. (1932) *Problems of Relative Growth*. Methuen, London.

I

IPCC (2014) *Climate Change 2014: Synthesis Report. Contribution of working groups I, II and III to the fifth assessment report of the intergovernmental panel on climate change* (eds: Core Writing Team, R.K. Pachauri and L.A. Meyer.). IPCC, Geneva.

J

- Jablonska, E. & Lamb, M.J. (2005) *Evolution in Four Dimensions: Genetic, Behavioral, and Symbolic Variation in the History of Life*. MIT press, Cambridge, Ms.
- Jablonska, E. & Lamb, M.J. (2007) Précis of Evolution in Four Dimensions. *Behavioral and Brain Sciences*, **30**, 353–365.
- Jensen, K., Mayntz, D., Toft, S., Clissold, F.J., Hunt, J., Raubenheimer, D. & Simpson, S.J. (2012) Optimal foraging for specific nutrients in predatory beetles. *Proceedings of the Royal Society B: Biological Sciences*, **279**, 2212–2218.
- Jeschke, J.M., Kopp, M. & Tollrian, R. (2002) Predator functional responses: discriminating between handling and digesting prey. *Ecological Monographs*, **72**, 95–112.
- Jeschke, J.M. & Tollrian, R. (2000) Density-dependent effects of prey defences. *Oecologia*, **123**, 391–396.
- Johnson, M., Diouris, M. & Le Pennec, M. (1994) Endosymbiotic bacterial contribution in the carbon nutrition of *Loripes lucinalis* (Mollusca: Bivalvia). *Symbiosis*, **17**, 1–13.
- Johnson, W.C., Thomas, L. & Adkisson, C.S. (1993) Dietary circumvention of acorn tannins by blue jays. *Oecologia*, **94**, 159–164.
- Jørgensen, B.B. (1982) Mineralization of organic matter in the sea bed - the role of sulphate reduction. *Nature*, **296**, 643–645.

K

- Kandylis, K. (1984) Toxicology of sulfur in ruminants: review. *Journal of Dairy Science*, **67**, 2179–2187.
- Katsanevakis, S., Thessalou-Legaki, M., Karlou-Riga, C., Lefkaditou, E., Dimitriou, E. & Verriopoulos, G. (2007) Information-theory approach to allometric growth of marine organisms. *Marine Biology*, **151**, 949–959.
- Kays, R., Crofoot, M.C., Jetz, W. & Wikelski, M. (2015) Terrestrial animal tracking as an eye on life and planet. *Science*, **348**, aaa2478, 1–19.
- Kicklighter, C.E., Fisher, C.R. & Hay, M.E. (2004) Chemical defense of hydrothermal vent and hydrocarbon seep organisms: a preliminary assessment using shallow-water consumers. *Marine Ecology Progress Series*, **275**, 11–19.
- Klaassen, M. & Ens, B.J. (1990) Is salt stress a problem for waders wintering on the Banc d'Arguin Mauritania? *Ardea*, **78**, 67–74.
- Kokko, H., Mappes, J. & Lindström, L. (2003) Alternative prey can change model-mimic dynamics between parasitism and mutualism. *Ecology Letters*, **6**, 1068–1076.
- Kotler, B.P. & Mitchell, W.A. (1995) The effect of costly information in diet choice. *Evolutionary Ecology*, **9**, 18–29.
- Kraan, C., Aarts, G., Piersma, T. & Dormann, C.F. (2013) Temporal variability of ecological niches: a study on intertidal macrobenthic fauna. *Oikos*, **122**, 754–760.
- Kraan, C., van Gils, J.A., Spaans, B., Dekkinga, A. & Piersma, T. (2010) Why Afro-Siberian red knots *Calidris canutus canutus* have stopped staging in the western Dutch Wadden Sea during southward migration. *Ardea*, **98**, 155–160.
- Kraan, C., van der Meer, J., Dekkinga, A. & Piersma, T. (2009a) Patchiness of macrobenthic invertebrates in homogenized intertidal habitats: hidden spatial structure at a landscape scale. *Marine Ecology Progress Series*, **383**, 211–224.
- Kraan, C., van Gils, J.A., Spaans, B., Dekkinga, A., Bijleveld, A.I., van Roomen, M., Kleefstra, R. & Piersma, T. (2009b) Landscape-scale experiment demonstrates that Wadden Sea intertidal flats are used to capacity by molluscivore migrant shorebirds. *Journal of Animal Ecology*, **78**, 1259–1268.
- Krebs, J.R. & Davies, N.B. (1997) *Behavioural Ecology: an Evolutionary Approach*. Sinauer Associates, Sunderland, MA.

- Krebs, J.R., Ryan, J.C. & Charnov, E.L. (1974) Hunting by expectation or optimal foraging? A study of patch use by chickadees. *Animal Behaviour*, **22, Part 4**, 953–964.
- Křivan, V. (1996) Optimal foraging and predator-prey dynamics. *Theoretical Population Biology*, **49**, 265–290.

L

- Laake, J., with contributions from E. Rakhimberdiev, B. Augustine (2012) Package "RMark": R Code for MARK Analysis, version 2.1.4.
- Laland, K.N. & Hoppitt, W. (2003) Do animals have culture? *Evolutionary Anthropology: Issues, News, and Reviews*, **12**, 150–159.
- Laland, K.N., Uller, T., Feldman, M.W., Sterelny, K., Müller, G.B., Moczek, A., Jablonka, E. & Odling-Smee, J. (2015) The extended evolutionary synthesis: its structure, assumptions and predictions. *Proceedings of the Royal Society B: Biological Sciences*, **282**.
- Lavielle, M. (2005) Using penalized contrasts for the change-point problem. *Signal Processing*, **85**, 1501–1510.
- Leadley, P., Pereira, H.M., Alkemade, R., Fernandez-Manjarrés, J.F., Proença, V., Scharlemann, J.P.W. & Walpole, M.J. (2010) Biodiversity scenarios: projections of 21st century change in biodiversity and associated ecosystem services. *Technical Series*, pp. 132. Secretariat of the Convention on Biological Diversity, Montreal.
- Lebreton, J.-D., Burnham, K.P., Clobert, J. & Anderson, D.R. (1992) Modeling survival and testing biological hypotheses using marked animals: a unified approach with case studies. *Ecological Monographs*, **62**, 67–118.
- Lessells, C.M. (1995) Putting resource dynamics into continuous input ideal free distribution models. *Animal Behaviour*, **49**, 487–494.
- Lewontin, R.C. (1983) Gene, organism and environment. *Evolution: from Molecules to Men* (ed. D.S. Bendall). Cambridge University Press, Cambridge.
- Lewontin, R.C. (2001) *The Triple Helix: Gene, Organism, and Environment*. Harvard University Press, Cambridge, MA.
- Leyrer, J. (2011) *Being at the Right Place at the Right Time: Interpreting the Annual Life Cycle of Afro-Siberian Red Knots*. PhD thesis. University of Groningen
- Leyrer, J., Lok, T., Brugge, M., Dekkinga, A., Spaans, B., van Gils, J.A., Sandercock, B.K. & Piersma, T. (2012) Small-scale demographic structure suggests preemptive behavior in a flocking shorebird. *Behavioral Ecology*, **23**, 1226–1233.
- Leyrer, J., Lok, T., Brugge, M., Spaans, B., Sandercock, B.K. & Piersma, T. (2013) Mortality within the annual cycle: seasonal survival patterns in Afro-Siberian Red Knots *Calidris canutus canutus*. *Journal of Ornithology*, **4**, 933–943.
- Leyrer, J., Spaans, B., Camara, M. & Piersma, T. (2006) Small home ranges and high site fidelity in red knots (*Calidris c. canutus*) wintering on the Banc d'Arguin, Mauritania. *Journal of Ornithology*, **147**, 376–384.
- Lima, S.L. & Dill, L.M. (1990) Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology*, **68**, 619–640.
- Lincango, M.P., Causton, C.E., Alvarez, C.C. & Jimenez-Uzcategui, G. (2011) Evaluating the safety of *Rodolia cardinalis* to two species of Galapagos finch; *Camarhynchus parvulus* and *Geospiza fuliginosa*. *Biological Control*, **56**, 145–149.
- Lindström, L., Alatalo, R.V., Lyytinen, A. & Mappes, J. (2001) Strong antiapostatic selection against novel rare aposematic prey. *Proceedings of the National Academy of Sciences USA*, **98**, 9181–9184.

M

- Ma, Z., Melville, D.S., Liu, J., Chen, Y., Yang, H., Ren, W., Zhang, Z., Piersma, T. & Li, B. (2014) Rethinking China's new great wall. *Science*, **346**, 912–914.
- MacArthur, R. (1955) Fluctuations of animal populations and a measure of community stability. *Ecology*, **36**, 533–536.
- MacArthur, R. & Levins, R. (1964) Competition, habitat selection, and character displacement in a patchy environment. *Proceedings of the National Academy of Sciences USA*, **51**, 1207–1210.

- MacArthur, R.H. & Pianka, E.R. (1966) On optimal use of a patchy environment. *American Naturalist*, **100**, 603–609.
- MacAvoy, S.E., Morgan, E., Carney, R.S. & Macko, S.A. (2008) Chemoautotrophic production incorporated by heterotrophs in Gulf of Mexico hydrocarbon seeps: An examination of mobile benthic predators and seep residents. *Journal of Shellfish Research*, **27**, 153–161.
- MacCurdy, R.B., Gabrielson, R.M. & Cortopassi, K.A. (2012) Automated wildlife radio tracking. In: *Handbook of Position Location: Theory, Practice and Advances* (eds. S.A. Zekavat & R.M. Buehrer). Wiley & Sons, Hoboken, NJ, 1129–1167
- Marsh, K.J., Wallis, I.R., McLean, S., Sorensen, J.S. & Foley, W.J. (2006) Conflicting demands on detoxification pathways influence how common brushtail possums choose their diets. *Ecology*, **87**, 2103–2112.
- Martin, G.R. (2017) *The Sensory Ecology of Birds*. Oxford University Press, Oxford.
- Mathot, K.J., Dekkinga, A. & Piersma, T. (2017) An experimental test of state-behaviour feedbacks: gizzard mass and foraging behaviour in red knots. *Functional Ecology*, doi: 10.1111/1365–2435.12827
- Mathot, K.J., Wright, J., Kempenaers, B. & Dingemanse, N. (2012) Adaptive strategies for managing uncertainty may explain personality-related differences in behavioural plasticity. *Oikos*, **121**.
- May, R.M. (1973) *Stability and Complexity in Model Ecosystems*. Princeton University Press.
- Mayntz, D., Raubenheimer, D., Salomon, M., Toft, S. & Simpson, S.J. (2005) Nutrient-specific foraging in invertebrate predators. *Science*, **307**, 111–113.
- Mayr, E. (1963) *Animal Species and Evolution*, Harvard University Press, Cambridge, MA.
- McFarland, D.J. (1977) Decision making in animals. *Nature*, **269**, 15–21.
- McNab, J.M. (1973) The avian caeca: a review. *World's Poultry Science Journal*, **29**, 251–263.
- McNair, J.N. (1982) Optimal giving-up times and the marginal value theorem. *American Naturalist*, **119**, 511–529.
- McNamara, J.M. & Houston, A.I. (1987) Partial preferences and foraging. *Animal Behaviour*, **35**, 1084–1099.
- McNamara, J.M. & Houston, A.I. (1994) The effect of a change in foraging options on intake rate and predation rate. *American Naturalist*, **144**, 978–1000.
- McNamara, J.M., Houston, A.I. & Collins, E.J. (2001) Optimality models in behavioral biology. *SIAM Review*, **43**, 413–466.
- McWilliams, S.R. & Karasov, W.H. (2014) Spare capacity and phenotypic flexibility in the digestive system of a migratory bird: defining the limits of animal design. *Proceedings of the Royal Society B: Biological Sciences*, **281**, 20140308.
- Moore, B.D. & Foley, W.J. (2005) Tree use by koalas in a chemically complex landscape. *Nature*, **435**, 488–490.
- Mouritsen, K.N. (1992) Predator avoidance in night-feeding dunlins *Calidris alpina*: a matter of concealment. *Ornis Scandinavica*, **23**, 195–198.
- Murdoch, W.W. (1969) Switching in general predators: experiments on predator specificity and stability of prey populations. *Ecological Monographs*, **39**, 335–354.
- Myers, J.P. (1980) Territoriality and flocking by buff-breasted sandpipers: variations in non-breeding dispersion. *Condor*, **82**, 241–250.
- Ma, Z., Melville, D.S., Liu, J., Chen, Y., Yang, H., Ren, W., Zhang, Z., Piersma, T. & Li, B. (2014) Rethinking China's new great wall. *Science*, **346**, 912–914.

N

- Nathan, R., Getz, W.M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D. & Smouse, P.E. (2008) A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences USA*, **105**, 19052–19059.
- Nebel, S., Piersma, T., van Gils, J.A., Dekkinga, A. & Spaans, B. (2000) Length of stopover, fuel storage and a sex-bias in the occurrence of two subspecies of red knots *Calidris c. canutus* and *C.c. islandica* in the Dutch Wadden Sea during southward migration. *Ardea*, **88**, 11.
- Noble, D. (2015) Conrad Waddington and the origin of epigenetics. *Journal of Experimental Biology*, **218**, 816–818.
- Nolte, D.L. & Provenza, F.D. (1991) Food preferences in lambs after exposure to flavours in milk. *Applied Animal Behaviour Science*, **32**, 381–389.
- Nolte, D.L., Provenza, F.D., R, C. & Panter, K.E. (1992) Garlic in the ovine fetal environment. *Physiology & Behavior*, **52**, 1091–1093.

O

- Ólafsson, E.B., Peterson, C.H. & Ambrose, W.G. (1995) Does recruitment limitation structure populations and communities of macro-invertebrates in marine soft sediments - the relative significance of pre- and post-settlement processes. *Oceanographic Literature Review*, **8**, 667.
- Olsson, O. & Holmgren, N.M.A. (1998) The survival-rate-maximizing policy for Bayesian foragers: wait for good news. *Behavioral Ecology*, **9**, 345–353.
- Olsson, J., Quevedo, M., Colson, C. & Svanbäck, R. (2007) Gut length plasticity in perch: into the bowels of resource polymorphisms. *Biological Journal of the Linnean Society*, **90**, 517–523.
- Onrust, J., de Fouw, J., Oudman, T., van der Geest, M., Piersma, T. & van Gils, J.A. (2013) Red Knot diet reconstruction revisited: context dependence revealed by experiments at Banc d'Arguin, Mauritania. *Bird Study*, **60**, 298–307.
- Oudman, T., Bijleveld, A.I., Kavelaars, M.M., Dekkinga, A., Cluderay, J., Piersma, T. & van Gils, J.A. (2016) Diet preferences as the cause of individual differences rather than the consequence. *Journal of Animal Ecology*, **85**, 1378–1388.
- Oudman, T., Bijleveld, A.I., Kavelaars, M., Cluderay, J., Piersma, T. & van Gils, J.A. (2016) Data from: Diet preferences as the cause of individual differences rather than the consequence. *Dryad Digital Repository*, <http://dx.doi.org/10.5061/dryad.2s8rr>.
- Oudman, T., Hin, V., Dekkinga, A. & van Gils, J.A. (2015) The effect of digestive capacity on the intake rate of toxic and non-toxic prey in an ecological context. *PLoS ONE*, **10**, e0136144.
- Oudman, T., Onrust, J., de Fouw, J., Spaans, B., Piersma, T. & van Gils, J.A. (2014) Digestive capacity and toxicity cause mixed diets in red knots that maximize intake rate. *American Naturalist*, **183**, 650–659.
- Oudman, T., Onrust, J., Spaans, B., Piersma, A.H. & van Gils, J.A. (2014) Data from: Digestive capacity and toxicity cause mixed diets in red knots that maximize energy intake rate. *Dryad Digital Repository*, <http://dx.doi.org/10.5061/dryad.5fp4g>.
- Oudman, T., Schekkerman, H., Kidee, A., van Roomen, M., Tentij, M. & Piersma, T. (2017) The waterbirds of Parc National du Banc d'Arguin: evaluation of all complete winter counts, workshop proceedings and an evaluation. *Report for Programme Towards a Rich Wadden Sea*.
- Owen-Smith, N. (1993) Evaluating optimal diet models for an African browsing ruminant, the kudu: how constraining are the assumed constraints? *Evolutionary Ecology*, **7**, 499–524.
- Owen-Smith, N. (1996) Circularity in linear programming models of optimal diet. *Oecologia*, **108**, 259–261.

P

- Patrick, S.C., Bearhop, S., Grémillet, D., Lescroël, A., Grecian, W.J., Bodey, T.W., Hamer, K.C., Wakefield, E., Le Nuz, M. & Votier, S.C. (2014) Individual differences in searching behaviour and spatial foraging consistency in a central place marine predator. *Oikos*, **123**, 33–40.
- Piersma, T. (1994) *Close to the Edge: Energetic Bottlenecks and the Evolution of Migratory Pathways in Knots*. PhD thesis, University of Groningen.
- Piersma, T. (2002) Energetic bottlenecks and other design constraints in avian annual cycles. *Integrative and Comparative Biology*, **42**, 51–67.
- Piersma, T. (2007) Using the power of comparison to explain habitat use and migration strategies of shorebirds worldwide. *Journal of Ornithology*, **148** (Suppl 1), S45–S59.
- Piersma, T. (2011) Flyway evolution is too fast to be explained by the modern synthesis: proposals for an 'extended' evolutionary research agenda. *Journal of Ornithology*, **152**, 151–159.
- Piersma, T. (2012) What is habitat quality? Dissecting a research portfolio on shorebirds. *Birds and Habitat: Relationships in Changing Landscapes*. (ed. R. Fuller). Cambridge University Press, Cambridge (UK), 383–407.
- Piersma, T. & Davidson, N.C. (1992) The migrations and annual cycles of five subspecies of knots in perspective. *Wader Study Group Bulletin*, **64**, 187–197.
- Piersma, T., de Goeij, P. & Tulp, I. (1993) An evaluation of intertidal feeding habitats from a shorebird perspective: towards comparisons between temperate and tropical mudflats. *Netherlands Journal of Sea Research*, **31**, 503–512.
- Piersma, T., Hoekstra, R., Dekkinga, A., Koolhaas, A., Wolf, P., Battley, P. & Wiersma, P. (1993) Scale and intensity of intertidal habitat use by knots *Calidris canutus* in the western Wadden sea in relation to food, friends and foes. *Netherlands Journal of Sea Research*, **31**, 331–357.

- Piersma, T., Koolhaas, A. & Dekkinga, A. (1993) Interactions between stomach structure and diet choice in shorebirds. *Auk*, **110**, 552–564.
- Piersma, T. & Lindström, Å. (1997) Rapid reversible changes in organ size as a component of adaptive behaviour. *Trends in Ecology & Evolution*, **12**, 134–138.
- Piersma, T., Lok, T., Chen, Y., Hassell, C.J., Yang, H.-Y., Boyle, A., Slaymaker, M., Chan, Y.-C., Melville, D.S., Zhang, Z.-W. & Ma, Z. (2016) Simultaneous declines in summer survival of three shorebird species signals a flyway at risk. *Journal of Applied Ecology*, **53**, 479–490.
- Piersma, T., MacCurdy, R.B., Gabrielson, R.M., Cluderay, J., Dekkinga, A., Spaulding, E., Oudman, T., Onrust, J., van Gils, J.A., Winkler, D.W. & Bijleveld, A.I. (2014) Fine-scale measurements of individual movements within bird flocks: the principles and three applications of TOA tracking. *Limosa*, **87**, 156–167.
- Piersma, T. & van Gils, J.A. (2011) *The Flexible Phenotype: A Body-centred Integration of Ecology, Physiology and Behaviour*. Oxford University Press, Oxford.
- Piersma, T., van Gils, J.A., de Goeij, P. & van der Meer, J. (1995) Holling's functional-response model as a tool to link the food-finding mechanism of a probing shorebird with its spatial distribution. *Journal of Animal Ecology*, **64**, 493–504.
- Pigliucci, M. (2001) *Phenotypic Plasticity: Beyond Nature and Nurture*. JHU Press, Baltimore, MD.
- Prater, A.J. (1972) The ecology of Morecambe Bay. III. The food and feeding habits of knot (*Calidris canutus* L.) in Morecambe Bay. *Journal of Applied Ecology*, **9**, 179–194.
- Prater, A.J., Merchant, J.H. & Vuorinen, J. (1977) *Guide to the Identification and Ageing of Holarctic Shorebirds*. BTO, Tring.
- Provenza, F.D. & Balph, D.F. (1987) Diet learning by domestic ruminants: theory, evidence and practical implications. *Applied Animal Behaviour Science*, **18**, 211–232.
- Provenza, F.D. & Cincotta, R.P. (1993) Foraging as a self-organizational learning process: accepting adaptability at the expense of predictability. *Diet Selection* (ed. R.N. Hughes), pp. 78–101. Blackwell Scientific Publications, London.
- Pulliam, H.R. (1975) Diet optimization with nutrient constraints. *American Naturalist*, **109**, 765–768.
- Pulliam, H.R. & Caraco, N. (1984) Living in groups: is there an optimal group size? In: *Behavioural Ecology: an Evolutionary Approach* (eds. J.R. Krebs & N.B. Davies). Sinauer Associates, Sunderland, MA, pp. 122–147.

Q

- Quainten, G., van Gils, J.A., Bocher, P., Dekkinga, A. & Piersma, T. (2010) Diet selection in a molluscivore shorebird across Western Europe: does it show short- or long-term intake rate-maximization? *Journal of Animal Ecology*, **79**, 53–62.

R

- Rapport, D.J. (1971) An optimization model of food selection. *American Naturalist*, **105**, 575–587.
- R Core Team (2015) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>.
- R Development Core Team (2013) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org>.
- Réale, D. & Dingemanse, N.J. (2010) Personality and individual social specialisation. In: *Social Behaviour: Genes, Ecology and Evolution* (eds. T. Székely, A.J. Moore & J. Komdeur), pp. 417–441. Cambridge University Press, Cambridge, UK.
- Réale, D., Dingemanse, N.J., Kazem, A.J.N. & Wright, J. (2010) Evolutionary and ecological approaches to the study of personality. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 3937–3946.
- Richter, C.P. (1943) Total self-regulatory functions in animals and human beings. *Harrey Lectures*, **38**, 63–103.
- Rita, H. & Ranta, E. (1998) Stochastic patch exploitation model. *Proceedings of the Royal Society of London B: Biological Sciences*, **265**, 309–315.
- Ritchie, M. (1988) Individual variation in the ability of Columbian ground squirrels to select an optimal diet. *Evolutionary Ecology*, **2**, 232–252.

- Rosenthal, O.D. & Berenbaum, M.R. (1992) *Herbivores: their Interactions with Secondary Plant Metabolites* (Second Edition). Vol. II. Academic Press Limited, San Diego, CA.
- S**
- Saikkonen, K., Faeth, S.H., Helander, M. & Sullivan, T.J. (1998) Fungal endophytes: A continuum of interactions with host plants. *Annual Review of Ecology and Systematics*, **29**, 319–343.
- Schmidt, K.A. (2000) Interactions between food chemistry and predation risk in fox squirrels. *Ecology*, **81**, 2077–2085.
- Schmidt, K.A., Brown, J.S. & Morgan, R.A. (1998) Plant defenses as complementary resources: a test with squirrels. *Oikos*, **81**, 130–142.
- Schmidt, K.A., Dall, S.R. & van Gils, J.A. (2010) The ecology of information: an overview on the ecological significance of making informed decisions. *Oikos*, **119**, 304–316.
- Schoener, T.W. (1971) Theory of feeding strategies. *Annual Review of Ecology and Systematics*, **2**, 369–404.
- Secor, S.M. & Diamond, J. (1995) Adaptive responses to feeding in Burmese pythons: pay before pumping. *Journal of Experimental Biology*, **198**, 1313–1325.
- Senner, N.R., Conklin, J.R. & Piersma, T. (2015) An ontogenetic perspective on individual differences. *Proceedings of the Royal Society of London B: Biological Sciences*, **282**, 20151050.
- Sherratt, T.N. (2003) State-dependent risk-taking by predators in systems with defended prey. *Oikos*, **103**, 93–100.
- Sherratt, T.N., Speed, M.P. & Ruxton, G.D. (2004) Natural selection on unpalatable species imposed by state-dependent foraging behaviour. *Journal of Theoretical Biology*, **228**, 217–226.
- Shrader, A.M., Kotler, B.P., Brown, J.S. & Kerley, G.I.H. (2008) Providing water for goats in arid landscapes: effects on feeding effort with regard to time period, herd size and secondary compounds. *Oikos*, **117**, 466–472.
- Sih, A., Bell, A. & Johnson, J.C. (2004) Behavioral syndromes: an ecological and evolutionary overview. *Trends in Ecology & Evolution*, **19**, 372–378.
- Sih, A. & Christensen, B. (2001) Optimal diet theory: when does it work, and when and why does it fail? *Animal Behaviour*, **61**, 379–390.
- Simpson, S.J., Sibly, R.M., Lee, K.P., Behmer, S.T. & Raubenheimer, D. (2004) Optimal foraging when regulating intake of multiple nutrients. *Animal Behaviour*, **68**, 1299–1311.
- Sims, D.W., Southall, E.J., Humphries, N.E., Hays, G.C., Bradshaw, C.J.A., Pitchford, J.W., James, A., Ahmed, M.Z., Brierley, A.S., Hindell, M.A., Morritt, D., Musyl, M.K., Righton, D., Shepard, E.L.C., Wearmouth, V.J., Wilson, R.P., Witt, M.J. & Metcalfe, J.D. (2008) Scaling laws of marine predator search behaviour. *Nature*, **451**, 1098–1102.
- Sitters, H.P., González, P.M., Piersma, T., Baker, A.J. & Price, D.J. (2001) Day and night feeding habitat of red knots in Patagonia: profitability versus safety? *Journal of Field Ornithology*, **72**, 86–95.
- Skelhorn, J. & Rowe, C. (2007) Predators' toxin burdens influence their strategic decisions to eat toxic prey. *Current Biology*, **17**, 1479–1483.
- Skelhorn, J. & Rowe, C. (2010) Birds learn to use distastefulness as a signal of toxicity. *Proceedings of the Royal Society B: Biological Sciences*, **277**, 1729–1734.
- Speed, M.P. (1993) When is mimicry good for predators? *Animal Behaviour*, **46**, 1246–1248.
- Speed, M.P. (2000) Warning signals, receiver psychology and predator memory. *Animal Behaviour*, **60**, 269–278.
- Spiegel, O., Leu, S.T., Bull, C.M. & Sih, A. (2017) What's your move? Movement as a link between personality and spatial dynamics in animal populations. *Ecology Letters*, **20**, 3–18.
- Stamps, J.A. (2003) Behavioural processes affecting development: Tinbergen's fourth question comes of age. *Animal Behaviour*, **66**, 1–13.
- Stamps, J.A. & Groothuis, T.G.G. (2010a) Developmental perspectives on personality: implications for ecological and evolutionary studies of individual differences. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **365**, 4029–4041.
- Stamps, J.A. & Groothuis, T.G.G. (2010b) The development of animal personality: relevance, concepts and perspectives. *Biological Reviews*, **85**, 301–325.
- Stamps, J.A. (2015) Individual differences in behavioural plasticities. *Biological Reviews*, [10.1111/bvr.12186](https://doi.org/10.1111/bvr.12186).

- Starck, J.M. (1999) Structural flexibility of the gastro-intestinal tract of vertebrates-implications for evolutionary morphology. *Zoologischer Anzeiger*, **238**, 87–102.
- Stephens, D.W. & Krebs, J.R. (1986) *Foraging Theory*. Princeton University Press, Princeton, NJ.
- Sutherland, W.J. (1983) Aggregation and the 'ideal free' distribution. *Journal of Animal Ecology*, **52**, 821–828.
- Sutherland, W.J. & Parker, G.A. (1985) Distribution of unequal competitors. In: *Behavioural Ecology: Ecological Consequences of Adaptive Behaviour* (eds. R.M. Smith & R.H. Smith). Blackwell Oxford, pp. 255–274.
- Switzer, P.V. (1993) Site fidelity in predictable and unpredictable habitats. *Evolutionary Ecology*, **7**, 533–555.

T

- Taylor, J.D. & Glover, E.A. (2006) Lucinidae (Bivalvia) - the most diverse group of chemosymbiotic molluscs. *Zoological Journal of the Linnean Society*, **148**, 421–438.
- Taylor, J.D., Glover, E.A., Smith, L., Dyal, P. & Williams, S.T. (2011) Molecular phylogeny and classification of the chemosymbiotic bivalve family Lucinidae (Mollusca: Bivalvia). *Zoological Journal of the Linnean Society*, **163**, 15–49.
- Templeton, C.N., Laland, K.N. & Boogert, N.J. (2014) Does song complexity correlate with problem-solving performance in flocks of zebra finches? *Animal Behaviour*, **92**, 63–71.
- Tilman, D. (1982) *Resource Competition and Community Structure*. Princeton University Press, Princeton, NJ.
- Tilman, D. (1994) Competition and biodiversity in spatially structured habitats. *Ecology*, **75**, 2–16.
- Tilman, D., Fargione, J., Wolff, B., D'Antonio, C., Dobson, A., Howarth, R., Schindler, D., Schlesinger, W.H., Simberloff, D. & Swackhamer, D. (2001) Forecasting agriculturally driven global environmental change. *Science*, **292**, 281–284.
- Tim Tinker, M., Guimarães, P.R., Novak, M., Marquitti, F.M.D., Bodkin, J.L., Staedler, M., Bentall, G. & Estes, J.A. (2012) Structure and mechanism of diet specialisation: testing models of individual variation in resource use with sea otters. *Ecology Letters*, **15**, 475–483.
- Toft, S. & Wise, D.H. (1999) Behavioral and ecophysiological responses of a generalist predator to single- and mixed-species diets of different quality. *Oecologia*, **119**, 198–207.
- Tregenza, T. (1995) Building on the ideal free distribution. In: *Advances in Ecological Research* (eds. M. Begon & A.H Fitter). Academic Press, London, pp. 253–307.
- Tunnicliffe, V. (1991) The biology of hydrothermal vents - ecology and evolution. *Oceanography and Marine Biology*, **29**, 319–407.

V

- Vahl, W.K., Lok, T., van der Meer, J., Piersma, T. & Weissing, F.J. (2005) Spatial clumping of food and social dominance affect interference competition among ruddy turnstones. *Behavioral Ecology*, **16**, 834–844.
- van Baalen, M., Křivan, V., van Rijn, P.C.J. & Sabelis, M.W. (2001) Alternative food, switching predators, and the persistence of predator-prey systems. *American Naturalist*, **157**, 512–524.
- van Bemmelen, R., Moe, B., Hanssen, S.A., Schmidt, N.M., Hansen, J., Lang, J., Sittler, B., Bollache, L., Tulp, I., Klaassen, R. & Gilg, O. (2017) Flexibility in otherwise consistent non-breeding movements of a long-distance migratory seabird, the long-tailed skua. *Marine Ecology Progress Series*, doi: 10.3354/meps12010.
- van de Kam, J., Ens, B.J., Piersma, T. & Zwarts, L. (2004) *Shorebirds. An illustrated behavioural ecology*. KNNV Publishers, Utrecht.
- van den Hout, P.J. (2010) *Struggle for Safety: Adaptive Responses of Wintering Waders to their Avian Predators*. PhD thesis, University of Groningen.
- van den Hout, P.J., Mathot, K.J., Maas, L.R.M. & Piersma, T. (2009) Predator escape tactics in birds: linking ecology and aerodynamics. *Behavioral Ecology*, **21**, 16–25.
- van den Hout, P.J., Piersma, T., ten Horn, J., Spaans, B. & Lok, T. (2016) Individual shifts toward safety explain age-related foraging distribution in a gregarious shorebird. *Behavioral Ecology*, arw173.

- van den Hout, P.J., Spaans, B. & Piersma, T. (2008) Differential mortality of wintering shorebirds on the Banc d'Arguin, Mauritania, due to predation by large falcons. *Ibis*, **150** (Suppl. 1), 219–230.
- van den Hout, P.J., van Gils, J.A., Robin, F., van der Geest, M., Dekkinga, A. & Piersma, T. (2014) Interference from adults forces young red knots to forage for longer and in dangerous places. *Animal Behaviour*, **88**, 137–146.
- van der Geest, M., Sall, A.A., Ely, S.O., Nauta, R.W., van Gils, J.A. & Piersma, T. (2014) Nutritional and reproductive strategies in a chemosymbiotic bivalve living in a tropical intertidal seagrass bed. *Marine Ecology Progress Series*, **501**, 113–126.
- van der Geest, M., van Gils, J.A., van der Meer, J., Olff, H. & Piersma, T. (2011) Suitability of calcine as an in situ growth marker in burrowing bivalves. *Journal of Experimental Marine Biology and Ecology*, **399**, 1–7.
- van der Heide, T., Bouma, T.J., van Nes, E.H., van de Koppel, J., Scheffer, M., Roelofs, J.G.M., van Katwijk, M.M. & Smolders, A.J.P. (2010) Spatial self-organized patterning in seagrasses along a depth gradient of an intertidal ecosystem. *Ecology*, **91**, 362–369.
- van der Heide, T., Govers, L.L., de Fouw, J., Olff, H., van der Geest, M., van Katwijk, M.M., Piersma, T., van de Koppel, J., Silliman, B.R., Smolders, A.J.P. & van Gils, J.A. (2012) A three-stage symbiosis forms the foundation of seagrass ecosystems. *Science*, **336**, 1432–1434.
- van der Heide, T., van Nes, E.H., Geerling, G.W., Smolders, A.J., Bouma, T.J. & van Katwijk, M.M. (2007) Positive feedbacks in seagrass ecosystems: implications for success in conservation and restoration. *Ecosystems*, **10**, 1311–1322.
- van der Meer, J., Beukema, J.J. & Dekker, R. (2001) Long-term variability in secondary production of an intertidal bivalve population is primarily a matter of recruitment variability. *Journal of Animal Ecology*, **70**, 159–169.
- van der Meer, J., Piersma, T. & Beukema, J.J. (2001) Population dynamics of benthic species on tidal flats: the possible roles of shorebird predation. In: *Ecological comparisons of sedimentary shores* (ed. K. Reise). Springer-Verlag, Berlin Heidelberg.
- van Gils, J.A. (2004) *Foraging Decisions in a Digestively Constrained Long-distance Migrant, the Red Knot (Calidris canutus)*. PhD thesis, University of Groningen.
- van Gils, J.A., Battley, P.F., Piersma, T. & Drent, R. (2005a) Reinterpretation of gizzard sizes of red knots world-wide emphasises overriding importance of prey quality at migratory stopover sites. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 2609–2618.
- van Gils, J.A., de Rooij, S.R., van Belle, J., van der Meer, J., Dekkinga, A., Piersma, T. & Drent, R. (2005b) Digestive bottleneck affects foraging decisions in red knots *Calidris canutus*. I. Prey choice. *Journal of Animal Ecology*, **74**, 105–119.
- van Gils, J.A., Dekkinga, A., Spaans, B., Vahl, W.K. & Piersma, T. (2005c) Digestive bottleneck affects foraging decisions in red knots *Calidris canutus*. II. Patch choice and length of working day. *Journal of Animal Ecology*, **74**, 120–130.
- van Gils, J.A., Dekkinga, A., van den Hout, P.J., Spaans, B. & Piersma, T. (2007) Digestive organ size and behavior of red knots (*Calidris canutus*) indicate the quality of their benthic food stocks. *Israel Journal of Ecology & Evolution*, **53**, 329–346.
- van Gils, J.A., Lisovski, S., Lok, T., Meissner, W., Ożarowska, A., de Fouw, J., Rakhimberdiev, E., Soloviev, M.Y., Piersma, T. & Klaassen, M. (2016) Body shrinkage due to Arctic warming reduces red knot fitness in tropical wintering range. *Science*, **352**, 819–821.
- van Gils, J.A., Piersma, T., Dekkinga, A. & Battley, P.F. (2006) Modelling phenotypic flexibility: an optimality analysis of gizzard size in red knots (*Calidris canutus*). *Ardea*, **94**, 409–420.
- van Gils, J.A., Piersma, T., Dekkinga, A. & Dietz, M.W. (2003a) Cost-benefit analysis of mollusc-eating in a shorebird. II. Optimizing gizzard size in the face of seasonal demands. *Journal of Experimental Biology*, **206**, 3369–3380.
- van Gils, J.A., Schenk, I.W., Bos, O. & Piersma, T. (2003b) Incompletely informed shorebirds that face a digestive constraint maximize net energy gain when exploiting patches. *American Naturalist*, **161**, 777–793.
- van Gils, J.A., Spaans, B., Dekkinga, A. & Piersma, T. (2006) Foraging in a tidally structured environment by red knots (*Calidris canutus*): ideal, but not free. *Ecology*, **87**, 1189–1202.

- van Gils, J.A., van der Geest, M., Jansen, E.J., Govers, L.L., de Fouw, J. & Piersma, T. (2012) Trophic cascade induced by molluscivore predator alters pore-water biogeochemistry via competitive release of prey. *Ecology*, **93**, 1143–1152.
- van Gils, J.A., van der Geest, M., Kraan, C., Folmer, E.O., Jansen, E.J. & Piersma, T. (2009) How the carrying capacity of the Wadden Sea regulates the number of wintering waders at Banc d'Arguin. *Limosa*, **82**, 134–140.
- van Gils, J.A., van der Geest, M., Leyrer, J., Oudman, T., Lok, T., Onrust, J., de Fouw, J., van der Heide, T., van den Hout, P.J., Spaans, B., Dekkinga, A., Brugge, M. & Piersma, T. (2013) Toxin constraint explains diet choice, survival and population dynamics in a molluscivore shorebird. *Proceedings of the Royal Society B: Biological Sciences*, **280**, 20130861.
- van Moorter, B., Rolandsen, C.M., Basille, M. & Gaillard, J.-M. (2016) Movement is the glue connecting home ranges and habitat selection. *Journal of Animal Ecology*, **85**, 21–31.
- van Roomen, M., Nagy, S., Foppen, R., Dodman, T., Citegetse, G. & Ndiaye, A. (2015) *Status of Coastal Waterbird Populations in the East Atlantic Flyway*. Program Rich Wadden Sea, Leeuwarden, Sovon, Nijmegen, Wetlands International, Wageningen, BirdLife International, Cambridge, Common Wadden Sea Secretariat, Wilhelmshaven.
- Verboven, N. & Piersma, T. (1995) Is the evaporative water loss of Knot *Calidris canutus* higher in tropical than in temperate climates? *Ibis*, **137**, 308–316.
- Verstraete, B., van Elst, D., Steyn, H., Van Wyk, B., Lemaire, B., Smets, E. & Dessein, S. (2011) Endophytic bacteria in toxic South African plants: identification, phylogeny and possible involvement in gousiekte. *PLoS ONE*, **6**, e19265.
- Villalba, J.J., Provenza, F.D. & Han, G. (2004) Experience influences diet mixing by herbivores: implications for plant biochemical diversity. *Oikos*, **107**, 100–109.

W

- Waddington, C.H. (1942) Canalization of development and the inheritance of acquired characters. *Nature*, **150**, 563–565.
- Waddington, C.H. (1953) Genetic assimilation of an acquired character. *Evolution*, **7**, 118–126.
- Waddington, C.H. (1957) *The Strategy of the Genes*. Allen and Unwin, London.
- Waddington, C.H. (1959) Evolutionary systems - animal and human. *Nature*, **183**, 1634–1638.
- Warnock, N. & Warnock, S. (1993) Attachment of radio-transmitters to sandpipers: review and methods. *Wader Study Group Bulletin*, **70**, 28–30.
- Warton, D.I. & Hui, F.K.C. (2011) The arcsine is asinine: the analysis of proportions in ecology. *Ecology*, **92**, 3–10.
- Weimerskirch, H. (2007) Are seabirds foraging for unpredictable resources? *Deep Sea Research Part II: Topical Studies in Oceanography*, **54**, 211–223.
- Werner, E.E. & Peacor, S.D. (2003) A review of trait-mediated indirect interactions in ecological communities. *Ecology*, **84**, 1083–1100.
- West-Eberhard, M.J. (1989) Phenotypic plasticity and the origins of diversity. *Annual Review of Ecology and Systematics*, **20**, 249–278.
- West-Eberhard, M.J. (2003) *Developmental Plasticity and Evolution*. Oxford University Press, New York, NY.
- Westoby, M. (1974) An analysis of diet selection by large generalist herbivores. *American Naturalist*, **108**, 290–304.
- Whelan, C.J. & Brown, J.S. (2005) Optimal foraging and gut constraints: reconciling two schools of thought. *Oikos*, **110**, 481–496.
- White, G.C. & Burnham, K.P. (1999) Program MARK: survival estimation from populations of marked animals. *Bird Study*, **46**, S120–S139.
- Whiteside, M.A., Sage, R. & Madden, J.R. (2015) Diet complexity in early life affects survival in released pheasants by altering foraging efficiency, food choice, handling skills and gut morphology. *Journal of Animal Ecology*, **84**, 1480–1489.
- Wiersma, P. & Piersma, T. (1994) Effects of microhabitat, flocking, climate and migratory goal on energy expenditure in the annual cycle of red knots. *Condor*, **96**, 257–279.

- Williams, B., Lovenburg, V., Huffard, C. & Caldwell, R. (2011) Chemical defense in pelagic octopus paralarvae: tetrodotoxin alone does not protect individual paralarvae of the greater blue-ringed octopus (*Hapalochlaena lunulata*) from common reef predators. *Chemoecology*, **21**, 131–141.
- Winchester, C.F. & Morris, M.J. (1956) Water intake rates of cattle. *Journal of Animal Science*, **15**, 722–740.
- Wolff, W.J., Duiven, A.G., Duiven, P., Esselink, P., Gueye, A., Meijboom, A., Moerland, G. & Zegers, J. (1993) Biomass of macrobenthic tidal flat fauna of the Banc d'Arguin, Mauritania. *Hydrobiologia*, **258**, 151–163.
- Wolff, W.J. & Smit, C. (1990) The Banc d'Arguin, Mauritania, as an environment for coastal birds. *Ardea*, **78**, 17–38.

Y

- Yamauchi, A. & Yamamura, N. (2005) Effects of defense evolution and diet choice on population dynamics in a one-predator-two-prey system. *Ecology*, **86**, 2513–2524.
- Yang, H.-Y., Chen, B., Ma, Z.-j., Hua, N., van Gils, J.A., Zhang, Z.-W. & Piersma, T. (2013) Economic design in a long-distance migrating molluscivore: how fast-fuelling red knots in Bohai Bay, China, get away with small gizzards. *Journal of Experimental Biology*, **216**, 3627–3636.

Z

- Zwarts, L. (1991) Seasonal variation in body weight of the bivalves *Macoma balthica*, *Scrobicularia plana*, *Mya arenaria* and *Cerastoderma edule* in the Dutch Wadden Sea. *Netherlands Journal of Sea Research*, **28**, 231–245.
- Zwarts, L. & Blomert, A.M. (1992) Why knot *Calidris canutus* take medium-sized *Macoma balthica* when 6 prey species are available. *Marine Ecology Progress Series*, **83**, 113–128.
- Zwarts, L., Blomert, A.-M. & Wanink, J.H. (1992) Annual and seasonal variation in the food supply harvestable by knot *Calidris canutus* staging in the Wadden Sea in late summer. *Marine Ecology Progress Series*, **83**, 129–139.
- Zwarts, L. & Wanink, J.H. (1993) How the food supply harvestable by waders in the Wadden Sea depends on the variation in energy density, body weight, biomass, burying depth and behaviour of tidal-flat invertebrates. *Netherlands Journal of Sea Research*, **31**, 441–476.