

University of Groningen

Alternative habitat

Lei, Weipan; Masero, José A.; Piersma, Theunis; Zhu, Bingrun; Yang, Hong Yan; Zhang, Zhengwang

Published in:
 Bird Conservation International

DOI:
[10.1017/S0959270917000508](https://doi.org/10.1017/S0959270917000508)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Lei, W., Masero, J. A., Piersma, T., Zhu, B., Yang, H. Y., & Zhang, Z. (2018). Alternative habitat: The importance of the Nanpu Saltpans for migratory waterbirds in the Chinese Yellow Sea. *Bird Conservation International*, 28(4), 549-566. <https://doi.org/10.1017/S0959270917000508>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Alternative habitat: the importance of the Nanpu Saltpans for migratory waterbirds in the Chinese Yellow Sea

WEIPAN LEI, JOSÉ A. MASERO, THEUNIS PIERSMA, BINGRUN ZHU,
HONG-YAN YANG and ZHENGWANG ZHANG

Summary

The natural coastal wetlands of the East Asian-Australasian Flyway (EAAF) are disappearing at alarming rates, leading to rapid declines of many populations of waterbirds in the most species-rich flyway in the world. The identification and assessment of possible alternative habitats that may buffer the loss of natural wetlands should, therefore, be a priority for the conservation of migratory waterbirds using this flyway. Coastal saltpans are functional wetlands that support large numbers of waterbirds worldwide. The Nanpu Saltpans in the northern Bohai Bay of the Yellow Sea in China are one of the largest (290 km²) saltpan complexes in the world. In this paper, we document the value of the Nanpu Saltpans for supporting waterbirds. The surveys, carried out from 2013 to 2016, included waterbird counts in the saltpans (93 km²) at high and low tide and on the adjacent natural tidal flats (57 km²) at low tide. Of the 89 waterbird species recorded, 27 had maximum counts exceeding the 1% threshold value of estimated flyway populations. The maximum counts of waterbirds in northward migration and southward migration in the Nanpu Saltpans were 96,000 and 93,500, respectively, including both foraging and roosting birds; these figures do not account for turnover, so the total number of birds using the site is likely to be higher. The maximum counts on the adjacent tidal flats at low tide amounted to 73,000 and 20,000 waterbirds during northward and southward migration, respectively, and most of them were foraging birds. In the boreal winter, few birds fed in the saltpans, but several thousand fed on the tidal flats. Waterbirds used the inland ponds (2.0–18.0 km from the intertidal area) mainly for feeding both during low tide and high tide and used the nearshore ponds (0.3–4.3 km from the intertidal area) mainly for high-tide roosting. Some species, such as Black-tailed Godwit *Limosa limosa*, Marsh Sandpiper *Tringa stagnatilis*, Pied Avocet *Recurvirostra avosetta*, and Black-winged Stilt *Himantopus himantopus*, occurred mainly in the saltpans; other species preferred tidal flats, such as Red Knot *Calidris canutus*, Great Knot *Calidris tenuirostris*, Bar-tailed Godwit *Limosa lapponica*, Eurasian Curlew *Numenius arquata*, Relict Gull *Larus relictus*, and Grey Plover *Pluvialis squatarola*. This study clearly demonstrates the joint ecological function of the Nanpu Saltpan complex and adjacent tidal flats as a key staging area for waterbirds in the EAAF, and as such both urgently warrant protected status.

Introduction

The East Asian-Australasian Flyway (EAAF) supports more waterbird species and is facing greater threats with more threatened species than any other flyway in the world (Kirby 2010, Wetlands International 2017). The shores of the Yellow Sea are a particularly important component of the EAAF, with internationally significant staging areas for at least 24 migratory

shorebird species (Barter 2002, Bamford et al. 2008, MacKinnon et al. 2012). However, the tidal flat ecosystems along the Yellow Sea are disappearing at an alarming rate: 28% of tidal flats existing in the 1980s had disappeared by the late 2000s, indicating an annual rate of loss of 1.2% (MacKinnon et al. 2012, Ma et al. 2014, Murray et al. 2014, Melville et al. 2016). Meanwhile, measurable correlated population and survival declines of the shorebirds and other waterbirds that rely on intertidal areas as feeding habitat were detected (Amano et al. 2010, Wilson et al. 2011, Piersma et al. 2016). These studies highlight the question of whether these waterbirds can use alternative habitats, for example human-made saltpans (Hua et al. 2015, Studds et al. 2017).

Anthropogenic wetlands, created or extensively modified by humans, are rapidly expanding and replacing natural wetlands worldwide (Czech and Parsons 2002, Ma et al. 2010). Although anthropogenic wetlands cannot completely replace the functions of natural ones as waterbird habitat (e.g. Ma et al. 2004, Desrochers et al. 2008, Choi et al. 2014, Sebastián-González and Green 2016), they can provide alternative or supplemental habitats for waterbirds. Examples of anthropogenic wetlands used by waterbirds include rice fields (Elphick 2000, Sanchez-Guzman et al. 2007), coastal grazing marshes (Milsom et al. 2000, Rhymer et al. 2010), extensive aquaculture ponds (Ma et al. 2002, Li et al. 2013, Navedo and Fernández in press), and coastal saltpans (Warnock and Takekawa 1995, Masero et al. 2000). Although the effective management of these anthropogenic wetlands can help to mitigate the adverse effects of natural wetland loss or degradation (reviewed by Ma et al. 2010, Navedo et al. 2017), in the EAAF many salt ponds, for example, are being converted to modern aquaculture or industrial land (Flaherty and Karnjanakesorn 1995, Sripanomyom et al. 2011).

In China, the total area of coastal saltpans increased from 1,639 km² in 1985 to 2,413 km² in 2010 and the total area of aquaculture ponds increased from 2,606 km² to 12,099 km² (Yao et al. 2016). However, during this century the area of coastal saltpans decreased owing to competing land uses driven by urbanisation and economic growth (Li et al. 2011).

Coastal saltpans are anthropogenic wetlands used for producing salt by the solar evaporation of seawater (Rocha et al. 2016). They are often classified as 'functional wetlands' with a high biological richness and support large numbers of shorebirds and other waterbird groups worldwide (e.g. Masero 2003, Sripanomyom et al. 2011, Houston et al. 2012, Green et al. 2015, Rogers et al. 2015). Their role in providing foraging and/or resting grounds for waterbirds varies according to species, time of year and tidal cycle. Many shorebirds feed on the tidal flats at low tide and move to adjacent supratidal saltpans to continue feeding or resting at high tide, while many others prefer to spend most of their time in the saltpans (Masero et al. 2000, Dias 2009). Likewise, waterbirds such as herons and gulls respond to the tide cycle in a similar way to shorebirds, and ducks and grebes use the supratidal ponds throughout the tide cycle (Warnock et al. 2002). Coastal saltpans also provide breeding grounds for waterbirds (Ramírez et al. 2011, 2012, Rocha et al. 2016).

The Nanpu Saltpan complex in Bohai Bay of the Yellow Sea, North China, is one of the largest coastal saltpans in the world, occupying an area of 290 km² (Figure 1). In 2010, the adjacent intertidal zone was documented as a critical foraging ground for 62% of the Red Knot *Calidris canutus*, and 23% of the Curlew Sandpiper *Calidris ferruginea* flyway populations during northward migration (Rogers et al. 2010, Yang et al. 2011). In 2012, in the Nanpu Saltpans, large numbers of Curlew Sandpipers, Marsh Sandpipers *Tringa stagnatilis*, Black-tailed Godwits *Limosa limosa*, and White-winged Terns *Chlidonias leucopterus* were counted in the northward migration, comprising up to 11%, 1%, 3%, and 4% of their flyway populations, respectively (Hassell et al. 2012). However, these counts were only performed during the northward migration in the supratidal ponds, and they did not cover the whole of the Nanpu Saltpans. Therefore, we do not know the relative importance of these supratidal ponds in relation to the natural tidal flats, and it is likely that the true peak numbers present in these saltpans are higher than the peak numbers reported previously.

Here, we investigate the role of the Nanpu Saltpans for waterbirds during the northward migration, southward migration and boreal winter. We report, for the first time, the waterbird numbers using the entire Nanpu Saltpans at both high and low tide, as well as on the adjacent




Figure 1. Map of Nanpu wetlands. The surveyed areas included inland and nearshore ponds and tidal flats. See legend and text for details.

tidal flats at low tide. These comparative counts will improve our understanding of how migratory waterbirds use the system and the current international importance of these unprotected salt pans for migratory waterbirds in the East Asian-Australasian Flyway.

Methods

Study area

The study area comprised most ponds available for waterbirds as feeding and/or roosting grounds in the Nanpu Salt pans (93 km²), as well as the adjacent tidal flats (57 km²) in the north of Bohai Bay (39°–12'N, 118°8'–28'E) and other supratidal anthropogenic habitats in the area such as shrimp and fish ponds (Figure 1). Similar to other industrial salt pans, the Nanpu Salt pans consist of shallow, interconnected pans of varying sizes separated by dikes. There are three types of ponds: storage, evaporation, and crystallization ponds (Figure 1). Briefly, seawater flows into the storage ponds with the rising tide and circulates through a series of evaporation ponds until it reaches many small crystallization ponds (Britton and Johnson 1987, Masero 2003). The evaporation ponds are operated in two ways: one maintains a high water level (60–100 cm) and one maintains

a low water level (40–60 cm) (salt worker Mr. Hu, pers. comm.). Shorebirds normally use water levels < 10 cm (Ntiamoa-Baidu *et al.* 1998). The low water level ponds can fall below 20 cm in the water transfer process, but the water in high water level ponds rarely falls to levels allowing waterbirds to feed. We focused our survey effort on the evaporation ponds because most waterbirds used these ponds during previous counts. We classified these ponds into two groups according to their distances to the tidal flats: inland ponds (86 km²; 2.4–18.0 km from the tidal flats) and nearshore ponds (Nanpu: 3.50 km², Zuidong: 3.64 km²; 0.3–4.3 km from the tidal flats). Due to saltpan management, the south inland ponds had restricted access in northward migration but free access in southward migration, and the north inland ponds had free access throughout the year (Figure 1). The tidal flats are 1–3 km wide at low tide and are completely submerged from about two hours before high tide (Yang *et al.* 2011). For the counts, the tidal flats were divided into four sectors: Beipu (14 km²), Nanpu (22 km²), Double Bridge (10 km²), and Zuidong (11 km²) (Figure 1).

Surveys

To survey the saltpans, we drove along the perimeter roads and counted from the roads or pond levees, while the tidal flats were surveyed from a dike close to the sea. Waterbirds were counted using telescopes (25–60 x magnification eyepieces) or binoculars (8 x 30). The schedule of waterbird counts and sample sizes are shown in Table 1. For more details, see Appendix S1 in the online supplementary materials.

Northward migration counts - Inland ponds were counted regularly at high and low tide (2014–2016); nearshore ponds were only counted at high tide (2015–2016) because few birds used them during low tide. Tidal flats were counted at low tide synchronously with the low tide counts in the inland ponds (2014–2016). Due to logistical constraints, on the tidal flats all waterbirds were counted in 2014, nine abundant or important shorebird species in 2015 and all shorebirds species in 2016. All counts in 2013 and a few counts in 2014 in inland ponds performed independently of the tidal cycle were also included (Table 1); i.e. a single count included the high-tide period but also part of the low-tide period. All inland pond counts during northward migration focused only on the northern inland ponds because the southern inland ponds had restricted access.

Southward migration and boreal winter counts - We counted the inland ponds and tidal flats in southward migration in 2013–2016 and in boreal winter of 2013. Due to logistical constraints, the tidal flats were counted before or after the counts on the inland ponds, and the counts in the saltpans were performed independently of the tidal cycle. Unfortunately, we did not count nearshore ponds during southward migration and boreal winter because we could not catch high tide in these ponds. The counts in the inland ponds also included the southern inland ponds in 2013 (southward migration and boreal winter) and in 2016 (southward migration).

During high tide, all tidal flats are submerged, so waterbirds have to move into saltpans for roosting or feeding (previous experience at the study site allowed us to know all large feeding and roosting sites). In each round of counts, the maximum number of a species in the study area was obtained from the high-tide counts in the saltpans or from the total number of the synchronised low-tide counts on tidal flats and in saltpans. To estimate a species' habitat usage, the number in the saltpans or on the tidal flats was divided by the maximum number in the study area (times 100%). Bird names followed the BirdLife Checklist Version 9.1 (<http://datazone.birdlife.org/species/taxonomy>).

Foraging activity in the saltpans

The foraging activity of waterbirds in the saltpans during northward migration from 2014 to 2016 was calculated at both the low- and high-tide periods, following Masero *et al.* (2000) and Masero and Pérez-Hurtado (2001). During the low- and high-tide counts we noted the activity of

Table 1. Waterbird numbers (mean \pm SE) in Nanpu Salt pans and adjacent tidal flats. Range (minimum and maximum counts), date of peak counts, shorebird numbers in relation to maximum abundance (%), number of waterbird species using the study area, count area (km²) and number of counts per season, respectively, also are shown. See text for details.

	2013	2014	2015	2016
NORTHWARD MIGRATION				
Whole salt pans at high tide	–	–	19,254 \pm 6,875 6,080 – 29,240 10 & 13 May 99%, 37 53, n = 3	25,014 \pm 2,697 19,820 – 29,860 25 ~ 26 Apr 98%, 47 57, n = 4
Whole salt pans at low tide	–	18,837 \pm 5,582 4,130 – 36,320 30 Apr ~ 1 May 99%, 43 41, n = 6	17,867 \pm 7,117 1,830 – 64,490 19 May 97.4%, 37 41, n = 8	13,464 \pm 3,737 8,130 – 24,000 17 ~ 19 Apr 96%, 37 41, n = 4
Inland ponds at high tide	–	21,680 \pm 6,743 7,520 – 39,950 29 ~ 30 Apr 99%, 36 41, n = 4	12,626 \pm 3,204 2,510 – 27,530 27 Apr 98%, 32 41, n = 7	17,011 \pm 2,885 10,230 – 25,100 17 ~ 18 Apr 96%, 39 41, n = 5
Inland ponds*	40,953 \pm 11,497 20,160 – 96,000 16 May 98.7%, 36 –, n = 6	17,746 \pm 5,389 1,290 – 48,590 4 May 100%, 43 –, n = 8	–	–
Exposed tidal flats	–	39,997 \pm 8,173 12,180 – 73,300 13 ~ 14 May 99.8%, 37 57, n = 6	42,079 \pm 8,714 33,360 – 50,800 4 ~ 5 May –, 9 57, n = 2	35,061 \pm 6,891 21,600 – 51,520 8 May, –, 34 57, n = 4
SOUTHWARD MIGRATION				
Inland ponds**	23,926 \pm 5,586 4,440 – 51,030 21~22 Aug 74.6%, 57 61, n = 8	45,198 \pm 8,919 18,860 – 93,500 23 ~ 24 Aug 51.5%, 53 44, n = 8	13,936 \pm 4,406 6,040 – 21,850 25 Sep 42.7%, 32 57, n = 4	25,656 \pm 4,888 20,770 – 30,540 23 Oct 6.7%, 35 69, n = 2
Exposed tidal flats***	3,381 \pm 887 940 – 5,190 22~23 Nov 56.5%, 29 46, n = 4	13,052 \pm 2,186 5,100 – 20,000 26~27 Sep 88%, 37 46, n = 6	8,377 8,377 23 Sep 70.9%, 24 46, n = 1	8,863 \pm 772 7,980 – 9,640 23 Oct 89.2%, 20 46, n = 2
BOREAL WINTER				
Inland ponds**	431 \pm 383 6 – 1,200 15 Dec 0%, 7 61, n = 3	–	–	–
Exposed tidal flats***	5,743 \pm 663 5,080 – 6,406 15 Dec 18.9%, 12 46, n = 2	–	–	–

* Inland ponds were surveyed independently of the tidal cycle and/or did not cover all ponds.

** Inland ponds were surveyed independently of the tidal cycle.

*** Zuidong tidal flats were excluded from the counts.

individuals as foraging or resting (including preening). For calculating the foraging activity in salt pans, the duration of the low-tide period in the salt pans was considered the same as the emersion period on the tidal flats during low tide; the rest, up to the completion of the tidal cycle, was considered to be high tide (tidal flats submerged period) (Masero and Pérez-Hurtado 2001). For a given species, the number of foraging individuals divided by the total number was its foraging percentage. The overall foraging percentage is based on the average of all species in each survey.

Data analyses

We described the abundance and richness of waterbirds in the Nanpu Salt pans and on the adjacent tidal flats. The abundance of waterbirds was expressed as the mean abundance (MA), maximum abundance (MaxA) and density (birds/km²). Waterbird richness was expressed as the number of species. To compare the relative use of the salt pans and tidal flats during low tide we used the densities of waterbirds. Many ponds were not used for feeding as the water levels were too high, so for calculating density in the salt pans, 58 km² of ponds were selected, usually with birds feeding or roosting, with approximately 62% of the total salt pans studied (93 km²). Note that not all of the 58 km² of the ponds were available for waterbirds during each survey, so this will lead to a conservative estimate of the density in salt pans. The area of the tidal flats used for the density calculation was 57 km².

Internationally important species are defined as those that are present in numbers exceeding 1% of their flyway populations according to the Convention on Wetlands (Ramsar Convention 1971, Yang *et al.* 2011). Waterbird flyway populations followed either Hansen *et al.* (2016) or Waterbird Population Estimates Fifth Edition (WPE5) (Wetlands International 2017).

To describe the community composition during each season and habitat, we calculated the percentage contributed by each species. Waterbirds that were too distant to identify to species level were not included. The northward migration community composition of the tidal flats was based only on the data of 2014, because in 2015 and 2016, not all waterbird species were counted.

After checking for normality by Kolmogorov-Smirnov tests, we used independent t-tests to test the differences between the densities in salt pans and tidal flats during low tide, and differences between mean foraging percentage in inland ponds and nearshore ponds during high tide. We used paired sample t-tests to test the total abundance in the salt pans (inland ponds or whole salt pans). In addition, we used non-parametric Mann-Whitney U-tests to compare the density of each species between the salt pans and tidal flats during low tide and Wilcoxon signed-rank tests to compare the abundance of each species between high tide and low tide in the salt pans (inland ponds or whole salt pans). Statistical analyses were carried out in IBM SPSS Statistics 20.0 for Windows (Armonk, NY: IBM Corp). All tests were two-tailed, and significance levels were set at $P < 0.05$. Values given are means \pm SE.

Results

Northward migration started in early March and ended in early June, with the abundance and richness of waterbird species reaching their peaks in mid-May. Some birds remained throughout June - presumably subadults or weak birds that were unable to migrate further north. Southward migration started in late June and lasted until late November, peaking in late August (Figure 2). After most birds had moved on in southward migration, some Dunlin *Calidris alpina*, Grey Plovers *Pluvialis squatarola*, and Eurasian Curlews *Numenius arquata* overwintered in the study area.

A total of 89 waterbird species were recorded in the study area, among which 13 species were threatened or 'Near Threatened'. Of these, 27 were recorded exclusively in salt pans and one exclusively in the adjacent tidal flats. In the salt pans, a total of 85 species occurred in inland ponds and 42 in nearshore ponds. During the northward migrations, the maximum number of waterbirds recorded in salt pans was 96,000 individuals on 16 May 2013 at high tide, and 64,490 on




Figure 2. Bird numbers (A) and species (B) of waterbirds in inland ponds and tidal flats of Nanpu wetlands. Data are the maximum count and the maximum number of species found in each month.

19 May 2014 at low tide. The maximum number of waterbirds recorded in salt pans during the southward migration was 93,500 individuals on 23–24 August 2014 (Table 1). At low tide, the peak number of waterbirds on the tidal flats occurred on 13–14 May 2014 and 26–27 September 2014, with 73,000 and 20,000 individuals, respectively. Among the waterbirds, shorebirds were the most abundant group of birds both in the Nanpu Salt pans and on the tidal flats (Table 1).

A total of 27 species exceeded the 1% threshold of the flyway populations (Table 2). Among them, 23 species exceeded the threshold in the salt pans (13 species only in salt pans); 14 species exceeded the threshold on the tidal flats (four species only on tidal flats); and 10 species exceeded the threshold in both habitats. Twenty-three species exceeded the threshold in northward migration; 17 species exceeded the threshold in southward migration and three species did so in boreal winter. Of these species, 20 were shorebirds.

The most abundant species in inland ponds were similar between high and low tides. The most abundant species in the inland ponds were different from the nearshore ponds, and the latter were similar to the tidal flats. The most abundant species were different between northward migration and southward migration, both in the salt pans and tidal flats (Appendix S2).

During northward migration, the density of waterbirds on the tidal flats was higher than in the salt pans during low tide ($P = 0.001$, $t = -4.01$, $n = 12$, $df = 22$). Species such as Red Knot and Great Knot *Calidris tenuirostris* had higher densities on the tidal flats than in the salt pans, whereas

Table 2. Waterbirds exceeding 1% of the flyway estimate population (FEP) in Nanpu Salt pans and adjacent tidal flats. Maximum abundance (MaxA) were obtained from the high tide and low tide counts in Nanpu Salt pans, low tide counts on tidal flats, or the sum of both habitats during low tide synchronized counts (the percentage of FEP is shown between parenthesis after MaxA numbers). Peak counts during northward and southward migration in both salt pans and tidal flats also are shown.

Specie name	Scientific name	MaxA (Percentage %)	FEP	Salt pans		Tidal flats	
				Northward migration	Southward migration	Northward migration	Southward migration
Curlew Sandpiper (NT)	<i>Calidris ferruginea</i>	61,890 ^a (69)	90,000 ^b	61,890	2,460	16,080	50
Red Knot (NT)	<i>Calidris canutus</i>	44,460 (40)	110,000 ^b	35,280	210	44,430	160
White-winged Tern	<i>Chlidonias leucopterus</i>	26,030 (3 ~ 26)	100,000 – 1,000,000	5,850	26,030	0	5
Black-headed Gull	<i>Larus ridibundus</i>	21,880 (2 ~ 22)	100,000 – 1,000,001	5,690	21,840	6	1,170
Common Shelduck	<i>Tadorna tadorna</i>	21,090 (14 ~ 21)	100,000 – 150,000	660	21,090	28	170
Red-necked Stint (NT)	<i>Calidris ruficollis</i>	20,590 ^a (4)	475,000 ^b	20,590	1,670	1,940	3
Black-tailed Godwit (NT)	<i>Limosa limosa</i>	17,480 (11)	160,000 ^b	17,480	5,248	14,040	1,350
Great Knot (EN)	<i>Calidris tenuirostris</i>	16,760 (4)	425,000 ^b	739 ⁰	12	16,764	420
Marsh Sandpiper	<i>Tringa stagnatilis</i>	15,850 (12)	130,000 ^b	15,850	12,360	270	410
Black-winged Stilt	<i>Himantopus himantopus</i>	15,190 (15 ~ 61)	25,000 – 100,000	712	15,190	4	1
Pied Avocet	<i>Recurvirostra avosetta</i>	14,250 (14 ~ 57)	25,000 – 100,000	3,440	14,250	360	800
Spotted Redshank	<i>Tringa erythropus</i>	13,490 (13 ~ 54)	25,000 – 100,000	860	13,490	11	140
Dunlin	<i>Calidris alpina</i>	10,500 (0.4 ~ 1.9)	553,900 – 2,674,900	6,640	800	10,180	10,500
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	9,470 (11)	85,000 ^b	9,470	8,380	210	140
Eurasian Curlew (NT)	<i>Numenius arquata</i>	8,270 (8)	100,000	1,250	44 ⁰	8,270	1,540
Kentish Plover	<i>Charadrius alexandrinus</i>	6,900 (7)	100,000	3,630	2,323	4,360	6,900
Relict Gull (VU)	<i>Larus relictus</i>	5,160 ^c (43)	12,000	160	1,430	2,000	3,000
Sanderling	<i>Calidris alba</i>	4,320 (14)	30,000 ^b	1,380	1	4,320	0
Black-necked Grebe	<i>Podiceps nigricollis</i>	3,810 (4 ~ 38)	10,000 – 100,000	1	3,810	0	0
Grey Plover	<i>Pluvialis squatarola</i>	3,620 (5)	80,000 ^b	930	10	3,620	1,380
Great Cormorant	<i>Phalacrocorax carbo</i>	2,480 (3 ~ 10)	25,000 – 100,000	1,060	2,480	0	130
Mew Gull	<i>Larus canus</i>	1,300 (1 ~ 5)	25,000 – 100,000	1,300	360	1	14
Broad-billed Sandpiper	<i>Calidris falcinellus</i>	730 (2)	30,000 ^b	76	200	730	1
Caspian Tern	<i>Hydroprogne caspia</i>	590 (0.6 ~ 2.4)	25,000 – 100,000	590	110	0	140
Ruddy Turnstone	<i>Arenaria interpres</i>	360 (1)	30,000 ^b	210	25	360	0
Asian Dowitcher (NT)	<i>Limnodromus semipalmatus</i>	260 (2)	14,000 ^b	56	107	260	22
Nordmann's Greenshank (EN)	<i>Tringa guttifer</i>	7 (1 ~ 2)	400 – 600	0	0	7	0

NT — Near Threatened, VU — Vulnerable, EN — Endangered, as listed by (IUCN 2015). ^a data from GFN; ^b FEP followed Hansen *et al.* (2016), other species followed WPE5.

^c MaxA counted during boreal winter on tidal flats.

Black-tailed Godwit, Pied Avocet *Recurvirostra avosetta*, and Black-winged Stilt *Himantopus himantopus* had higher densities in the salt pans than on the tidal flats (Appendix S3).

The number of birds in the salt pans during high tide was higher than during low tide ($P = 0.002$, $t = -5.18$, $n = 7$, $df = 6$), but the number of birds on the inland ponds did not differ between high and low tide ($P = 0.119$, $t = 1.68$, $n = 13$, $df = 12$). Few birds used nearshore ponds during low tide (one count recorded 814 birds on 9 May 2016). When comparing the number in all the salt pans during high and low tide, Red Knot ($P = 0.018$, $Z = -2.37$), Great Knot ($P = 0.018$, $Z = -2.37$), Black-headed Gull *Larus ridibundus* ($P = 0.028$, $Z = -2.20$), Eurasian Curlew ($P = 0.043$, $Z = -2.02$), and Bar-tailed Godwit *Limosa lapponica* ($P = 0.028$, $Z = 2.20$; all $n = 7$) were more abundant during high tide. For the inland ponds only, Black-tailed Godwit ($P = 0.023$, $Z = -2.27$), Common Shelduck *Tadorna tadorna* ($P = 0.034$, $Z = -2.12$), Dunlin ($P = 0.028$, $Z = -2.20$), Red Knot ($P = 0.021$, $Z = -2.31$), Red-necked Stint *Calidris ruficollis* ($P = 0.019$, $Z = -2.35$), and White-winged Tern ($P = 0.028$, $Z = -2.20$; all $n = 13$) were more abundant during high tide. Black-tailed Godwit, Curlew Sandpiper, Dunlin, Marsh Sandpiper, Pied Avocet, Red Knot, Red-necked Stint also were abundant during low tide (> 500 birds on average). During high tide, species like Red Knot and Great Knot were mainly present in nearshore ponds (Figure 3).

During northward migration, most waterbirds in the inland ponds were actively feeding (except for Great Cormorant *Phalacrocorax carbo* which were just roosting), while the waterbirds (mainly shorebirds) in the nearshore ponds were roosting during high tide (Figure 4). However, Great




Figure 3. Bird numbers of the main species using inland and nearshore ponds during northward migration. High tide and low tide surveys were conducted in consecutive days or in days as close as possible. CAST= Caspian Tern, for other acronyms, see legend in Appendix S2. See text for methodological details.




Figure 4. Feeding percentage of waterbirds in inland and nearshore ponds during northward migration. CAST = Caspian Tern, for other acronyms, see legend in Appendix S2. See text for methodological details.

Cormorants were found feeding in a storage pond in southward migration, which were dry in northward migration. The mean foraging percentage (average of each count) in the inland ponds was high during both high tide ($80.7 \pm 3.9\%$, $n = 8$) and low tide ($91.2 \pm 0.8\%$, $n = 8$). The foraging rate in the nearshore ponds is lower than in the inland ponds during high tide ($10.4 \pm 4.2\%$, $n = 5$, $P < 0.001$, $t = -11.69$, $df = 11$).

The percentage of birds using different habitats and during different tides grouped into three categories (Figure 5). White-winged Tern, Pied Avocet, Marsh Sandpiper, Spotted Redshank *Tringa erythropus*, Black-winged Stilts, Common Shelduck, Black-tailed Godwit, Black-headed Gull, and Caspian Tern *Hydroprogne caspia* mainly used salt pans. Bar-tailed Godwit, Grey Plover, Eurasian Curlew, Great Knot, and Red Knot mainly used tidal flats, and they also used nearshore ponds during high tide. The remaining species occurred in both habitats (Figure 5).

Discussion

This study demonstrates the important role of the Nanpu Salt pans for migratory waterbirds using the East Asian-Australasian Flyway. Large numbers of many waterbird species used the Nanpu Salt pans as feeding and/or roosting sites during both the northward migration and southward migrations. A total of 89 species was recorded, of which 27 occurred in numbers of international importance, with 13 threatened or 'Near Threatened' species. Salt pans are not only supplementary feeding grounds for some waterbirds during high tide but also major feeding grounds for some species throughout the tidal cycle.

Habitat use

There are no natural wetlands in Nanpu, except the tidal flats, and all supratidal natural habitats had been converted into salt pans or aquaculture ponds. Both tidal flats and salt pans can provide feeding grounds for waterbirds; however, during high tide, when tidal flats are totally submerged, only salt pans can provide roosting and foraging grounds.

We showed that salt pans and tidal flats use patterns by waterbirds varied among species and seasonally. Thus, both habitats were used heavily by waterbirds during migration, but only a few species used Nanpu as a wintering area. The abundance of waterbirds recorded on southward




Figure 5. Percentage of birds using salt pans or tidal flats during northward migration at low or high tide. In these foraging behaviour data, there were no Great Knot and Grey Plover recorded during low tide, no Bar-tailed Godwit during both low tide and high tide in inland ponds, no White-winged Tern, Caspian Tern, and Spotted Redshank in nearshore ponds during high tide. The foraging percentages of Great Knot, Sanderling, Eurasian Curlew, and Common Shelduck in nearshore ponds were zero. CAST= Caspian Tern, for other acronyms, see legend in Appendix B. See text for methodological details.

migration 2015 and 2016 was lower than on southward migration 2013 and 2014, perhaps because there were fewer surveys in 2015 and 2016 than in 2013 and 2014. We did not count the nearshore ponds during southward migration and boreal winter, so the number of waterbirds in salt pans in both seasons may be higher than we recorded.

Migratory shorebirds were the most abundant group during northward migration. On southward migration, the percentage of shorebirds was lower than on northward migration, because the most abundant species on northward migration such as Red Knot, Great Knot and Curlew Sandpiper seldom used the study area on southward migration. We also noted during the counts that Pied Avocet, Black-winged Stilt, Kentish Plover, Common Redshank *Tringa totanus*, Little Tern *Sterna albifrons*, Common Tern *Sterna hirundo*, Gull-billed Tern *Gelochelidon nilotica*, and Eastern Spot-billed Duck *Anas zonorhyncha* breed in Nanpu Salt pans.

There are three hypotheses on why shorebirds feed in salt pans adjacent to natural intertidal areas: the 'supplementary food hypothesis', the 'preference hypothesis', and the 'disturbance hypothesis' (Masero *et al.* 2000). Although the shellfish harvest on tidal flats may affect the foraging activities of waterbirds (Navedo and Masero 2007), the number of shellfish collectors was few (Yang *et al.* 2016) and the tidal flats seem large enough for birds to feed free from humans (pers. obs.). Therefore, this hypothesis should be rejected.

The use of salt pans by Red Knots, Relict Gull *Larus relictus*, Dunlin, and Sanderling was consistent with the 'supplementary food hypothesis', as most birds used the tidal flats as foraging grounds and only fed in the salt pans at high tide. Other species such as Great Knot, Eurasian Curlew, Grey Plover and Bar-tailed Godwit almost only used the salt pans as roosting sites during high tide.

The 'preference hypothesis' seems to explain the foraging patterns of White-winged Tern, Pied Avocet, Marsh Sandpiper, Spotted Redshank, Black-winged Stilt, Common Shelduck, Black-tailed Godwit, and Black-headed Gull as they foraged in the inland ponds throughout the tidal cycle. The habitat use patterns of Curlew Sandpiper, Red-necked Stint, Sharp-tailed Sandpiper *Calidris acuminata*, and Kentish Plover were variable because many individuals fed on tidal flats during low tide, but many others also fed in salt pans during both high tide and low tide.

The findings of previous studies on the foraging use of coastal salt pans in other parts of the world were generally similar (e.g. Velasquez and Hockey 1992, Masero *et al.* 2000, Warnock *et al.* 2002, Dias 2009). Black-tailed Godwit, Black-winged Stilt, Pied Avocet, Spotted Redshank, Kentish Plover (Masero *et al.* 2000, Dias 2009), Black-necked Stilt *Himantopus mexicanus*, and American Avocet *Recurvirostra americana* (Warnock *et al.* 2002) preferred salt pans for foraging, while Grey Plover, Ruddy Turnstone *Arenaria interpres*, Bar-tailed Godwit, and Red Knot used salt pans as supplementary feeding grounds (Masero *et al.* 2000, Dias 2009).

The number and behaviour of waterbirds in inland ponds and nearshore ponds showed that these two groups of ponds had different functions for waterbirds in daytime. Inland ponds were mainly used as feeding grounds, whereas nearshore ponds were mainly used as high-tide roosting sites by shorebirds. High-tide roost sites are important to waterbirds because roosting sites can constrain their use of forage areas (Dias *et al.* 2006, Rogers *et al.* 2006b). These constraints are particularly relevant in the context of Nanpu, since as stated previously, this area is densely populated and there are few sites available for roosting. We found a large number of shorebirds roosting in the nearshore ponds at high tide during daylight. These ponds are also used as high-tide roosting sites at night (Chris Hassell pers. comm.). Distance is an important factor affecting high-tide roosting site selection, and shorebirds using these ponds would be minimising the costs of flying from tidal flats to roost sites (Rogers *et al.* 2006a).

That salt pans may buffer the loss of natural habitats is supported by a large amount of evidence (Masero and Pérez-Hurtado 2001, Masero 2003, Sripanomyom *et al.* 2011, Dias *et al.* 2014). The use patterns of salt pans and tidal flats by waterbirds found in this study indicate that the buffer role of salt pans varies among species and relies on the extent to which they use the salt pans. For the species preferring to feed in salt pans, such as Black-tailed Godwits and Black-winged Stilts, the loss of natural tidal flats could be largely compensated if the conditions in salt pans are suitable for feeding. For the species that use both salt pans and tidal flats, such as Curlew Sandpiper and Red-necked Stint, the loss of natural tidal flats can be partly compensated. This is important for those species that rely heavily on the Yellow Sea and whose flyway populations are dramatically declining (Studds *et al.* 2017). However, for the species that seldom feed in the salt pans, such as Great Knot and Red Knot, the role of salt pans for feeding is limited, although they are important high-tide roosting sites.

International importance

Our findings confirmed the international importance of the Nanpu Salt pans for migrating waterbirds in the East Asian-Australasian Flyway. The Nanpu Salt pans provided feeding and roosting sites for large numbers of waterbirds and supported species of conservation concern. Thirteen threatened or 'Near Threatened' species on the IUCN Red List were recorded in salt pans or on tidal flats. One is Critically Endangered (Spoon-billed Sandpiper *Calidris pygmaea*). Four are 'Endangered' (Great Knot, Nordmann's Greenshank *Tringa guttifer*, Far Eastern Curlew *Numenius madagascariensis*, and Oriental White Stork *Ciconia boyciana*). Two are 'Vulnerable' (Relict Gull and Saunders' Gull *Chroicocephalus saundersi*). Six are 'Near Threatened' (Curlew Sandpiper, Red-necked Stint, Red Knot, Black-tailed Godwit, Eurasian Curlew, and Asian Dowitcher *Limnodromus semipalmatus*). The number of Great Knot, Nordmann's Greenshank, Relict Gull, Curlew Sandpiper, Black-tailed Godwit, and Eurasian Curlew also exceeded 1% of their flyway populations (Hansen *et al.* 2016, Wetlands International 2017). The Nanpu tidal flats and salt pans were listed in China's top 21 priority coastal wetlands for waterbird conservation (Xia *et al.* 2017).

From 1993 to 2012, 10 shorebird taxa that refuel on the Yellow Sea tidal mudflats have declined by 65% (Studds *et al.* 2017). These taxa include species that also showed a declining trend in this study, such as Curlew Sandpiper. The latter has been listed as a 'Critically Endangered' species in Australia under its Environment Protection and Biodiversity Conservation Act 1999 and as 'Near Threatened' on the IUCN Red List (IUCN 2017). This species has suffered an 80.5% population

decline over the last three generations (a generation time of 7.6 years), with an annual rate of decline of 7.5% (Garnett *et al.* 2011, BirdLife International 2016a, Studds *et al.* 2017). The maximum number of Curlew Sandpipers in the Nanpu Saltpans was 61,890 (May 16, 2013) recorded in a single pond (3.27 km²) (Hassell *et al.* 2013), which comprised approximately 69% of the flyway population (Hansen *et al.* 2016). Together with 20,580 Red-necked Stints and other species that fed in the same ponds, this single pond held more than 90,000 birds (Hassell *et al.* 2013). However, the maximum percentages of the estimated flyway population of Curlew Sandpiper recorded in the saltpans were only 26% (23,700 birds) in 2014, 39% (35,270) in 2015, and 18% (16,570) in 2016. Although four years of data are too short to detect population declines, given the high natural variability of shorebird numbers from year to year, the decline of the Curlew Sandpiper in this area is still of concern. This is because the environmental factors in the saltpans and tidal flats were similar for four years, and a decline was also found in other species, such as Red-necked Stint, Black-tailed Godwit, and Red Knot. Flyway populations of these species have declined sharply in recent decades (Studds *et al.* 2017, Wetlands International 2017).

The maximum count of Black-tailed Godwit in the Nanpu Saltpans was 17,480 (22–23 April 2013), comprising approximately 11% of the flyway population (Wetlands International 2017). The global population of Black-tailed Godwits may have declined by approximately 25% since 1990 (BirdLife International 2016b). The maximum count of Black-tailed Godwits in northward migration declined during the study period, with 14,790 in 2015 and 10,180 in 2016.

Approximately 34,200 Red Knots (90% were the subspecies *piersmai*) were recorded feeding or roosting in a single pond (3.30 km²) on 29 May 2013, an estimated 51% of the world population of the subspecies *piersmai* (Hassell *et al.* 2013). The summer survival of Red Knot has sharply declined since 2012, and coastal intertidal habitat loss in the Yellow Sea is the most likely cause (Piersma *et al.* 2016). The maximum number of Red Knot in the study area decreased from 44,430 (2014) to 30,750 (2015) and 15,340 (2016), suggesting that the summer survival of Red Knot is probably still declining. The flyway population of Red Knots in the study area sharply increased from 13% in 2007 to 62% in 2010 due to tidal flat loss in the surrounding area (Yang *et al.* 2011), and the recent decrease suggests that their prospects are not good.

Conservation implications

Despite its conservation importance, the birds and their habitats in the Nanpu Saltpans have not been properly protected to date. Habitat loss and human disturbance are increasing in recent years in our study area. For instance, in the adjacent Tianjin Municipality, and Fengnan, Tangshan City, Hebei Province, more than 100 km² of saltpans have been lost since Barter *et al.* (2001) identified the international importance of this area for shorebirds (Melville *et al.* 2016). Since 2010, at least 20 km² in the Nanpu Saltpans have been converted to industrial land. In addition, another large project plans to use 100 km² of the Nanpu Saltpans (Hebei Daily 2015). Moreover, construction of a new road across the Nanpu Saltpans starting in 2015 has cut through many ponds, including the one with 96,000 birds on 16 May 2013. This has caused sharp declines of waterbirds in these ponds (pers. obs.). Therefore, an urgent need exists to prevent further loss of the remaining saltpans and adjacent tidal flats. Other options include restoring lost habitats and modifying existing artificial coastal habitats to maximise shorebird foraging and roosting opportunities (Studds *et al.* 2017). However, we do not suggest building more saltpans to replace natural feeding habitats because the saltpans attract different species to the tidal flats and this habitat is not suitable for all (Masero 2003).

Today, some of the most important areas for shorebirds include coastal saltpans (Rogers *et al.* 2015). It has been suggested that the presence of coastal saltpans could increase the carrying capacity of some estuarine systems for non-breeding shorebirds (Batty 1992, Masero *et al.* 2000). However, across the world, large saltpans are disappearing or are being abandoned due to changing economics driven by industrialisation, urbanisation and competing land uses (Masero 2003). Saltpans do not only produce salt (see review by Rocha *et al.* 2012). In the Nanpu Saltpans,

storage and evaporation ponds are used for aquaculture, while in many other evaporation ponds the crustacean brine shrimp *Artemia* is harvested. All these activities can benefit local people. However, many ponds have high water levels to meet the needs of salt production (especially the high-water level evaporation ponds) or aquaculture, which limits their use by foraging shorebirds. Environmental conditions in the salt pans can affect the availability and accessibility of prey for waterbirds, so the available foraging area within the salt pans can vary several-fold (Dias 2009). Therefore, salt pans need waterbird-friendly management to maintain their importance as feeding habitats for shorebirds (Rocha *et al.* 2017). For example, during the migration season, water levels and the available foraging area of the ponds should be manipulated whenever possible (Dias *et al.* 2014). For shorebirds, the best water level would be 0–10 cm, and for non-shorebirds 30–40 cm (Lei 2017). Salinity should be maintained in several ranges, especially 20–60 ppt (for fish and fish-eating birds) and ~140 ppt for shorebirds, waterfowl, gulls and terns (Warnock *et al.* 2002).

This study clearly demonstrates the joint ecological function of The Nanpu Saltpan complex and adjacent tidal flats as a key staging area for waterbirds in the EAAF, and both urgently warrant protected status. Further studies in Nanpu complex should also focus on the potential impact of high-voltage power lines, since collision killed many birds in Nanpu Salt pans (pers. obs.), as well as on illegal eggs collection of breeding shorebirds such as Kentish Plover and Pied Avocet (Que *et al.* 2015, Lei 2017).

Supplementary Material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S0959270917000508>

Acknowledgements

This research was supported by the National Natural Science Foundation of China (Zhengwang Zhang, grant number 31572288), (Hong-yan Yang, grant number 31301885); and the Grant from WWF-China Programme. Theunis Piersma acknowledges funding from WWF-Netherlands and the Netherlands Organization for Scientific Research (NWO). We thank volunteers, Katarzyna Kucharska, Tong Mu, Jian Zhao and Peter Crighton's great help. We thank the Global Flyway Network team (Chris Hassell, Adrian Boyle, Matthew Slaymaker, Bob Loos, supported by WWF-Netherlands and BirdLife Netherlands) who helped survey the tidal flats. Thanks to Jason Loghry, Jin Liu, Ginny Chen, Katherine Leung, Chao Yan, Fuxing Wu, Yu Liu, Pinjia Que, Yanjing Chang, Jiayu Liu, Ying Chen, David Melville, Jan Erik Nilsen, Terry Townshend, Per Alstrom, and friends from GSWJ birdwatching groups who helped with the surveys. Thanks also to Yajing Chang and Lei Guan who helped to produce the map. Professor Zhijun Ma reviewed an early version of the manuscript and gave valuable suggestions. Terry Townshend helped revise the language for the figures, table legends and appendices. We would also like to thank P. Atkinson and two anonymous referees for their comments on earlier versions of the manuscript.

References

- Amano, T., Szekely, T., Koyama, K., Amano, H. and Sutherland, W. J. (2010) A framework for monitoring the status of populations: An example from wader populations in the East Asian–Australasian flyway. *Biol. Conserv.* 143: 2238–2247.
- Bamford, M., Watkins, D., Bancroft, W., Tischler, G. and Wahl, J. (2008) *Migratory shorebirds of the East Asian–Australasian flyway: Population estimates and internationally important sites*. Canberra, Australia: Wetlands International, Oceania.
- Barter, M. (2002) *Shorebirds of the Yellow Sea: importance, threats and conservation status*. Canberra, Australia: Wetlands International–Oceania. (International Wader Studies 12).

- Barter, M., Li, Z. W. and Xu, J. L. (2001) Shorebird numbers on the Tianjin Municipality coast in May 2000. *Stilt* 39: 29.
- Batty, L. (1992) The wader communities of a saline and an intertidal site on the Ria Formosa, Portugal. *Wader Study Group Bull.* 66: 66–72.
- BirdLife International. (2016a) Species factsheet: *Calidris ferruginea*. Downloaded from <http://www.birdlife.org> on 15/08/2016.
- BirdLife International. (2016b) Species factsheet: *Limosa limosa*. Downloaded from <http://www.birdlife.org> on 12/03/2016.
- Britton, R. and Johnson, A. (1987) An ecological account of a Mediterranean salina: the Salin de Giraud, Camargue (S. France). *Biol. Conserv.* 42: 185–230.
- Choi, C. Y., Gan, X. J., Hua, N., Wang, Y. and Ma, Z. J. (2014) The habitat use and home range analysis of dunlin (*Calidris alpina*) in Chongming Dongtan, China and their conservation implications. *Wetlands* 34: 255–266.
- Czech, H. A. and Parsons, K. C. (2002) Agricultural wetlands and waterbirds: A review. *Waterbirds* 25: 56–65.
- Desrochers, D. W., Keagy, J. C. and Cristol, D. A. (2008) Created versus natural wetlands: Avian communities in Virginia salt marshes. *Ecoscience* 15: 36–43.
- Dias, M. P. (2009) Use of salt ponds by wintering shorebirds throughout the tidal cycle. *Waterbirds* 32: 531–537.
- Dias, M. P., Granadeiro, J. P., Lecoq, M., Santos, C. D. and Palmeirim, J. M. (2006) Distance to high-tide roosts constrains the use of foraging areas by dunlins: Implications for the management of estuarine wetlands. *Biol. Conserv.* 131: 446–452.
- Dias, M. P., Lecoq, M., Moniz, F. and Rabaga, J. E. (2014) Can human-made saltpans represent an alternative habitat for shorebirds? Implications for a predictable loss of estuarine sediment flats. *Environ. Manage.* 53: 163–171.
- Elphick, C. S. (2000) Functional equivalency between rice fields and seminatural wetland habitats. *Conserv. Biol.* 14: 181–191.
- Flaherty, M. and Karnjanakesorn, C. (1995) Marine shrimp aquaculture and natural resource degradation in Thailand. *Environ. Manage.* 19: 27–37.
- Garnett, S., Szabo, J. and Dutson, G. (2011) *The action plan for Australian birds 2010*. Canberra, Australia: CSIRO Publishing.
- Green, J. M. H., Sripanomyom, S., Giam, X. and Wilcove, D. S. (2015) The ecology and economics of shorebird conservation in a tropical human - modified landscape. *J. Appl. Ecol.* 52: 1483–1491.
- Hansen, B. D., Fuller, R. A., Watkins, D., Rogers, D. I., Clemens, R. S., Newman, M., Woehler, E. J., and Weller, D. R. (2016) *Revision of the East Asian-Australasian Flyway population estimates for 37 listed migratory shorebird species*. Melbourne: BirdLife Australia.
- Hassell, C., Boyle, A., Slaymaker, M. and Chan, Y. C. (2012) Red Knot northward migration through Bohai Bay, China, field trip report April & May 2012. Broome: Global Flyway Network.
- Hassell, C., Boyle, A., Slaymaker, M., Chan, Y. C. and Piersma, T. (2013) Red Knot northward migration through Bohai Bay, China, field trip report April & May 2013. Broome: Global Flyway Network.
- Hebei Daily (2015) Beijing and Hebei jointly issued 'The industry development plan of Beijing (Caofeidian) Modern Industry Development Experimental Zone'. http://hbrb.hebnews.cn/html/2015-09/26/content_73633.htm (In Chinese) (accessed on 1 October 2015).
- Houston, W., Black, R., Elder, R., Black, L. and Segal, R. (2012) Conservation value of solar salt ponds in coastal tropical eastern Australia to waterbirds and migratory shorebirds. *Pac. Conserv. Biol.* 18: 100–122.
- Hua, N., Tan, K., Chen, Y. and Ma, Z. J. (2015) Key research issues concerning the conservation of migratory shorebirds in the Yellow Sea region. *Bird Conserv. Internatn.* 25: 38–52.
- IUCN (2017) *The IUCN Red List of Threatened Species. Version 2017.2*. www.iucnredlist.org (accessed on 1 April 2017).
- Kirby, J. (2010) *Review 2-Review of Current Knowledge of Bird Flyways, Principal Knowledge Gaps and Conservation Priorities*. CMS Scientific Council: Flyway Working Group Reviews.
- Lei, W. P. (2017) *Studies on the waterbirds used saltpans in north of Bohai Bay*. PhD thesis. Beijing: Beijing Normal University.

- Li, D. L., Chen, S. H., Lloyd, H., Zhu, S. Y., Shan, K., and Zhang, Z. W. (2013) The importance of artificial habitats to migratory waterbirds within a natural/artificial wetland mosaic, Yellow River Delta, China. *Bird Conserv. Internatn.* 23: 184–198.
- Li, S., Zhao, S., Zhou, X. and Li, H. (2011) Current environmental situation analysis on the Changlu Saltworks in North China. *Tianjin Science & Technology*: 114–116. (In Chinese).
- Ma, Z. J., Cai, Y. T., Li, B. and Chen, J. K. (2010) Managing wetland habitats for waterbirds: An international perspective. *Wetlands* 30: 15–27.
- Ma, Z. J., Jing, K., Tang, S. M. and Chen, J. K. (2002) Shorebirds in the eastern intertidal areas of Chongming Island during the 2001 northward migration. *Stilt* 41: 6–10.
- Ma, Z. J., Li, B., Zhao, B., Jing, K., Tang, S. M. and Chen, J. K. (2004) Are artificial wetlands good alternatives to natural wetlands for waterbirds? A case study on Chongming Island, China. *Biodivers. Conserv.* 13: 333–350.
- Ma, Z. J., Melville, D. S., Liu, J. G., Chen, Y., Yang, H. Y., Ren, W. W., Zhang, Z. W., Piersma, T. and Li, B. (2014) Rethinking China's new great wall. *Science* 346: 912–914.
- MacKinnon, J., Verkuil, Y. and Murray, N. (2012) *IUCN situation analysis on East and Southeast Asian intertidal habitats, with particular reference to the Yellow Sea (including the Bohai Sea)*. Gland, Switzerland and Cambridge, UK: IUCN. (Occasional paper of the IUCN species survival commission 47).
- Masero, J. A. (2003) Assessing alternative anthropogenic habitats for conserving waterbirds: salinas as buffer areas against the impact of natural habitat loss for shorebirds. *Biodivers. Conserv.* 12: 1157–1173.
- Masero, J. A. and Pérez-Hurtado, A. (2001) Importance of the supratidal habitats for maintaining overwintering shorebird populations: how redshanks use tidal mudflats and adjacent saltworks in southern Europe. *The Condor* 103: 21–30.
- Masero, J. A., Perez-Hurtado, A., Castro, M., and Arroyo, G. M. (2000) Complementary use of intertidal mudflats and adjacent salinas by foraging waders. *Ardea* 88: 177–191.
- Melville, D. S., Chen, Y. and Ma, Z. J. (2016) Shorebirds along the Yellow Sea coast of China face an uncertain future - a review of threats. *Emu* 116: 100–110.
- Milsom, T. P., Langton, S. D., Parkin, W. K., Peel, S., Bishop, J. D., Hart, J. D. and Moore, N. P. (2000) Habitat models of bird species' distribution: an aid to the management of coastal grazing marshes. *J. Appl. Ecol.* 37: 706–727.
- Murray, N. J., Clemens, R. S., Phinn, S. R., Possingham, H. P. and Fuller, R. A. (2014) Tracking the rapid loss of tidal wetlands in the Yellow Sea. *Front. Ecol. Environ.* 12: 267–272.
- Navedo, J. G. and Fernández, G. (in press) Use of semi-intensive shrimp farms as alternative foraging areas by migratory shorebird populations in tropical areas. *Bird Conserv. Internatn.* <https://doi.org/10.1017/S0959270918000151>
- Navedo, J. G., Fernández, G., Valdivia, N., Drever, M. C. and Masero, J. A. (2017) Identifying management actions to increase foraging opportunities for shorebirds at semi-intensive shrimp farms. *J. Appl. Ecol.* 54: 567–576.
- Navedo, J. G. and Masero, J. A. (2007) Measuring potential negative effects of traditional harvesting practices on waterbirds: a case study with migrating curlews. *Anim. Conserv.* 10: 88–94.
- Ntiamoa-Baidu, Y., Piersma, T., Wiersma, P., Poot, M., Battley, P. and Gordon, C. (1998) Water depth selection, daily feeding routines and diets of waterbirds in coastal lagoons in Ghana. *Ibis* 140: 89–103.
- Piersma, T., Lok, T., Chen, Y., Hassell, C. J., Yang, H. Y., Boyle, A., Slaymaker, M., Chan, Y. C., Melville, D. S. and Zhang, Z. W. (2016) Simultaneous declines in summer survival of three shorebird species signals a flyway at risk. *J. Appl. Ecol.* 53: 479–490.
- Que, P. J., Chang, Y. J., Eberhart-Phillips, L., Liu, Y., Szekely, T. and Zhang, Z. W. (2015) Low nest survival of a breeding shorebird in Bohai Bay, China. *J. Ornithol.* 156: 297–307.
- Ramírez, F., Abdennadher, A., Sanpera, C., Jover, L., Wassenaar, L. I. and Hobson, K. A. (2011) Assessing waterbird habitat use in coastal evaporative systems using stable isotopes ($\delta^{13}\text{C}$, $\delta^{15}\text{N}$ and δD) as environmental tracers. *Estuarine, Coastal and Shelf Science* 92: 217–222.

- Ramírez, E., Navarro, J., Afán, I., Hobson, K. A., Delgado, A. and Forero, M. G. (2012) Adapting to a changing world: unraveling the role of man-made habitats as alternative feeding areas for slender-billed gull (*Chroicocephalus genei*). *PLoS one* 7: e47551.
- Ramsar Convention (1971) *Convention on Wetlands of International Importance especially as Waterfowl Habitat. Ramsar (Iran), 2 February 1971*. UN Treaty Series No. 14583.
- Rhymer, C. M., Robinson, R. A., Smart, J. and Whittingham, M. J. (2010) Can ecosystem services be integrated with conservation? A case study of breeding waders on grassland. *Ibis* 152: 698–712.
- Rocha, A., Fonseca, D., Masero, J. A. and Ramos, J. A. (2016) Coastal saltpans are a good alternative breeding habitat for Kentish plover *Charadrius alexandrinus* when umbrella species are present. *J. Avian Biol.* 47: 824–833.
- Rocha, A. R., Ramos, J. A., Paredes, T. and Masero, J. A. (2017) Coastal saltpans as foraging grounds for migrating shorebirds: an experimentally drained fish pond in Portugal. *Hydrobiologia* 790: 141–155.
- Rocha, R. D. M., Costa, D. F., Lucena-Filho, M. A., Bezerra, R. M., Medeiros, D. H., Azevedo-Silva, A. M., Araujo, C. N. and Xavier-Filho, L. (2012) Brazilian solar saltworks-ancient uses and future possibilities. *Aquat. Biosyst.* 8: 8.
- Rogers, D. I., Battley, P. F., Piersma, T., Gils, Van, J. A. and Rogers, K. G. (2006a) High-tide habitat choice: insights from modelling roost selection by shorebirds around a tropical bay. *Anim. Behav.* 72: 563–575.
- Rogers, D. I., Piersma, T. and Hassell, C. J. (2006b) Roost availability may constrain shorebird distribution: Exploring the energetic costs of roosting and disturbance around a tropical bay. *Biol. Conserv.* 133: 225–235.
- Rogers, D. I., Stamation, K., Loyn, R. H. and Menkhorst, P. (2015) *Literature review: management of non-tidal ponds for shorebirds*. Heidelberg, Victoria: Arthur Rylah Institute for Environmental Research, Department of Environment, Land, Water and Planning.
- Rogers, D. I., Yang, H. Y., Hassell, C. J., Boyle, A. N., Rogers, K. G., Chen, B., Zhang, Z. W. and Piersma, T. (2010) Red knots (*Calidris canutus piersmai* and *C. c. rogersi*) depend on a small threatened staging area in Bohai Bay, China. *Emu* 110: 307–315.
- Sanchez-Guzman, J. M., Moran, R., Masero, J. A., Corbacho, C., Costillo, E., Villegas, A. and Santiago-Quesada, F. (2007) Identifying new buffer areas for conserving waterbirds in the Mediterranean basin: the importance of the rice fields in Extremadura, Spain. *Biodivers. Conserv.* 16: 3333–3344.
- Sebastián-González, E. and Green, A. J. (2016) Reduction of avian diversity in created versus natural and restored wetlands. *Ecography* 39: 1176–1184.
- Sripanomyom, S., Round, P. D., Savini, T., Trisurat, Y. and Gale, G. A. (2011) Traditional salt-pans hold major concentrations of overwintering shorebirds in Southeast Asia. *Biol. Conserv.* 144: 526–537.
- Studds, C. E., Kendall, B. E., Murray, N. J., Wilson, H. B., Rogers, D. I., Clemens, R. S., Gosbell, K., Hassell, C. J., Jessop, R., Melville, D. S., Milton, D. A., Minton, C. D. T., Possingham, H. P., Riegen, A. C., Straw, P., Woehler, E. J. and Fuller, R. A. (2017) Rapid population decline in migratory shorebirds relying on Yellow Sea tidal mudflats as stopover sites. *Nat. Commun.* 8: 14895.
- Velasquez, C. R. and Hockey, P. A. R. (1992) The importance of supratidal foraging habitats for waders at a south temperate estuary. *Ardea* 80: 243–253.
- Warnock, N., Page, G. W., Ruhlen, T. D., Nur, N., Takekawa, J. Y. and Hanson, J. T. (2002) Management and conservation of San Francisco Bay salt ponds: Effects of pond salinity, area, tide, and season on Pacific Flyway waterbirds. *Waterbirds* 25: 79–92.
- Warnock, S. E. and Takekawa, J. Y. (1995) Habitat preferences of wintering shorebirds in a temporally changing environment: Western Sandpipers in the San Francisco Bay estuary. *The Auk*: 920–930.
- Wetlands International (2017) *Waterbird Population Estimates*. Retrieved from wpe.wetlands.org on 5 Jul 2017.
- Wilson, H. B., Kendall, B. E., Fuller, R. A., Milton, D. A. and Possingham, H. P. (2011) Analyzing variability and the rate of decline of migratory shorebirds in Moreton Bay, Australia. *Conserv. Biol.* 25: 758–766.

- Xia, S., Yu, X., Millington, S., Liu, Y., Jia, Y., Wang, L., Hou, X. and Jiang, L. (2017) Identifying priority sites and gaps for the conservation of migratory waterbirds in China's coastal wetlands. *Biol. Conserv.* 210: 72–82.
- Yang, H. Y., Chen, B., Barter, M., Piersma, T., Zhou, C. F., Li, F. S. and Zhang, Z. W. (2011) Impacts of tidal land reclamation in Bohai Bay, China: ongoing losses of critical Yellow Sea waterbird staging and wintering sites. *Bird Conserv. Internatn.* 21: 241–259.
- Yang, H. Y., Chen, B., Piersma, T., Zhang, Z. W. and Ding, C. Q. (2016) Molluscs of an intertidal soft-sediment area in China: Does overfishing explain a high density but low diversity community that benefits staging shorebirds? *J. Sea. Res.* 109: 20–28.
- Yao, Y. C., Ren, C. Y., Wang, Z. M., Wang, C. and Deng, P. Y. (2016) Monitoring of salt ponds and aquaculture ponds in the coastal zone of China in 1985 and 2010. *Wetland Sci.* 14: 874–882. (In Chinese).

WEIPAN LEI^{1,2,3}, JOSÉ A. MASERO², THEUNIS PIERSMA^{3,4}, BINGRUN ZHU^{1,3},
HONG-YAN YANG^{3,4}, ZHENGWANG ZHANG^{1*}

¹Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life Sciences, Beijing Normal University, Beijing, 100875, China.

²Department de Anatomy, Cell Biology and Zoology, University of Extremadura, Avenida de Elvas s/n, 06071 Badajoz, Spain.

³Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, PO Box 11103, 9700 CC Groningen, The Netherlands.

⁴NIOZ Royal Netherlands Institute for Sea Research, Department of Coastal Systems and Utrecht University, PO Box 59, 1790 AB Den Burg, Texel, The Netherlands.

* Author for correspondence; mail: zzw@bnu.edu.cn

Received 6 January 2017; revision accepted 3 October 2017;

Published online 17 September 2018