

University of Groningen

Morphological and digestive adjustments buffer performance

Zhang, Shou-Dong; Ma, Zhijun; Choi, Chi-Yeung; Peng, He-Bo; Melville, David S.; Zhao, Tian-Tian; Bai, Qing-Quan; Liu, Wen-Liang; Chan, Ying-Chi; van Gils, Jan A.

Published in:
 Ecology and Evolution

DOI:
[10.1002/ece3.5013](https://doi.org/10.1002/ece3.5013)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Zhang, S-D., Ma, Z., Choi, C-Y., Peng, H-B., Melville, D. S., Zhao, T-T., Bai, Q-Q., Liu, W-L., Chan, Y-C., van Gils, J. A., & Piersma, T. (2019). Morphological and digestive adjustments buffer performance: How staging shorebirds cope with severe food declines. *Ecology and Evolution*, 9(7), 3868-3878. <https://doi.org/10.1002/ece3.5013>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.



Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

ORIGINAL RESEARCH

Morphological and digestive adjustments buffer performance: How staging shorebirds cope with severe food declines

Shou-Dong Zhang^{1,2,3} | Zhijun Ma¹  | Chi-Yeung Choi^{4,5,6}  | He-Bo Peng^{2,3} | David S. Melville⁷ | Tian-Tian Zhao¹ | Qing-Quan Bai⁸ | Wen-Liang Liu⁹ | Ying-Chi Chan^{2,3}  | Jan A. van Gils²  | Theunis Piersma^{2,3} 

¹Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, Coastal Ecosystems Research Station of the Yangtze River Estuary, Shanghai Institute of Eco-Chongming (SIEC), Fudan University, Shanghai, China

²Department of Coastal Systems, NIOZ Royal Netherlands Institute for Sea Research and Utrecht University, Texel, The Netherlands

³Rudi Drent Chair in Global Flyway Ecology, Conservation Ecology Group, Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen, Groningen, The Netherlands

⁴School of Biological Sciences, University of Queensland, Brisbane, Queensland, Australia

⁵Centre for Integrative Ecology, School of Life & Environmental Sciences, Deakin University, Geelong, Victoria, Australia

⁶School of Environmental Science and Engineering, Southern University of Science and Technology, Shenzhen, China

⁷Global Flyway Network, Nelson, New Zealand

⁸Forestry Bureau of Dandong, Dandong, China

⁹School of Ecological and Environmental Sciences, East China Normal University, Shanghai, China

Correspondence

Zhijun Ma, Ministry of Education Key Laboratory for Biodiversity Science and Ecological Engineering, Coastal Ecosystems Research Station of the Yangtze River Estuary, Shanghai Institute of Eco-Chongming (SIEC), Fudan University, Shanghai, China.
Email: zhijunm@fudan.edu.cn

Funding information

The National Key Research and Development Program of China, Grant/Award Number: 2018YFC1406402; National Natural Science Foundation of China, Grant/Award Number: 31071939, 31572280 and 31830089; the KNAW China Exchange Programme grant, Grant/Award Number: 530-5CDP16; the CSC scholarship, Grant/Award Number: 201706100078

Abstract

Organisms cope with environmental stressors by behavioral, morphological, and physiological adjustments. Documentation of such adjustments in the wild provides information on the response space in nature and the extent to which behavioral and bodily adjustments lead to appropriate performance effects. Here we studied the morphological and digestive adjustments in a staging population of migrating Great Knots *Calidris tenuirostris* in response to stark declines in food abundance and quality at the Yalu Jiang estuarine wetland (northern Yellow Sea, China). At Yalu Jiang, from 2011 to 2017 the densities of intertidal mollusks, the food of Great Knots, declined 15-fold. The staple prey of Great Knots shifted from the relatively soft-shelled bivalve *Potamocorbula laevis* in 2011–2012 to harder-shelled mollusks such as the gastropod *Umbonium thomasi* in 2016–2017. The crushing of the mollusks in the gizzard would require a threefold to 11-fold increase in break force. This was partially resolved by a 15% increase in gizzard mass which would yield a 32% increase in shell processing capacity. The consumption of harder-shelled mollusks was also accompanied by reliance on regurgitates to excrete unbreakable parts of prey, rather than the usual intestinal voidance of shell fragments as feces. Despite the changes in digestive morphology and strategy, there was still an 85% reduction in intake rate in 2016–2017 compared with 2011–2012. With these morphological and digestive

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

adjustments, the Great Knots remaining faithful to the staging site to a certain extent buffered the disadvantageous effects of dramatic food declines. However, compensation was not complete. Locally, birds will have had to extend foraging time and use a greater daily foraging range. This study offers a perspective on how individual animals may mitigate the effects of environmental change by morphological and digestive strategies and the limits to the response space of long-distance migrating shorebirds in the wild.

KEYWORDS

East Asian-Australasian Flyway, energetics, food decline, Great Knot *Calidris tenuirostris*, phenotypic flexibility, prey quality, regurgitates

1 | INTRODUCTION


When animals encounter problems in achieving a positive energy balance, they may move away (Piersma, 2012), adjust behavior (Sydeman et al., 2001), or change relevant morphological (Grant & Grant, 2017), and correlated physiological (i.e., “physiomorphic,” Oudman et al., 2016) aspects of the phenotype (Piersma & van Gils, 2011). Bodily adjustments are usually studied under controlled experimental conditions (Dekinga, Dietz, Koolhaas, & Piersma, 2001; van Gils, Piersma, Dekinga, & Dietz, 2003). Here we provide field evidence for phenotypic adjustment in a highly site-faithful and long-lived, long-distance migratory shorebird species confronted with dramatic changes in their food supply at their main staging site.

Long-distance migratory shorebirds have provided several landmark examples of phenotypic flexibility (Piersma, 2002), including extensive body remodeling during migration (Vézina, Williams, Piersma, & Morrison, 2012). During the staging episodes prior to departure on long-distance flight, digestive and exercise organs grow and shrink in adaptive ways as the fuel stores are built up (Piersma & Gill, 1998; Piersma, Gudmundsson, & Lillendahl, 1999). Of particular interest, here are the digestive organs. Shorebirds may

regress their digestive organs (also some other visceral organs) in the course of long-distance migratory flights (Battley et al., 2000), but need to upregulate them during fueling (Battley, Dekinga, et al., 2001; Battley, Dietz, et al., 2001; Battley et al., 2000; Landys-Ciannelli, Piersma, & Jukema, 2003; Lindström & Piersma, 1993).

We studied Great Knots *Calidris tenuirostris*, an endangered endemic shorebird species in the East Asian-Australasian Flyway, seasonally commuting between main nonbreeding grounds in Northwest Australia and breeding grounds in the highlands of eastern Siberia, with halfway staging areas on the East Asian coast (Battley, Dekinga, et al., 2001; Battley, Dietz, et al., 2001; Battley et al., 2000; Lisovski, Gosbell, Hassell, & Minton, 2016; Ma et al., 2013). Except for their time on the tundra breeding grounds, Great Knots feed on mollusks on intertidal mudflats (Choi et al., 2017; Tulp & de Goeij, 1994). Shelled prey items are swallowed whole and crushed in the muscular gizzard, the shell fragments being evacuated from the gut as shell-rich feces (one-way stream, Figure 1). It takes longer to process hard-shelled than soft-shelled prey, and thus, the digestive bottleneck due to consumption of hard-shelled prey will cause a decrease in food intake rate (van Gils, Battley, Piersma, & Drent, 2005; van Gils et al., 2003).

FIGURE 1 Summary of the factors contributing to different feeding performance of Great Knot. Depending on prey quality (flesh/shell ratio) and shell hardness (break force), Great Knots may have smaller or larger gizzards. Generally, crushed shells are evacuated through the gut as feces (one-way stream), while Great Knots can also void unbreakable parts of ingested shells as pellets (two-way stream)



The decline in annual survival and population size of the Red Knot *Calidris canutus*, a sister species of the Great Knot, was earlier explained by food shortages at their main northward staging area due to over-harvesting and low food availability at their wintering ground (Baker et al., 2004; van Gils et al., 2016). The recent declines in survival and population sizes of Great Knots have been explained by the loss and degradation of coastal staging habitat in the Yellow Sea region due to land claim, pollution, invasive species, and over-exploitation of marine resources (Hua, Tan, Chen, & Ma, 2015; Melville, Chen, & Ma, 2016; Piersma et al., 2016). Following the destruction of Saemangeum, South Korea, by reclamation (Moores, Rogers, Rogers, & Hansbro, 2016), the Yalu Jiang estuarine wetland (hereafter YLJ, Figure 2) became the main known staging area of Great Knots in the Yellow Sea region (Choi, Battley, Potter, Rogers, & Ma, 2015; Riegen, Vaughan, & Rogers, 2014; Zhang et al., 2018). Despite there having been little loss of intertidal habitat over the past two decades at YLJ, the bivalve *Potamocorbula laevis*, one of the dominant benthic invertebrate species and the main prey of Great Knots (Choi et al., 2017), has dramatically declined in abundance since 2013 (Zhang et al., 2018).

Assessments of the degree to which such phenotypically flexible adjustments help animals in real-life situations require field studies

where the changing aspects of both the target species and their immediate environment including their food are described well enough (Battley, Dekinga, et al., 2001; Battley, Dietz, et al., 2001; Battley et al., 2000; Piersma, 2011). Here we assess how Great Knots staging at YLJ in 2011–2017 coped with the changing abundance and digestive quality of their food in terms of internal morphology and attributes of the digestion process (Figure 1). We analyzed food availability, composition, diet selection, and food intake rates in two years before and two years after the dramatic decline of *P. laevis*. We compared the quality of the different mollusk prey over the study period and examined changes in gizzard mass and the way that the birds coped with unbreakable parts of their prey. This study highlights how morphological and digestive strategies may change under resource limitation and offers a perspective on how a species can mitigate such effects (Figure 1).

2 | MATERIALS AND METHODS

2.1 | Study site

This study was carried out during March–May 2011–2017, at YLJ (39°40′–39°58′N, 123°34′–124°07′E; Figure 2), which is close to




FIGURE 2 Location of Yalu Jiang estuarine wetland in the north part of the Yellow Sea, China. The dots show all the 104 macrobenthos stations sampled in 2013, 2015, 2016, and 2017. The 36 dots within the dashed-line square in the central area show the macrobenthos stations sampled in 2011 and 2014; in 2012, we sampled the 36 sites within the dashed-line square in the middle part together with the 12 sites within the dashed-line rectangle in the east part

the national boundary between China and North Korea (Choi et al., 2017; Zhang et al., 2018). About 44,000 Great Knots (Zhang et al., 2018) stage here for nearly 2 months during northward migration (Choi et al., 2015; Ma et al., 2013; Riegen et al., 2014). From 2011–2012 to 2016–2017, the numbers of Great Knots at YLJ decreased by 29% (Zhang et al., 2018). The seasonal pattern of occurrence also changed. In the early years, numbers built up from late March to early April, to then remain steady until the time of northward departures from 14–21 May. However, departures substantially advanced in 2017, with numbers already declining from mid-April. Some birds might move to other staging sites because it is too early to depart to the breeding grounds (Zhang et al., 2018). In 2015, a large number of Great Knots may have moved about 160 km westward from YLJ to Gaizhou, on the east of Liaodong Bay, Bohai (Melville, Peng, et al., 2016), where *P. laevis* were plentiful at the time (HBP and YCC unpublished data).

Tides at YLJ are typically semi-diurnal, with an average tidal amplitude of 4.5 m and a maximum of 6.9 m (Wang, Ren, & Zhu, 1986). Local movements of Great Knots are strongly affected by the tides. They forage on the intertidal flats at low tide and roost in aquaculture ponds inside the seawall when intertidal flats are submerged (Zhang et al., 2018). Bivalves (especially superabundant *P. laevis*) have previously been identified as the staple food of Great Knots at YLJ (Choi et al., 2017).

2.2 | Diet composition and food availability

To measure food availability, we sampled potential prey (macrobenothos, especially mollusks living in the soft sediments) once a month from March to May in 2011, 2012, 2015, 2016, and 2017, and April and May in 2013 and 2014. We established 16 sampling transects across the intertidal flats with a total of 104 sampling stations spaced 500 m apart (Figure 2). All the stations (104 sampling stations) were sampled in 2013, 2015, 2016, and 2017, but we focused on the middle parts (36 sampling stations) of the study area in 2011, 2012, and 2014 as this zone is the main foraging area of Great Knots (Choi et al., 2017). There is no significant difference in macrobenthos species and ash-free dry mass (AFDM) between the middle parts of sampling stations and the total 104 sampling stations (t tests, $p > 0.2$ for both macrobenthos species and AFDM in all the years in 2013, 2015, 2016, and 2017) (Zhang et al., 2018). At each station, one core sample (diameter 15.5 cm, 5 cm in depth) was collected and then washed through a 0.5 mm sieve. We considered all bivalves and gastropods from the top 5 cm of sediment as potential food for Great Knots which have an average bill length of about 45 mm (Tulp & de Goeij, 1994). All soft-bodied organisms were soaked in 5% formalin for at least 72 hr before being stored in 70% ethanol, and hard-bodied organisms including large bivalves, gastropods, and crabs were kept frozen. Organisms were identified to the finest practicable taxonomic level (Supporting Information Table S1) with size (the longest measurement) measured (to 0.01 mm) in the laboratory (see Zhang et al., 2018). For each taxon, we selected complete individuals of different lengths to determine AFDM (Zhang et al., 2018).

We measured the height of the left hinge for different shell lengths of all bivalve species (Dekinga & Piersma, 1993; Yang et al., 2013), and the width of the last whorl of the columella for different sizes of *Umbonium thomasi*. The species-specific relationships between size (the longest measurement) and AFDM, AFDM, and DM_{shell} (shell dry mass), size (the longest measurement) and height of left hinge or columella width were established using regression analysis (polynomial, e-exponential, logarithmic, and power regression) for each taxonomic group (Bom et al., 2018). Models with the highest Pearson correlation coefficient were selected. Regression models were established for each year for the two dominant bivalves, *P. laevis* and *Macra veneriformis*, while samples from all years were pooled for other groups in the regression models due to small sample sizes (Zhang et al., 2018).

Food retention time in the digestive tracts of sandpipers is short (e.g., 20–50 min for red knots, Piersma, 1994). During low tide, we followed flocks of Great Knot at foraging sites to collect droppings after birds had been feeding at a site for more than 30 min (to ensure the droppings were produced by Great Knots feeding on prey from that foraging site) (Choi et al., 2017). We collected a total of 2,712 droppings (1,093 in 2011, 856 in 2012, 117 in 2016 and 646 in 2017) and 439 pellets (184 in 2016 and 255 in 2017). Each dropping and pellet were placed in a separate plastic bag and stored at -20°C .

In the laboratory, droppings and pellets were dried at 60°C for 72 hr, and then sifted through a 0.3 mm sieve. The shell fragments from the droppings and pellets were sorted to species and measured to the nearest 0.1 mm using an Olympus SZX7 dissecting microscope: the height of unbroken hinges for bivalve species, the width of the last whorl of the columella for *U. thomasi*, and the shell length for undigested individuals. The weight of shell fragments (no columella present) and the columella of *U. thomasi* in droppings and pellets were weighed to the nearest 0.1 mg. To determine the size composition of ingested organisms contained in the dropping and pellet samples, we used the regression of left hinge height against shell length for bivalve species, and the regression between columella width and total width in *U. thomasi* (Supporting Information Table S3).

At YLJ, Great Knots mainly consumed *P. laevis*, but also took other bivalves and gastropods, all available prey being within the top 5 cm of sediment (Choi et al., 2017). Based on the analysis of feces and pellets, the maximum sizes of prey taken by Great Knots were determined (Zwarts & Blomert, 1992). We defined potential (“harvestable”) food as follows: *P. laevis* (less than 25 mm, divided into small size: <10 mm and large size: >10 mm) and *Moerella iridescens* less than 25 mm, *M. veneriformis* less than 23 mm, *U. thomasi* less than 15 mm, *Nassarius variciferus* and *Nassarius festivus* less than 12 mm, other smaller proportions of bivalves and gastropods less than 20 mm.

2.3 | Intake rate, prey quality, and shell hardness

To record bird behavior, a focal bird was chosen randomly from a flock of foraging birds and watched for 5-min, using a $\times 20$ –60

telescope. Before the start of each 5-min observation bout, the date, time, and location were noted. During each observation bout, activities such as pecks, probes, items swallowed and interference with other individuals were recorded on digital voice recorders (2011 and 2012), while we used digital video cameras to record all behaviors in 2016 and 2017. The digital sound files were transcribed using JWWatcher 1.0 (Blumstein, Daniel, & Evans, 2006), which allowed us to quantify the time a bird spent on different activities. We used BORIS (Friard & Gamba, 2016) to transcribe behaviors on video. We identified ingested prey as bivalve, crab, gastropod, ghost shrimp, razor clam, polychaeta, sea anemone, or unknown.

Based on the composition of the prey size and the regression of size-species AFDM/DM_{shell} (Supporting Information Tables S1 and S2), we estimated ingested AFDM and DM_{shell} for Great Knots feeding on a specific food item. Based on the species and numbers of prey taken by Great Knots, and the relationship of size-species AFDM/DM_{shell} (Supporting Information Tables S1 and S2), we calculated AFDM/DM_{shell} intake rate per unit time. We evaluated prey quality based on flesh/shell ratio and shell hardness based on the break force of shells. The flesh/shell ratio was calculated using AFDM and DM_{shell} intake by each individual (AFDM/DM_{shell}). A higher flesh/shell ratio (van Gils, de Rooij, et al., 2005) indicates higher prey quality. A fixed digital force gauge (HP-20 and HP-300, Yueqing Ai Li Instrument, China) was used to measure the break force (N) of four major prey of Great Knots (*P. laevis*, *M. veneriformis*, *M. iridescens*, and *U. thomasi*) (Bom et al., 2018; Yang et al., 2013). We regressed break force on prey size trying polynomial, e-exponential, logarithmic, and power regression. Models with the largest Pearson correlation coefficient were selected.

2.4 | Gizzard size

Some Great Knots drowned accidentally in fishing nets set on the intertidal mudflat. In April–May 2011–2012 and 2016–2017, a total of 48 dead birds (22 and 26, respectively) were collected and used opportunistically for the study. The birds were collected, the feathers dried with a hairdryer and weighed. Carcasses were sealed in airtight plastic bags and stored at -20°C . In laboratory, carcasses were dissected following the procedures of Piersma et al. (1999). The fresh mass of the gizzard was weighed to the nearest 0.1 g. The contents of the intestines and gizzard were washed separately through a 0.3 mm sieve and then dried at 60°C for 72 hr. The fragments were separated and identified. The shell fragments and the columella of *U. thomasi* in intestinal and gizzard contents were weighed to the nearest 0.1 mg.

2.5 | Data analysis

We used Ivlev's electivity index (E_i , Ivlev, 1961) to quantify the prey selection by Great Knots: $E_i = (r_i - p_i)/(r_i + p_i)$, where r_i is the percentage of prey species i in the diet (number of species/total number of preys in the diet $\times 100$), p_i is the percentage of this species on the mudflat. E_i ranges from -1 to 1 interval, values from -1 to 0 meaning

negative selection, values from 0 to 1 meaning positive selection, and 0 meaning no selection.

Linear models controlling the effect of date were used to compare gizzard size (dry mass, g) between the period before (2011 and 2012 combined) and after food changes (2016 and 2017 combined). We calculated the changes of shell processing capacity based on the quadratic relationship between gizzard mass and shell processing capacity (van Gils, Piersma, Dekinga, & Battley, 2006). The percentage by weight of columella of *U. thomasi* in feces, pellets, intestines, and gizzard were compared using analysis of variance (ANOVA) followed by Fisher's least significant difference (LSD). Differences in intake rate of AFDM, DM_{shell}, and prey quality between the two periods were also compared using ANOVA followed by LSD.

Logarithmic transformation was used when the data were not normally distributed. When modeling the functional response between intake rate and prey density, we took account of the variation in prey quality using a modified form of Holling's disk equation: $IR = a \times N / (1 + a \times N \times h / Q)$, where IR is the average intake rate (mg AFDM/s) for each year, N is the average prey density (ind/m²) for each year, and Q is the average prey quality (AFDM/DM_{shell}) for each year. We estimated a (searching efficiency) and h (prey handling time) using the nonlinear least square (nls) function in R version 3.5.2. By multiplying handling time h by the inverse of prey quality Q , we included time lost to digestion in the standard type II functional response model (similar to eq. 5 in Jeschke, Kopp, & Tollrian, 2002). The significance level was set at 0.05. Statistical analyses were carried out in SPSS 20.0 unless referenced otherwise.

3 | RESULTS

Prey available to Great Knots declined dramatically from 2011 to 2017. The density of small *P. laevis* (<10 mm) declined by 98.5%, *P. laevis* (>10 mm) declined by 99.8%, while densities of the gastropod *U. thomasi* increased 57 times and the bivalve *M. iridescens* increased seven times, respectively (Figure 3a). The main prey of Great Knots in 2011 was *P. laevis* (of total prey consumed 62% for small sized and 34% for large sized) and in 2012 (8% for small sized and 88% for large sized). In 2016, the main prey was *U. thomasi* (89.6%), and in 2017 *U. thomasi* (37%) and *M. veneriformis* (35%) (Figure 3b). In 2011 and 2012, Great Knots exhibited positive selection for small *P. laevis*, while they selected large *P. laevis* in 2016 and 2017 (Figure 3c). In 2016, Great Knots selected *U. thomasi* while rejecting *M. veneriformis*, but in 2017, they selected *M. veneriformis* (Figure 3c). Great Knots exhibited negative selection for small *P. laevis* in 2016 and 2017 (Figure 3c).

Break force of the four main prey species increased with shell length (Supporting Information Figure S2). The break forces of modal prey size were 4.0 N (*P. laevis*, 7.86 mm) in 2011, 12.0 N (*P. laevis*, 12.91 mm) in 2012, 45.2 N (*U. thomasi*, 11.61 mm) in 2016, and 35.4 N (*U. thomasi*, 10.37 mm) in 2017 (Supporting Information Figure S2). The break force required to crush the main prey in 2016




FIGURE 3 (a) Food availability (ind/m²) and (b) diet composition (percentage of individual prey in feces and pellets combined) of Great Knots from 2011 to 2017; (c) Electivity index (E_i) of Great Knots for different prey in 2011, 2012, 2016, and 2017. E_i in the interval of [-1, 0] means negative selection, (0, 1] positive selection, and 0 no selection




FIGURE 4 Comparison of gizzard mass of Great Knots in 2011–2012 ($n = 22$, open circles and dashed line) and 2016–2017 ($n = 27$, filled circles and solid line). (a) The equation in 2011–2012: Gizzard mass = $7.20 + 0.04 \times$ capture date (Julian day: 1 Mar = 1)–1.53; in 2016–2017: Gizzard mass = $7.20 + 0.04 \times$ capture date. (b) Gizzard mass was significantly different ($F = 8.716$, $p = 0.005$) between the two periods. Diamonds show the mean value of gizzard mass, horizontal lines show the median, the upper and lower edges of the box plots represent the first and the third quartiles, bars represent the 95% confidence interval. (c) Photographs of gizzards in 2011–2012 (bottom) and in 2016–2017 (top)

and 2017 was 9–11 times greater than that in 2011 and 3–4 times greater than that in 2012.

After controlling the effect of date, gizzard mass significantly increased ($F = 8.72$, $p = 0.005$) after the diet shift (a 15% increase from 7.86 ± 0.31 g in 2011–2012 to 9.43 ± 0.41 g in 2016–2017, Figure 4). This would equal a 32% increase in shell processing capacity.

The consumption of hard-shelled mollusks in 2016–2017 was accompanied by a greater reliance on regurgitation (rather than intestinal evacuation as feces) to excrete unbreakable parts of prey such as the columella of *U. thomasi* (Figure 1). The weight percentages

of columella in feces, intestinal contents, pellets, and gizzard contents were 0.71 ± 0.44 , 0.00 ± 0.00 , 27.18 ± 1.96 , and 35.47 ± 5.34 , respectively (Figure 5). There was no significant difference in the weight percentage of columella in feces and intestinal contents, and in pellets and gizzard contents ($p > 0.05$ for both), respectively (Figure 5). However, the weight percentage of columella in feces and intestinal contents was significantly lower than that in pellets and gizzard contents ($p < 0.001$, Figure 5), suggesting that—unlike other parts of the shell—the columella was not passed through the entire digestive tract.




FIGURE 5 The weight percentage of columella remains in feces, intestinal contents, pellets, and gizzard contents. The three asterisks refer to a significant difference in the columella percent (%) between the combination of faces and intestinal contents and the combination of pellets and gizzard contents ($F = 40.88$, $p < 0.001$). Diamonds show the mean value, horizontal lines show the median, the upper and lower edges of the box plots represent the first and third quartiles, the bars represent the 95% confidence interval. Open circles indicate outliers. The top-left photograph shows feces of a Great Knot on the mudflat and the top-right a pellet. The letters a and b denote significant differences detected by LSD test

After the shift from *P. laevis* in 2011–2012 to a predominantly gastropod diet in 2016–2017, the intake rates expressed as AFDM/s and DM_{shell}/s significantly decreased ($F = 91.73$, $p < 0.001$ and $F = 36.94$, $p < 0.001$, respectively) (Supporting Information Figure S1a,b). The intake rate of AFDM decreased by 86%, and the intake rate of DM_{shell} decreased by 77% (Supporting Information Figure S1a,b). The prey quality significantly differed among the 4 years ($F = 8.56$, $p < 0.001$), with the highest in 2011 and the lowest in 2017. There was no significant difference in prey quality in 2011 and 2012. However, that prey quality in 2016 and 2017 was significantly lower than in 2011 ($p < 0.05$) and prey quality in 2017 was also significantly lower than in 2012 ($p < 0.05$, Figure 6a), suggest a serious decrease in prey quality in the course of this study. Similarly, shell hardness of prey significantly differed among the 4 years ($F = 456.54$, $p < 0.001$), with the hardest in 2016 and the softest in 2011. Shell hardness in 2016 was 4.0 and 2.7-fold harder than that in 2011 and 2012, respectively; shell hardness in 2017 was 2.8 and 2.0-fold harder than that in 2011 and 2012, respectively (Figure 6b).

The functional response relationship between intake rate and prey density for the year-specific prey qualities (Figure 6c) showed




FIGURE 6 (a) Quality (AFDM/ DM_{shell}), and (b) shell hardness (break force, N) of prey of Great Knots in 2011, 2012, 2016, and 2017, and (c) the relationship between intake rate of ash-free dry mass (AFDM/s) and prey density. Diamonds show the mean, horizontal lines show the median, the upper and lower edges of the box plots represent the first and third quartiles, and the bars represent the 95% confidence interval (a, b) or standard error (c). The open circles indicate outliers (a, b). Different letters denote significant differences detected by LSD test (a, b). The curves in (c) (the dotted line, dashed lines, dotted-dashed line, and solid line represent 2011, 2012, 2016, and 2017, respectively) show the relationships between intake rate of ash-free dry mass (AFDM/s) and prey density for different prey qualities expressed as: $IR = a \times N / (1 + a \times N \times h / Q)$, where IR, N, and Q are, respectively, the average intake rate (mg AFDM/s), prey density (ind/ m^2), and average prey quality (AFDM/ DM_{shell}) for each year; a (searching efficiency) = $6.5 \times 10^{-3} m^2/s$, h (prey handling time) = $7.3 \times 10^{-2} s/mg DM_{shell}$. At a given prey density, intake rate increases with prey quality

that the difference in intake rates between the two periods was explained by changes in both prey density and prey quality. The latter was especially evident from the comparison between 2011, when small *P. laevis* were abundant and large *P. laevis* were scarce, and 2012, when the reverse situation occurred.

4 | DISCUSSION

This study yields an interesting example of the morphological and digestive adjustments by which animals cope with environmental change (Abrahms et al., 2018; Colles, Liow, & Prinzing, 2009; van Gils, Battley, et al., 2005). After a dramatic food decline at a staging site, Great Knots changed the composition of their diet composition and prey selection to include prey types that were physically more difficult to process. They could do so because they simultaneously adjusted both the features of the digestive organ (gizzard mass) and the way of dealing with indigestible matter (voiding the hardest shell fragments through regurgitates rather than as feces).

The decline of food availability was mainly due to an almost complete loss of *P. laevis*. *P. laevis* occurred at high densities in 2011 and 2012, accounting for more than 95% of the total macrobenthos (Choi et al., 2017; Zhang et al., 2018). The small *P. laevis* are relatively easy to crush in the gizzard (Yang et al., 2013), thus birds exhibited a high processing rate and thus a high intake rate. Indeed, Great Knots exhibited positive selection for small *P. laevis* in 2011 and 2012. By 2016 and 2017, small *P. laevis* had all but disappeared and *M. iridescens* only occurred at low densities (Figure 3a). However, there were still a few large *P. laevis* available, which were selected by Great Knots (Figure 3c).

According to symmorph design rules, organs should grow to a size to satisfy but not exceed the requirements (Taylor & Weibel, 1981). Digestive organs have a high energy cost per unit mass (Rolfe & Brown, 1997; Scott & Evans, 1992), and the amount and metabolic cost can increase basal metabolic rate (BMR) (McKechnie, 2008; Piersma, 2002; Piersma et al., 1996; Vézina, Love, Lessard, & Williams, 2009). This is suggested to be one of the reasons that migratory birds ready for take-off on long-distance flights avoid oversized digestive organs (Piersma & Gill, 1998; van Gils, Battley, et al., 2005; Yang et al., 2013). However, this is also a function of ecological context as Red Knots, when forced to eat hard-shelled food generally enlarge their gizzard, to reduce it when soft food is available (Dekinga et al., 2001; van Gils et al., 2003). At YLJ, the prey quality decreased from 2011–2012 to 2016–2017, which means birds would need to process more shell material for similar amounts of energy gained. However, the shell of prey consumed in 2016 and 2017 required larger break force to be crushed than those in 2011 and 2012 (Figure 6b). The larger the break force, the longer the processing time (Yang et al., 2013). Thus, Great Knots feeding on hard-shelled mollusks will be digestively constrained by the amount of prey processed per unit time (Piersma, Koolhaas, & Dekinga, 1993; van Gils, Battley, et al., 2005; van Gils et al., 2003). As a consequence, the AFDM intake rate of Great Knots not only decreased with prey density, but also decreased with prey quality and shell hardness (Figure 6c). Great Knots at YLJ increased gizzard mass by 15% which, if similar to Red Knots, would have resulted in an increase of the shell processing rate by 32% (van Gils et al., 2006). This, however, will not be sufficient to compensate for the overall reduction in food availability and quality.

Shorebirds usually evacuate crushed shell material from the gut as feces (Battley & Piersma, 2005). The regurgitation of pellets, which occurs in Tringids but has not often been reported in Calidrid sandpipers (Dekinga & Piersma, 1993; Fedrizzi, Carlos, & Campos, 2016; Worrall, 1984), coincided with the strongly increased break forces required for ingested prey in 2016 and 2017. It appears that Great Knots were unable to crush the extremely hard calcified columella of *U. thomasi*. The potential risk of the columella, with its pointed apex and relatively sharp whorls, damaging the digestive tract (see Piersma et al., 1993; van Gils, Battley, et al., 2005; Yang et al., 2013; pers. obs.), may explain the change from a one-way stream to a two-way stream for ballast evacuation (see Figure 1).

The long-term AFDM intake rate in 2016 and 2017 was only 14%–30% of that in 2011 and 2012 (Figure 6a). As a consequence, to achieve the daily body mass gain at YLJ necessary to migrate, Great Knots would need to forage for 13–16 hr per day in 2016–2017 rather than less than five hours in 2011–2012 (Supporting Information Appendix S1). The mudflats at YLJ are exposed for an average of 18.6 hr a day during spring tides and 21.0 hr a day during neap tides (C.Y. Choi et al., unpublished data). However, Great Knots eating very hard-shelled *U. thomasi* were observed to take long pauses of over one hour (pers. obs., see van Gils, Battley, et al., 2005). We interpret this as time necessary for digestion and/or the preparation and voidance of regurgitates. It thus appears likely that birds were rather challenged to have enough foraging time to satisfy the energy requirements, especially at spring tides.

Staging numbers of Great Knots at YLJ declined by a third between 2011/2012 and 2016/2017 (Zhang et al., 2018), implying that some birds were not able to fully compensate for the severe reduction in both prey quantity and quality. Considering the effects on survival of the dramatic losses and degradation of intertidal habitats in the Yellow Sea (Piersma et al., 2016), migrating Great Knots appear to be short of alternative high-quality staging habitats. The decrease in body mass gain rates of Great Knots at the study site could well lead to cause downstream effects on survival and breeding success (Piersma et al., 2016; Senner, Conklin, & Piersma, 2015). The long-term studies on birds and their food conditions at the staging and nonbreeding sites which spawned this contribution, seem critical if we are serious in trying to understand the causes of decline and to responsively manage the key habitats of endangered species such as Great Knots.

ACKNOWLEDGMENTS

We thank the Dandong Yalu Jiang Estuary Wetland National Nature Reserve for facilitating our fieldwork. We are indebted to Xiaojing Gan, Peter Brakels, Peter Crighton, Rosanne Michielsen, Jason Loghry, Dezhong Xing, Yongjun Sui, Haoying Huang, Yi Yang, Yayi Huang, Julia Melville, Kun Tan, and Jianping Yu for field assistance. We also thank Jingyao Niu, Shaoping Zang, Yishu Zhu, and Youjia Li for helping us process macrozoobenthos samples, behavior videos and measuring break force. This study was financially supported

by the National Key Research and Development Program of China (2018YFC1406402), National Natural Science Foundation of China (31830089, 31572280, and 31071939), the CSC scholarship (201706100078) and the KNAW China Exchange Programme grant (530-5CDP16) awarded to TP.

CONFLICT OF INTEREST

None declared.

AUTHORS' CONTRIBUTIONS

Z.M, T.P., J.A.v.G., and S.D.Z. conceived the ideas and designed the experiment; S.D.Z., C.Y.C., H.B.P., D.S.M., Q.Q.B., T.T.Z., and Y.C.C. collected the data in the field; S.D.Z., C.Y.C., and W.L.L. processed the samples and recordings with help from H.B.P. and Y.C.C.; S.D.Z. analyzed the data; S.D.Z., T.P., and Z.M. led the writing of the manuscript. All authors contributed to the drafts and approved the final version.

DATA ACCESSIBILITY

Relevant data will be available via Dryad: <https://doi.org/10.5061/dryad.9021d2v>.

ORCID

Zhijun Ma  <https://orcid.org/0000-0001-7459-9448>

Chi-Yeung Choi  <https://orcid.org/0000-0001-9829-7460>

Ying-Chi Chan  <https://orcid.org/0000-0002-7183-4411>

Jan A. van Gils  <https://orcid.org/0000-0002-4132-8243>

Theunis Piersma  <https://orcid.org/0000-0001-9668-466X>

REFERENCES

- Abrahms, B., Hazen, E. L., Bograd, S. J., Brashares, J. S., Robinson, P. W., Scales, K. L., ... Costa, D. P. (2018). Climate mediates the success of migration strategies in a marine predator. *Ecology Letters*, 21, 63–71. <https://doi.org/10.1111/ele.12871>
- Baker, A. J., Gonzalez, P. M., Piersma, T., Niles, L. J., de Lima Serraro do Nascimento, I., Atkinson, P. W., ... Aarts, G. (2004). Rapid population decline in red knots: Fitness consequences of decreased refuelling rates and late arrival in Delaware Bay. *Proceedings of the Royal Society B: Biological Sciences*, 271, 875–882. <https://doi.org/10.1098/rspb.2003.2663>
- Battley, P. F., Dekinga, A., Dietz, M. W., Piersma, T., Tang, S., & Hulsman, K. (2001). Basal metabolic rate declines during long-distance migratory flight in great knots. *Condor*, 103, 838–845.
- Battley, P. F., Dietz, M. W., Piersma, T., Dekinga, A., Tang, S. X., & Hulsman, K. (2001). Is long-distance bird flight equivalent to a high-energy fast? Body composition changes in freely migrating and captive fasting great knots. *Physiological and Biochemical Zoology*, 74, 435–449. <https://doi.org/10.1086/320432>
- Battley, P. F., & Piersma, T. (2005). Adaptive interplay between feeding ecology and features of the digestive tract in birds. In J. M. Starck, & T. Wang (Eds.), *Physiological and ecological adaptations to feeding in vertebrates* (pp. 201–228). Enfield, NH: Science Publishers.
- Battley, P. F., Piersma, T., Dietz, M. W., Tang, S. X., Dekinga, A., & Hulsman, K. (2000). Empirical evidence for differential organ reductions during trans-oceanic bird flight. *Proceedings of the Royal Society B: Biological Sciences*, 267, 191–195. <https://doi.org/10.1098/rspb.2000.0986>
- Blumstein, D. T., Daniel, J. C., & Evans, C. S. (2006). JWwatcher (Version 1.0). <http://www.jwatcher.ucla.edu/>
- Bom, R. A., de Fouw, J., Klaassen, R. H. G., Piersma, T., Lavaleye, M. S. S., Ens, B. J., ... van Gils, J. A. (2018). Food web consequences of an evolutionary arms race: Molluscs subject to crab predation on intertidal mudflats in Oman are unavailable to shorebirds. *Journal of Biogeography*, 45, 342–354. <https://doi.org/10.1111/jbi.13123>
- Choi, C. Y., Battley, P. F., Potter, M. A., Ma, Z. J., Melville, D. S., & Sukkaewmanee, P. (2017). How migratory shorebirds selectively exploit prey at a staging site dominated by a single prey species. *Auk*, 134, 76–91. <https://doi.org/10.1642/AUK-16-58.1>
- Choi, C. Y., Battley, P. F., Potter, M. A., Rogers, K. G., & Ma, Z. (2015). The importance of Yalu Jiang coastal wetland in the north Yellow Sea to Bar-tailed Godwits *Limosa lapponica* and Great Knots *Calidris tenuirostris* during northward migration. *Bird Conservation International*, 25, 53–70. <https://doi.org/10.1017/S0959270914000124>
- Colles, A., Liow, L. H., & Prinzing, A. (2009). Are specialists at risk under environmental change? Neoecological, paleoecological and phylogenetic approaches. *Ecology Letters*, 12, 849–863. <https://doi.org/10.1111/j.1461-0248.2009.01336.x>
- Dekinga, A., Dietz, M. W., Koolhaas, A., & Piersma, T. (2001). Time course and reversibility of changes in the gizzards of red knots alternately eating hard and soft food. *Journal of Experimental Biology*, 204, 2167–2173.
- Dekinga, A., & Piersma, T. (1993). Reconstructing diet composition on the basis of faeces in a mollusc-eating wader, the Knot *Calidris canutus*. *Bird Study*, 40, 144–156. <https://doi.org/10.1080/00063659309477140>
- Fedrizzi, C. E., Carlos, C. J., & Campos, A. A. (2016). Annual patterns of abundance of Nearctic shorebirds and their prey at two estuarine sites in Ceará, NE Brazil, 2008–2009. *Wader Study*, 123, 122–135. <https://doi.org/10.18194/ws.00036>
- Friard, O., & Gamba, M. (2016). BORIS: A free, versatile open-source event-logging software for video/audio coding and live observations. *Methods in Ecology and Evolution*, 7, 1325–1330. <https://doi.org/10.1111/2041-210X.12584>
- Grant, B. R., & Grant, P. R. (2017). Watching speciation in action. *Science*, 355, 910–911. <https://doi.org/10.1126/science.aam6411>
- Hua, N., Tan, K., Chen, Y., & Ma, Z. J. (2015). Key research issues concerning the conservation of migratory shorebirds in the Yellow Sea region. *Bird Conservation International*, 25, 38–52. <https://doi.org/10.1017/S0959270914000380>
- Ivlev, V. S. (1961). *Experimental ecology of the feeding of fishes*. New Haven, CT: Yale University Press.
- Jeschke, J. M., Kopp, M., & Tollrian, R. (2002). Predator functional responses: Discriminating between handling and digesting prey. *Ecological Monographs*, 72, 95–112. <https://doi.org/10.2307/3100087>
- Landys-Ciannelli, M. M., Piersma, T., & Jukema, J. (2003). Strategic size changes of internal organs and muscle tissue in the bar-tailed godwit during fat storage on a spring stopover site. *Functional Ecology*, 17, 151–159. <https://doi.org/10.1046/j.1365-2435.2003.00715.x>
- Lindström, Å., & Piersma, T. (1993). Mass changes in migrating birds: The evidence for fat and protein storage re-examined. *Ibis*, 135, 70–78. <https://doi.org/10.1111/j.1474-919X.1993.tb02811.x>
- Lisovski, S., Gosbell, K., Hassell, C., & Minton, C. (2016). Tracking the full annual-cycle of the great knot *Calidris tenuirostris*, a long-distance migratory shorebird of the East Asian-Australasian Flyway. *Wader Study*, 123, 177–189. <https://doi.org/10.18194/ws.00048>

- Ma, Z., Hua, N., Peng, H., Choi, C., Battley, P. F., Zhou, Q., ... Tang, C. (2013). Differentiating between stopover and staging sites: Functions of the southern and northern Yellow Sea for long-distance migratory shorebirds. *Journal of Avian Biology*, 44, 504–512. <https://doi.org/10.1111/j.1600-048X.2013.00213.x>
- McKechnie, A. E. (2008). Phenotypic flexibility in basal metabolic rate and the changing view of avian physiological diversity: A review. *Journal of Comparative Physiology B*, 178, 235–247. <https://doi.org/10.1007/s00360-007-0218-8>
- Melville, D. S., Chen, Y., & Ma, Z. J. (2016). Shorebirds along the Yellow Sea coast of China face an uncertain future: A review of threats. *Emu*, 116, 100–110. <https://doi.org/10.1071/MU15045>
- Melville, D. S., Peng, H. B., Chan, Y. C., Bai, Q. Q., He, P., Tan, K., ... Ma, Z. J. (2016). Gaizhou, Liaodong Bay, Liaoning Province, China – A site of international importance for Great Knot *Calidris tenuirostris* and other shorebirds. *Stilt*, 69–70, 57–61.
- Moores, N., Rogers, D. I., Rogers, K., & Hansbro, P. M. (2016). Reclamation of tidal flats and shorebird declines in Saemangeum and elsewhere in the Republic of Korea. *Emu*, 116, 136–146. <https://doi.org/10.1071/MU16006>
- Oudman, T., Bijleveld, A. I., Kavelaars, M. M., Dekinga, A., Cludera, J., Piersma, T., & van Gils, J. A. (2016). Diet preferences as the cause of individual differences rather than the consequence. *Journal of Animal Ecology*, 85, 1378–1388. <https://doi.org/10.1111/1365-2656.12549>
- Piersma, T. (1994). *Close to the edge: Energetic bottlenecks and the evolution of migratory pathways in knots*. PhD thesis, University of Groningen, The Netherlands.
- Piersma, T. (2002). Energetic bottlenecks and other design constraints in avian annual cycles. *Integrative and Comparative Biology*, 42, 51–67. <https://doi.org/10.1093/icb/42.1.51>
- Piersma, T. (2011). Why marathon migrants get away with high metabolic ceilings: Towards an ecology of physiological restraint. *Journal of Experimental Biology*, 214, 295–302. <https://doi.org/10.1242/jeb.046748>
- Piersma, T. (2012). What is habitat quality? Dissecting a research portfolio on shorebirds. In R. J. Fuller (Ed.), *Birds and habitat: Relationships in changing landscapes* (pp. 383–407). Cambridge, UK: Cambridge University Press.
- Piersma, T., Bruinzeel, L., Drent, R., Kersten, M., van der Meer, J., & Wiersma, P. (1996). Variability in basal metabolic rate of a long-distance migrant shorebird (Red Knot, *Calidris canutus*) reflects shifts in organ sizes. *Physiological Zoology*, 69, 191–217. <https://doi.org/10.1086/physzool.69.1.30164207>
- Piersma, T., & Gill, R. E. (1998). Guts don't fly: Small digestive organs in obese bar-tailed godwits. *Auk*, 115, 196–203. <https://doi.org/10.2307/4089124>
- Piersma, T., Gudmundsson, G. A., & Lilliendahl, K. (1999). Rapid changes in the size of different functional organ and muscle groups during refueling in a long distance migrating shorebird. *Physiological and Biochemical Zoology*, 72, 405–415. <https://doi.org/10.1086/316680>
- Piersma, T., Koolhaas, A., & Dekinga, A. (1993). Interactions between stomach structure and diet choice in shorebirds. *Auk*, 110, 552–564. <https://doi.org/10.2307/4088419>
- Piersma, T., Lok, T., Chen, Y., Hassell, C. J., Yang, H.-Y., Boyle, A., ... Ma, Z. (2016). Simultaneous declines in summer survival of three shorebird species signals a flyway at risk. *Journal of Applied Ecology*, 53, 479–490. <https://doi.org/10.1111/1365-2664.12582>
- Piersma, T., & van Gils, J. A. (2011). *The flexible phenotype: A body-centred integration of ecology, physiology, and behaviour*. Oxford, UK: Oxford University Press.
- Riegen, A. C., Vaughan, G. R., & Rogers, K. D. (2014). *Yalu Jiang Estuary Shorebird Survey Report 1999–2010*. Yalu Jiang Estuary Nature Reserve, China and Miranda Naturalists' Trust, New Zealand.
- Rolfe, D. F. S., & Brown, G. C. (1997). Cellular energy utilization and molecular origin of standard metabolic rate in mammals. *Physiological Reviews*, 77, 731–758. <https://doi.org/10.1152/physrev.1997.77.3.731>
- Scott, I., & Evans, P. R. (1992). The metabolic output of avian (*Sturnus vulgaris*, *Calidris alpina*) adipose tissue liver and skeletal muscle: Implications for BMR/body mass relationships. *Comparative Biochemistry and Physiology A. Comparative Physiology*, 103, 329–332. [https://doi.org/10.1016/0300-9629\(92\)90589-1](https://doi.org/10.1016/0300-9629(92)90589-1)
- Senner, N. R., Conklin, J. R., & Piersma, T. (2015). An ontogenetic perspective on individual differences. *Proceedings of the Royal Society B. Biological Sciences*, 282, 20151050. <https://doi.org/10.1098/rspb.2015.1050>
- Sydeman, W. J., Hester, M. M., Thayer, J. A., Gress, F., Martin, P., & Buffa, J. (2001). Climate change, reproductive performance and diet composition of marine birds in the southern California Current system, 1969–1997. *Progress in Oceanography*, 49, 309–329. [https://doi.org/10.1016/S0079-6611\(01\)00028-3](https://doi.org/10.1016/S0079-6611(01)00028-3)
- Taylor, C. R., & Weibel, E. R. (1981). Design of the mammalian respiratory system. I. Problem and strategy. *Respiration Physiology*, 44, 1–10. [https://doi.org/10.1016/0034-5687\(81\)90073-6](https://doi.org/10.1016/0034-5687(81)90073-6)
- Tulp, I., & de Goeij, P. (1994). Evaluating wader habitats in Roebuck Bay (North-western Australia) as a springboard for Northbound migration in waders, with a focus on Great Knots. *Emu*, 94, 78–95. <https://doi.org/10.1071/MU9940078>
- van Gils, J. A., Battley, P. F., Piersma, T., & Drent, R. (2005). Reinterpretation of gizzard sizes of red knots world-wide emphasises overriding importance of prey quality at migratory stopover sites. *Proceedings of the Royal Society B. Biological Sciences*, 272, 2609–2618. <https://doi.org/10.1098/rspb.2005.3245>
- van Gils, J. A., de Rooij, S. R., van Belle, J., van der Meer, J., Dekinga, A., Piersma, T., & Drent, R. (2005). Digestive bottleneck affects foraging decisions in red knots *Calidris canutus*. I. Prey choice. *Journal of Animal Ecology*, 74, 105–119. <https://doi.org/10.1111/j.1365-2656.2004.00903.x>
- van Gils, J. A., Lisovski, S., Lok, T., Meissner, W., Ozarowska, A., de Fouw, J., ... Klaassen, M. (2016). Body shrinkage due to Arctic warming reduces red knot fitness in tropical wintering range. *Science*, 352, 819–821. <https://doi.org/10.1126/science.aad6351>
- van Gils, J. A., Piersma, T., Dekinga, A., & Battley, P. F. (2006). Modelling phenotypic flexibility: An optimality analysis of gizzard size in Red Knots (*Calidris canutus*). *Ardea*, 94, 409–420.
- van Gils, J. A., Piersma, T., Dekinga, A., & Dietz, M. W. (2003). Cost-benefit analysis of mollusc-eating in a shorebird. II. Optimizing gizzard size in the face of seasonal demands. *Journal of Experimental Biology*, 206, 3369–3380. <https://doi.org/10.1242/jeb.00546>
- Vézina, F., Love, O. P., Lessard, M., & Williams, T. D. (2009). Shifts in metabolic demands in growing altricial nestlings illustrate context-specific relationships between BMR and body composition. *Physiological and Biochemical Zoology*, 82, 248–257. <https://doi.org/10.1086/597548>
- Vézina, F., Williams, T. D., Piersma, T., & Morrison, R. I. G. (2012). Phenotypic compromises in a long-distance migrant during the transition from migration to reproduction in the High Arctic. *Functional Ecology*, 26, 500–512. <https://doi.org/10.1111/j.1365-2435.2011.01955.x>
- Wang, Y., Ren, M., & Zhu, D. (1986). Sediment supply to the continental shelf by the major rivers of China. *Journal of the Geological Society*, 143, 935–944. <https://doi.org/10.1144/gsjgs.143.6.0935>
- Worrall, D. H. (1984). Diet of the Dunlin *Calidris alpina* in the Severn Estuary. *Bird Study*, 31, 203–212. <https://doi.org/10.1080/00063658409476842>

- Yang, H. Y., Chen, B., Ma, Z. J., Hua, N., van Gils, J. A., Zhang, Z. W., & Piersma, T. (2013). Economic design in a long-distance migrating molluscivore: How fast-fuelling red knots in Bohai Bay, China, get away with small gizzards. *Journal of Experimental Biology*, 216, 3627–3636. <https://doi.org/10.1242/jeb.083576>
- Zhang, S.-D., Ma, Z., Choi, C.-Y., Peng, H.-B., Bai, Q.-Q., Liu, W.-L., ... Piersma, T. (2018). Persistent use of a shorebird staging site in the Yellow Sea despite severe declines in food resources implies a lack of alternatives. *Bird Conservation International*, 28, 534–548. <https://doi.org/10.1017/S0959270917000430>
- Zwarts, L., & Blomert, A. M. (1992). Why knot *Calidris canutus* take medium-sized *Macoma balthica* when six prey species are available. *Marine Ecology Progress Series*, 83, 113–128. <https://doi.org/10.3354/meps083113>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Zhang S-D, Ma Z, Choi C-Y, et al. Morphological and digestive adjustments buffer performance: How staging shorebirds cope with severe food declines. *Ecol Evol*. 2019;9:3868–3878. <https://doi.org/10.1002/ece3.5013>