

University of Groningen

Giant Arcuate Lupus Vulgaris with Rapid Progression

Parajuli, Niraj; Jha, Harendra Kumar; Jonkman, Marcel Franciscus

Published in:
Journal of nepal medical association

DOI:
[10.31729/jnma.4453](https://doi.org/10.31729/jnma.4453)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Parajuli, N., Jha, H. K., & Jonkman, M. F. (2019). Giant Arcuate Lupus Vulgaris with Rapid Progression: A Case Report. *Journal of nepal medical association*, 57, 275-277. [218]. <https://doi.org/10.31729/jnma.4453>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Giant Arcuate Lupus Vulgaris with Rapid Progression: A Case Report

Niraj Parajuli,¹ Harendra Kumar Jha,² Marcel Franciscus Jonkman³

¹Department of Dermatology and Venereology, National Academy of Medical Sciences, Bir Hospital, Kathmandu, Nepal, ²Chitwan Hospital, Bharatpur, Nepal, ³Department of Dermatology, University Medical Center, Groningen, Netherlands.

ABSTRACT

Lupus vulgaris is the commonest form of cutaneous tuberculosis. It is a chronic and slowly progressive disease. It can be transmitted either through hematogenous or lymphatic spread but most commonly through contiguous extension. There are many reports on different form of lupus vulgaris but there are only a few reports on large sized lupus vulgaris. Here, we report a case of 75-year-old man with a giant lupus vulgaris rapidly progressing in just two year's duration.

Keywords: *cutaneous tuberculosis; lupus vulgaris; Nepal.*

INTRODUCTION

Lupus vulgaris is the most common form of cutaneous tuberculosis which is chronic and progressive in nature.¹ This form of cutaneous tuberculosis is found in individual with moderate immunity and a strong tuberculin sensitivity. A characteristic feature of lupus vulgaris is an extremely chronic course with slow but steady growth of the lesions over a period of many years, or even decades.

We present here a case of giant arcuate shaped lupus vulgaris, rapidly progressing over a period of just two years.

CASE REPORT

A 75 years old male came to dermatology out-patient with a single large atrophic, hypopigmented to skin colored plaque on antero-lateral chest wall for the last two years. According to the patient, the lesion started as few asymptomatic erythematous papules that coalesced to form plaque in the lateral part of left chest. This plaque increased gradually leaving a healed atrophic center. The borders were covered with thick crusts. There was no history of cough, fever, night sweats, or weight loss. There was no history of tuberculosis or malignancy in other family members. Patient was non-

smoker and non-alcoholic beverage drinker.

On examination, a single large plaque on left anterior chest wall covering the whole of the left chest superiorly from clavicular line involving the entire left chest up to the subcostal region and extending posteriorly to infrascapular region, sparing the left axilla. The approximate size of the plaque was 30 x 30 cm² (Figure 1). It was arcuate in shape with clearing of the adjacent areas with few areas of atrophy. The parasternal and subcostal margin of the plaque had thick adherent crusts on an erythematous base. There were areas of scarring and atrophy in the lesion. The plaque was non-tender and diascopy was negative for apply jelly nodules. The corresponding lymph nodes were not enlarged.

Complete blood counts were within normal limits but a raised erythrocyte sedimentation rate of 105mm/hr was noted. Mantoux test performed showed an induration of 20 mm. Chest x-ray showed no abnormalities. Sputum for acid-fast bacilli did not show any acid-fast bacilli. Liver function test and renal function test were all within normal limits.

Correspondence: Dr. Niraj Parajuli, Department of Dermatology and Venereology, National Academy of Medical Sciences, Bir Hospital, Kathmandu, Nepal. Email: drnirajparajuli@gmail.com, Phone: +977-9841777800.



Figure 1. Clinical photograph showing a single large plaque on the whole of left chest.

An incisional biopsy done from the plaque showed pseudo-epitheliomatous hyperplasia, neutrophilic abscess formation in the corneal layer. Underlying dermis showed ill-defined epithelioid cell granuloma, langhans giant cells and a surrounding inflammatory infiltrates consisting of plasma cells, lymphocytes and neutrophilic micro abscesses and fibrosis (Figure 2a and 2b).




Figure 2a. HPE shows pseudo-epitheliomatous hyperplasia, neutrophilic abscess formation in the corneal layer. Underlying dermis shows ill-defined epithelioid cell granuloma (40x magnification, H & E stain).




Figure 2b. HPE showing epithelioid cell granuloma, Langhans giant cells, surrounding mixed inflammatory infiltrate consisting of plasma cells and lymphocytes (100x magnifications, H & E stain).

The patient was diagnosed as a case of lupus vulgaris and was started on anti-tubercular treatment category 1 and was advised for follow-up.

DISCUSSION

Cutaneous tuberculosis results from a chronic infection by *Mycobacterium tuberculosis*, *M. bovis* and occasionally by the Calmette-Guerin bacillus. The clinical manifestations are variable which depends on the interaction of several factors including the site of infection and the host's immunity.²

Lupus vulgaris a chronic and progressive form of cutaneous tuberculosis. Lesions are generally solitary and found on the head and neck region. The clinical presentation can be of several different clinical appearances.³ It can be transmitted either through hematogenous or lymphatic spread but most commonly through contiguous extension. Rarely, it can also occur from exogenous inoculation or on post- BCG scar.⁴ The common locations of lupus vulgaris are the lower half of the body involving legs, thighs, buttocks, and feet.⁵ The different variants of lupus vulgaris includes plaque, ulcerative, mutilating, vegetative, papular or nodule and hypertrophic form. A prominent feature of lupus vulgaris is atrophic scarring, with or without ulceration.⁶ Typically, the histopathology shows tuberculoid granulomas composed of lymphocytes, plasma cells, epithelioid and giant cells, scant or absent central caseation, in the superficial dermis. The epidermis is usually hyperplastic, but may be atrophic or ulcerated. Stain for acid fast bacilli is usually negative. A characteristic feature of lupus vulgaris is its extremely chronic course. In this reported case, the patient had a

large plaque within a short period of just over 2 years. The largest reported case of lupus vulgaris in literature is from India with size of 60x45 cm and 40x 50cm.⁷

Consent: [JNMA Case Report Consent Form](#) was signed by the patient and the original is attached with the patient's chart.

Conflict of Interest: None.

REFERENCES

1. Sehgal VN, Srivastava G, Bajaj P, Sengal R. Re-infection (secondary) inoculation cutaneous tuberculosis. *Int J Dermatol.* 2001 Mar;40(3):205-9. [[PubMed](#)]
2. dos Santos JB, Figueiredo AR, Ferraz CE, de Oliveira MH, da Silva PG, de Medeiros VLS. Cutaneous tuberculosis: epidemiologic, etiopathogenic and clinical aspects - Part I. *An Bras Dermatol.* 2014;89(2):219-29. [[PubMed](#) | [FullText](#)]
3. Turan E, Yurt N, Yesilova Y, Celik OI. Lupus vulgaris diagnosed after 37 years: a case of delayed diagnosis. *Dermatol Online J.* 2012;18(5):13. [[PubMed](#) | [FullText](#)]
4. Kanwar AJ. Lupus vulgaris following BCG vaccination. *Int J Dermatol.* 1988;27:525-6. [[PubMed](#)]
5. Kumar B, Muralidhar S. Cutaneous tuberculosis: a twenty-year prospective study. *Int J Tuberc Lung Dis.* 1999 Jun;3(6):494-500. [[PubMed](#)]
6. Jain VK, Aggarwal K, Jain S, Singh S. Hypertrophic lupus vulgaris: An unusual presentation. *Indian J Dermatol.* 2009;54:287-9. [[PubMed](#) | [FullText](#)]
7. Sacchidanand S, Sharavana S, Mallikarjun M, Nataraja HV. Giant lupus vulgaris: A rare presentation. *Indian Dermatol Online J.* 2012 Jan-Apr;3(1):34-6. [[PubMed](#) | [FullText](#)]

© The Author(s) 2018.

This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>