

University of Groningen

Night-time activity forecast by season and weather in a longitudinal design

Wahnschaffe, Amely; Nowozin, Claudia; Rath, Andreas; Floessner, Theresa; Appelhoff, Stefan; Muench, Mirjam; Kunz, Dieter

Published in:
International Psychogeriatrics

DOI:
[10.1017/S1041610217001235](https://doi.org/10.1017/S1041610217001235)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wahnschaffe, A., Nowozin, C., Rath, A., Floessner, T., Appelhoff, S., Muench, M., & Kunz, D. (2017). Night-time activity forecast by season and weather in a longitudinal design: natural light effects on three years' rest-activity cycles in nursing home residents with dementia. *International Psychogeriatrics*, 29(12), 2071-2080. <https://doi.org/10.1017/S1041610217001235>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Night-time activity forecast by season and weather in a longitudinal design – natural light effects on three years' rest-activity cycles in nursing home residents with dementia

Amely Wahnschaffe,^{1,2} Claudia Nowozin,^{1,2} Andreas Rath,³ Theresa Floessner,⁴ Stefan Appelhoff,^{1,2} Mirjam Münch^{1,2} and Dieter Kunz^{1,2}

¹Charité – Universitätsmedizin Berlin, Sleep Research & Clinical Chronobiology, Institute of Physiology, Berlin, Germany

²St. Hedwig-Hospital, Clinic for Sleep and Chronomedicine, Berlin, Germany

³SplittFennert Nursing Home, Berlin, Germany

⁴University of Groningen, Groningen Institute for Evolutionary Life Sciences, Chronobiology Unit, Groningen, The Netherlands

ABSTRACT

Background: Night-time agitation is a frequent symptom of dementia. It often causes nursing home admission and has been linked to circadian rhythm disturbances. A positive influence of light interventions on night-time agitation was shown in several studies. The aim of our study was to investigate whether there is a long-term association between regional weather data (as indicator for daylight availability) and 24-hour variations of motor activity.

Methods: Motor activity of 20 elderly nursing home residents living with dementia was analyzed using recordings of continuously worn wrist activity monitors over a three-year period. The average recording duration was 479 ± 206 days per participant (mean \pm SD). Regional cloud amount and day length data from the local weather station (latitude: $52^{\circ}56'N$) were included in the analysis to investigate their effects on several activity variables.

Results: Nocturnal rest, here defined as the five consecutive hours with the least motor activity during 24 hours (L5), was the most predictable activity variable per participant. There was a significant interaction of night-time activity with day length and cloud amount ($F_{1,1174} = 4.39$; $p = 0.036$). Night-time activity was higher on cloudy short days than on clear short days ($p = 0.007$), and it was also higher on cloudy short days than on cloudy long days ($p = 0.032$).

Conclusions: The need for sufficient *zeitgeber* (time cue) strength during winter time, especially when days are short and skies are cloudy, is crucial for elderly people living with dementia. Activity forecast by season and weather might be a valuable approach to anticipate adequately complementary use of electrical light and thereby foster lower night-time activity.

Key words: rest-activity cycle, agitation, dementia, light, season, weather, circadian

Introduction

Night-time agitation, a symptom complex that is part of different subtypes of dementia, includes confusion, agitation, aggressive behavior, aimless wandering, and disturbances of the sleep–wake cycle (Burns *et al.*, 1990; Finkel, 2000). Because

the need for night care also impairs sleep duration and quality of caregiving relatives, night-time agitation is a frequent cause for nursing home placement and high care costs (Sanford, 1975; Spira *et al.*, 2012). Pharmacological interventions show inconsistent results and bear the risk of side effects especially in the elderly (Declercq *et al.*, 2013; Mahlberg *et al.*, 2004; Seitz *et al.*, 2011). Both sleep disturbances in old age (Campbell *et al.*, 1988; Satlin *et al.*, 1991; van Someren *et al.*, 1993) and night-time agitation in dementia were reported to be linked to a dysregulation of the circadian system that causes disturbances in sleep–wake regulation (Witting *et al.*, 1990;

Correspondence should be addressed to: Dr rer. medic. Amely Wahnschaffe, Research Group: Sleep & Clinical Chronobiology, Institute of Physiology (CBF), Charité – Universitätsmedizin Berlin, Grosse Hamburger Strasse 5–11, 10115 Berlin, Germany. Phone: +49 30 23112901; Fax: +49 30 23112903. Email: amely.wahnschaffe@gmx.de. Received 10 Jan 2017; revision requested 24 Feb 2017; revised version received 28 May 2017; accepted 7 Jun 2017. First published online 10 July 2017.

Cohen-Mansfield *et al.*, 1995; van Someren *et al.*, 1996; Altimus *et al.*, 2010; Bedrosian and Nelson, 2012).

The circadian system in humans is governed by a master clock located in the suprachiasmatic nuclei (SCN) that modulates a system of peripheral internal clocks. Light is the main *zeitgeber* (“time cue,” a term coined by Jürgen Aschoff, according to Oxford English Dictionary 2016 “a rhythmically occurring natural phenomenon which acts as a cue in the regulation of the body’s circadian rhythms,” see also Lockley, 2009) of the circadian timing system, predominantly conveyed via melatonin-expressing retinal ganglion cells (with greatest sensitivity at 480 nm); these cells are also modulated by rod and cone input (Altimus *et al.*, 2010), which project directly to the SCN via the retinohypothalamic tract (Foster, 2005). The master clock in the SCN and peripheral biological clocks thus stay aligned with the astronomical 24 hours day with its natural dichotomy of light and dark in seasonally changing proportions and its spectral transitions at dusk and dawn.

With age, the synchronization between internal and external clocks tends to weaken and the amplitudes of circadian rhythms (e.g. body temperature, melatonin, sleep propensity) flatten (Sellix *et al.*; Van Gool and Mirmiran, 1986). In dementia, the decrease of circadian strength is even more severe (Liu *et al.*, 2000; Harper *et al.*, 2005; Wu *et al.*, 2007). Potential physiological reasons for these changes are a functional disruption in the SCN starting in the early stages of dementia (Liu *et al.*, 2000; Wu *et al.*, 2007), a decline of melatonin production related to pineal calcification (Mahlberg *et al.*, 2008) and age-related changes in the functioning of the eye with less light reaching the retina (Turner *et al.*, 2010). Insufficient exposure to the main *zeitgeber* light adds to this weakening of the circadian system in old age and dementia (Mishima *et al.*, 2001). Natural light could thus serve as an ideal model for the design of dynamic lighting systems that use different light intensities and spectral compositions in the course of 24 hours (e.g. Figueiro, 2008).

Some studies aiming to stabilize circadian rhythms with electrical light interventions reported positive influences on night-time agitation (e.g. Lovell *et al.*, 1995; Van Someren *et al.*, 1997a; Mishima *et al.*, 1998; Skjerve *et al.*, 2004; Burns *et al.*, 2009; Wahnschaffe *et al.* 2017). Natural lighting conditions change with season and weather. Thus, season and weather characteristics could also add to or compensate for an existing lack of natural light exposure for entraining circadian rhythms in the elderly. Studies focusing on links between weather and circadian parameters in

the elderly (Obayashi *et al.*, 2012) or people with dementia (Figueiro *et al.*, 2012) are rare. A recent study showed that higher actual light exposures (measured with wrist worn luxmeters) had positive impacts on emotions, alertness, and life quality in nursing home residents living with dementia (Münch *et al.* 2017). Understanding the impact of season and weather on rest-activity cycles in nursing home residents with dementia could enhance the development of architectural environments and compensatory electrical lighting systems or treatment with exogenous melatonin (e.g. Riemersma-van der Lek *et al.*, 2008).

Therefore, the objective of the present study was to determine in a retrospective and explorative analysis the impact of day length and cloud amount as season- and weather-dependent natural light parameters on rest-activity cycles of 20 elderly nursing home residents living with dementia. We hypothesized that day length and cloud amount as major determinants of outside day light patterns affect rest-activity cycles in residents with dementia.

Methods

We investigated the influence of season and weather on 24-hour rest-activity rhythms in nursing home residents living with dementia. Therefore, we retrospectively analyzed a data set that was collected to assess the effects of a dynamic lighting system (Wahnschaffe *et al.*, 2017).

Setting

The nursing home (“Splitt-Fennert”, Berlin, Germany, latitude 52.506 N) is a small institution, specialized on care of residents living with dementia. At the time of this data collection, there were 34 residents living there. It is located in an old villa with large windows made off standard glass, allowing high daylight fluxes to enter the residents’ bedrooms and common rooms (about 90% of natural light are transmitted through standard glass). Windows could be covered with curtains, when residents went to sleep or when they felt blinded by direct sunlight. During daylight hours, windows remained mostly uncovered. Most residents spent the majority of their waking time in the common living room or outside in the garden, if the weather allowed it (information from personal communication with nursing home staff). Accordingly, a high amount of natural light exposure across seasons was warranted for all participating residents.

Electrical lighting consisted of ordinary room light from ceiling mounted and standard lamps

equipped with incandescent light bulbs (2700 K). This kind of illumination is known to be between 70 and 170 lx at the eye level and was switched on/off according to individual needs. One of the nursing home's care guidelines is to only use a minimum of sedative medication, which reduces its possible confounding effect in the study. Psychopharmacological medication in the study sample included anti-depressants (three residents received citalopram, three mirtazapine, one venlafaxine), neuroleptics (three risperidone, three pipamperone, one melperone), antidementives (one memantine, two donepezil), and only one resident received bencodiazepines (oxacepam). Dosages ranged in the lower third of recommended amounts.

Participants

Inclusion criteria for the study were being a resident in the assigned nursing home and having a diagnosis of dementia. Exclusion criteria were bilateral blindness and a lack of tolerance to wear an actigraph as expressed verbally or non-verbally by efforts to get rid of the device or signs of stress when it was put on. The study was conducted in accordance with the Declaration of Helsinki; legal representatives of all participants had given their written, informed consent. Only residents with reliable activity data (see below) for more than 120 days were included in the analysis. Twenty residents were included in the analysis (19 female and 1 male), their mean age being 83.8 years (\pm 8.8 SD) at the beginning of the study. The dementia diagnoses were made based on ICD-10 criteria and included: Alzheimer's disease ($n = 9$); vascular dementia ($n = 2$); fronto-temporal dementia ($n = 1$); Korsakoff's syndrome ($n = 1$) and dementia subtypes, which were not further specified ($n = 7$). Regarding psychiatric comorbidity, there were three residents with a concomitant diagnosis of depression. Two residents had cataract replacements, one patient had a retinal detachment in one eye, and one patient was visually blind in one eye.

Activity data

Rest-activity cycles were continuously recorded over a three-year period between July 21, 2009 and June 17, 2012 by activity monitors (Actiwatch, Cambridge Neurotechnology, UK), worn on the non-dominant wrist. Rest-activity data was recorded in 2-minute intervals. Every three months, data were downloaded onto a hard drive. Data processing was performed semi-manually by an independent scorer. The 24-hour days with missing data for less than 3 hours were edited with the 24-hour average, and 24-hour days with more than

3 hours of missing data were excluded from the analysis, similarly as it was done by Bromundt *et al.* (2012). A non-parametric circadian rhythm analysis (Sokolove and Bushnell, 1978; Van Someren *et al.*, 1999) was performed on edited rest-activity data with the software Sleep Analysis 7 (Software Package Cambridge Neurotechnology, UK, Version 7.3) and the following parameters were calculated: (1) relative amplitude (RA), defined as the difference between the average hourly movement duration over the most active uninterrupted 10-hour period (day) and the least active uninterrupted 5-hour period (night) of the average 24-hour pattern, normalized with the sum of day and night activity to a scale from 0 to 1; (2) inter-daily stability (IS): quantifying the invariability between the days ("strength of coupling to *zeitgebers* (=time cues)") (Van Someren *et al.*, 1999); scale 0 to 1; the higher, the more stable); (3) intra-daily variability (IV): quantifying the frequency and extent of transitions between rest and activity ("fragmentation of the rhythm"; scale 1 to > 2 ; the higher, the more fragmented (Van Someren *et al.*, 1999)); (4) mean hourly movement duration during the uninterrupted 5-hour period with the lowest activity level in the average 24-hour pattern (L5), with high L5 indicating nocturnal restlessness; (5) clock time at the beginning of L5 (L5 onset); (6) mean hourly movement duration during the uninterrupted 10-hour period with the highest activity level in the average 24-hour pattern (M10); (7) clock time at the beginning of M10 (M10 onset; for formulas, see Van Someren *et al.*, 1997b; 1999) based on forward-moving averages from seven-day intervals (i.e. the value on a certain date represents the parameters calculated from the interval including the current day and the following six days), as recommended for this software.

Weather data

Daily regional weather data (Berlin Tegel Airport, Germany, latitude 52.506 N, 6.3 km linear distance from the nursing home) was downloaded from the webpage of the German Meteorological Services (DWD, 2015) for the time span of activity data acquisition.

The variable "cloud amount" was included in further analysis as determining variable defined as the proportion of the sky that is completely covered by clouds. Cloud amount is expressed as daily arithmetic mean of three measurements at fixed times: 7:30 hours, 14:30 hours, and 21.30 hours (DWD, 2015). Its assessment is conducted by a human observer and is reported on a scale between 0 and 8; 0 is indicating that there are no clouds at all; 4, that maximally 50% of the sky is covered

Table 1. Descriptive statistics of circadian rest-activity variables

ACTIVITY PARAMETER	MEAN	SD
IS	0.42	0.10
IV	1.28	0.34
RA	0.68	0.14
L5	1542	1233
M10	7895	5161
L5 onset (h)	22:13	1:56
M10 onset (h)	9:01	1:26

Values are reported in arbitrary units, except for L5 and M10 onsets.

Abbreviations: SD = standard deviation; IS = interdaily stability (strength of coupling to *zeitgebers*; the higher, the more stable), IV = intradaily variability (fragmentation of the circadian rest-activity rhythm; the higher, the more fragmented); RA = relative amplitude (L5/M10 ratio), L5 = activity during the 5-hour period with the lowest activity during 24 hours; M10 = activity during the 10-hour period with the highest activity during 24 hours; L5 onset = start time of L5; M10 onset = start time of M10.

by clouds; and 8 that the whole sky is covered by clouds. Because sunshine duration correlates strongly and inversely with cloud amount (Pearson correlation in our data set: $r = -0.835$, $p < 0.0001$), sunshine duration was not entered in the analysis as relevant separate variable. The variable “day length” was derived from a commercial website (Gerding, 2015) for the latitude of the nursing home and verified with the Berlin communal weather web page (BerlinOnline, 2015). Day length is defined as the duration from sunrise until sunset in hours (Oxford Dictionary, 2016).

Statistical analyses

All statistical analyses were conducted with PASW Statistics 18 (2009, SPSS Inc., Hong Kong). We first calculated descriptive statistics for sample and activity data (see Table 1). Second, we explored the association of cloud amount and day length with the different rest-activity variables to identify relevant parameters for further analysis. To this end, we performed for each participant linear multiple regression analyses with weather parameters (cloud amount, day length) as predictors and rest-activity parameters (IS, IV, and RA indicating circadian rhythm characteristics; L5 indicating nocturnal restlessness) as dependent variables (see Table 2). To rule out random day-by-day variations and include preceding time periods (taking into account light history and slow and cumulative effects of light on circadian rhythms), we smoothed weather parameters in 14-day backward-moving averages. For activity parameters, we choose the corresponding forward-moving seven-day-based averages. We chose backward-moving averages

for weather and forward-moving averages for activity because we presuppose that prior weather influences the following activity and never the other way round. We only included every seventh value in the analysis to avoid overestimation of associations by combining the same values several times as a consequence of the moving average data due to the seven-day intervals. To control for potential subgroup differences between dementia subtypes, we exploratively compared intra-individual linear regressions (cloud amount predicting rest-activity variables as described above) between residents with a diagnosis of Alzheimer’s disease and those with other dementia subtype diagnoses in an independent sample *t*-test.

After selecting L5 as the rest-activity variable with the highest intra-individual regression coefficient as predictor of cloud amount, we plotted a sequence chart of L5 in relation to cloud amount and day length. Then we calculated *post-hoc* Pearson correlations between date and activity parameters (L5, M10) to describe development over time after visual analysis of the graph as well as between cloud amount and L5 to test the observed correspondences for significance. Last, a linear mixed-model analysis was performed using cloud amount and day length both as dichotomous variables (cloudy vs. clear days; long vs. short days), by applying a median split on all available days. For significant interactions, *t*-tests for dependent samples were used to assess differences in the outcome measures between four fixed subgroups: short cloudy days, long cloudy days, short clear days, and long clear days.

Results

The time intervals of included rest-activity data ranged between 140 and 723 days (mean \pm SD: 479 ± 206 days) per participant. Mean values of rest-activity variables (\pm SD) are displayed in Table 1 and were for IS: $0.42 (\pm 0.10)$; for IV: $1.28 (\pm 0.34)$; and for RA: $0.68 (\pm 0.14)$; L5: $1542 (\pm 1233)$; M10 = $7895 (\pm 5161)$; L5 onset (h): $22:13 (\pm 1:56)$; and M10 onset (h): $9:01 (\pm 1:26)$. The onset times indicate that M10 predominantly occurred during daytime and L5 during night-time.

The multiple regression analyses for predicting rest-activity-related variables (IS, IV, L5, and RA) with light-related weather parameters (cloud amount and day length) per participant showed a high inter-individual variance of regression coefficients (see Table 2). There were no significant differences in regression coefficients between subgroups of residents, i.e. participants living with Alzheimer’s disease and those having other

Table 2. Intra-individual linear regression analyses

LINEAR REGRESSION (R^2)	PREDICTORS: CLOUD AMOUNT, DAY LENGTH			
	DEPENDENT VARIABLE			
SUBJECT	IS	IV	L5	RA
1	0.03	0.03	0.00	0.03
2	0.01	0.17	0.17	0.02
4	0.07	0.45	0.35	0.08
5	0.06	0.11	0.50	0.14
6	0.53	0.02	0.38	0.22
7	0.06	0.01	0.02	0.05
8	0.00	0.19	0.04	0.02
9	0.03	0.03	0.02	0.14
10	0.03	0.00	0.00	0.02
11	0.05	0.06	0.08	0.02
12	0.01	0.02	0.10	0.18
14	0.02	0.02	0.42	0.26
15	0.02	0.00	0.06	0.01
16	0.00	0.02	0.07	0.00
21	0.03	0.04	0.07	0.19
23	0.11	0.03	0.34	0.11
29	0.1	0.04	0.10	0.01
30	0.05	0.10	0.08	0.01
31	0.01	0.09	0.08	0.00
32	0.01	0.00	0.05	0.01
Mean (SD)	0.06(0.11)	0.07 (0.11)	0.15 (0.15)	0.08 (0.08)
Number of $p < .05$	1	5	11	4

The 5 hours with the least consecutive motor activity during 24 hours (L5), indicative for nocturnal rest, was the most predictable activity variable per participant. Dependent variables: individual activity parameters (seven-day forward-moving averages). Predictors: cloud amount and day length (14-day backward-moving averages). Only every seventh day was included in the analysis to avoid overestimation of associations caused by moving averages. Significant results ($p < 0.05$) are marked in bold.

Abbreviations: IS = interdaily stability (strength of the coupling to *zeitgebers*; the higher, the more stable); IV = intradaily variability (fragmentation of the rhythm; the higher, the more fragmented); RA = relative amplitude (relative ratio of day and night activity); L5 = activity during the 5-hour period with the lowest activity during 24 hours.

dementia subtype diagnoses ($p > 0.05$). Therefore, the analyses were performed with the entire group of study participants within one sample.

Intra-individual regression coefficients showed that cloud amount and day length were significant predictors for IS in only two participants, for IV in five participants, and for RA in four participants (Table 2). On the other hand, L5 was significantly predicted by cloud amount in 11 ($0.074 \geq R^2 \leq 0.504$; $p < 0.05$) out of 20 participants. Thus, the rest-activity parameter L5 had the highest proportion of significant predictions by cloud amount and day length and the highest overall mean \pm SD ($R^2 = 0.146 \pm 0.153$) across participants. L5 was therefore chosen as the dependent variable for further analysis. Figure 1 shows the time course of mean L5 (as an indicator of nocturnal restlessness) in relation to cloud amount and day length over the study period ($n = 20$). The time courses of the two variables mean L5 and cloud amount show visually some similarities with co-occurrence of peaks and troughs. They correlate moderately but significantly ($r = 0.206$,

$p = 0.015$). A decline of L5 over time is observable and is reflected in a significant negative correlation of L5 with date ($r = -0.524$; $p < 0.01$). This decline might be attributed to a general decline in activity due to illness progression because M10 also declines with time (M10 vs. date: $r = -0.676$; $p < 0.01$).

For L5, the mixed-model analysis showed a significant interaction between day length and cloud amount ($p = 0.036$; $F_{1,1174} = 4.39$; Figure 2). There were no significant main effects of cloud amount or day length. *Post-hoc* tests showed that night-time activity was higher during cloudy short days when compared to clear short days ($T_{19} = 3.1$; $p = 0.007$). Night-time activity was also higher during cloudy short days when compared to cloudy long days ($T_{17} = -2.3$; $p = 0.032$).

Discussion

The aim of the study was to analyze possible effects of season and weather on rest-activity patterns in




Figure 1. Time course of nocturnal restlessness (L5) is related to cloud amount and day length. Thick black line (left y-axis) represents the time course of L5 (30-day centered moving average; $n = 20$ nursing-home residents with dementia; 140–723 days of activity recordings); thin black lines indicate \pm standard errors. Grey line (right y-axis) represents time course of cloud amount (30-day backward-moving average). Day length is represented dichotomously as short (<12:08 hours; light grey rectangle) versus long days (>12:08 hours; white rectangles). The dates (month, year) are noted on the horizontal x-axis. Time courses of cloud amount and L5 are significantly associated ($r = 0.206, p < 0.05$). Abbreviations and definitions: L5 = averaged movement counts per hour (arbitrary unit) during the 5-hour period with the lowest activity during 24 hours indicating nocturnal restlessness; cloud amount = size of the part of the sky that is completely covered by clouds reported in eighths.




Figure 2. Nocturnal restlessness is influenced by natural light conditions. Bars represent L5 group means calculated from individual means in subgroups; error bars represent standard errors. In the mixed-model analysis with nocturnal restlessness (L5) as dependent variable there was a significant interaction with day length and cloud amount ($F_{1,173,8} = 4.39; p = 0.036$; both entered in the model as dichotomous variables determined by median split and expressed as cloudy versus clear days and short versus long days, respectively; $n = 20$ nursing-home residents with dementia; 140–723 days of activity recordings). * indicates significant differences in *post-hoc* *t*-tests for dependent samples ($p < 0.05$). Abbreviations: L5 = averaged movement counts per hour (arbitrary unit) during the 5-hour period with the lowest activity during 24 hours indicating nocturnal restlessness.

nursing home residents with dementia. Activity during the 5 hours with the lowest activity within 24 hours (L5) turned out to be the rest-activity parameter that showed the highest association with cloud amount and day length in the explorative analysis. These 5 hours predominantly occurred during night time (L5 onset mean \pm SD = 22:13 \pm 1:56 hours) and might be an objective indicator of the clinical observation of nocturnal restlessness as well as a proxy for night-time agitation in dementia (Burns *et al.*, 1990; Finkel, 2000). Our results showed that season- and weather-dependent natural lighting conditions had an influence on nocturnal restlessness. Cloud amount and day length interacted in their impact on night-time activity: only when there was already little natural light available regarding intensity (cloud amount) or exposure duration (day length) did a further reduction of natural light dose play a role.

Several attempts to stabilize rest-activity cycles of nursing home residents and home-dwelling elderly with dementia through interventions with light therapy or enhanced ambient light yielded inconsistent results. However, most studies reported a stabilizing effect of light therapy on rest-activity patterns (Satlin *et al.*, 1992; Mishima *et al.*, 1994; Ancoli-Israel *et al.*, 2002, 2003; Dowling *et al.*, 2005), on agitation (Lovell *et al.*, 1995; Mishima *et al.*, 1998; Skjerve *et al.*, 2004; Burns *et al.*, 2009), and on sleep (Lyketsos *et al.*, 1999; Fetveit and Bjorvatn, 2005). Light-intervention studies are mostly conducted in the winter months to yield clearer results by avoiding the influencing effects of natural light. Our results suggest that daylight should also be taken into account as a potentially effective intervention. Cloud amount could be a still-underestimated interacting variable in light-intervention studies because from our results, it seems relevant for nocturnal activity (which is at least partly circadian regulated), especially during the winter months.

Seasonal variations in circadian rhythms of hormones, mood, and activity are often reported for northern latitudes with variable daylight lengths but not for latitudes close to the equator with lower day length variability (e.g. Laakso *et al.*, 1994; Park *et al.*, 2007; Friberg *et al.*, 2012). Former studies on day length and weather influences on activity only address overall activity levels and not the circadian pattern of activity (e.g. Feinglass *et al.*, 2011; Klenk *et al.*, 2011; Goodman *et al.*, 2012). We found only two studies directly addressing daylight and weather effects on circadian parameters in elderly and persons with dementia. Obayashi and colleagues reported from a field study with a sample of 192 elderly a positive association of duration of bright daylight exposure (>1,000 lx) and amount of

urinary 6-sulfatoxy melatonin excretion (Obayashi *et al.*, 2012). Another field study with people with dementia showed lower rates of daylight exposure and a lower amplitude of the circadian-driven rest-activity rhythm in winter (Figueiro *et al.*, 2012). Other studies report a generally low level of light exposure in nursing-home residents with dementia (Ancoli-Israel *et al.*, 1997; Shochat *et al.*, 2000). The potentially lower circadian amplitude with lower light exposure (Figueiro *et al.*, 2012; Obayashi *et al.*, 2012) suggests that more daylight would always improve rest-activity cycles. It was therefore surprising that our findings suggest that beyond a certain level of light exposure, additional light does not lead to further improvements. Several causes could lead to the observation: (1) in the summer, the investigated nursing-home residents were not only exposed to daylight through the windows for longer periods of time but also spent more time outside in the garden so that they were more activated during the day and could receive relevant circadian light input even during cloudy days. Accordingly, light exposure through the windows of the nursing home on cloudy winter days may have been of much lower efficacy for circadian rhythmicity; (2) seasonal and weather-dependent differences in daily activity and habitual bedtimes might add to these effects; (3) there might have been additional disturbing influence of indoor evening light on short days. (4) a yet to be specified "saturation effect" might be responsible for the fact that the received light dose was high enough to have a positive influence on nocturnal restlessness at some point so that it could not be enhanced by more light exceeding this dose. Accordingly, it was shown earlier that the gain in phase shifting effects of light diminishes with growing exposure duration (Rimmer *et al.*, 2000; Beersma *et al.*, 2009). This effect could be caused by an interaction of response reduction under light and response restoration in darkness (Comas *et al.*, 2007); (5) an additional explanation of the fact that clear days did not significantly add to the reduction of nocturnal restlessness on long (summer) days could be the high temperatures occurring especially on such days. High ambient temperatures might cause physical discomfort, high blood pressure, and sleep disturbances especially in old age, which could counteract benefits from light effects.

There are some limitations to our study. One is the small sample size of 20 participants, which was accepted to allow for the extended data acquisition period of three years. Additionally, the sample was, while homogenous in terms of age, diagnosis, and living circumstances, heterogeneous regarding mobility and eye functioning of participants. The small sample size and its heterogeneity might also

be the reason that possible main effects of day length and cloud amount did not yield significance. It was not controlled for potentially moderating variables like social activities. The explorative character of rest-activity parameter selection and the *post-hoc* hypothesis testing determine the preliminary character of our results, which should be replicated in another sample with predefined parameters. Furthermore, we did not record systematic subjective reports on behavior that would have allowed drawing conclusions on the clinical relevance of the observed variability in night-time activity.

To conclude, the results showed lower nocturnal restlessness in nursing-home residents with dementia during summer seasons or after clearer days in fall/winter/spring. The observed higher nocturnal restlessness after short and cloudy days may indicate a weakness of light as *zeitgeber* for the circadian system, especially during time periods with low natural light levels and exposure durations. For the planning of light interventions, this would imply that complementary light exposure to electrical lighting could be highly efficient in winter and on cloudy days.

Conflict of interest

None.

Description of authors' roles

D. Kunz and A. Wahnschaffe had developed concept and design of the study. C. Nowozin, A. Wahnschaffe, and A. Rath conducted data acquisition. M. Münch, A. Wahnschaffe, T. Floessner, and S. Appelhoff analyzed and interpreted the data. A. Wahnschaffe drafted the paper and all authors revised it with substantial contributions and approved the final version.

Acknowledgments

We thank the residents of the Splitt-Fennert nursing home and their relatives for participating in the study. We thank the team for their help with collecting data and the specialist for seismology Prof Dr Frederik Tilmann for his data analysis advice (because long-term activity patterns can sometimes resemble seismic wave patterns). We thank Prof Dr Andrea Rodenbeck, Dr Frederik Bes, Dr Alexandra Papakonstantinou, and Jan de Zeeuw for their helpful comments on an earlier version of the paper and Julia Ritter for proofreading this document. The data collection

phase of the study was financially supported by the German Federal Ministry of Economics (FKZ 0319057L) as preparation for a later intervention study.

References

- Altimus, C. M. et al.** (2010). Rod photoreceptors drive circadian photoentrainment across a wide range of light intensities. *Nature Neuroscience*, 13, 1107–1112. doi: [10.1038/nn.2617](https://doi.org/10.1038/nn.2617)
- Ancoli-Israel, S. et al.** (1997). Variations in circadian rhythms of activity, sleep, and light exposure related to dementia in nursing-home patients. *Sleep*, 20, 18–23.
- Ancoli-Israel, S. et al.** (2003). Effect of light on agitation in institutionalized patients with severe Alzheimer disease. *The American Journal of Geriatric Psychiatry*, 11, 194–203.
- Ancoli-Israel, S., Martin, J. L., Kripke, D. F., Marler, M. and Klauber, M. R.** (2002). Effect of light treatment on sleep and circadian rhythms in demented nursing home patients. *Journal of the American Geriatric Society*, 50, 282–289. doi: [jgs50060](https://doi.org/jgs50060)
- Bedrosian, T. A. and Nelson, R. J.** (2012). Sundowning syndrome in aging and dementia: research in mouse models. *Experimental Neurology*, 243, 67–73.
- Beersma, D. G. et al.** (2009). The progression of circadian phase during light exposure in animals and humans. *Journal of Biological Rhythms*, 24, 153–160. doi: [10.1177/0748730408330196](https://doi.org/10.1177/0748730408330196).
- Berlin Online.** (2015). BerlinOnline Stadtportal GmbH & Co. KG. Available at: <http://www.berlin.de/wetter/>; last accessed 18 March 2015.
- Bromundt, V. et al.** (2012). Circadian sleep-wake cycles, well-being, and light therapy in borderline personality disorder. *Journal of Personality Disorders*, 27, 680–696. doi: [10.1521/pedi_2012_26_057](https://doi.org/10.1521/pedi_2012_26_057)
- Burns, A., Allen, H., Tomenson, B., Duignan, D. and Byrne, J.** (2009). Bright light therapy for agitation in dementia: a randomized controlled trial. *International Psychogeriatrics*, 21, 711–721. doi: [S1041610209008886](https://doi.org/S1041610209008886)
- Burns, A., Jacoby, R. and Levy, R.** (1990). Psychiatric phenomena in Alzheimer's disease. IV: disorders of behaviour. *British Journal of Psychiatry*, 157, 86–94.
- Campbell, S. S., Kripke, D. F. Gillin, J. C. and Hrubovcak, J. C.** (1988). Exposure to light in healthy elderly subjects and Alzheimer's patients. *Physiology and Behavior*, 42, 141–144.
- Cohen-Mansfield, J., Werner, P. and Freedman, L.** (1995). Sleep and agitation in agitated nursing home residents: an observational study. *Sleep*, 18, 674–680.
- Comas, M., Beersma, D. G., Spoelstra, K. and Daan, S.** (2007). Circadian response reduction in light and response restoration in darkness: a “skeleton” light pulse PRC study in mice (*Mus musculus*). *Journal of Biological Rhythms*, 22, 432–444. doi: [22/5/432](https://doi.org/22/5/432)
- Declercq, T. et al.** (2013). Withdrawal versus continuation of chronic antipsychotic drugs for behavioural and psychological symptoms in older people with dementia. *Cochrane Database Systematic Reviews*, (3), CD007726. doi: [10.1002/14651858.CD007726.pub2](https://doi.org/10.1002/14651858.CD007726.pub2)

- Dowling, G. A. et al.** (2005). Effect of morning bright light treatment for rest-activity disruption in institutionalized patients with severe Alzheimer's disease. *International Psychogeriatrics*, 17, 221–236.
- DWD.** (2015). Deutscher wetterdienst. Available at: www.dwd.de; last accessed 18 March 2015.
- Feinglass, J. et al.** (2011). The effects of daily weather on accelerometer-measured physical activity. *Journal of Physical Activity and Health*, 8, 934–943.
- Fetveit, A. and Bjorvatn, B.** (2005). Bright-light treatment reduces actigraphic-measured daytime sleep in nursing home patients with dementia: a pilot study. *The American Journal of Geriatric Psychiatry*, 13, 420–423. doi: [10.1176/appi.ajgp.13.5.420](https://doi.org/10.1176/appi.ajgp.13.5.420)
- Figueiro, M. G.** (2008). A proposed 24 h lighting scheme for older adults. *Lighting Research and Technology*, 40, 153–160.
- Figueiro, M. G., Hamner, R., Higgins, P., Hornick, T. and Rea, M. S.** (2012). Field measurements of light exposures and circadian disruption in two populations of older adults. *Journal of Alzheimers Disease*, 31, 711–715. doi: [10.3233/JAD-2012-120484](https://doi.org/10.3233/JAD-2012-120484)
- Finkel, S.** (2000). Introduction to behavioural and psychological symptoms of dementia (BPSD). *International Journal of Geriatric Psychiatry*, 15, S2–S4. doi: [10.1002/\(SICI\)1099-1166\(200004\)15:1+<S2::AID-GPS159>3.0.CO;2-3 \[pii\]](https://doi.org/10.1002/(SICI)1099-1166(200004)15:1+<S2::AID-GPS159>3.0.CO;2-3 [pii])
- Foster, R. G.** (2005). Neurobiology: bright blue times. *Nature*, 433, 698–699. doi: [10.1038/433698a](https://doi.org/10.1038/433698a)
- Friborg, O., Bjorvatn, B., Amponsah, B. and Pallesen, S.** (2012). Associations between seasonal variations in day length (photoperiod), sleep timing, sleep quality and mood: a comparison between Ghana (5 degrees) and Norway (69 degrees). *Journal of Sleep Research*, 21, 176–184. doi: [10.1111/j.1365-2869.2011.00982.x](https://doi.org/10.1111/j.1365-2869.2011.00982.x)
- Gerding, G.** (2015). Sonnenaufgang-Sonnenuntergang. Available at: www.sonnenaufgang-sonnenuntergang.de; last accessed 18 March 2015.
- Goodman, A., Paskins, J. and Mackett, R.** (2012). Day length and weather effects on children's physical activity and participation in play, sports, and active travel. *Journal of Physical Activity and Health*, 9, 1105–1116.
- Harper, D. G. et al.** (2005). Disturbance of endogenous circadian rhythm in aging and Alzheimer disease. *The American Journal of Geriatric Psychiatry*, 13, 359–368. doi: [10.1176/appi.ajgp.13.5.359](https://doi.org/10.1176/appi.ajgp.13.5.359)
- Klenk, J., Buchele, G., Rapp, K., Franke, S. and Peter, R.** (2011). Walking on sunshine: effect of weather conditions on physical activity in older people. *Journal of Epidemiology and Community Health*, 66, 474–476. doi: [10.1136/jech.2010.128090](https://doi.org/10.1136/jech.2010.128090)
- Laakso, M. L., Porkka-Heiskanen, T., Alila, A., Stenberg, D. and Johansson, G.** (1994). Twenty-four-hour rhythms in relation to the natural photoperiod: a field study in humans. *Journal of Biological Rhythms*, 9, 283–293.
- Liu, R. Y. et al.** (2000). Decreased vasopressin gene expression in the biological clock of Alzheimer disease patients with and without depression. *Journal of Neuropathology and Experimental Neurology*, 59, 314–322.
- Lockley, S. W.** (2009). Circadian rhythms: influence of light in humans. In R. S. Larry (ed.), *Encyclopedia of Neuroscience* (pp 971–988). Oxford, UK: Academic Press.
- Lovell, B. B., Ancoli-Israel, S. and Gevirtz, R.** (1995). Effect of bright light treatment on agitated behavior in institutionalized elderly subjects. *Psychiatry Research*, 57, 7–12. doi: [016517819502550G](https://doi.org/10.1651/7819502550G)
- Lyketsos, C. G., Lindell Veiel, L., Baker, A. and Steele, C.** (1999). A randomized, controlled trial of bright light therapy for agitated behaviors in dementia patients residing in long-term care. *International Journal of Geriatric Psychiatry*, 14, 520–525. doi: [10.1002/\(SICI\)1099-1166\(199907\)14:7<520::AID-GPS983>3.0.CO;2-M](https://doi.org/10.1002/(SICI)1099-1166(199907)14:7<520::AID-GPS983>3.0.CO;2-M)
- Mahlberg, R. et al.** (2008). Pineal calcification in Alzheimer's disease: an in vivo study using computed tomography. *Neurobiology of Aging*, 29, 203–209.
- Mahlberg, R., Kunz, D., Sutej, I., Kuhl, K. P. and Hellweg, R.** (2004). Melatonin treatment of day-night rhythm disturbances and sundowning in Alzheimer disease: an open-label pilot study using actigraphy. *Journal of Clinical Psychopharmacology*, 24, 456–459.
- Mishima, K. et al.** (1994). Morning bright light therapy for sleep and behavior disorders in elderly patients with dementia. *Acta Psychiatrica Scandinavica*, 89, 1–7.
- Mishima, K., Hishikawa, Y. and Okawa, M.** (1998). Randomized, dim light controlled, crossover test of morning bright light therapy for rest-activity rhythm disorders in patients with vascular dementia and dementia of Alzheimer's type. *Chronobiology International*, 15, 647–654.
- Mishima, K., Okawa, M., Shimizu, T. and Hishikawa, Y.** (2001). Diminished melatonin secretion in the elderly caused by insufficient environmental illumination. *Journal of Clinical Endocrinology and Metabolism*, 86, 129–134. doi: [10.1210/jcem.86.1.7097](https://doi.org/10.1210/jcem.86.1.7097)
- Münch, M. et al.** (2017) Bright light delights: effects of daily light exposure on emotions, rest-activity cycles, sleep and melatonin secretion in severely demented patients. *Current Alzheimer Research*. doi: [10.2174/1567205014666170523092858](https://doi.org/10.2174/1567205014666170523092858)
- Obayashi, K. et al.** (2012). Positive effect of daylight exposure on nocturnal urinary melatonin excretion in the elderly: a cross-sectional analysis of the HEIJO-KYO study. *Journal of Clinical Endocrinology and Metabolism*, 97, 4166–4173. doi: [10.1210/jc.2012-1873](https://doi.org/10.1210/jc.2012-1873)
- Oxford Dictionaries** (2016). Available at: <http://www.oxforddictionaries.com>; last accessed 15 March 2016.
- Park, D. H., Kripke, D. F. and Cole, R. J.** (2007). More prominent reactivity in mood than activity and sleep induced by differential light exposure due to seasonal and local differences. *Chronobiology International*, 24, 905–920. doi: [10.1080/07420520701669677](https://doi.org/10.1080/07420520701669677)
- Riemersma-van der Lek, R. F., Swaab, D. F., Twisk, J., Hol, E. M., Hoogendijk, W. J. and Van Someren, E. J.** (2008). Effect of bright light and melatonin on cognitive and noncognitive function in elderly residents of group care facilities: a randomized controlled trial. *JAMA*, 299, 2642–2655.
- Rimmer, D. W. et al.** (2000). Dynamic resetting of the human circadian pacemaker by intermittent bright light.

- American Journal of Physiology: Regular, Integrative and Comparative Physiology*, 279, R1574–R1579.
- Sanford, J. R.** (1975). Tolerance of debility in elderly dependants by supporters at home: its significance for hospital practice. *British Medical Journal*, 3, 471–473.
- Satlin, A. et al.** (1991). Circadian locomotor activity rhythms in Alzheimer's disease. *Neuropsychopharmacology*, 5, 115–126.
- Satlin, A., Volicer, L., Ross, V., Herz, L. and Campbell, S.** (1992). Bright light treatment of behavioral and sleep disturbances in patients with Alzheimer's disease. *American Journal of Psychiatry*, 149, 1028–1032.
- Seitz, D. P. et al.** (2011). Antidepressants for agitation and psychosis in dementia. *Cochrane Database Systematic Reviews*, doi: [10.1002/14651858.CD008191.pub2](https://doi.org/10.1002/14651858.CD008191.pub2)
- Sellix, M. T. et al.** Aging differentially affects the re-entrainment response of central and peripheral circadian oscillators. *Journal of Neuroscience*, 32, 16193–16202. doi: [10.1523/JNEUROSCI.3559-12.2012](https://doi.org/10.1523/JNEUROSCI.3559-12.2012)
- Shochat, T., Martin, J., Marler, M. and Ancoli-Israel, S.** (2000). Illumination levels in nursing home patients: effects on sleep and activity rhythms. *Journal of Sleep Research*, 9, 373–379.
- Skjerve, A. et al.** (2004). Improvement in behavioral symptoms and advance of activity acrophase after short-term bright light treatment in severe dementia. *Psychiatry and Clinical Neuroscience*, 58, 343–347. doi: [10.1111/j.1440-1819.2004.01265.x](https://doi.org/10.1111/j.1440-1819.2004.01265.x)
- Sokolove, P. G. and Bushell, W. N.** (1978). The chi square periodogram: its utility for analysis of circadian rhythms. *Journal of Theoretical Biology*, 72, 131–160.
- Spira, A. P. et al.** (2012). Objectively measured sleep quality and nursing home placement in older women. *Journal of the American Geriatric Society*, 60(7) 1237–1243.
- Turner, P. L., Van Someren, E. J. and Mainster, M. A.** (2010). The role of environmental light in sleep and health: effects of ocular aging and cataract surgery. *Sleep Medicine Reviews*, 14, 269–280.
- Van Gool, W. A. and Mirmiran, M.** (1986). Aging and circadian rhythms. *Progressive Brain Research*, 70, 255–277.
- Van Someren, E. J. et al.** (1996). Circadian rest-activity rhythm disturbances in Alzheimer's disease. *Biological Psychiatry*, 40, 259–270. doi: [0006-3223\(95\)00370-3](https://doi.org/0006-3223(95)00370-3)
- Van Someren, E. J. et al.** (1999). Bright light therapy: improved sensitivity to its effects on rest-activity rhythms in Alzheimer patients by application of nonparametric methods. *Chronobiology International*, 16, 505–518.
- Van Someren, E. J., Kessler, A., Mirmiran, M. and Swaab, D. F.** (1997a). Indirect bright light improves circadian rest-activity rhythm disturbances in demented patients. *Biological Psychiatry*, 41, 955–963. doi: [S0006-3223\(97\)89928-3](https://doi.org/S0006-3223(97)89928-3)
- Van Someren, E. J., Lijzenga, C., Mirmiran, M. and Swaab, D. F.** (1997b). Long-term fitness training improves the circadian rest-activity rhythm in healthy elderly males. *Journal of Biological Rhythms*, 12, 146–156.
- Van Someren, E. J., Mirmiran, M. and Swaab, D. F.** (1993). Non-pharmacological treatment of sleep and wake disturbances in aging and Alzheimer's disease: chronobiological perspectives. *Behavioural Brain Research*, 57, 235–253.
- Wahnschaffe, A. et al.** (2017). Implementation of dynamic lighting in a nursing home: impact on agitation but not on rest-activity patterns. *Current Alzheimer Research*. doi: [10.2174/1567205014666170608092411](https://doi.org/10.2174/1567205014666170608092411).
- Witting, W., Kwa, I. H., Eikelenboom, P., Mirmiran, M. and Swaab, D. F.** (1990). Alterations in the circadian rest-activity rhythm in aging and Alzheimer's disease. *Biological Psychiatry*, 27, 563–572. doi: [0006-3223\(90\)90523-5](https://doi.org/0006-3223(90)90523-5)
- Wu, Y. H., Zhou, J. N., Van Heerikhuizen, J., Jockers, R. and Swaab, D. F.** (2007). Decreased MT1 melatonin receptor expression in the suprachiasmatic nucleus in aging and Alzheimer's disease. *Neurobiology of Aging*, 28, 1239–1247. doi: [10.1016/j.neurobiolaging.2006.06.002](https://doi.org/10.1016/j.neurobiolaging.2006.06.002)