

University of Groningen

**Corrigendum to: "Using MCMC chain outputs to efficiently estimate Bayes factors" (vol 55, pg 367, 2011)**

Morey, Richard D.; Rouder, Jeffrey N.; Pratte, Michael S.; Speckman, Paul L.

*Published in:*  
Journal of Mathematical Psychology

*DOI:*  
[10.1016/j.jmp.2017.03.001](https://doi.org/10.1016/j.jmp.2017.03.001)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2017

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Morey, R. D., Rouder, J. N., Pratte, M. S., & Speckman, P. L. (2017). Corrigendum to: "Using MCMC chain outputs to efficiently estimate Bayes factors" (vol 55, pg 367, 2011). *Journal of Mathematical Psychology*, 79, 131-131. <https://doi.org/10.1016/j.jmp.2017.03.001>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Contents lists available at ScienceDirect

## Journal of Mathematical Psychology

journal homepage: [www.elsevier.com/locate/jmp](http://www.elsevier.com/locate/jmp)

## Corrigendum

## Corrigendum to “Using MCMC chain outputs to efficiently estimate Bayes factors” [J. Math. Psychol. 55 (2011) 367–378]

Richard D. Morey<sup>a,\*</sup>, Jeffrey N. Rouder<sup>b</sup>, Michael S. Pratte<sup>b</sup>, Paul L. Speckman<sup>c</sup><sup>a</sup> University of Groningen, Psychometrics and Statistics, Grote Kruisstraat 2/1, 9712NS Groningen, The Netherlands<sup>b</sup> University of Missouri, Department of Psychological Sciences, 210 McAlester Hall, Columbia, MO 65201, United States<sup>c</sup> University of Missouri, Department of Statistics, 146 Middlebush Hall, Columbia, MO 65201, United States

## ARTICLE INFO

## Article history:

Available online 19 April 2017

The authors would like to publicise an error on page 372, section 4.1 of the article “Using MCMC chain outputs to efficiently estimate Bayes factors”. There is a mistake in the definition of the inverse gamma density function  $p(g | r)$ : every appearance of  $r$  should be replaced by  $r/\sqrt{2}$ . The correct density function is

$$p(g | r) = r(2\pi)^{-\frac{1}{2}} g^{-\frac{3}{2}} \exp\left\{-\frac{r^2}{2g}\right\}, \quad g, r > 0.$$

To the best of the authors' knowledge, this oversight did not affect any other aspect of the manuscript. The authors thank Jorge Tendeiro and Don van Ravenzwaaij for calling their attention to this error.

DOI of original article: <http://dx.doi.org/10.1016/j.jmp.2011.06.004>.

\* Correspondence to: School of Psychology, Cardiff University, United Kingdom.

E-mail addresses: [moreyr@cardiff.ac.uk](mailto:moreyr@cardiff.ac.uk), [r.d.morey@rug.nl](mailto:r.d.morey@rug.nl) (R.D. Morey).