

University of Groningen

Factors associated with workarounds in Bar-Code-assisted medication administration in hospitals

BCMA Study Group

Published in:
Journal of Clinical Nursing

DOI:
[10.1111/jocn.15217](https://doi.org/10.1111/jocn.15217)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

BCMA Study Group (2020). Factors associated with workarounds in Bar-Code-assisted medication administration in hospitals. *Journal of Clinical Nursing*, 29(13-14), 2239-2250. <https://doi.org/10.1111/jocn.15217>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Factors associated with workarounds in barcode-assisted medication administration in hospitals

Willem van der Veen PhD, Researcher¹ | Katja Taxis PharmD, PhD, Professor¹ |
Hans Wouters PhD² | Hester Vermeulen PhD, RN, Professor³ | David W. Bates MD,
PhD, Professor⁴ | Patricia M. L. A. van den Bemt PharmD, PhD, Professor⁵ |
the BCMA Study Group

¹Unit Pharmacotherapy, Epidemiology & Economics, Faculty of Science and Engineering, University of Groningen, Groningen, The Netherlands

²General Practitioners Research Institute, University of Groningen, Groningen, The Netherlands

³Department of IQ Healthcare, Scientific Center for Quality of Healthcare, Radboud Institute of Health Sciences, Nijmegen, The Netherlands

⁴Harvard Medical School and Brigham and Women's Hospital, Boston, MA, USA

⁵Department Clinical Pharmacy and Pharmacology, University Medical Center Groningen, Groningen, The Netherlands

Correspondence

Willem van der Veen,
Unit Pharmacotherapy, Epidemiology & Economics, Faculty of Science and Engineering, University of Groningen, Antonius Deusinglaan 1 9713 AV Groningen, The Netherlands.
Email: willem.van.der.veen@rug.nl

Abstract

Aims and objectives: To identify that workarounds (defined as “informal temporary practices for handling exceptions to normal procedures or workflow”) by nurses using information technology potentially compromise medication safety. Therefore, we aimed to identify potential risk factors associated with workarounds performed by nurses in Barcode-assisted Medication Administration in hospitals.

Background: Medication errors occur during the prescribing, distribution and administration of medication. Errors could harm patients and be a tragedy for both nurses and medical doctors involved. Interventions to prevent errors have been developed, including those based on information technology. To cope with shortcomings in information technology-based interventions as Barcode-assisted Medication Administration, nurses perform workarounds. Identification of workarounds in information technology is essential to implement better-designed software and processes which fit the nurse workflow.

Design: We used the data from our previous prospective observational study, performed in four general hospitals in the Netherlands using Barcode techniques, to administer medication to inpatients.

Methods: Data were collected from 2014–2016. The disguised observation was used to gather information on potential risk factors and workarounds. The outcome was a medication administration with one or more workarounds. Logistic mixed models were used to determine the association between potential risk factors and workarounds. The STROBE checklist was used for reporting our data.

Results: We included 5,793 medication administrations among 1,230 patients given by 272 nurses. In 3,633 (62.7%) of the administrations, one or more workarounds were observed. In the multivariate analysis, factors significantly associated with workarounds were the medication round at 02 p.m.–06 p.m. (adjusted odds ratio [OR]: 1.60,

BCMA Study Group members are listed in Appendix.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Journal of Clinical Nursing* published by John Wiley & Sons Ltd

95% CI: 1.05–2.45) and 06 p.m.–10 p.m. (adjusted OR: 3.60, 95% CI: 2.11–6.14) versus the morning shift 06 a.m.–10 a.m., the workdays Monday (adjusted OR: 2.59, 95% CI: 1.51–4.44), Wednesday (adjusted OR: 1.92, 95% CI: 1.2–3.07) and Saturday (adjusted OR: 2.24, 95% CI: 1.31–3.84) versus Sunday, the route of medication, nonoral (adjusted OR: 1.28, 95% CI: 1.05–1.57) versus the oral route of drug administration, the Anatomic Therapeutic Chemical classification-coded medication “other” (consisting of the irregularly used Anatomic Therapeutic Chemical classes [D, G, H, L, P, V, Y, Z]) (adjusted OR: 1.49, 95% CI: 1.05–2.11) versus Anatomic Therapeutic Chemical class A (alimentary tract and metabolism), and the patient–nurse ratio $\geq 6-1$ (adjusted OR: 5.61, 95% CI: 2.9–10.83) versus $\leq 5-1$.

Conclusions: We identified several potential risk factors associated with workarounds performed by nurses that could be used to target future improvement efforts in Barcode-assisted Medication Administration.

Relevance to clinical practice: Nurses administering medication in hospitals using Barcode-assisted Medication Administration frequently perform workarounds, which may compromise medication safety. In particular, nurse workload and the patient–nurse ratio could be the focus for improvement measures as these are the most clearly modifiable factors identified in this study.

KEYWORDS

Barcode-assisted Medication Administration, health information technology, medication safety, nurse-performed-workarounds, nurse workload, quality of care

1 | INTRODUCTION

Nurses are in the front-line caring for hospital inpatients, which involves a multitude of complex tasks, including the administration of prescribed medication to patients. Nurses perform the last step in the multifaceted process performed by several health professionals of prescribing and administering medication, capable of protecting patients from medication errors. Therefore, they have a crucial role in medication safety.

Electronic Barcode-assisted Medication Administration (BCMA) systems have been developed to help nurses to reduce medication administration errors (Hutton, Ding, & Wellman, 2017; Poon et al., 2010). The BCMA system checks the information obtained by scanning both the barcode on the medication package and the bar-coded patient wristband against the information provided by the hospital's electronic prescribing system. An electronic alert is given in case of a mismatch. In some BCMA systems, nurses also have a personal barcode so that the individuals administering the medication are registered automatically. Several studies have shown a substantial decrease in medication administration errors following the implementation of BCMA systems in hospitals (Berdot et al., 2016; Hassink, Essenberg, Roukema, & van den Bemt, 2013; Helmons, Wargel, & Daniels, 2009; Maaskant et al., 2015; Poon et al., 2010).

What does this paper contribute to the wider global clinical community

- Nurses administering medication using Barcode-assisted Medication Administration frequently perform workarounds.
- Potential risk factor associated with these workarounds was the nurse workload.
- Especially, the nurse workload found in infrequently performed nursing procedures and the patient–nurse ratio could be the focus for improvement measures in medication administration using Barcode-assisted Medication Administration.

However, Information Technology (IT) based interventions as BCMA are not always used as instructed and required, and nurses perform workarounds. Workarounds (defined as “informal temporary practices for handling exceptions to normal procedures or workflow”) by nurses using IT potentially compromise medication safety (van der Veen, van den Bemt, Wouters, et al., 2017). Until now, the factors that force nurses to carry out workarounds were unclear.

1.1 | Aim

This study aimed to identify potential risk factors associated with nurse workarounds in BCMA in order to explore why nurses perform workarounds.

2 | BACKGROUND

Medication errors occur during the prescribing, distribution and administration of medication (Krahenbuhl-Melcher et al., 2007; Lisby, Nielsen, & Mainz, 2005). Errors may not only harm patients but could also be a tragedy for healthcare workers as nurses. Several interventions to prevent errors have been developed, including those based on information technology (IT). IT-based interventions are most promising because they have the potential to structure and standardise processes as prescribing and administration of medication (Bates, 2000; Kaushal & Bates, 2002; Seidling & Bates, 2016). In practice, IT-based interventions in health care, such as BCMA, are not always used as instructed and required or fit the daily workflow (Cheung et al., 2014; van der Veen, van den Bemt, Wouters, et al., 2017). Nurses adopt so-called workarounds (Ash, Berg, & Coiera, 2004; Koppel, Wetterneck, Telles, & Karsh, 2008a; Rack, Dudjak, & Wolf, 2012) to cope with IT-based interventions as BCMA in daily practice. Workarounds were defined as "informal temporary practices for handling exceptions to normal workflow" (Kobayashi, Fussell, Xiao, & Seagull, 2005). While performing workarounds, the nurse workflow can be changed, either once, temporarily or even over an extended period, but identity, purpose or construction of the system remains unchanged (Alter, 2014). End users of BCMA such as the nursing staff occasionally practice workarounds to deal with perceived issues, which may relate to lack of confidence in technology, the time that using this technology takes or other issues relating to hardware, programming, screen design, user knowledge or communication problems (Debono et al., 2013; Koppel, 2014; Patterson, 2018). Examples of workarounds performed by nurses using BCMA are as follows: not scanning at all, scanning patient's barcodes remotely (i.e. not the actual wristbands of patients), ignoring signals or alerts or scanning of medication for multiple patients at the same time. Identification of workarounds in IT is essential to implement better-designed software and processes which fit the nurse workflow. As seen in qualitative research (Cresswell, Mozaffar, Lee, Williams, & Sheikh, 2016), workarounds can improve efficiency, but may also create new risks and compromise the safety and effectiveness of patient care. A lack of coherence between the wishes and expectations of healthcare providers, patients and technological capabilities could be the cause of this. Suggested is that, to avoid this, healthcare workers such as nurses, pharmacists and medical doctors should already be involved in the implementation phase of new IT-based systems (Blijleven, Koelemeijer, Wetzels, & Jaspers, 2017; Koppel, Smith, Blythe, & Kothari, 2015; van der Veen, de Gier, van der Schaaf, Taxis, & van den Bemt, 2013). In our previous study in four hospitals in the Netherlands, we showed that workarounds

are associated with medication administration errors (van der Veen, van den Bemt, Wouters, et al., 2017). Other research identified circumstantial factors for performing workarounds but focused mainly on the usability of the BCMA system (Debono et al., 2013; Holden, Rivera-Rodriguez, Faye, Scanlon, & Karsh, 2013; Patterson, Rogers, Chapman, & Render, 2006). Our data (van der Veen, van den Bemt, Wouters, et al., 2017) provide the opportunity to study quantitatively (instead of qualitatively) the potential risk factors associated with workarounds in IT-based intervention BCMA. This may be considered as a first step to develop interventions intended to reduce the frequency of nurse workarounds in the use of BCMA in hospitals.

3 | DESIGN

We used the data (van der Veen, van den Bemt, Wouters, et al., 2017) of our multicentre prospective observational study conducted in adult patients (aged 18 years and older) admitted to four hospitals in the Netherlands using BCMA in the medication administration process (at the time of planning our study, eight hospitals in the Netherlands were using BCMA-based systems). The research project was started in 2014, and enrolment was completed at the end of 2016. Both a detailed version of the study protocol (van der Veen, van den Bemt, Bijlsma, de Gier, & Taxis, 2017) and the analysis of the association between workarounds and medication errors (van der Veen, van den Bemt, Wouters, et al., 2017) have been published before. The study was registered in the "Dutch trial register" with trial ID NTR4355. Data were anonymised following Dutch privacy legislation (van der Veen, van den Bemt, Wouters, et al., 2017).

3.1 | Ethics

The regional medical ethics committee (in Dutch: "Regionale Toetsingscommissie Patientgebonden Onderzoek RTPO") approved the study protocol on 22 May 2014.

4 | METHODS

The STROBE checklist for reporting cohort, case-control and cross-sectional studies was used (File S1).

4.1 | Setting

All four included hospitals operated electronic prescribing of medication and BCMA, each using different software for both the prescribing and the BCMA part. After scanning the barcode of both the patient and the medication, the BCMA system checked whether the patient and the medication matched the physician's prescription.

To facilitate the scanning of medication, pharmacy technicians dispensed unit-dose barcode-labelled medication. Medication rounds

were scheduled on the following times: 6 a.m.–10 a.m., 10 a.m.–2 p.m., 6 p.m.–8 p.m. and 8 p.m.–10 p.m. Per the medication round, one nurse was responsible for medication administration. Nurse trainees were supervised by registered nurses (van der Veen, van den Bemt, Wouters, et al., 2017).

4.2 | Participants

Patients on participating nursing wards of four Dutch hospitals using BCMA to administer medication were included; patients aged 17 years and younger were excluded (van der Veen, van den Bemt, Bijlsma, et al., 2017). These were four out of eight hospitals using BCMA in the Netherlands at the time of planning the study. We included wards from the following areas: cardiology, pulmonary diseases, geriatrics, internal medicine, neurology, surgery and orthopaedics. The observers (three undergraduate students from the schools of pharmacy of the universities of Groningen and Utrecht in the Netherlands, who were all well trained in the technique of observation and who all had to pass an examination in order to be appointed as an observer) accompanied nurses working on the selected wards during the selected drug rounds. The observed nurses were aware of the fact that they were being observed, but not about the precise nature of the data, which were collected (disguised observation).

All the nurses agreed to be observed. Based on Dutch regulation, nurses in training were not responsible for nursing actions but worked under the supervision of a registered nurse who has the final responsibility for the actions of the nurse in training. For this reason, we did not distinguish between a registered nurse and a training nurse in the final analysis.

4.3 | Data collection

The disguised observation method (Dean & Barber, 1999, 2001; Smeulders, Hoekstra, van Dijk, Overkamp, & Vermeulen, 2013) was used to collect data on potential risk factors and workarounds (van der Veen, van den Bemt, Bijlsma, et al., 2017). In practice, each observer accompanied the nurse who administered the medication using BCMA and observed the administration of each medication to the patient. The following observation schedule was followed: at least three rounds were observed each day of the week, with a weekly minimum of 21 medication administration rounds. Details of the drug administration to the patient were documented using a case report form (van der Veen, van den Bemt, Bijlsma, et al., 2017). In case the observer noticed a potentially dangerous error, he or she intervened for ethical reasons, while retaining these observations in the data set. Incomplete observations were excluded.

4.4 | Definition and classification

We defined workarounds using the definition of Kobayashi as “informal temporary practices for handling exceptions to the normal

workflow” (Kobayashi et al., 2005). Workarounds were classified as described earlier (van der Veen, van den Bemt, Bijlsma, et al., 2017) using a self-developed classification system and observation form based on the research of Koppel (Koppel et al., 2008). Workarounds were classified as procedural-based, related to patient identification, the scanning process, the computers or scanner alert signals, and other workflow procedures, or nurse-work-related. To determine whether a workaround took place, the observers compared their observation records after each medication administration round to the hospital or ward procedures and local guidelines on the BCMA process (van der Veen, van den Bemt, Bijlsma, et al., 2017).

4.5 | Outcome measure and potential risk factors

The proportion of medication administrations with one or more workarounds was the main outcome measure. Potential risk factors associated with workarounds were selected based on the research of Van den Bemt (van den Bemt, 2006; van den Bemt, Idzinga, Robertz, Kormelink, & Pels, 2009), Schimmel (Schimmel, Becker, van den Bout, Taxis, & van den Bemt, 2011), Driscoll (Driscoll et al., 2018), Aiken (Aiken et al., 2014), Spetz (Spetz, Donaldson, Aydin, & Brown, 2008), Donaldson and Shapiro (Donaldson & Shapiro, 2010) and Wise (Wise, 2016). The following factors were included to analyse their association with workarounds: general characteristics (ward type, time of medication round, day of the week, patient age and gender), medication characteristics (percentage barcoded medication, route of administration, i.e. oral vs. nonoral), the first level of the Anatomic Therapeutic Chemical (ATC) classification code system (Anonymous, 2012, 2017) (the ATC code system is an international drug classification scheme, aimed to categorise the active ingredients of drugs according to the organ or system on which they act and their therapeutic, pharmacological and chemical properties, developed by the World Health Organization [WHO] (Table 2)), nurse characteristics (work experience [≤ 24 months, > 24 months]), nurse workload characteristics during the medication shift (i.e. the number of medications per patient [1, 2, ≥ 3]) and the patient–nurse ratio (i.e. the number of beds occupied by patients divided by the number of registered nurses on that ward during one shift).

4.6 | Statistical analysis

The association between potential risk factors and workarounds was analysed using logistic mixed models using a similar statistical approach as in our previous study (van der Veen, van den Bemt, Wouters, et al., 2017). To take into account the potential dependency of observations (i.e. more than one observation was made for each nurse), a random intercept at the ward and the nurse level was included in the models. Owing to observed multicollinearity between the training of the nurse (student nurse vs. registered nurse) and the work experience (≤ 24 months vs. > 24 months) of the nurse, we only included work experience as a variable in the model. The type of hospital (general vs. training hospital) corresponded

TABLE 1 Characteristics of study hospitals ($N = 4$) and nurses ($N = 272$)

Characteristics	Category	N	%
Hospitals ($n = 4$)			
Location	Rural area	2	50
	Urban area	2	50
Number of beds	200–400	1	25
	401–600	2	50
	601–800	1	25
Hospital type	General	3	75
	Teaching	1	25
Hospital BCMA experience	2–4 years	1	25
	4–6 years	2	50
	6–8 years	1	25
Nurses ($n = 272$)			
Gender	Male	24	8.8
	Female	248	91.2
Education level	Student nurse	33	12.1
	Registered nurse	236	86.7
	Unknown ^a	3	1.2
Experience as a registered nurse	≤24 months	36	15.3
	>24 months	198	83.9
	Unknown	2	0.8
Registered nurse BCMA experience	≤6 months	28	11.9
	>6 months	206	87.3
	Unknown	2	0.8
Nursing ward	Cardiology	39	14.3
	Pulmonary diseases	29	10.7
	Geriatrics	15	5.5
	Internal medicine	53	19.5
	Neurological diseases	35	12.9
	Surgical diseases	60	22.1
	Orthopaedics	30	11.0
	Other type of nursing ward	11	4.0
	Mean patient–nurse ratio on wards during medication administration shifts	Hospital 1	5
Hospital 2		4	
Hospital 3		7	
Hospital 4		6	

^aThese 3 nurses were responsible for 48 medication administrations, 25 without a workaround and 23 with a workaround.

with the percentage of medication supplied with a barcode (<99% vs. ≥99%). Therefore, we did not include the hospital type in the analysis. First, univariate analyses were performed in which we examined the factors individually. Subsequently, multivariate analysis was performed, including the percentage of barcoded medication, type of nursing department, the day of the week, time of the medication round, ATC classes, the number of drugs per round, route

TABLE 2 Anatomic Therapeutic Chemical (ATC) classification system

ATC Code	Drugs related to organ system or use
A	Alimentary tract and metabolism
B	Blood and blood-forming organs
C	Cardiovascular system
D	Dermatological medication
G	Genitourinary system and sex hormones
H	Systemic hormonal preparations, excluding sex hormones and insulins
J	Anti-infective for systemic use
L	Antineoplastic and immunomodulating agents
M	Muscular–skeletal system
N	Nervous system
P	Antiparasitic products, insecticides and repellents
R	Respiratory system
S	Sensory organs, eye, nose, ear
V	Various drugs
Y	Not supplied
Z	Not relevant

of administration and the patient–nurse ratio as the independent variables. Mixed model analyses were carried out using MLWIN version 6.3. All other analyses were carried out using SPSS version 23.0, similar to our protocol and our previous study (van der Veen, van den Bemt, Bijlsma, et al., 2017; van der Veen, van den Bemt, Wouters, et al., 2017).

5 | RESULTS

We observed 6,021 medication administrations in patients admitted to four hospitals, of which 228 (3.8%) were excluded because they were incomplete. The included 5,793 medication administrations were given to 1,230 inpatients by 272 nurses. In 3,633, medication administrations (62.7%), one or more workarounds, were observed (van der Veen, van den Bemt, Wouters, et al., 2017). The characteristics of the study hospitals and the nurses are presented in Table 1.

Procedural workarounds (as not scanning at all) were most common ($n = 1,307$, 36%). Other workarounds observed were patient scanning-related (as no barcode wristband on the patient; $n = 1,017$, 28%) and medication scanning-related (including scanning before actual administration of medication, scanning medication for more than one patient at a time, and ignoring computer or scanner alerts; $n = 400$, 11%). The observers did not have to intervene during the observation period.

In the multivariate analysis, factors significantly associated with workarounds were the medication round at 02 p.m.–06 p.m. (adjusted OR: 1.60, 95% CI: 1.05–2.45) and 06 p.m.–10 p.m. (adjusted OR: 3.60, 95% CI: 2.11–6.14) versus the morning shift 06 a.m.–10

TABLE 3 Unadjusted and adjusted associations between factors and workarounds in barcode-assisted medication administration

Factors	Factor	No WA (N)	%	WA (N)	%	Unadjusted OR ^b	95% CI	Adjusted OR ^c	95% CI	
General factors	Ward type	Cardiology	341	5.89	682	11.77	Ref. ^a	—	Ref. ^a	—
		Pulmonary diseases	380	6.56	278	4.80	0.30	0.08–1.13	0.29	0.07–1.29
		Geriatrics	159	2.74	122	1.93	0.34	0.07–1.58	0.82	0.14–4.84
	Internal medicine	406	7.01	611	10.55	0.83	0.30–2.31	1.31	0.40–4.25	
		Neurological diseases	219	3.78	425	7.34	0.96	0.28–3.30	0.91	0.23–3.60
	Surgical diseases	406	7.01	1,008	17.40	0.90	0.32–2.51	1.45	0.45–4.70	
		Orthopaedics	153	2.64	447	7.72	1.08	0.29–4.06	0.77	0.18–3.33
	Other type of nursing ward, for example day care	96	1.66	60	1.04	0.39	0.05–2.95	0.90	0.09–8.7	
		06 a.m.–10 a.m.	1509	26.05	1775	30.64	Ref. ^a	—	Ref. ^a	—
	Time of medication run	10 a.m.–02 p.m.	98	1.69	160	2.76	1.57	1.01–2.42	1.57	0.93–2.66
		02 p.m.–06 p.m.	472	8.15	472	8.15	1.57	1.11–2.22	1.60	1.05–2.45
	Day of the week	06 p.m.–10 p.m.	81	1.40	1,226	21.16	8.79	6.39–12.1	3.60	2.11–6.14
		Sunday	159	2.74	374	6.46	Ref. ^a	—	Ref. ^a	—
		Monday	228	3.94	504	8.07	1.91	1.24–2.96	2.59	1.51–4.44
		Tuesday	360	6.21	572	9.87	1.43	0.94–2.17	1.34	0.79–2.26
Wednesday		377	6.51	681	11.76	1.19	0.80–1.75	1.92	1.20–3.07	
Thursday		405	6.99	723	12.48	1.57	1.02–2.39	1.58	0.95–2.62	
Friday		290	5.01	331	5.71	1.20	0.75–1.91	1.66	0.92–3.00	
Patient age ^d	Saturday	305	5.26	374	6.46	1.57	1.02–2.42	2.24	1.31–3.84	
	<74 years of age	1,072	18.51	1855	32.02	Ref. ^a	—	Ref. ^a	—	
Patient gender	≥74 years of age	1,088	18.78	1778	30.69	1.02	0.90–1.15	0.95	0.82–1.1	
	Women	1,037	17.90	1,657	28.60	Ref. ^a	—	Ref. ^a	—	
Medication factors	Men	1,123	19.39	1,976	34.11	0.89	0.79–1.01	0.89	0.78–1.03	
	% barcoded medication	713	12.31	815	14.07	Ref. ^a	—	Ref. ^a	—	
Route of administration	<99%	1,447	24.98	2,818	48.64	0.65	0.28–1.53	1.22	0.47–3.15	
	Oral medication	1,831	31.61	2,951	50.94	Ref. ^a	—	Ref. ^a	—	
	Nonoral route ^e	329	5.68	682	11.77	1.28	1.10–1.49	1.28	1.05–1.57	

(Continues)

TABLE 3 (Continued)

Factors	Factor	No WA (N)	%	WA (N)	%	Unadjusted OR ^b	95% CI	Adjusted OR ^c	95% CI	
ATC ^f code (first level)	ATC A	556	9.60	757	13.07	Ref. ^a	—	Ref. ^a	—	
	ATC B	182	3.14	381	6.58	1.14	0.92–1.42	0.99	0.76–1.28	
	ATC C	479	8.27	620	10.70	0.96	0.81–1.13	0.96	0.79–1.15	
	ATC J	67	1.16	187	3.23	1.59	1.17–2.16	1.42	0.99–2.04	
	ATC M	68	1.17	104	1.80	1.13	0.80–1.59	1.12	0.77–1.63	
	ATC N	530	9.15	1,095	18.90	1.16	0.99–1.36	1.01	0.84–1.21	
	ATC R	119	2.05	235	4.06	1.17	0.90–1.51	0.89	0.64–1.23	
	ATC S	86	1.48	110	1.90	0.98	0.71–1.35	0.85	0.58–1.24	
	Infrequently used classes (D, G, H, L, P, V, Y, Z) ^g	73	1.26	144	2.49	1.48	1.07–2.04	1.49	1.05–2.11	
	Nurse factors									
Work experience	<24 months	355 ^{hr}	6.18	541 ⁱ	9.42	Ref. ^a	—	Ref. ^a	—	
	≥24 months	1,780 ^{hr}	30.98	3,069 ⁱ	53.42	1.30	0.86–1.96	1.03	0.6–1.76	
Workload factors										
	Drugs per round per patient	1	52	0.90	187	3.23	Ref. ^a	—	Ref. ^a	—
		2	130	2.24	367	6.34	0.92	0.65–1.30	0.97	0.63–1.50
	≥3	1978	34.14	3,079	53.15	0.81	0.60–1.10	0.96	0.66–1.41	
	Patient–nurse ratio	≤5 to 1	1755	30.30	1,412	24.37	Ref. ^a	—	Ref. ^a	—
		≥6 to 1	405	6.99	2,221	38.34	12.21	8.14–18.31	5.61	2.90–10.83

^aReference category.

^bOdds ratio.

^cAdjusted for all variables shown in table.

^dAge of 74 is median age.

^eNumbers nonoral routes; inhalation 414, parenteral 240, sublingual 118, eye drops 69, dermal drugs 56, other route 114.

^fAnatomic Therapeutic Chemical (ATC) classification.

^gOther, infrequently used ATC classes D, G, H, L, P, V, Y, Z (in which we observed a total of 217 administrations, range 2–75).

^h25 missing values from 3 nurses.

ⁱ23 missing values from 3 nurses.

a.m., the workdays Monday (adjusted OR: 2.59, 95% CI: 1.51–4.44), Wednesday (adjusted OR: 1.92, 95% CI: 1.2–3.07), Saturday (adjusted OR: 2.24, 95% CI: 1.31–3.84) versus Sunday, the route of medication, nonoral (adjusted OR: 1.28, 95% CI: 1.05–1.57) versus the oral route of drug administration, the Anatomic Therapeutic Chemical (ATC) classification-coded (Table 2) medication “other” (consisting of infrequently used ATC classes [D, G, H, L, P, V, Y, Z]) (adjusted OR: 1.49, 95% CI: 1.05–2.11) versus ATC class A (alimentary tract and metabolism) and the patient–nurse ratio, $\geq 6-1$ (adjusted OR: 5.61, 95% CI: 2.90–10.83) versus $\leq 5-1$ (Table 3). Factors not significantly associated with workarounds were the ward type, the patient age and gender, the percentage of barcoded medication, the number of medications per patient and the nurse work experience. Observers did not record the level of education of three nurses. They were responsible for 48 administrations (23 with a workaround and 25 of them without a workaround). Those figures did not significantly change the workaround percentage (62.8% workarounds instead of 62.7%).

6 | DISCUSSION

Potential risk factors associated with workarounds were the day of the week, the timing of the medication administration, the route of administration, the administration of medication from irregularly used ATC classes and the patient–nurse ratio. Other factors, such as the percentage of barcoded medication and work experience, were not associated with workarounds. These results can be used to help target efforts to reduce the frequency of workarounds in the future.

Procedures should be reviewed critically to ensure that nonorally administered medication can be administered correctly using the BCMA system. Furthermore, nurses need to be well trained to perform infrequent nursing procedures. The association of the nonoral route of administration with workarounds may have several causes. For example, the dermal application, as well as inhalation, is often left to the patient self-administering this medication. This may enhance the risk of workarounds because nurses may forget to scan such medication. Another example is a parenteral medication that needs handling to make it ready to administer: the original vial with infusion powder may contain a barcode, but the infusion bag with the added drug may not be barcoded. The handling of infrequently used medication (as expressed by the ATC class “other”) may lead to workarounds because of the nurses not being familiar with administering this medication.

A higher patient–nurse ratio was also associated with workarounds. This finding is in line with other studies finding associations between the number of nursing staff and quality of care for hospitalised patients (Ball, Murrells, Rafferty, Morrow, & Griffiths, 2014; Driscoll et al., 2018; Goedhart, van Oostveen, & Vermeulen, 2017; Wise, 2016). Death rates in British hospitals with nurses caring for six or fewer patients were 20% lower than in hospitals with nurses caring for ten or more patients (Griffiths, Ball, Murrells, Jones, & Rafferty, 2016). Little is known about the optimal patient–nurse ratio, and ratios may vary by time of day and patient awareness. In California, the USA, rules require a patient–nurse ratio of one nurse to every five

patients (Donaldson & Shapiro, 2010). In our study, the work pressure may also have led to nurses leaving out time-consuming steps such as scanning patients or medications (van Onzenoort et al., 2008).

Workarounds were associated with the time of the medication round and particular days. Workarounds seem to be more likely on busy weekdays versus the relatively quiet Sunday. Also, workarounds were more likely on the rounds scheduled during the afternoon and evening. This may also be due to the busier parts of the day, leading to nurse workarounds to save time. Our findings emphasise the need to review the patient–nurse ratio, work schedules and medication-related workload per day of the week and per shift to ensure the safe use of the system. Nurse managers are responsible for a positive work environment and for planning an adequate balance between patients and available nursing care (van Oostveen, Braaksma, & Vermeulen, 2014).

Interestingly, we found several factors such as work experience of nurses and the percentage of barcoded medications not associated with workarounds. Work experience of nurses has been found to be associated with the quality of performance (Blegen, Vaughn, & Goode, 2001) but obviously did not play a role in how nurses dealt with recently introduced IT-based systems as barcode administration. Efforts to prevent workarounds should be therefore targeted to all nursing staff involved, independent of their work experience. Another reason could be found in our dichotomous analysis of the nurse experience. A further refinement of the work experience would possibly have shed light on this item.

Our finding that the percentage of barcoded medication was a factor not associated with workarounds is noteworthy. In the Netherlands, about 80% of the medication in hospitals is barcoded by the vendor (single-cell medication). Especially, liquids, eye drops and eardrops and ointments are not provided with single-cell barcodes. For reasons of patient safety, a lot of effort is taken by hospitals to apply their barcodes to individual medication without those vendor-supplied barcodes. Since the percentage of barcodes was not associated as a factor in carrying out workarounds by nurses, current efforts seem to be sufficient in this respect.

6.1 | Strengths and limitations

A strength of our study is that we quantitatively assessed the association of potential risk factors with workarounds in a large sample from multiple institutions using a robust method of data collection. The multicentre design of the study enhances the generalisability of our data.

Our study has some limitations, as well. Despite disguised observation being considered as the “golden standard” of data collection in medication administration error studies (Barker, Flynn, & Pepper, 2002; Dean & Barber, 2001; Westbrook & Ampt, 2009), observation bias cannot be excluded: observers may become tired and therefore less precise. We educated the observers intensively and trained them to stay nearby to the nurses managing the medication. Only a limited quantity (228, 3.8%) of observations had to be excluded due to the incompleteness of the observation. Furthermore, the observer may have influenced the nurses, but this phenomenon known

as “Hawthorne effect” is reported to be small (Gale, 2004). Also, observers may have missed some workarounds. An ethical question that can be raised in our research is whether observers have the right to observe persons who are not fully aware of the nature of the data that are collected. However, in addition to the national permission for this study, every participating hospital was informed (both the board of directors and ward management) as well and received copies of the research protocol and the nationwide approval of our research, and no single objection was noted. Other limitations were that all four hospitals had BCMA software systems from different vendors (van der Veen, van den Bemt, Bijlsma, et al., 2017), and we observed only nurses from internal medicine and surgical wards and patients aged 18 years and older. Finally, we based the selection of potential risk factors on literature. Hence, we cannot rule out that we have missed some factors. Exploring nurses' beliefs and attitudes using BCMA may reveal additional user aspects, as has been shown in a study on double-checking procedures (Schwappach, Pfeiffer, & Taxis, 2016). Using the Australian Work Observation Method by Activity Timing (WOMBAT) technique (Westbrook, 2009; Westbrook & Ampt, 2009) may be another way to gain a better understanding of the underlying causes of some of the factors (Westbrook, Duffield, Li, & Creswick, 2011).

6.2 | Further research

Our results suggest that workload may be an important cause of workarounds. One example of a workload reducing intervention could be the introduction of dedicated personnel—such as pharmacy technicians—who are solely responsible for medication administration. Pharmacy technicians are trained to handle medication as the main part of their daily work, in contrast to nurses for whom medication administration is only a part of their daily routine. In addition to this, pharmacy technicians, given the nature of the work in the pharmacy, might be better trained in the use of technology in general. Research from both the USA and the UK shows the feasibility of medication administration to hospitalised patients by pharmacy technicians (Keers, 2017; Pedersen, Schneider, & Scheckelhoff, 2012).

On the other hand, this could be costly, and pharmacy technicians would have less of a sense of the patient and their conditions. Cluster randomised controlled trials (cluster RCTs) (Harris et al., 2018) or interrupted time series (Westbrook, Raban, Lehnbohm, & Li, 2016) are needed to demonstrate the effectiveness of this intervention. Further research should also include qualitative research methods, for example interviewing nurses responsible for medication administration using IT-based interventions as BCMA, exploring further the causes for workarounds.

7 | CONCLUSIONS

Nurses administering medication using BCMA frequently performed workarounds. Potential risk factors associated with these workarounds

were the administration of nonoral drugs, medication from ATC classes that were infrequently given, the day of the week, the time of the medication round and the patient–nurse ratio. Especially, the nurse workload reflected in the day of the week, time of the medication round and patient–nurse ratio could be the focus for improvement measures as these are the most clearly modifiable factors identified in this study.

8 | RELEVANCE TO CLINICAL PRACTICE

Our study has identified several potential risk factors of workarounds, which may compromise medication safety. Especially, the patient–nurse ratio and certain—potential busier—moments of the day and week were associated with performing nurse workarounds. These factors could be the focus for improvement measures aimed at reducing the workload per nurse.

AUTHOR CONTRIBUTION

Willem van der Veen, Patricia M.L.A. van den Bemt and Katja Taxis each contributed equally to the design, the analysis and the interpretation of the data of the study, drafted the manuscript and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. Hans Wouters, Hester Vermeulen and David W. Bates made each equally substantial contributions to the analysis and interpretation of the data of the study and the drafting of the final manuscript. All authors read and approved the final manuscript and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

ETHICAL APPROVAL

The study is approved by the “Regionale Toetsingscommissie Patientgebonden Onderzoek RTPO” (22 May 2014, reference number RTPO 920a). The study was registered in the “Dutch trial register” with trial ID NTR4355.

ORCID


Willem van der Veen  <https://orcid.org/0000-0003-4603-8736>

Katja Taxis  <https://orcid.org/0000-0001-8539-2004>

Hans Wouters  <https://orcid.org/0000-0001-7711-0476>

Hester Vermeulen  <https://orcid.org/0000-0002-0743-2979>

David W. Bates  <https://orcid.org/0000-0001-6268-1540>

Patricia M. L. A. van den Bemt  <https://orcid.org/0000-0003-1418-5520>

REFERENCES

- Aiken, L. H., Sloane, D. M., Bruyneel, L., Van den Heede, K., Griffiths, P., Busse, R., ... Sermeus, W. (2014). Nurse staffing and education and hospital mortality in nine European countries: A retrospective observational study. *Lancet (London, England)*, 383(9931), 1824–1830. [https://doi.org/10.1016/S0140-6736\(13\)62631-8](https://doi.org/10.1016/S0140-6736(13)62631-8)

- Alter, S. (2014). Theory of workarounds. *Communications of the Association for Information Systems*, 34(55), 1041–1066. <https://doi.org/10.17705/1CAIS.03455>
- Anonymous (2012). WHO collaborating centre for drug statistics methodology, Oslo, 2012. *Guidelines for ATC classification and DDD assignment 2013*. Retrieved from https://www.whooc.no/filearchive/publications/1_2013guidelines.pdf
- Anonymous (2017). WHO collaborating centre for drug statistics methodology, international language for drug utilization research. Retrieved from <https://www.whooc.no/>
- Ash, J. S., Berg, M., & Coiera, E. (2004). Some unintended consequences of information technology in health care: The nature of patient care information system-related errors. *Journal of the American Medical Informatics Association: JAMIA*, 11(2), 104–112. <https://doi.org/10.1197/jamia.M1471>
- Ball, J. E., Murrells, T., Rafferty, A. M., Morrow, E., & Griffiths, P. (2014). 'Care left undone' during nursing shifts: Associations with workload and perceived quality of care. *BMJ Quality and Safety*, 23(2), 116–125. <https://doi.org/10.1136/bmjqs-2012-001767>
- Barker, K. N., Flynn, E. A., & Pepper, G. A. (2002). Observation method of detecting medication errors. *American Journal of Health-System Pharmacy*, 59(23), 2314–2316. <https://doi.org/10.1093/ajhp/59.23.2314>
- Bates, D. W. (2000). Using information technology to reduce rates of medication errors in hospitals. *BMJ (Clinical Research Ed.)*, 320(7237), 788–791.
- Berdot, S., Roudot, M., Schramm, C., Katsahian, S., Durieux, P., & Sabatier, B. (2016). Interventions to reduce nurses' medication administration errors in inpatient settings: A systematic review and meta-analysis. *International Journal of Nursing Studies*, 53, 342–350. <https://doi.org/10.1016/j.ijnurstu.2015.08.012>
- Blegen, M. A., Vaughn, T. E., & Goode, C. J. (2001). Nurse experience and education: Effect on quality of care. *Journal of Nursing Administration*, 31(1), 33–39. <https://doi.org/10.1097/00005110-200101000-00007>
- Blijleven, V., Koelemeijer, K., Wetzels, M., & Jaspers, M. (2017). Workarounds emerging from electronic health record system usage: Consequences for patient safety, effectiveness of care, and efficiency of care. *JMIR Human Factors*, 4(4), e27. <https://doi.org/10.2196/humanfactors.7978>
- Cheung, K. C., van der Veen, W., Bouvy, M. L., Wensing, M., van den Bemt, P. M., & de Smet, P. A. (2014). Classification of medication incidents associated with information technology. *Journal of the American Medical Informatics Association: JAMIA*, 21(e1), e63–e70. <https://doi.org/10.1136/amiajnl-2013-001818>
- Cresswell, K. M., Mozaffar, H., Lee, L., Williams, R., & Sheikh, A. (2016). Workarounds to hospital electronic prescribing systems: A qualitative study in English hospitals. *BMJ Quality and Safety*, 26(7), 542–551. <https://doi.org/10.1136/bmjqs-2015-005149>
- Dean, B. S., & Barber, N. D. (1999). A validated, reliable method of scoring the severity of medication errors. *American Journal of Health-System Pharmacy*, 56(1), 57–62. <https://doi.org/10.1093/ajhp/56.1.57>
- Dean, B., & Barber, N. (2001). Validity and reliability of observational methods for studying medication administration errors. *American Journal of Health-System Pharmacy*, 58(1), 54–59. <https://doi.org/10.1093/ajhp/58.1.54>
- Debono, D. S., Greenfield, D., Travaglia, J. F., Long, J. C., Black, D., Johnson, J., & Braithwaite, J. (2013). Nurses' workarounds in acute healthcare settings: A scoping review. *BMC Health Services Research*, 13, 175–6963-13-175. <https://doi.org/10.1186/1472-6963-13-175>
- Donaldson, N., & Shapiro, S. (2010). Impact of California mandated acute care hospital nurse staffing ratios: A literature synthesis. *Policy, Politics and Nursing Practice*, 11(3), 184–201. <https://doi.org/10.1177/1527154410392240>
- Driscoll, A., Grant, M. J., Carroll, D., Dalton, S., Deaton, C., Jones, I., ... Astin, F. (2018). The effect of nurse-to-patient ratios on nurse-sensitive patient outcomes in acute specialist units: A systematic review and meta-analysis. *European Journal of Cardiovascular Nursing*, 17(1), 6–22. <https://doi.org/10.1177/1474515117721561>
- Gale, E. A. (2004). The Hawthorne studies-a fable for our times? *QJM: Monthly Journal of the Association of Physicians*, 97(7), 439–449. <https://doi.org/10.1093/qjmed/hch070>
- Goedhart, N. S., van Oostveen, C. J., & Vermeulen, H. (2017). The effect of structural empowerment of nurses on quality outcomes in hospitals: A scoping review. *Journal of Nursing Management*, 25(3), 194–206. <https://doi.org/10.1111/jonm.12455>
- Griffiths, P., Ball, J., Murrells, T., Jones, S., & Rafferty, A. M. (2016). Registered nurse, healthcare support worker, medical staffing levels and mortality in English hospital trusts: A cross-sectional study. *British Medical Journal Open*, 6(2), e008751–2015-008751. <https://doi.org/10.1136/bmjopen-2015-008751>
- Harris, T., Kerry, S., Victor, C., Illiffe, S., Ussher, M., Fox-Rushby, J., ... Cook, D. (2018). A pedometer-based walking intervention in 45- to 75-year-olds, with and without practice nurse support: The PACE-UP three-arm cluster RCT. *Health Technology Assessment (Winchester, England)*, 22(37), 1–274. <https://doi.org/10.3310/hta22370>
- Hassink, J. J., Essenberg, M. D., Roukema, J. A., & van den Bemt, P. M. (2013). Effect of bar-code-assisted medication administration on medication administration errors. *American Journal of Health-System Pharmacy*, 70(7), 572–573. <https://doi.org/10.2146/ajhp120257>
- Helmons, P. J., Wargel, L. N., & Daniels, C. E. (2009). Effect of bar-code-assisted medication administration on medication administration errors and accuracy in multiple patient care areas. *American Journal of Health-System*, 66(13), 1202–1210. <https://doi.org/10.2146/ajhp080357>
- Holden, R. J., Rivera-Rodriguez, A. J., Faye, H., Scanlon, M. C., & Karsh, B. T. (2013). Automation and adaptation: Nurses' problem-solving behavior following the implementation of bar coded medication administration technology. *Cognition, Technology and Work (Online)*, 15(3), 283–296. <https://doi.org/10.1007/s10111-012-0229-4>
- Hutton, K., Ding, Q., & Wellman, G. (2017). The effects of bar-coding technology on medication errors: A systematic literature review. *Journal of Patient Safety*, 1–15. <https://doi.org/10.1097/PTS.0000000000000366>
- Kaushal, R., & Bates, D. W. (2002). Information technology and medication safety: What is the benefit? *Quality and Safety in Health Care*, 11(3), 261–265. <https://doi.org/10.1136/qhc.11.3.261>
- Keers, R. N. (2017). *Evaluation of pharmacy TECHnician supported MEDication administration rounds (TECHMED) on reducing omitted doses: A pilot randomised controlled trial and process evaluation in a university teaching hospital*. (No. CPRG2-TECHMED vs.3).
- Kobayashi, M., Fussell, S., Xiao, Y., & Seagull, J. (2005). *Work coordination, workflow, and workarounds in a medical context. CHI 2005 late breaking results* (pp. 1561–1564). New York, NY: ACM Press.
- Koppel, R. (2014). Bar-coded medication administration brings trade-offs even with humane automation and lean approaches. *Joint Commission Journal on Quality and Patient Safety/Joint Commission Resources*, 40(8), 339–340.
- Koppel, R., Smith, S., Blythe, J., & Kothari, V. (2015). Workarounds to computer access in healthcare organizations: You want my password or a dead patient? *Studies in Health Technology and Informatics*, 208, 215–220.
- Koppel, R., Wedterneck, T., Telles, J. L., & Karsh, B. T. (2008). Workarounds to barcode medication administration systems: Their occurrences, causes, and threats to patient safety. *Journal of the American Medical Informatics Association: JAMIA*, 15(4), 408–423. <https://doi.org/10.1197/jamia.M2616>
- Krahenbuhl-Melcher, A., Schlienger, R., Lampert, M., Haschke, M., Drewe, J., & Krahenbuhl, S. (2007). Drug-related problems in hospitals: A review of the recent literature. *Drug Safety*, 30(5), 379–407.

- Lisby, M., Nielsen, L. P., & Mainz, J. (2005). Errors in the medication process: Frequency, type, and potential clinical consequences. *International Journal for Quality in Health Care*, 17(1), 15–22.
- Maaskant, J. M., Vermeulen, H., Apampa, B., Fernando, B., Ghaleb, M. A., Neubert, A., & Soe, A. (2015). Interventions for reducing medication errors in children in hospital. *Cochrane Database of Systematic Reviews*, (3), CD006208. <https://doi.org/10.1002/14651858.CD006208.pub3>
- Patterson, E. S. (2018). Workarounds to intended use of health information technology: A narrative review of the human factors engineering literature. *Human Factors*, 60(3), 281–292. <https://doi.org/10.1177/0018720818762546>
- Patterson, E. S., Rogers, M. L., Chapman, R. J., & Render, M. L. (2006). Compliance with intended use of bar code medication administration in acute and long-term care: An observational study. *Human Factors*, 48(1), 15–22. <https://doi.org/10.1518/001872006776412234>
- Pedersen, C. A., Schneider, P. J., & Scheckelhoff, D. J. (2012). ASHP national survey of pharmacy practice in hospital settings: Dispensing and administration—2011. *American Journal of Health-System Pharmacy*, 69(9), 768–785. <https://doi.org/10.2146/ajhp110735>
- Poon, E. G., Keohane, C. A., Yoon, C. S., Dittmore, M., Bane, A., Levtzion-Korach, O., ... Gandhi, T. K. (2010). Effect of bar-code technology on the safety of medication administration. *New England Journal of Medicine*, 362(18), 1698–1707. <https://doi.org/10.1056/NEJMs0907115>
- Rack, L. L., Dudjak, L. A., & Wolf, G. A. (2012). Study of nurse workarounds in a hospital using bar code medication administration system. *Journal of Nursing Care Quality*, 27(3), 232–239. <https://doi.org/10.1097/NCQ.0b013e318240a854>
- Schimmel, A. M., Becker, M. L., van den Bout, T., Taxis, K., & van den Bemt, P. M. (2011). The impact of type of manual medication cart filling method on the frequency of medication administration errors: A prospective before and after study. *International Journal of Nursing Studies*, 48(7), 791–797. <https://doi.org/10.1016/j.ijnurstu.2010.12.007>
- Schwappach, D. L., Pfeiffer, Y., & Taxis, K. (2016). Medication double-checking procedures in clinical practice: A cross-sectional survey of oncology nurses' experiences. *British Medical Journal Open*, 6(6), e011394. <https://doi.org/10.1136/bmjopen-2016-011394>
- Seidling, H. M., & Bates, D. W. (2016). Evaluating the impact of health IT on medication safety. *Studies in Health Technology and Informatics*, 222, 195–205.
- Smeulders, M., Hoekstra, M., van Dijk, E., Overkamp, F., & Vermeulen, H. (2013). Interruptions during hospital nurses' medication administration rounds. *Nursing Reports*, 3(1), 4. <https://doi.org/10.4081/nursrep.2013.e4>
- Spetz, J., Donaldson, N., Aydin, C., & Brown, D. S. (2008). How many nurses per patient? Measurements of nurse staffing in health services research. *Health Services Research*, 43(5 Pt 1), 1674–1692. <https://doi.org/10.1111/j.1475-6773.2008.00850.x>
- van den Bemt, P. M. L. A. (2006). Drug-related problems: Definitions and classifications. *EJHP Practice*, 12(suppl), 10–12.
- van den Bemt, P. M., Idzinga, J. C., Robertz, H., Kormelink, D. G., & Pels, N. (2009). Medication administration errors in nursing homes using an automated medication dispensing system. *Journal of the American Medical Informatics Association: JAMIA*, 16(4), 486–492. <https://doi.org/10.1197/jamia.M2959>
- van der Veen, W., de Gier, H. J., van der Schaaf, T., Taxis, K., & van den Bemt, P. M. (2013). Risk analysis and user satisfaction after implementation of computerized physician order entry in Dutch hospitals. *International Journal of Clinical Pharmacy*, 35(2), 195–201. <https://doi.org/10.1007/s11096-012-9727-y>
- van der Veen, W., van den Bemt, P. M., Bijlsma, M., de Gier, H. J., & Taxis, K. (2017). Association between workarounds and medication administration errors in bar code-assisted medication administration: Protocol of a multicenter study. *JMIR Research Protocols*, 6(4), e74. <https://doi.org/10.2196/resprot.7060>
- van der Veen, W., van den Bemt, P. M. L. A., Wouters, H., Bates, D. W., Twisk, J. W. R., de Gier, H. J., ... BCMA Study Group (2017). Association between workarounds and medication administration errors in bar-code-assisted medication administration in hospitals. *Journal of the American Medical Informatics Association: JAMIA*, 25(4), 385–392. <https://doi.org/10.1093/jamia/ocx077>
- van Onzenoort, H. A., van de Plas, A., Kessels, A. G., Veldhorst-Janssen, N. M., van der Kuy, P. H., & Neef, C. (2008). Factors influencing bar-code verification by nurses during medication administration in a Dutch hospital. *American Journal of Health-System Pharmacy*, 65(7), 644–648. <https://doi.org/10.2146/ajhp070368>
- van Oostveen, C. J., Braaksma, A., & Vermeulen, H. (2014). Developing and testing a computerized decision support system for nurse-to-patient assignment: A multimethod study. *Computers, Informatics, Nursing: CIN*, 32(6), 276–285. <https://doi.org/10.1097/CIN.000000000000056>
- Westbrook, J. I. (2009). Development and testing of an observational method for detecting medication administration errors using information technology. *Studies in Health Technology and Informatics*, 146, 429–433.
- Westbrook, J. I., & Ampt, A. (2009). Design, application and testing of the work observation method by activity timing (WOMBAT) to measure clinicians' patterns of work and communication. *International Journal of Medical Informatics*, 78(Suppl 1), S25–S33. <https://doi.org/10.1016/j.ijmedinf.2008.09.003>
- Westbrook, J. I., Duffield, C., Li, L., & Creswick, N. J. (2011). How much time do nurses have for patients? A longitudinal study quantifying hospital nurses' patterns of task time distribution and interactions with health professionals. *BMC Health Services Research*, 11, 319. <https://doi.org/10.1186/1472-6963-11-319>
- Westbrook, J. I., Raban, M. Z., Lehnbohm, E. C., & Li, L. (2016). The precise observation system for the safe use of medicines (POSSUM): An approach for studying medication administration errors in the field. *Studies in Health Technology and Informatics*, 228, 629–633.
- Wise, J. (2016). Higher nurse to patient ratio is linked to reduced risk of inpatient death. *BMJ (Clinical Research Ed.)*, 352, i797. <https://doi.org/10.1136/bmj.i797>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: van der Veen W, Taxis K, Wouters H, Vermeulen H, Bates DW, van den Bemt PMLA; the BCMA Study Group. Factors associated with workarounds in barcode-assisted medication administration in hospitals. *J Clin Nurs*. 2020;29:2239–2250. <https://doi.org/10.1111/jocn.15217>

APPENDIX**BCMA Study Group**

Michiel Duyvendak, PharmD, PhD, Hospital Pharmacist, Department of Hospital Pharmacy, Antonius Hospital, Sneek, the Netherlands. Karen Oude Luttikhuis, MSc, PharmD, Goorse Apotheek, Goor, the Netherlands. Johannes J. W. Ros, MSc, PharmD, Hospital Pharmacist, Department of Clinical Pharmacy, Gelre Hospitals, Apeldoorn, the Netherlands. Erwin C. Vasbinder, PharmD, Ph.D.,

Hospital Pharmacist, Department of Hospital Pharmacy, Groene Hart Hospital, Gouda, the Netherlands. Maryam Atrafi, MSc, Utrecht University, school of Pharmacy, Utrecht, the Netherlands. Bjorn Brasse, MSc, PharmD, Hospital Pharmacist, Department of Hospital Pharmacy, Elkerliek Hospital, Helmond, the Netherlands. Iris Mangelaars, MSc, PharmD, Utrecht University, School of Pharmacy, Utrecht, the Netherlands.