

University of Groningen

## Comparative transcriptome analyses of adzuki bean weevil (*Callosobruchus chinensis*) response to hypoxia and hypoxia/hypercapnia

Cui, S. F.; Wang, L.; Ma, L.; Wang, Y. L.; Qiu, J. P.; Liu, Zh Ch; Geng, X. Q.

*Published in:*  
Bulletin of entomological research

*DOI:*  
[10.1017/S0007485318000512](https://doi.org/10.1017/S0007485318000512)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2019

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Cui, S. F., Wang, L., Ma, L., Wang, Y. L., Qiu, J. P., Liu, Z. C., & Geng, X. Q. (2019). Comparative transcriptome analyses of adzuki bean weevil (*Callosobruchus chinensis*) response to hypoxia and hypoxia/hypercapnia. *Bulletin of entomological research*, 109(2), 266-277. <https://doi.org/10.1017/S0007485318000512>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Comparative transcriptome analyses of adzuki bean weevil (*Callosobruchus chinensis*) response to hypoxia and hypoxia/hypercapnia

S.F. Cui<sup>1,2</sup>, L. Wang<sup>2</sup>, L. Ma<sup>3</sup>, Y.L. Wang<sup>2</sup>, J.P. Qiu<sup>2</sup>,  
Zh.Ch. Liu<sup>2\*</sup> and X.Q. Geng<sup>2\*</sup>

<sup>1</sup>School of Grain Science and Technology, Jiangsu University of Science and Technology, Zhenjiang 212004, China: <sup>2</sup>Department of Resources and Environment, School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai 200240, China: <sup>3</sup>Behavioral & Physiological Ecology (BPE) Group, Groningen Institute for Evolutionary Life Sciences, University of Groningen, Nijenborgh 7, 9747 AG Groningen, Netherlands

## Abstract

Stored product insects show high adaption to hypoxia and hypercapnia, but the underlying mechanism is still unclear. Herein, a comparative transcriptome on 4<sup>th</sup> adzuki bean weevil (*Callosobruchus chinensis*) instar larvae was studied to clarify the response mechanisms to hypoxia (HA) and hypoxia/hypercapnia (HHA) using NextSeq500 RNA-Seq. Transcript profiling showed a significant difference in HA or HHA exposure both quantitatively and qualitatively. Compared with control, 631 and 253 genes were significantly changed in HHA and HA, respectively. Comparing HHA with HA, 1135 differentially expressed genes (DEGs) were identified. The addition of hypercapnia made a complex alteration on the hypoxia response of bean weevil transcriptome, carbohydrate, energy, lipid and amino acid metabolism were the most highly enriched pathways for genes significantly changed. In addition, some biological processes that were not significantly enriched but important were also discussed, such as immune system and signal transduction. Most of the DEGs related to metabolism both in HHA and HA were up-regulated, while the DEGs related to the immune system, stress response or signal transduction were significantly down-regulated or suppressed. This research reveals a comparatively full-scale result in adzuki bean weevil hypoxia and hypoxia/hypercapnia tolerance mechanism at transcription level, which might provide new insights into the genomic research of this species.

**Keywords:** adzuki bean weevil, comparative transcriptome, hypoxia, hypercapnia

(Accepted 21 May 2018; First published online 12 July 2018)

## Introduction

A modified atmosphere (MA) with low oxygen (hypoxia) and high carbon dioxide (hypercapnia) at biologically achievable levels have been demonstrated to be cost-effective and

environmentally friendly in the control of stored product insects and to maintain grain quality. MA could be achieved biologically in a barn bio-system or storage container by purging with CO<sub>2</sub> or N<sub>2</sub> and then sealing it hermetically, resulting in the suppression of insect feeding, growth, development, population expansion, and reproduction (Cheng *et al.*, 2013; García-Lara *et al.*, 2013; Iturralde-García *et al.*, 2016; Yan *et al.*, 2016). Generally, an MA with hypoxia (<5% oxygen) and hypercapnia (7.5–19.2% carbon dioxide) have been successful for the control of some storage pests (Conyers & Bell, 2007; Azab *et al.*, 2013; Wong-Corral *et al.*, 2013). However,

\*Author for correspondence  
Phone: 86-21-34208239 and 86-021-34208042  
Fax: 86-34205877 and 86-21-34206142  
E-mail: zhchl@sjtu.edu.cn and xqgeng@sjtu.edu.cn

many insect species from Coleoptera show high tolerance under similar situations, and they could even recover from hours to days of treatments. The physiological responses of the insects to MAs have been studied for years. As early as 1986, people had found that adults of *Cryptolestes ferrugineus*, *Sitophilus granaries*, *Tribolium castaneum*, and *Oryzaephilus surinamensis* could live under MA with low oxygen (1–1.6%) and increased carbon dioxide (10–30%) for nearly 7 days (Krishnamurthy *et al.*, 1986). In 1990, the hypoxia- and hypercapnia-selected *T. castaneum* was found to be capable of developing slowly under MAs (Donahaye, 1990). Recently, 4<sup>th</sup> wild-type *Callosobruchus maculatus* instar larvae could recover to normal development from more than 20 days of exposure to hypoxia and hypercapnia (Cheng *et al.*, 2012). Although some remarkable observations have been made in recent years, the mechanism of insect tolerance to hypoxia and hypercapnia is not fully understood. For instance, a number of insects moderate ventilator responses (e.g., changes in spiracles opening, convective gas, and tracheoles fluid levels) for adequate oxygen to maintain aerobic metabolism under hypoxia in the short term (seconds–hours), leading to appropriate levels of internal  $P_{CO_2}$  and pH. The concentration of carbon dioxide strongly affects the function of insect ventilation, and a fall in tracheal  $P_{CO_2}$  reduces the ventilation rate (Emekci *et al.*, 2002; Schmitz & Harrison, 2004; Matthews & White, 2011). An  $O_2$ -sensing system, including soluble guanylyl cyclases (e.g., Gyc-89 Da, Gyc-89 Db, and Gyc-88E) acting as molecular  $O_2$  sensors, are critical for mediating behavioral responses to hypoxia (Vermehren-Schmaedick *et al.*, 2010). In the long-term (hours–days), insects change morphological responses, including changes in tracheoles, respiratory and open tracheal system (Wingrove & O'farrell, 1999; Jha *et al.*, 2015). In these processes, the energy demand and metabolism rate in high hypoxia/hypercapnia-tolerant species (e.g., *Drosophila* and *C. maculatus*) was suppressed, shifting from aerobic to anaerobic metabolism, along with the changes in signal transduction (e.g., Notch, Toll/Imd, and Wnt pathways), cellular structure, and muscular contraction at the transcript level (Zhou *et al.*, 2008b; Chi *et al.*, 2011; Gersten *et al.*, 2014). Hypoxia-inducible factors (HIFs) are considered the key gene regulators. Under hypoxic conditions, HIFs would stimulate the expression of downstream genes related to immunity, apoptosis, development, stress responses and aging (Majmundar *et al.*, 2010).

Most of the previous researches have focused on the tolerance mechanism of insect under hypoxia, but fewer studies have investigated the response of insect under hypoxia/hypercapnia, let alone hypoxia/hypercapnia tolerant mechanism. *C. maculatus* larvae mortality was significantly increased when hypercapnia was added in hypoxia situation, but the larvae survived under hypoxia/hypercapnia grew faster than those under hypoxia only. As previously reported, hypercapnia decreased *Drosophila* resistance to bacterial infection by suppressing induction of specific antimicrobial peptides, but it does not induce a response characteristic of hypoxia, oxidative stress or heat shock, nor does it decrease a flies lifespan or cell viability (Helenius *et al.*, 2009; Selfridge *et al.*, 2016). The majority of metabolic genes in adult flies under 13%  $CO_2$  for 24 h were up-regulated, while over half of fertility- and immune-related genes were down-regulated (Selfridge *et al.*, 2016), which was contrary to the response of hypoxia-selected *Drosophila* (3rd larvae and adults) or *C. maculatus* larvae under 3%  $O_2$  + 17%  $CO_2$  for 24 h. Most of the

commonly up-regulated genes were involved in immunity, and the down-regulated genes were involved in metabolism (Zhou *et al.*, 2008a; Chi *et al.*, 2011). The transcript profile of *Litopenaeus vannamei* was altered under hypercapnia and hypoxic conditions, which showed the energetic cost for stabilizing translational machinery, mobilizing energy stores, and down-regulating ubiquitin/proteasomal degradation (Selfridge *et al.*, 2016).

Adzuki bean weevil, *Callosobruchus chinensis* (Coleoptera: Bruchidae), one of the most important destructive Bruchids, causes severe damage to stored legume grains and mainly inhabits tropical and subtropical regions of the world (Tuda *et al.*, 2005; Duan *et al.*, 2014; Kedia *et al.*, 2015; Duan *et al.*, 2016). Previous studies have paid more attention to this pest control strategies of MA, but little is known about this mechanism of action (Mohapatra *et al.*, 2015). According to our previous research, bean weevil shows high adaptability to hypoxia even hypoxia/hypercapnia (Cui *et al.*, 2017). To further reveal bean weevil hypoxia and hypoxia/hypercapnia tolerance mechanism, we used NextSeq500 RNA-Seq to investigate the differences in gene expression profiles among 4th adzuki bean weevil larvae under hypoxia, hypoxia/hypercapnia and normoxia. Here, we supposed that hypercapnia not only enhanced effects of hypoxia on gene expression in bean weevil but also produced different results.

## Materials and methods

### *Insect rearing and MA treatment*

Adzuki bean weevils were maintained on *Ormosia* seeds at a density of two or three insects per seed in glass bottles in a normoxic climate box at  $28 \pm 1^\circ C$  and 65% relative humidity under a 16 h:8 h light:dark photoperiod regime. We determined the developmental stages as previously described, using head capsule size as a standard (Cheng *et al.*, 2012). The developmental substages of egg, 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup>, and 4<sup>th</sup> instar, pupae, and adult from egg lying was 4, 8, 12, 16, 21, 23 and 23.5 d, respectively. When the larvae reached the middle of the fourth instar stage (16 days old), 20 larvae with 10 infested seeds (two larvae per seed) were transferred to specifically designed flat-bottomed flasks, with an inlet rubber hose connected to a mixed air cylinder (2%  $O_2$  + 98%  $N_2$  [hypoxia, HA] or 2%  $O_2$  + 18%  $CO_2$  + 80%  $N_2$  [hypercapnia/hypoxia, HHA] and an outlet rubber hose connected to a bubbler with silicone oil), and the mixed gas needed to be pre-humidified to nearly 68% r.h. by passing through a bubbling container containing a saturated solution of potassium iodide. The mixed gas was delivered into the flasks from the inlet rubber hose controlled by a gas reducing valve and flowmeter until the air concentration of the flasks became stable, which was verified using a head space analyzer (58,143,389; Danensor-checkpoint; Hergesteillon); then, the pressure was regulated to 1 atm and the gas flow rate to  $10 \text{ ml min}^{-1}$ . The pre-mixed gas was purchased in the form of pressurized cylinders from Shanghai He sheng Specialty Gases, and the concentration was verified (Donahaye & Navarro, 2000; Riudavets *et al.*, 2009; Cheng *et al.*, 2015). Control larvae were transferred to another flat-bottomed flask, then it was sealed with gauze to allow atmospheric air to diffuse, maintaining the concentrations of 20%  $O_2$  + 0%  $CO_2$  + 80%  $N_2$ , 68% r.h. in the phyton. Each treatment group consisted of three replicates. Larvae were removed from the seeds 24 h later. Collected

samples were immediately frozen in liquid nitrogen and stored at  $-80^{\circ}\text{C}$  for later RNA extraction.

#### *RNA extraction and cDNA library construction and sequencing*

Nine samples from the HHA and HA treatments and the control were included. Each sample contained a pool of 20 4th instar larvae, homogenized together in a liquid nitrogen for total RNA extraction, and RNeasy Plus Total RNA extraction reagent (Takara Bio, Shiga, Japan) was used following the manufacturer's instructions. Residual genomic DNA was removed with an RNA clean RNA kit (Bioteck, Beijing, China) according to the manufacturer's instructions and then dissolved in RNase-free water. The integrity of the total RNA was measured using an Agilent 2100 Bioanalyzer system using the RNA 6000 Nano kit according to the manufacturer's protocols (Agilent Technologies, Palo Alto, CA, USA). The high-quality RNA (RIN values  $>9.0$ ) was used to construct the cDNA library, which was sequenced using an Illumina instrument following the NextSeq500 RNA-Seq experimental procedure with a TruSeq RNA Sample Prep Kit (Illumina, San Diego, CA, USA).

Poly (A) mRNA was isolated from the total RNA and further purified by combining with oligo-dT magnetic beads. The mRNA was mixed with a fragmentation buffer to obtain 200–300 bp fragments, which were used as templates to synthesize first-strand cDNA with random primers, and double-strand cDNA was transformed from the first-strand cDNA using RNase H and DNA polymerase I. The double-strand cDNA fragments (200–300 bp) were purified using the QIA quick polymerase chain reaction (PCR) Extraction Kit (Qiagen, Valencia, CA, USA), end-repaired with 3'–5' exonuclease and polymerase, and ligated with sequencing adapters through the A and T complementary base pairing. AMPure XP beads (Beckman Coulter, Shanghai, China) were used to remove unsuitable fragments, and the sequencing library was constructed by PCR amplification. To prevent an overestimation of the mRNA copy number, the cycle number of cDNA libraries was strictly controlled. The multiplexed cDNA libraries were checked using PicoGreen (Quantifluor<sup>TM</sup>-ST fluorometer E6090; Promega, Madison, WI, USA) and fluorospectrophotometry (P7589 Quant-iT PicoGreen dsDNA Assay Kit; Invitrogen, Carlsbad, CA, USA) and quantified with the Agilent2100 Bioanalyzer and Agilent High Sensitivity DNA Kit (5067–4626). The synthesized cDNA libraries were normalized to 10 nM, diluted, quantified to 4–5 pM, and sequenced using the Illumina NextSeq<sup>TM</sup> 500 platform.

#### *De novo transcriptome assembly and bioinformatics analysis*

Raw reads from nine samples were collected and filtered by removing unsuitable reads as follows: adapter sequences with at least 10 bp overlap were removed using Cutadapt (ver. 1.2.1) (Martin, 2011), and reads with low average quality ( $Q \leq 20$ ) or a final length  $<50$  were trimmed off. The high-quality sequences were assembled into contigs and transcripts using Trinity (R20140717, kmer 25 bp) with the default parameters (kmer = 25 and minimum contig length = 48), and the longest transcript in each cluster was selected as unigenes for subsequent analyses (Altschul *et al.*, 1990; Haas *et al.*, 2013). Unigenes were submitted to a TBLASTX search in the non-redundant (NR) protein database to identify homology. Annotations with E-values  $<10^{-5}$  were retained and the top-hit species were indicated. In addition to NR, the Evolutionary

Genealogy of Genes, Non-supervised Orthologous Groups (eggNOG), and Gene Ontology (GO) databases were also used to identify homology, annotate the unigene distribution, and analyze biological behaviors, respectively (Ashburner *et al.*, 2000; Powell *et al.*, 2014). The Kyoto Encyclopedia of Genes and Genomes (KEGG) database was also used for annotation and to certify the functions of the unigenes in metabolic pathways (Kanehisa *et al.*, 2004). All sequences were mapped to unigenes using Bowtie (ver. 2.2.4, default parameters) (Langdon, 2015). The reads per kilobase per million mapped reads (RPKM) were normalized and identified using the DESeq 1.18 package in the R software, using an adjusted  $P$ -value of 0.05 as the threshold (Anders & Huber, 2010). Genes with a relative RPKM ratio  $>2$  or  $<0.5$  and  $P < 0.05$  were considered to be differentially expressed genes (DEGs). The Vann program was used to analyze the relationships of the genes in the different treatments to detect overlapping DEGs. The DEGs were also presented visually in Volcano Plot using the GGplot2 package in the R software. The enrichment analysis used to identify the functions of the DEGs was performed using GOSlim and the KEGG pathway database.

#### *Identification of DEGs by q-PCR*

Quantitative real-time PCR was performed using the total RNA extracted as described previously to confirm the expression of the putative DEGs from the transcriptome. cDNA was reverse-transcribed with 1  $\mu\text{g}$  total RNA using a PrimeScript<sup>TM</sup> RT reagent Kit with gDNA Eraser (Takara Bio). Selected genes were amplified using a SYBR Premix Ex Taq<sup>TM</sup> II Kit (Takara Bio), the DyNmo Flash SYBR Green qPCR Kit (Thermo Scientific, Rockford, IL, USA), and the CFX connect Real-Time System (Bio-Rad, Hercules, CA, USA) machine.  $\beta$ -actin was chosen as the reference to normalize the mRNA level between treatment and control samples. Three biological replicates were performed for each treatment, with three technical repeats for each biological replicate. A melting curve analysis was performed, and the specificity of the amplification and fold-change value of selected genes was calculated using the  $2^{-\Delta\Delta\text{Ct}}$  method. When the control  $\Delta\text{Ct}$  is greater than the treatment  $\Delta\text{Ct}$ , the  $2^{-\Delta\Delta\text{Ct}}$  value is  $<1$ . The negative inverse  $2^{-\Delta\Delta\text{Ct}}$  is the fold-change reduction in expression; otherwise, the  $2^{-\Delta\Delta\text{Ct}}$  value is  $>1$ , and the original  $2^{-\Delta\Delta\text{Ct}}$  value is used as the fold change increase from the treatment (Schmittgen & Livak, 2008).

## Results

#### *Transcriptome sequencing and assembly*

A mean total of  $5.83 \times 10^7$ ,  $5.27 \times 10^7$ , and  $5.18 \times 10^7$  raw reads were obtained from the libraries of samples under HA, HHA, and NOR, respectively. The mean read length was 150 bp. After filtering the low-quality sequences (Q20), nearly  $5.37 \times 10^7$ ,  $5.20 \times 10^7$ , and  $4.66 \times 10^7$  reads were useful under HHA, HA and NOR, respectively (fig. 1a), while 94.13% of reads ( $5.07 \times 10^7$ ) and 92.84% of bases ( $7.3 \times 10^9$ ) remained for the *de novo* assembly, and the information of databases is presented in fig. 1b.

A total of 8,874,226 contigs with an average length of 209 bp were obtained, and among them, 197,813 contigs were longer than 209 bp, 710,845 bp were longer than 111 bp, the percent of GC was 44.94%. A total of 223,694 transcripts with an average length of 506 bp were obtained, and among




Fig. 1. Summary of sequence data. (a) Reads, (b) bases (c) the length of the transcript.

them, 41,247 transcripts were longer than 594 bp, 17,1930 transcripts were longer than 238 bp, and the percent of GC was 43.55%. A total of 41,110 unigenes with an average length of 816 bp were obtained, and among them, 6427 unigenes were longer than 1438 bp, 27,713 unigenes were longer than 312 bp, and the percent of GC was 46.28% (table 1). The number of unigenes decreases as its length increases (fig. 1c).

#### Unigene functional annotation

After being assembled, all unigenes were submitted to the NR, GO, Kegg Orthology (KO), eggNOG, and Swiss-Prot databases for functional annotation using BLSTTX. Among the annotated unigenes, 93.44% were similar to genes in eggNOG, which explores the ancestry of a protein in 26

categories; 65.89% of unigenes were annotated with GO, which is a standard database for classifying gene function; 13.72% of unigenes were annotated with KO; and 80.94% of unigenes were functionally annotated by Swiss-Prot. A total of 65.89% of unigenes were annotated into GO categories, whereas only 13% were annotated into KO categories (table 2).

#### Gene expression profiling in HHA and HA

Compared with NOR, a total of 631 DEGs (345 up-regulated, 286 down-regulated) were found in HHA, and 253 DGEs (74 up-regulated, 179 down-regulated) were found in HA. 1135 DEGs (940 up-regulated, 195 down-regulated) were obtained from the comparison of HHA and HA, and the details are listed completely in table S1. Based

Table 1. Summary of the assembly from sequencing for adzuki bean weevil.

	Contig	Transcript	Unigene
Total Length (bp)	185,525,298	113,207,377	33,551,294
Sequence Number	887,426	223,694	41,110
Max. Length (bp)	44,489	44,465	38,164
Mean Length (bp)	209.06	506.08	816.13
N50 (bp)	209	594	1438
N50 Sequence No.	197,813	41,247	6427
N90 (bp)	111	238	312
N90 Sequence No.	710,845	171,930	27,713
GC%	44.94	43.55	46.28

Table 2. The summary of unigene function annotation with different databases.

Database	Number	Percentage (%)
NR	41,111	100
GO	27,090	65.89
KEGG	5640	13.72
eggNOG	38,415	93.44
Swissprot	33,274	80.94
In all database	4113	10

on GO enrichment, more than ten biological processes were found to be significantly affected by both HHA and HA: molecular function (i.e., structural molecular activity and RNA binding), cellular components (i.e., translation and symbiosis) and organization processes (i.e., membrane organization and lyase activity) were the most significantly enriched categories (fig. 2b and table S2).

Based on KO enrichment, most of the metabolic pathways were significantly affected under HHA and HA ( $P < 0.05$ ), and the pathways of carbohydrate metabolism, energy, lipid metabolism were the top three significantly enriched categories by DEGs, followed by the pathways of energy metabolism, xenobiotics biodegradation, and metabolism. The processes of amino acid metabolism, transport and catabolism, and endocrine and digestive systems were significantly enriched in HHA, the process of folding, sorting, and degradation were enriched in HA. Due to the synthetic effect of hypoxia and hypercapnia, the metabolic process was also the most affected category, followed by the biological cellular process and molecular function of binding categories (fig. 2b and table S3). Some of the important processes underlying hypoxia and hypercapnia response in bean weevil were presented below.

#### *Important pathways related to hypoxia and hypercapnia resistant to bean weevil*

Carbohydrate metabolism was the most highly significant enriched pathway in both treatments. Sixty-nine DEGs was found for carbohydrate metabolism in HHA ( $P < 0.05$ ), six DEGs were found in HA ( $P < 0.05$ ) and 78 DEGs were obtained from the comparison of HHA to HA (fig. 2b and table S3). These DEGs related were widely distributed in branched metabolic pathways, especially the pathways of glycolysis/gluconeogenesis, the citrate cycle (TCA cycle), pentose phosphate, pentose, and glucuronate interconversions and pyruvate metabolism (table S4). Most of DEGs related to these pathways were up-regulated in HHA, except for *MINPPI*

and *LDH*, respectively encoding multiple inositol-polyphosphate phosphatases and L-lactate dehydrogenase (fig. 3a).

Lipid metabolism was the most significantly enriched pathway in HHA (28 DEGs,  $P < 4.36E-16$ ) but not in HA (0 DEGs) (fig. 2b and table S3), and most of the pathways in HHA were up-regulated, except for *PNLIP* encoding pancreatic triacylglycerol lipase (fig. 3b). These genes were widely distributed in branched metabolic pathways, especially fatty acid degradation, glycerolipid, glycerophospholipid, and sphingolipid metabolism (table S4).

Amino acid metabolism was the third mostly enriched category in HHA (51 DEGs,  $P < 0.05$ ), while it was not significantly enriched in HA (1 DEG,  $P > 0.05$ ), relative to NOR (fig. 2b and table S3). These DEGs were widely present in almost every amino acid metabolism branches, such as the pathway of alanine, aspartate and glutamate metabolism, arginine and proline metabolism, tyrosine metabolism, and valine, leucine and isoleucine degradation (table S4), most of DEGs related were up-regulated, except for *CHDH* encoding glutaryl-CoA dehydrogenase (fig. 3c).

Energy metabolism was significantly enriched by DEGs both in HHA and HA relative to NOR. Twenty-six DEGs were found in HHA ( $P < 0.05$ ), while only eight DEGs were in HA ( $P < 0.05$ ) (fig. 2b and table S3). Among them, 12 and 4 DEGs were assigned to oxidative phosphorylation (OXPHOS) in HHA and HA, respectively. Almost all of the related genes were up-regulated. Comparing HHA with HA, energy metabolism was also remarkably enriched, and most of the related genes were up-regulated (fig. 3d and table S4).

The GO enrichment analysis also showed that most DEGs are related to the processes of the immune system and the response to stress was suppressed in HHA or HA, except *PGRP2*, *SODM1*, and *PRGP*, respectively encoding peptidoglycan-recognition, superoxide dismutase [Mn]1 and peptidoglycan-recognition protein were up-regulated (table S4). The alteration in expression of DEGs related to signal transduction was presented in fig. 3e. In HA, the increase of 3 genes encoding those key enzymes such as small subunit ribosomal protein S6e(R-S6e), GAPDH, HSP90A, and R-S6e, besides these 3 genes, another 6 genes encoding PDHB, GAPDH, PEPCCK, and cyclic AMP-responsive element-binding protein 3(CREB3) were detected in HHA, most of which were up-regulated, except *GAPDH*.

#### *q-PCR validation*

Twenty-three genes were selected randomly to confirm the accuracy of the RNA-Seq analysis by q-PCR. The results showed an agreement between the q-PCR and RNA-Seq results in paired comparisons of HA and NOR, HHA and




Fig. 2. Summary of significantly enriched categories based on GOSlims (a) and KOSlims (b). GO and KO categories are shown on the  $y$ -axis, the distribution of DEGs is presented as  $\log_{10}(P\text{-value})$  calculated with the principle of hypergeometrics on the  $x$ -axis,  $P < 0.05$ .




Fig. 3. The alteration in expression of DEGs related to Carbohydrate metabolism (a), Amino acid metabolism (b), Lipid metabolism (c), oxidative phosphorylation (d) and signal transduction (e).






Fig. 4. The correlation between the transcript level and q-PCR certification of genes obtained from pairwise comparisons of (a) HA and NOR, (b) HHA and NOR, (c) HHA and HA.

NOR, and HHA and HA (fig. 4). For example, the expression of *SARDH*, which encodes sarcosine dehydrogenase and is a key regulatory enzyme in the glycine, serine, and threonine metabolic pathway, was up-regulated 5-fold (q-PCR) or 4.55-fold (RNA-Seq) after 24 h of exposure to HHA, and 1.65-fold (q-PCR) or 1.92-fold (RNA-Seq) after 24 h of exposure to HA compared with those of the NOR treatment. The q-PCR primers used are presented in table S6.

### Discussion

A modified atmosphere (MA) with hypoxia and hypercapnia is known to be a useful method for stored-product insect control, which has a great effect on the development of *Callosobruchus* species. The transcriptome profiles of *C. maculatus* in response to hypoxia/hypercapnia had been reported before (3% O<sub>2</sub> + 17% CO<sub>2</sub> for 4 h, 24 h) (Chi *et al.*, 2011). Our RNA-Seq results provide more depth and coverage of the gene expression of *C. chinensis* in different air situations (2% O<sub>2</sub>, 2% O<sub>2</sub> + 18% CO<sub>2</sub>) for 24 h. the current study represents our first attempt to investigate the molecular response of *C. chinensis* to hypoxia and hypercapnia. We found that genes related to primary metabolism and energy metabolism in bean weevil were highly significantly changed due to hypoxia and hypercapnia, and the presence of hypercapnia enhanced hypoxia effects on gene expression (Zhou *et al.*, 2008a; Chi *et al.*, 2011).

Most of DEGs related to carbohydrate metabolism (e.g., glycolysis, TCA cycle, pyruvate metabolism, and pentose phosphate) were down-regulated in hypoxia, while most of DEGs were up-regulated in the presence of hypercapnia (fig. 5a). For example, the decrease of *GAPDH*, *ALDO* respectively, encoding glyceraldehyde 3-phosphate dehydrogenase and fructose-bisphosphate aldolase, class I in glycolysis in HA, might suggest anaerobic metabolism suppressed. The increase of DEGs (e.g., *DLAT*, *PDHB*, *PDHA*, *PC*, *MDH2*, *SDHB*, and *CS*) involved in TCA cycle and glycolysis (e.g., *LDH* and *PK*) in HHA, might lead to the increase of energy supply but the disturbance of internal acid-base balance (Nicolas & Sillans, 1989; Harrison, 2001; Förster & Hetz, 2010). Furthermore, the increase of DEGs involved in pyruvate metabolism and pentose phosphate pathway in HHA might be useful for energy supply (Feala *et al.*, 2009; Ahn *et al.*, 2013). It might be the reason for larvae growing faster but the higher death rate in HHA than those in HA (Zhou *et al.*, 2008a; Ragland *et al.*, 2010; Chi *et al.*, 2011; Cheng *et al.*, 2012).

Lipid metabolism is not only nutrient reserves when insect metabolic depression under extreme environment, but also serve as cell structural components, growth regulators and precursors of neurotransmitters. In our paper, bean weevil lipid metabolism was significantly affected by hypoxia/hypercapnia, while it significantly changed in fungi, *C. neoformans*, and *Drosophila* under hypoxia only (Donahaye & Navarro, 2000; Zhou *et al.*, 2008a; Gleason *et al.*, 2011). Our results were similar to *C. maculatus* under hypoxic/hypercapnic (Donahaye & Navarro, 2000; Chi *et al.*, 2011; Ahn *et al.*, 2013). It suggests that stored product insects (i.e., bean weevil, cowpea bruchid) have much higher hypoxia tolerance. In hypoxia/hypercapnia, some DEGs encoding the rate-limiting enzymes in lipid metabolism were up-regulated (fig. 5b). For example, long-chain-fatty-acid-CoA ligase gene (*ACSBG*) in the initial stage were up-regulated and acetyl-CoA C-acetyltransferase gene (*atoB*) in the final stage of fatty acid degradation were down-regulated, which might lead to the activation of fatty acid degradation. The increase of *PNLTP* in glycerolipid metabolism was helpful for the mobilization of triglyceride (TG) reserves for energy sources and water balance, while caused TG to be exhausted in a short time, resulting in insect death due to desiccation (Donahaye & Navarro, 2000). The genes of glycerol-3-phosphate O-acyltransferase 3/4 (*GPAT3* 4), glycerol-3-phosphate dehydrogenase (*GLPA*), and glycerol-3-phosphate dehydrogenase (NAD<sup>+</sup>) (*GPD1*) in sphingomyelin metabolism were up-regulated, which might lead to brain lipid precursors transferring into the processes of signal transduction, resulting in insect neurological disorder; The increase of glucuronosyltransferase gene (*UGT*) might inactivate ecdysteroid in steroid hormone biosynthetic pathway, leading to insect diapause (O'Reilly & Miller, 1989; Lin *et al.*, 2016).

Amino acid metabolism was most significantly affected by bean weevil under hypoxia/hypercapnia. Most of DEGs related were up-regulated, for example, the genes of homogentisate 1,2-dioxygenase (*TAT*), aspartate aminotransferase, mitochondrial (*GOT2*), and kynurenine/2-amino adipate aminotransferase (*KAT2*) and 4-hydroxyphenylpyruvate dioxygenase gene (*HPD*) were up-regulated, which might make amino acids lose more important molecular functions than the energy supply (fig. 5c). The decrease of *TAT* and *HPD* in tyrosine metabolism might lead to the decrease of dopamine which is important for neurotransmission, curricular sclerotization, and melanization in insects (Manekar *et al.*, 2005; Marmaras & Lampropoulou, 2009); The increase of 4-aminobutyrate aminotransferase gene (*ABAT*) might lead



to the decrease of GABA, which is the principle synaptic transmission inhibitor both in the central nervous system and at nerve/muscle junctions in insects (Nicolas & Sillans, 1989; Ramputh & Bown, 1996; Duncan & Newton, 2000; Rash and Hodgson, 2002).

Most of DEGs related to oxidative phosphorylation (OPHOS), except for the decrease of cytochrome c oxidase subunit 3 gene (*COX3*), were found to be significantly up-regulated due to hypoxia or hypercapnia. Cytochrome c oxidase subunit (COX) is critical for the utilization of O<sub>2</sub> in process of mitochondrial respiration. COX has three largest subunits (COX1, COX2, and COX3). In this paper, *COX1* was up-regulated and *COX3* was down-regulated, which might influence the catalytic activity of COX and the utilization of O<sub>2</sub> in bean weevil under the hypoxic environment (Khalimonchuk & Rödel, 2005). This phenomenon was similar to the response of Tibetan migratory locust but opposite to the response of *Epiblema scudderiana* and *Eurosta solidaginis* under a similar situation (McMullen & Storey, 2008; Zhang *et al.*, 2013). The alteration in expressions of DEGs encoding NADP-producing enzymes (e.g., *NDUFA12*, *NDUFAS8* and *NDUFB11*) and succinate dehydrogenase (e.g., *SDHA* and *SDHB*) were up-regulated in hypoxia/hypercapnia. Unlike *Ephesthia cautella* and *Drosophila*, these enzymes were suppressed under pure carbon dioxide (Duncan & Newton, 2000; Colinet & Renault, 2012).

In addition, most of the genes related to immunity, stress response, and signal transduction were significantly suppressed in bean weevil under hypoxia and hypoxia/hypercapnia, except for a few genes that were up-regulated. For example, *PGRP2* and *PGRP-SC2* involved in pathways of Toll-like receptor signaling and immune deficiency were up-regulated (Dziarski & Gupta, 2006). *SODM1* and *CATA*, which are important for protecting cells from ROS damage, were up-regulated (Guzy *et al.*, 2005; Maroz *et al.*, 2008). *AAT2* and *RXRA* involved in oxidative phosphorylation,  $\beta$ -oxidation of fatty acids, were also up-regulated (Aprille, 1988; Huss *et al.*, 2001). These genes up-regulated might be helpful for bean weevil surviving under severe hypoxia and hypoxia/hypercapnia.

### Conclusion

We elucidated the differences in transcriptional profiles of the 4<sup>th</sup> adzuki bean weevil larvae under hypoxic, hypoxic/hypercapnic, and normal oxygen environments. Several differences between insect response to hypoxia and hypoxia/hypercapnia have been found, such as the regulation of primary metabolism, signal transduction pathways, immune system. We suppose that these multiple changes might contribute to the high hypoxia even hypoxia/hypercapnia tolerant in bean weevil. It might complement our previous metabolomics work on bean weevil under the same situation. These studies are critical for revealing the mechanism of hypoxia and hypoxia/hypercapnia tolerance in stored product insects, although further verification experiment should be done in future.

### Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S0007485318000512>.

### Acknowledgements

This study was supported by the Key Construction Program of the National '985' Project (985, Grant No.

TS0320215002) and Agri-X (Grant No. Agri-X 2016007) from Shanghai Jiao Tong University, Shanghai international scientific and technological cooperation foundation (Grant No.17310742000) and Zhenjiang scientific and technological foundation for social development (Grant No.SH2017053). The authors sincerely thank the experts from Shanghai Personal Biotechnology Co., Ltd for RNA-Seq technical support.

### References

- Ahn, J.-E., Zhou, X., Dowd, S.E., Chapkin, R.S. & Zhu-Salzman, K. (2013) Insight into hypoxia tolerance in cowpea bruchid: metabolic repression and heat shock protein regulation via hypoxia-inducible factor 1. *PLoS ONE* **8**, e57267.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.
- Anders, S. & Huber, W. (2010) Differential expression analysis for sequence count data. *Genome Biology* **11**, R106.
- Aprille, J. (1988) Regulation of the mitochondrial adenine nucleotide pool size in liver: mechanism and metabolic role. *The FASEB Journal* **2**, 2547–2556.
- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S. & Eppig, J.T. (2000) Gene Ontology: tool for the unification of biology. *Nature Genetics* **25**, 25–29.
- Azab, M., Darwish, A., Mohamed, R. & Sanad, M. (2013) Comparative efficacy of controlled atmospheres against two stored product insects. *Journal of Crop Protection* **2**, 343–353.
- Cheng, W., Lei, J., Ahn, J.-E., Liu, T.-X. & Zhu-Salzman, K. (2012) Effects of decreased O<sub>2</sub> and elevated CO<sub>2</sub> on survival, development, and gene expression in cowpea bruchids. *Journal of Insect Physiology* **58**, 792–800.
- Cheng, W., Lei, J., Ahn, J.-E., Wang, Y., Lei, C. & Zhu-Salzman, K. (2013) CO<sub>2</sub> enhances effects of hypoxia on mortality, development, and gene expression in cowpea bruchid, *Callosobruchus maculatus*. *Journal of Insect Physiology* **59**, 1160–1168.
- Cheng, W., Lei, J., Fox, C.W., Johnston, J.S. & Zhu-Salzman, K. (2015) Comparison of life history and genetic properties of cowpea bruchid strains and their response to hypoxia. *Journal of Insect Physiology* **75**, 5–11.
- Chi, Y.H., Ahn, J.-E., Yun, D.-J., Lee, S.Y., Liu, T.-X. & Zhu-Salzman, K. (2011) Changes in oxygen and carbon dioxide environment alter gene expression of cowpea bruchids. *Journal of Insect Physiology* **57**, 220–230.
- Colinet, H. & Renault, D. (2012) Metabolic effects of CO<sub>2</sub> anaesthesia in *Drosophila melanogaster*. *Biology Letters* **8**, 1050–1054.
- Conyers, S. & Bell, C. (2007) A novel use of modified atmospheres: storage insect population control. *Journal of Stored Products Research* **43**, 367–374.
- Cui, S., Wang, L., Qiu, J., Liu, Z. & Geng, X. (2017) Comparative metabolomics analysis of *Callosobruchus chinensis* larvae under hypoxia, hypoxia/hypercapnia and normoxia. *Pest Management Science* **73**, 1267–1276.
- Donahaye, E. (1990) The potential for stored-product insects to develop resistance to modified atmospheres. pp. 989–996 in *Proceedings of the Proceedings of the Fifth International Working Conference on Stored-Product Protection*. 9–14 September 1990, Bordeaux, France.
- Donahaye, E. & Navarro, S. (2000) Comparisons of energy reserves among strains of *Tribolium castaneum* selected for

- resistance to hypoxia and hypercarbia, and the unselected strain. *Journal of Stored Products Research* **36**, 223–234.
- Duan, C., Li, W., Zhu, Z., Li, D., Sun, S. & Wang, X. (2016) Genetic differentiation and diversity of *Callosobruchus chinensis* collections from China. *Bulletin of Entomological Research* **106**, 124–134.
- Duan, C.X., Li, D.D., Sun, S.L., Wang, X.M. & Zhu, Z.D. (2014) Rapid development of microsatellite markers for *Callosobruchus chinensis* using Illumina paired-end sequencing. *PLoS ONE* **9**, e95458.
- Duncan, F.D. & Newton, R.D. (2000) The use of the anaesthetic, enflurane, for determination of metabolic rates and respiratory parameters in insects, using the ant, *Camponotus maculatus* (Fabricius) as the model. *Journal of Insect Physiology* **46**, 1529–1534.
- Dziarski, R. & Gupta, D. (2006) The peptidoglycan recognition proteins (PGRPs). *Genome Biology* **7**, 1–13.
- Emekci, M., Navarro, S., Donahaye, E., Rindner, M., Azrieli, A., (2002) Respiration of *Tribolium castaneum* (Herbst) at reduced oxygen concentrations. *Journal of Stored Products Research* **38**, 413–425.
- Förster, T.D. & Hetz, S.K. (2010) Spiracle activity in moth pupae—the role of oxygen and carbon dioxide revisited. *Journal of Insect Physiology* **56**, 492–501.
- Feala, J.D., Coquin, L., Zhou, D., Haddad, G.G., Paternostro, G. & McCulloch, A.D. (2009) Metabolism as means for hypoxia adaptation: metabolic profiling and flux balance analysis. *BMC Systems Biology* **3**, 91.
- García-Lara, S., Ortiz-Islas, S. & Villers, P. (2013) Portable hermetic storage bag resistant to *Prostephanus truncatus*, *Rhyzopertha dominica*, and *Callosobruchus maculatus*. *Journal of Stored Products Research* **54**, 23–25.
- Gersten, M., Zhou, D., Azad, P., Haddad, G.G. & Subramaniam, S. (2014) Wnt pathway activation increases hypoxia tolerance during development. *PLoS ONE* **9**, e103292.
- Gleason, J.E., Corrigan, D.J., Cox, J.E., Reddi, A.R., McGinnis, L. A. & Culotta, V.C. (2011) Analysis of hypoxia and hypoxia-like states through metabolite profiling. *PLoS ONE* **6**, e24741.
- Guzy, R.D., Hoyos, B., Robin, E., Chen, H., Liu, L., Mansfield, K. D., Simon, M.C., Hammerling, U. & Schumacker, P.T. (2005) Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing. *Cell Metabolism* **1**, 401–408.
- Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., Couger, M.B., Eccles, D., Li, B. & Lieber, M. (2013) De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols* **8**, 1494–1512.
- Harrison, J.F. (2001) Insect acid-base physiology. *Annual Review of Entomology* **46**, 221–250.
- Helenius, I.T., Krupinski, T., Turnbull, D.W., Gruenbaum, Y., Silverman, N., Johnson, E.A., Sporn, P.H., Sznajder, J.I. & Beitel, G.J. (2009) Elevated CO<sub>2</sub> suppresses specific *Drosophila* innate immune responses and resistance to bacterial infection. *Proceedings of the National Academy of Sciences* **106**, 18710–18715.
- Huss, J.M., Levy, F.H. & Kelly, D.P. (2001) Hypoxia inhibits the peroxisome proliferator-activated receptor  $\alpha$ /retinoid X receptor gene regulatory pathway in cardiac myocytes a mechanism for  $\alpha$ 2-dependent modulation of mitochondrial fatty acid oxidation. *Journal of Biological Chemistry* **276**, 27605–27612.
- Iturralde-García, R.D., Borboa-Flores, J., Cinco-Moroyoqui, F.J., Riudavets, J., Del Toro-Sánchez, C.L., Rueda-Puente, E.O., Martínez-Cruz, O. & Wong-Corral, F.J. (2016) Effect of controlled atmospheres on the insect *Callosobruchus maculatus* Fab. in stored chickpea. *Journal of Stored Products Research* **69**, 78–85.
- Jha, A.R., Zhou, D., Brown, C.D., Kreitman, M., Haddad, G.G. & White, K.P. (2015) Shared genetic signals of hypoxia adaptation in *Drosophila* and in high-altitude human populations. *Molecular Biology and Evolution* **33**, 501–517.
- Kanehisa, M., Goto, S., Kawashima, S., Okuno, Y. & Hattori, M. (2004) The KEGG resource for deciphering the genome. *Nucleic Acids Research* **32**, D277–D280.
- Kedia, A., Prakash, B., Mishra, P.K., Singh, P. & Dubey, N.K. (2015) Botanicals as eco friendly biorational alternatives of synthetic pesticides against *Callosobruchus* spp. (Coleoptera: Bruchidae) – a review. *Journal of Food Science and Technology* **52**, 1239–1257.
- Khalimonchuk, O. & Rödel, G. (2005) Biogenesis of cytochrome c oxidase. *Mitochondrion* **5**, 363–388.
- Krishnamurthy, T.S., Spratt, E.C. & Bell, C.H. (1986) The toxicity of carbon dioxide to adult beetles in low oxygen atmospheres. *Journal of Stored Products Research* **22**, 145–151.
- Langdon, W.B. (2015) Performance of genetic programming optimised Bowtie2 on genome comparison and analytic testing (GCAT) benchmarks. *BioData Mining* **8**, 1.
- Lin, T., Cai, Z., Wu, H. & Luo, L. (2016) Changes in midgut gene expression following *Bacillus thuringiensis* (Bacillales: Bacillaceae) infection in *Monochamus alternatus* (Coleoptera: Cerambycidae). *Florida Entomologist* **99**, 60–66.
- Majmunder, A.J., Wong, W.J. & Simon, M.C. (2010) Hypoxia-inducible factors and the response to hypoxic stress. *Molecular Cell* **40**, 294–309.
- Manekar, A., Godshalwar, M., Rewale, M. & Singh, M. (2005) Effect of tyrosine on central nervous system of marine male crab, scylla serrata: a biochemical study. *Journal of Cell and Tissue Research* **5**, 345–348.
- Marmaras, V.J. & Lampropoulou, M. (2009) Regulators and signalling in insect haemocyte immunity. *Cellular Signalling* **21**, 186–195.
- Maroz, A., Kelso, G.F., Smith, R.A., Ware, D.C. & Anderson, R.F. (2008) Pulse radiolysis investigation on the mechanism of the catalytic action of Mn(II)–pentaazamacrocyclic compounds as superoxide dismutase mimetics. *The Journal of Physical Chemistry A* **112**, 4929–4935.
- Martin, M. (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet. Journal* **17**, 10–12.
- Matthews, P.G. & White, C.R. (2011) Regulation of gas exchange and haemolymph pH in the cockroach *Nauphoeta cinerea*. *Journal of Experimental Biology* **214**, 3062–3073.
- McMullen, D.C. & Storey, K.B. (2008) Mitochondria of cold hardy insects: responses to cold and hypoxia assessed at enzymatic, mRNA and DNA levels. *Insect Biochemistry and Molecular Biology* **38**, 367–373.
- Mohapatra, D., Kar, A. & Giri, S.K. (2015) Insect pest management in stored pulses: an overview. *Food and Bioprocess Technology* **8**, 239–265.
- Nicolas, G. & Sillans, D. (1989) Immediate and latent effects of carbon dioxide on insects. *Annual Review of Entomology* **34**, 97–116.
- O'Reilly, D.R. & Miller, L.K. (1989) A baculovirus blocks insect molting by producing ecdysteroid UDP-glucosyl transferase. *Science* **245**, 1110–1112.
- Powell, S., Forslund, K., Szklarczyk, D., Trachana, K., Roth, A., Huerta-Cepas, J., Gabaldón, T., Rattei, T., Creevey, C. &

- Kuhn, M.** (2014) eggNOG v4. 0: nested orthology inference across 3686 organisms. *Nucleic Acids Research* **42**, D231–D239.
- Ragland, G.J., Denlinger, D.L. & Hahn, D.A.** (2010) Mechanisms of suspended animation are revealed by transcript profiling of diapause in the flesh fly. *Proceedings of the National Academy of Sciences* **107**, 14909–14914.
- Ramputh, A.-I. & Bown, A.W.** (1996) Rapid [ $\gamma$ ]-aminobutyric acid synthesis and the inhibition of the growth and development of oblique-banded leaf-roller larvae. *Plant Physiology* **111**, 1349–1352.
- Rash, L.D. & Hodgson, W.C.** (2002) Pharmacology and biochemistry of spider venoms. *Toxicon* **40**, 225–254.
- Riudavets, J., Castañé, C., Alomar, O., Pons, M.J. & Gabarra, R.** (2009) Modified atmosphere packaging (MAP) as an alternative measure for controlling ten pests that attack processed food products. *Journal of Stored Products Research* **45**, 91–96.
- Schmittgen, T.D. & Livak, K.J.** (2008) Analyzing real-time PCR data by the comparative CT method. *Nature protocols* **3**, 1101–1108.
- Schmitz, A. & Harrison, J.F.** (2004) Hypoxic tolerance in air-breathing invertebrates. *Respiratory Physiology & Neurobiology* **141**, 229–242.
- Selfridge, A.C., Cavadas, M.A., Scholz, C.C., Campbell, E.L., Welch, L.C., Lecuona, E., Colgan, S.P., Barrett, K.E., Sporn, P.H. & Sznajder, J.I.** (2016) Hypercapnia suppresses the HIF-dependent adaptive response to hypoxia. *Journal of Biological Chemistry* **291**, 11800–11808.
- Tuda, M., Chou, L.-Y., Niyomdham, C., Buranapanichpan, S. & Tateishi, Y.** (2005) Ecological factors associated with pest status in *Callosobruchus* (Coleoptera: Bruchidae): high host specificity of non-pests to Cajaninae (Fabaceae). *Journal of Stored Products Research* **41**, 31–45.
- Vermehren-Schmaedick, A., Ainsley, J.A., Johnson, W.A., Davies, S.-A. & Morton, D.B.** (2010) Behavioral responses to hypoxia in *Drosophila* larvae are mediated by atypical soluble guanylyl cyclases. *Genetics* **186**, 183–196.
- Wingrove, J.A. & O'farrell, P.H.** (1999) Nitric oxide contributes to behavioral, cellular, and developmental responses to low oxygen in *Drosophila*. *Cell* **98**, 105–114.
- Wong-Corral, F.J., Castañé, C. & Riudavets, J.** (2013) Lethal effects of CO<sub>2</sub>-modified atmospheres for the control of three Bruchidae species. *Journal of Stored Products Research* **55**, 62–67.
- Yan, Y., Williams, S.B., Baributsa, D. & Murdock, L.L.** (2016) Hypoxia treatment of *Callosobruchus maculatus* females and its effects on reproductive output and development of progeny following exposure. *Insects* **7**, 26.
- Zhang, Z.-Y., Chen, B., Zhao, D.-J. & Kang, L.** (2013) Functional modulation of mitochondrial cytochrome c oxidase underlies adaptation to high-altitude hypoxia in a Tibetan migratory locust. *Proceedings of the Royal Society of London B: Biological Sciences*, 280:20122758.
- Zhou, D., Xue, J., Lai, J.C., Schork, N.J., White, K.P. & Haddad, G.G.** (2008a) Mechanisms underlying hypoxia tolerance in *Drosophila melanogaster*: hairy as a metabolic switch. *PLoS Genetics* **4**, e1000221.
- Zhou, X.R., Horne, I., Damcevski, K., Haritos, V., Green, A. & Singh, S.** (2008b) Isolation and functional characterization of two independently-evolved fatty acid  $\Delta$ 12-desaturase genes from insects. *Insect Molecular Biology* **17**, 667–676.