

University of Groningen

Infection control in dental health care during and after the SARS-CoV-2 outbreak

Volgenant, Catherine M. C.; Persoon, Ilona F.; de Ruijter, Rolf A. G.; de Soet, J. J. (Hans)

Published in:
 Oral diseases

DOI:
[10.1111/odi.13408](https://doi.org/10.1111/odi.13408)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
 Publisher's PDF, also known as Version of record

Publication date:
 2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Volgenant, C. M. C., Persoon, I. F., de Ruijter, R. A. G., & de Soet, J. J. H. (2021). Infection control in dental health care during and after the SARS-CoV-2 outbreak. *Oral diseases*, 27(S3), 674-683. <https://doi.org/10.1111/odi.13408>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Infection control in dental health care during and after the SARS-CoV-2 outbreak

Catherine M. C. Volgenant¹  | Ilona F. Persoon¹  | Rolf A. G. de Ruijter² | J. J. (Hans) de Soet¹ 

¹Department of Preventive Dentistry, Academic Centre of Dentistry Amsterdam (ACTA), University of Amsterdam and Vrije Universiteit Amsterdam, Amsterdam, The Netherlands

²Expert Group Behavioral and Contemplative Dentistry of the University Medical Center Groningen/Center for Dentistry and Oral Hygiene, Rijksuniversiteit Groningen, Groningen, The Netherlands

Correspondence

J. J. (Hans) de Soet, Department of Preventive Dentistry, ACTA, Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands.
Email: infectioncontrol@acta.nl

Abstract

COVID-19 is an emerging infectious disease caused by the widespread transmission of the coronavirus SARS-CoV-2. Some of those infected become seriously ill. Others do not show any symptoms, but can still contribute to transmission of the virus. SARS-CoV-2 is excreted in the oral cavity and can be spread via aerosols. Aerosol generating procedures in dental health care can increase the risk of transmission of the virus. Due to the risk of infection of both dental healthcare workers and patients, additional infection control measures for all patients are strongly recommended when providing dental health care. Consideration should be given to which infection control measures are necessary when providing care in both the current situation and in the future.

KEYWORDS

dentistry, infection control, infectious disease transmission, Public Health Dentistry, SARS-CoV-2

1 | INTRODUCTION

In recent months, the world has been confronted with an outbreak of the SARS-CoV-2 virus (Rabi, Al Zoubi, Kasasbeh, Salameh, & Al-Nasser, 2020), resulting in the *coronavirus disease 2019*; COVID-19 (Khan et al., 2020). This coronavirus has a single-stranded positive RNA chain, surrounded by a capsid (Adhikari et al., 2020; Ashour, Elkhatib, Rahman, & Elshabrawy, 2020). In most cases, the virus causes mild-to-severe respiratory complaints (Chen et al., 2020; Huang, Wang, et al., 2020). Since the population did not have contact with this virus before, no herd immunity against the virus has been acquired among the population. In addition, the virus is highly contagious, with one person infected with SARS-CoV-2 infecting on average 2–3 people (reproduction number R_0) (Liu, Gayle, Wilder-Smith, & Rocklöv, 2020; Sun et al., 2020). This results in a potentially

large group of infected individuals. By isolating these infected individuals and applying sufficient infection control measures, the R_0 will decrease. When the R_0 is less than 1, an outbreak will extinguish spontaneously (Chen, 2020; Heesterbeek & Dietz, 1996). However, when infected individuals show little or no symptoms, they are likely to be missed and that they will be exempted from isolation. This can contribute to the further dissemination of the virus (Munster, Koopmans, van Doremalen, van Riel, & de Wit, 2020).

The SARS-CoV-2 virus uses the membrane-bound angiotensin-converting enzyme 2 (ACE2) receptor of the host to penetrate the cells (Xu et al., 2020). The membrane-bound ACE2 receptor is mainly found on cells of the mucosal tissues, such as the dorsum of the tongue and salivary glands (Liu et al., 2011; Xu et al., 2020). Hence, saliva has been suggested for non-invasive diagnostics of this virus (Khurshid, Asiri, & Al Wadaani, 2020; Sabino-Silva, Jardim, & Siqueira, 2020). The mucosal membrane of the oral cavity, respiratory tract and eyes is an important *portal of entry* for this virus

Volgenant and Persoon contributed equally to this manuscript.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Oral Diseases* published by John Wiley & Sons Ltd



(Adhikari et al., 2020; Zhang, Zhang, & Wang, 2020). This *portal of entry* also serves as a reservoir from which transmission may occur to other individuals, for example coughing or sneezing (Adhikari et al., 2020). Now the outbreak has become a pandemic and no drugs or vaccines are yet available, infection control measures are the only option for the time being to slow down the number of new infections (Lai et al., 2020). These precautions include social distancing and isolation which prompted to provide only limited emergency (dental) health care during the beginning of the outbreak in most countries (Farooq & Ali, 2020; Guo, Zhou, Liu, & Tan, 2020; Izzetti, Nisi, Gabriele, & Graziani, 2020; Meng, Hua, & Bian, 2020b; Peng et al., 2020; Spagnuolo, De Vito, Rengo, & Tatullo, 2020). In time, dental health care should be slowly upscaled beyond emergency care in order to prevent and treat oral diseases. This paper summarises the infection control measures for the dental healthcare setting in relation to SARS-CoV-2, based on the currently available scientific evidence.

2 | INFECTION CONTROL IN DENTAL HEALTH CARE

Infection control in dental health care is about calculating risks; it is not possible to completely exclude all risks (Volgenant & de Soet, 2018). At first, infection control guidelines in dental health care aimed to prevent transmission of blood-borne diseases (Kohn et al., 2003). Unique to the dental healthcare setting is the profound production of aerosols during most treatment procedures. Aerosols are liquid or solid particles in the air, which can be responsible for the transfer of micro-organisms (Zemouri, de Soet, Crielaard, & Laheij, 2017). Considering the risk of aerosol transmission in dental health care, most patients are considered healthy and thus less strict aerosol precautions have to be taken compared to general healthcare settings (Siegel, Rhinehart, Jackson, Chiarello, & Committee, 2007). As a result, dental health care has always been provided in such a way that it has a relatively limited effect on feasibility and costs. Current guidelines in dental health care worldwide strive for *optimal* and feasible rather than *maximal* precautions.

3 | TRANSMISSION OF CORONAVIRUS SARS-CoV-2

It is suggested that four categories of transmission of SARS-CoV-2 occur: symptomatic transmission (direct transmission from a COVID-19 patient), presymptomatic transmission (direct transmission from a SARS-CoV-2-positive person without symptoms yet), asymptomatic transmission (direct transmission from a SARS-CoV-2-positive person who never develops symptoms) and environmental transmission (indirect transmission which is not traceable to an index patient) (Ferretti et al., 2020). Consequently, during the COVID-19 pandemic, patients in the dental healthcare setting cannot be considered healthy even when they are not experiencing

symptoms. Transmission of the SARS-CoV-2 virus has been reported from 12.6% of the patients before they showed symptoms (Du et al., 2020). Presymptomatic transmission from an index patient has been estimated to account for up to 44% of total transmissions and even occurred up to 2 days before symptoms were experienced (He et al., 2020). The time between infection and symptom onset (incubation period) of COVID-19 is on average 5.1 days (CI 95% 4.5–5.8), up to a period of 11.5 days (CI 95% 8.2–15.6) (Lauer et al., 2020). Numerous publications reported on transmission during the incubation period of COVID-19 (Huang, Xia, Chen, Shan, & Wu, 2020; Tong et al., 2020; Yu, Zhu, Zhang, & Han, 2020).

Several studies described individual cases of asymptomatic individuals (Arons et al., 2020; Chan et al., 2020; Hoehl et al., 2020; Kimball, 2020; Luo et al., 2020; Pan et al., 2020; Zou et al., 2020), as well as transmission from an asymptomatic index (Bai et al., 2020; Rothe et al., 2020). The viral load did not differ between symptomatic and asymptomatic carriers (Zou et al., 2020). In addition, mild symptoms or loss of smell or taste are also recognised as possible symptoms of infection with SARS-CoV-2 (Bai et al., 2020; Hu et al., 2020; Lai et al., 2020), making it difficult to identify a true asymptomatic individual. There could be a potential large reservoir of individuals in the population (Verity et al., 2020) that can contribute to rather hidden ongoing transmission (Munster et al., 2020).

The case fatality rate of this virus, reported in China, is 3%–11% (Rajgor, Lee, Archuleta, Bagdasarian, & Quek, 2020). In addition, often only mild symptoms occur (Kluytmans et al., 2020; Verity et al., 2020). Infected persons can thus continue to participate in the population, making it likely that dental healthcare workers (DHCWs) will come into contact with SARS-CoV-2-positive patients (Cheng et al., 2020). However, the case fatality rate is based on the number of deaths per confirmed case. The infection fatality rate of the virus has been estimated for the whole population of China at 0.66% (Verity et al., 2020). It is likely that both the infection fatality rate and the case fatality rate will change in the coming period when more tests will be performed.

4 | TRANSMISSION OF SARS-CoV-2 IN DENTAL HEALTH CARE

The transmission of the virus mainly occurs via respiratory droplets and faecal shedding (Liu, Ning, Chen, Guo, Liu, et al., 2020; Ngoc et al., 2020; Ong et al., 2020). These respiratory droplets are excreted from the oral cavity and pharynx, for example by speech, and usually do not reach more than 1.5–2 m (Ai & Melikov, 2018; Bischoff, Swett, Leng, & Peters, 2013). When coughing and sneezing, aerosols are also generated, in which the aerosols remain in the air for some time (Jones & Brosseau, 2015). Aerosols are able to reach beyond the social distancing instructions of 1.5–2 m, for example due to aerodynamic effects (Bischoff et al., 2013; Blocken, Malizia, van Druenen, & Marchal, 2020) and transmission of the SARS-CoV-2 virus via aerosols has been suggested (Liu, Ning, Chen, Guo, Liu, et al., 2020). Although aerosols do not play a major role in transmission



of SARS-CoV-2 in most daily activities, the situation is different in the dental clinic. Water in combination with compressed air used for coolant and spraying causes aerosols which become contaminated with micro-organisms from the oral cavity (Zemouri et al., 2017). DHCWs operate at a distance of 60 cm or less from a patient's oral cavity. A recent study indicates that the largest microbiological contamination within the dental healthcare clinic takes place within 1 m from the oral cavity, via both splashes and aerosols (Zemouri et al., 2020). In medical practice, transmission of SARS-CoV-2 virus via aerosols is suggested in addition to transmission of the virus via droplets (Ong et al., 2020; Wax & Christian, 2020). It appears that different types of coronaviruses can already be detected in aerosols produced by exhalation (Leung et al., 2020). Moreover, a SARS-CoV-2-positive patient has many virus particles in his saliva and on the dorsum of the tongue (To et al., 2020; Xu et al., 2020). This suggests that aerosols generated during dental healthcare treatment in these individuals can also contain SARS-CoV-2 and thereby transmit the virus to the DHCWs (Figure 1).

Even after completing treatment, aerosols are suspended in the air within the treatment room, with heavier and larger particles settling faster (Bennett et al., 2000; Nikitin, Petrova, Trifonova, & Karpova, 2014). Settling occurs on all horizontal surfaces, after which these can act as a vehicle for transmission of the SARS-CoV-2

virus via indirect contact (Figure 1). Viable virus was still detectable on, for example, plastic surfaces after 72 hr, up to 7 days (van Doremalen et al., 2020). Regardless of the modes of transmission, the minimal infectious dose of SARS-CoV-2 has not yet been established. Therefore, irrespective of the level of contamination, all surfaces contaminated with aerosol or touched by patients should be regarded as potentially contaminated.

5 | INFECTION CONTROL MEASURES

The dental healthcare setting can be an important route for transmission of airborne or drop-related infectious diseases, both for the dental team and the patient (Volgenant & de Soet, 2018) (Figure 1). Whenever new infectious diseases emerge, infection control protocols within the dental health care are modified accordingly (Monaghan, 2016; Smales & Samaranyake, 2003).

Protection against work-related infectious diseases can be implemented at different hierarchal levels (Figure 2). The National Institute for Occupational Safety and Health (NIOSH) in the United States conducts research and makes recommendations to prevent work-related diseases. Measures that intervene at a level closer to the source of the virus are generally more effective and provide





FIGURE 1 Transmission of SARS-CoV-2 can occur via direct contact, indirect contact and via air (droplets/aerosols). This can take place from patients to the DHCWs and vice versa, and reciprocal between patients or DHCWs. This applies to symptomatic, presymptomatic and asymptomatic individuals

FIGURE 2 Controlling exposure to occupational risks is aimed when protecting DHCWs. The hierarchy shown here is used to determine on which level feasible and effective solutions can be implemented in the dental clinic (adapted from National Institute for Occupational Safety and Health (NIOSH, 2015) of the Centre for Disease Control and Prevention (CDC), United States)



more protection than measures closer to the healthcare workers. The starting point of these necessary measures is that possibly all patients may be SARS-CoV-2 positive, although not symptomatic yet, and that SARS-CoV-2 is transmitted when providing dental health care, mainly via aerosols. Hence, possible adjustments to the regular protocols and the considerations involved are discussed below.

5.1 | Elimination of the reservoir

The infectious reservoir can be eliminated by preventing contact with an infected patient. Many guidelines in infection control in dental health care are based on this principle (Kohn et al., 2003). Infected patients are assumed to be too ill to visit the dental clinic or, as a result of the anamnesis, elective care is postponed. During the SARS-CoV-2 outbreak, dental health care is limited to providing urgent care in most countries (Farooq & Ali, 2020; Izzetti et al., 2020; Meng, Hua, & Bian, 2020a).

- Part of the demand of dental health care can be met by telephone, email or videoconferences (Guo, Wu, & Xie, 2020; Meng et al., 2020b). A detailed patient history, possibly accompanied by photographs or video conferencing, can aid the primary management of dental emergencies. Urgent care can be managed by prescribing analgesia, antiseptics or as a last option antibiotics (Ather, Patel, Ruparel, Diogenes, & Hargreaves, 2020). These modern techniques can also be applied when providing preventive dental health care (Darwish, 2020; Mallineni et al., 2020).
- In an area where widespread transmission has not been established, triage can aid in estimating risks of transmission of SARS-CoV-2. Individuals with a travel history to areas with ongoing transmission or individuals with recent exposure to SARS-CoV-2-positive individuals should be considered as a high risk for serving as a source of transmission. Moreover, individuals showing signs and symptoms of COVID-19 (e.g. coughing or a fever; see above) should also be considered as high risk for transmission. DHCWs

who comply with criteria as described above should be considered as high risk and therefore should not be present in the dental clinic.

- For detecting patients to who care can be provided with limited risk of transmission with SARS-CoV-2, triage is applicable (Alharbi, Alharbi, & Alqaidi, 2020; Ayebare, Flick, Okware, Bodo, & Lamorde, 2020; Izzetti et al., 2020; Prati, Pelliccioni, Sambri, Chersoni, & Gandolfi, 2020). Dental health care for patients showing signs and symptoms of COVID-19 should be limited to urgent care and can only be provided in a clinic with full protective measures. DHCWs have to keep in mind that triage is currently unable to differentiate asymptomatic or presymptomatic patients from unaffected individuals. Reliable, simple and cheap rapid tests can assist in determining to who dental health care can be provided without additional measures (Khurshid et al., 2020).
- When providing care is indispensable, elimination of (secondary) infectious reservoirs is essential in preventing transmission. Both hand hygiene and hygiene of surfaces have always been important measures against the spread of viruses in society and health care; this applies to SARS-CoV-2 as well (Lotfinejad, Peters, & Pittet, 2020; Lynch, Mahida, Oppenheim, & Gray, 2020; Nicolaidis, Avraam, Cueto-Felgueroso, González, & Juanes, 2020; Ran et al., 2020). It is important to realise that DHCWs should prevent touching their own face, both with and without personal protective equipment (PPE) (Elder, Sawyer, Pallerla, Khaja, & Blacker, 2014). Cleaning removes the virus mechanically and disinfection inactivates the virus. Surfaces that may be contaminated with SARS-CoV-2 can effectively be disinfected within 1 min by applying at least 62% alcohol, 0.5% hydrogen peroxide or 1,000 ppm (0.1%) sodium hypochlorite (Kampf, Todt, Pfaender, & Steinmann, 2020). Effective disinfection with different alcohol-based hand rub formulations and lower dilutions of alcohol has also been reported (Kratzel et al., 2020).
- The procedures for cleaning, disinfection and sterilisation of instruments can be performed as described in regularly applicable guidelines in dentistry (Kohn et al., 2003). However, mechanical



cleaning is strongly recommended (automated washer disinfectors) to prevent transmission by, for example, splashing during cleaning. Cleaning and/or disinfection should also include all horizontal surfaces in the treatment room and all other items and locations in the clinic that could have been touched by the patient (Kampf, Scheithauer, Lemmen, Saliou, & Suchomel, 2020).

5.2 | Engineering controls: isolating DHCWs from the hazard

- The air in the treatment room after an aerosol generating procedure should be regarded as contaminated. Dispersion of the virus throughout the dental clinic should be avoided, even though it is currently unknown whether the amount of virus particles in the air after an aerosol generating procedure in dental health care can exceed the infectious dose. Therefore, working under negative air pressure would be preferable (Cheong & Phua, 2006). Clean air will be drawn from less contaminated areas towards the treatment room. The active exhaust flow from the contaminated treatment room leads to removal of possible pathogens from the air.
- In most dental clinics, working under negative air pressure is not possible. Sufficient ventilation in the room (Meng et al., 2020a) will dilute the virus load (Stockwell et al., 2019). On the one hand, mechanical ventilation, possibly enhanced, can significantly increase the expulsion of air. On the other hand, natural ventilation can be improved by active ventilation and, if possible, create a draught through the room (Escombe, Ticona, Chávez-Pérez, Espinoza, & Moore, 2019). Research data on the required duration of ventilation regarding SARS-CoV-2 in the dental clinic is not yet available. A case report suggested the spread of SARS-CoV-2 virus particles via droplet transmission prompted by air-conditioned ventilation (Lu et al., 2020). Therefore, potentially infected air should not be transported to people in the vicinity of the clinic.
- Some procedures do not require direct patient contact, for example scheduling of appointments. Indicating at which distance interaction is recognised as safe may be considered or installing physical barriers at the front desk, for example clear partitions. The interior of the dental clinic should be assessed and if necessary rearranged to allow for maintaining a safe distance, for example rearranging the waiting area.

5.3 | Administrative controls: changing the way DHCWs organise their work

- The routing within the dental clinic should be arranged in such way that both DHCWs and patients are able to maintain distance from each other when DHCWs are not wearing PPE. Social distancing between DHCWs should also be maintained when not caring for a patient, for example when changing clothes or during breaks.

- Many infection control measures require changes in behaviour (Kretzer & Larson, 1998). Therefore, extra attention to (the behaviour of) the team is imperative and should be aimed at creating awareness to the adjusted procedures in order to prevent contamination between DHCWs. It is important to provide them with appropriate information, education and training and to provide sufficient resources to promote the behavioural changes.
- During dental treatment, the virus load in aerosols can be reduced by applying a leakproof rubber dam (Cochran, Miller, & Sheldrake, 1989; Rørslett Hardersen, Enersen, Kristoffersen, Ørstavik, & Sunde, 2019; Samaranayake, Reid, & Evans, 1989). The work field should be disinfected after the application of rubber dam. Furthermore, apart from reducing the microbial load from aerosols, rubber dam can also contribute to reducing splashes (Dahlke et al., 2012).
- Aerosol dispersion should be minimised by adjusting dental treatment procedures, for example by using hand instruments instead of water-cooled instruments or ultrasonic cleaning devices (Harrel, Barnes, & Rivera-Hidalgo, 1998). In addition, adequate saliva as well as aerosol extraction using high volume evacuation is important to minimise aerosol production (Devker et al., 2012; Narayana, Mohanty, Sreenath, & Vidhyadhari, 2016). Procedures that provoke gag reflexes or coughing should be avoided if possible (Meng et al., 2020a).
- Thirty minutes after aerosol formation, virus particles and bacteria can still be detected in the air of the treatment room (Bennett et al., 2000; Nikitin et al., 2014). Transmission to unprotected DHCWs in between treatments as well as to the next patient should be prevented. Alternatively, to waiting at least 30 min between patients, sufficient ventilation may be applied (more information under Engineering controls).
- Recent publications suggested that rinsing the oral cavity with hydrogen peroxide (1% H₂O₂) may be useful in reducing the risk of transmission of SARS-CoV-2 via aerosols (Ather et al., 2020; Peng et al., 2020). However, since the viral load is high in the throat, in the nose, on the tongue and in the saliva (Liu et al., 2011; Xu et al., 2020; Zou et al., 2020), the oral cavity will soon be recontaminated after rinsing. Povidine-iodine has been suggested to be useful for both oral and nasal disinfection against SARS-CoV-2 (Kirk-Bayley, Challacombe, Sunkaraneni, & Combes, 2020; Loftus, Dexter, Parra, & Brown, 2020; Rørslett Hardersen et al., 2019). A systematic literature review reported that rinsing with all kinds of other orally applied disinfectants reduces the microbiological load in aerosols generated during dental healthcare procedures (Marui et al., 2019), but it is unclear whether this reduction is clinically relevant for prevention of SARS-CoV-2 transmission. In vitro studies on chlorhexidine showed that it insufficiently inactivates SARS-CoV-2 (G. Kampf, Scheithauer, et al., 2020).

5.4 | Protection of the DHCW with PPE

- Since the respiratory tract is the main *portal of entry* of the virus, the respiratory tract should be shielded (Jin et al., 2020). Therefore,



the recommendation is to wear respiratory protection during aerosol generating procedures in patients infected with SARS-CoV-2 (WHO, 2020). These respiratory protective devices (filtering half masks: FFP-2/ N95/ KN95) filter particles significantly more effectively and have a better fit compared to regular medical face masks (type IIR, fluid resistant). PPE should protect the patient as well as fellow DHCWs against the micro-organisms exhaled by the user. Therefore, a mask with an exhalation valve should not be worn in dental health care, as it does not protect against splashes and respiratory micro-organisms from the user are released via the valve. It is essential that PPE complies with international standards for example European standard EN 149:2001 + A1: 2009 for respirators (British_Standards_Institution, 2011, 2019). In research on protection against fine particles, N95-equivalent respirators showed 9% total leakage, whereas for medical face masks leakage was 22%–35% (Steinle et al., 2018). It should be noted that medical face masks are designed to protect the patient against the exhaled air from the DHCW and do not protect DHCWs against aerosols. In a systematic review, the use of respirators was compared with medical face masks and was not associated with a lower risk of laboratory-confirmed influenza. The authors therefore suggested that these respirators should not be recommended for general public or non-high-risk medical staff, who are not in close contact with influenza patients or suspected patients (Long et al., 2020). Moreover, the effectiveness of respirators strongly depends on the proper intended use (Noti et al., 2012). The use of respirators significantly reduces the risks, but does not completely eliminate them. Clinical studies on the efficacy of masks in dentistry concerning virus protection have not been performed yet. The availability and prioritisation of PPE may influence which protection may be used within a dental clinic.

- The mucous membranes of the eyes are also a possible *portal of entry* (Adhikari et al., 2020). Therefore, goggles or a face shield should be used during treatment. The advantage of a face shield is its protection of mask from splashes (Lindsley, Noti, Blachere, Szalajda, & Beezhold, 2014).
- Transmission via surfaces like clothing can be prevented by careful behaviour (no touch) or by wearing a splash-proof long-sleeved apron over standard protective clothing. This apron should be considered contaminated after an aerosol generating treatment and should not be touched during treatment and should be discarded immediately after leaving the treatment room. All skin and other body parts left uncovered when wearing PPE should be carefully covered (wearing water resistant caps) or cleaned and/or disinfected afterwards (shoes, hair). An intact skin serves as a proper barrier against the SARS-CoV-2 virus, but can also serve as a vector for transmission. Hence, hygiene of DHCWs other than their hands is also required.

6 | FUTURE PERSPECTIVES FOR DENTAL HEALTH CARE

The first phase of the outbreak required providing only urgent dental health care. Gradually, elective dental health care can be

resumed, only on the condition of applying additional infection control measures combined with a risk assessment per patient. DHCWs and patients are at an increased risk. The modified procedures described in this paper aim towards a *maximal* effect instead of an *optimal* effect. These additional precautions have been effective in preventing new infections with SARS-CoV-2 when providing emergency dental care in China (Meng et al., 2020a, 2020b).

In Europe, several standard guidelines for infection control in dental health care have already been adapted to the SARS-CoV-2 pandemic. So far, these guidelines assume a reliable triage as well as a negligible number of asymptomatic or presymptomatic patients. Currently, insufficient original research is available on both the virus (e.g. on its infectivity and the minimal infection dose) and the specific risks of aerosol generating dental procedures. Given this uncertainty, the duty of care towards patients should be balanced with both the safety of DHCWs and patients, as well as the limited resources of both time and materials (especially respirators) (Kampf, Scheithauer, et al., 2020) and finally the economic consequences. In most countries, decisions regarding these issues are made by governmental institutions and are beyond the scope of this paper. However, the authors stress the importance on being open to all DHCWs to what extent the risks are unknown.

When gradually scaling down additional preventive measures, risks and feasibility should be carefully balanced. The pandemic, possibly followed by postpandemic outbreaks, will likely remain present for a prolonged period of time (Kissler, Tedijanto, Goldstein, Grad, & Lipsitch, 2020). Since it is impossible to perform social distancing in dental health care, it is required to continuously consider which precautions are needed, possibly for the coming years. Especially regarding the risk assessment of aerosol generated during dental health care, future research should focus on determining which measures are adequate for providing safe care.

ACKNOWLEDGEMENTS

The authors are very grateful to the many colleagues from the University Medical Center Groningen and the Academic Centre of Dentistry Amsterdam who provided us with advice while writing this article.

CONFLICT OF INTEREST

None to declare.

AUTHOR CONTRIBUTIONS

Catherine Volgenant: Conceptualization; Writing-original draft. **Ilona F. Persoon:** Conceptualization; Writing-original draft. **Rolf A.G. de Ruijter:** Conceptualization; Writing-review & editing. **J. J. de Soet:** Conceptualization; Writing-original draft.

ORCID

Catherine M. C. Volgenant  <https://orcid.org/0000-0002-4049-2914>



Ilona F. Persoon  <https://orcid.org/0000-0002-9773-3774>

J. J. (Hans) de Soet  <https://orcid.org/0000-0002-9964-0205>

REFERENCES

- Adhikari, S. P., Meng, S., Wu, Y. J., Mao, Y. P., Ye, R. X., Wang, Q. Z., ... Zhou, H. (2020). Epidemiology, causes, clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-19) during the early outbreak period: A scoping review. *Infectious Diseases of Poverty*, 9(1), 29. <https://doi.org/10.1186/s40249-020-00646-x>
- Ai, Z., & Melikov, A. K. (2018). Airborne spread of expiratory droplet nuclei between the occupants of indoor environments: A review. *Indoor Air*, 28(4), 500–524. <https://doi.org/10.1111/ina.12465>
- Alharbi, A., Alharbi, S., & Alqaidi, S. (2020). Guidelines for dental care provision during the COVID-19 pandemic. *The Saudi Dental Journal*, 32(4), 181–186. <https://doi.org/10.1016/j.sdentj.2020.04.001>
- Arons, M. M., Hatfield, K. M., Reddy, S. C., Kimball, A., James, A., Jacobs, J. R., ... Jernigan, J. A. (2020). Presymptomatic SARS-CoV-2 infections and transmission in a skilled nursing facility. *New England Journal of Medicine*, <https://doi.org/10.1056/NEJMoa2008457>
- Ashour, H. M., Elkhatib, W. F., Rahman, M. M., & Elshabrawy, H. A. (2020). Insights into the recent 2019 novel coronavirus (SARS-CoV-2) in light of past human coronavirus outbreaks. *Pathogens*, 9(3), 186. Retrieved from <https://www.mdpi.com/2076-0817/9/3/186> <https://doi.org/10.3390/pathogens9030186>
- Ather, A., Patel, B., Ruparel, N. B., Diogenes, A., & Hargreaves, K. M. (2020). Coronavirus disease 19 (COVID-19): Implications for clinical dental care. *Journal of Endodontics*, 46(5), 584–595. <https://doi.org/10.1016/j.joen.2020.03.008>
- Ayebare, R. R., Flick, R., Okware, S., Bodo, B., & Lamorde, M. (2020). Adoption of COVID-19 triage strategies for low-income settings. *The Lancet Respiratory Medicine*, 8(4), e22. [https://doi.org/10.1016/S2213-2600\(20\)30114-4](https://doi.org/10.1016/S2213-2600(20)30114-4)
- Bai, Y., Yao, L., Wei, T., Tian, F., Jin, D. Y., Chen, L., & Wang, M. (2020). Presumed asymptomatic carrier transmission of COVID-19. *JAMA*, 323(14), 1406. <https://doi.org/10.1001/jama.2020.2565>
- Bennett, A. M., Fulford, M. R., Walker, J. T., Bradshaw, D. J., Martin, M. V., & Marsh, P. D. (2000). Microbial aerosols in general dental practice. *British Dental Journal*, 189(12), 664–667. <https://doi.org/10.1038/sj.bdj.4800859>
- Bischoff, W. E., Swett, K., Leng, I., & Peters, T. R. (2013). Exposure to influenza virus aerosols during routine patient care. *The Journal of Infectious Diseases*, 207(7), 1037–1046. <https://doi.org/10.1093/infdis/jis773>
- Blocken, B., Malizia, F., van Druenen, T., & Marchal, T. (2020). *Towards aerodynamically equivalent COVID19 1.5 m social distancing for walking and running*. Eindhoven, the Netherlands: Eindhoven University of Technology. Retrieved from <http://www.urbanphysics.net/COVID19.html> (version 21 April 2020).
- British_Standards_Institution. (2011). 149: 2001+ A1: 2009 Respiratory protective devices. Filtering half masks to protect against particles. Requirements, testing, marking. London, UK: British Standards Institution (BSI).
- British_Standards_Institution. (2019). 14683:2019 Medical face masks. Requirements and test methods. London, UK: British Standards Institution (BSI).
- Chan, J.-F.-W., Yuan, S., Kok, K.-H., To, K.-K.-W., Chu, H., Yang, J., ... Yuen, K.-Y. (2020). A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: A study of a family cluster. *The Lancet*, 395(10223), 514–523. [https://doi.org/10.1016/S0140-6736\(20\)30154-9](https://doi.org/10.1016/S0140-6736(20)30154-9)
- Chen, J. (2020). Pathogenicity and transmissibility of 2019-nCoV-A quick overview and comparison with other emerging viruses. *Microbes and Infection*, 22(2), 69–71. <https://doi.org/10.1016/j.micinf.2020.01.004>
- Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F., Han, Y., ... Zhang, L. (2020). Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: A descriptive study. *The Lancet*, 395(10223), 507–513. [https://doi.org/10.1016/S0140-6736\(20\)30211-7](https://doi.org/10.1016/S0140-6736(20)30211-7)
- Cheng, H.-Y., Jian, S.-W., Liu, D.-P., Ng, T.-C., Huang, W.-T., & Lin, H.-H. (2020). High transmissibility of COVID-19 near symptom onset. *medRxiv*. <https://doi.org/10.1101/2020.03.18.20034561>
- Cheong, K. W. D., & Phua, S. Y. (2006). Development of ventilation design strategy for effective removal of pollutant in the isolation room of a hospital. *Building and Environment*, 41(9), 1161–1170. <https://doi.org/10.1016/j.buildenv.2005.05.007>
- Cochran, M. A., Miller, C. H., & Sheldrake, M. A. (1989). The efficacy of the rubber dam as a barrier to the spread of microorganisms during dental treatment. *The Journal of the American Dental Association*, 119(1), 141–144. <https://doi.org/10.14219/jada.archive.1989.0131>
- Dahlke, W. O., Cottam, M. R., Herring, M. C., Leavitt, J. M., Ditmyer, M. M., & Walker, R. S. (2012). Evaluation of the spatter-reduction effectiveness of two dry-field isolation techniques. *The Journal of the American Dental Association*, 143(11), 1199–1204. <https://doi.org/10.14219/jada.archive.2012.0064>
- Darwish, S. (2020). COVID-19 considerations in dental care. *Dental Update*, 47(4), 287–302. <https://doi.org/10.12968/denu.2020.47.4.287>
- Devker, N., Mohitey, J., Vibhute, A., Chouhan, V. S., Chavan, P., Malagi, S., & Joseph, R. (2012). A study to evaluate and compare the efficacy of preprocedural mouthrinsing and high volume evacuator attachment alone and in combination in reducing the amount of viable aerosols produced during ultrasonic scaling procedure. *The Journal of Contemporary Dental Practice*, 13(5), 681–689. <https://doi.org/10.5005/jp-journals-10024-1209>
- Du, Z., Xu, X., Wu, Y., Wang, L., Cowling, B. J., & Meyers, L. A. (2020). The serial interval of COVID-19 from publicly reported confirmed cases. *Emerging Infectious Diseases*, 26(6). <http://dx.doi.org/10.3201/eid2606.200357>
- Elder, N. C., Sawyer, W., Pallerla, H., Khaja, S., & Blacker, M. (2014). Hand hygiene and face touching in family medicine offices: A Cincinnati Area Research and Improvement Group (CARInG) network study. *The Journal of the American Board of Family Medicine*, 27(3), 339–346. <https://doi.org/10.3122/jabfm.2014.03.130242>
- Escombe, A. R., Ticona, E., Chávez-Pérez, V., Espinoza, M., & Moore, D. A. J. (2019). Improving natural ventilation in hospital waiting and consulting rooms to reduce nosocomial tuberculosis transmission risk in a low resource setting. *BMC Infectious Diseases*, 19(1), 88. <https://doi.org/10.1186/s12879-019-3717-9>
- Farooq, I., & Ali, S. (2020). COVID-19 outbreak and its monetary implications for dental practices, hospitals and healthcare workers. *Postgraduate Medical Journal*, postgradmedj-2020-137781, <https://doi.org/10.1136/postgradmedj-2020-137781>
- Ferretti, L., Wymant, C., Kendall, M., Zhao, L., Nurtay, A., Abeler-Dörner, L., ... Fraser, C. (2020). Quantifying SARS-CoV-2 transmission suggests epidemic control with digital contact tracing. *Science*, 368(6491), eabb6936. <https://doi.org/10.1126/science.abb6936>
- Guo, H., Zhou, Y., Liu, X., & Tan, J. (2020). The impact of the COVID-19 epidemic on the utilization of emergency dental services. *Journal of Dental Sciences*, <https://doi.org/10.1016/j.jds.2020.02.002>
- Guo, J., Wu, H., & Xie, H. (2020). Letter to the Editor: How to deal with suspended oral treatment during the COVID-19 epidemic. *Journal of Dental Research*, 0022034520920169. <https://doi.org/10.1177/0022034520920169>
- Harrel, S. K., Barnes, J. B., & Rivera-Hidalgo, F. (1998). Aerosol and splatter contamination from the operative site during ultrasonic scaling.



- The Journal of the American Dental Association*, 129(9), 1241–1249. <https://doi.org/10.14219/jada.archive.1998.0421>
- He, X., Lau, E. H. Y., Wu, P., Deng, X., Wang, J., Hao, X., ... Leung, G. M. (2020). Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nature Medicine*, 26(5), 672–675. <http://dx.doi.org/10.1038/s41591-020-0869-5>
- Heesterbeek, J., & Dietz, K. (1996). The concept of R_0 in epidemic theory. *Statistica Neerlandica*, 50(1), 89–110.
- Hoehl, S., Rabenau, H., Berger, A., Kortenbusch, M., Cinatl, J., Bojkova, D., ... Naujoks, F. (2020). Evidence of SARS-CoV-2 infection in returning travelers from Wuhan, China. *New England Journal of Medicine*, 382(13), 1278–1280. <https://doi.org/10.1056/NEJMc2001899>
- Hu, Z., Song, C., Xu, C., Jin, G., Chen, Y., Xu, X., ... Zheng, Y. (2020). Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Science China Life Sciences*, 63(5), 706–711.
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., ... Cao, B. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*, 395(10223), 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5)
- Huang, R., Xia, J., Chen, Y., Shan, C., & Wu, C. (2020). A family cluster of SARS-CoV-2 infection involving 11 patients in Nanjing, China. *The Lancet Infectious Diseases*, 20(5), 534–535. [https://doi.org/10.1016/S1473-3099\(20\)30147-X](https://doi.org/10.1016/S1473-3099(20)30147-X)
- Izzetti, R., Nisi, M., Gabriele, M., & Graziani, F. (2020). COVID-19 transmission in dental practice: Brief review of preventive measures in Italy. *Journal of Dental Research*, 0022034520920580. <https://doi.org/10.1177/0022034520920580>
- Jin, Y., Yang, H., Ji, W., Wu, W., Chen, S., Zhang, W., & Duan, G. (2020). Virology, epidemiology, pathogenesis, and control of COVID-19. *Viruses*, 12(4), 372. <https://doi.org/10.3390/v12040372>
- Jones, R. M., & Brosseau, L. M. (2015). Aerosol transmission of infectious disease. *Journal of Occupational and Environmental Medicine*, 57(5), 501–508. <https://doi.org/10.1097/jom.0000000000000448>
- Kampf, G., Scheithauer, S., Lemmen, S., Saliou, P., & Suchoemel, M. (2020). COVID-19-associated shortage of alcohol-based hand rubs, face masks, medical gloves and gowns; proposal for a risk-adapted approach to ensure patient and healthcare worker safety. *Journal of Hospital Infection*. <https://doi.org/10.1016/j.jhin.2020.04.041>
- Kampf, G., Todt, D., Pfaender, S., & Steinmann, E. (2020). Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *Journal of Hospital Infection*, 104(3), 246–251. <https://doi.org/10.1016/j.jhin.2020.01.022>
- Khan, S., Siddique, R., Shereen, M. A., Ali, A., Liu, J., Bai, Q., ... Xue, M. (2020). The emergence of a novel coronavirus (SARS-CoV-2), their biology and therapeutic options. *Journal of Clinical Microbiology*, JCM.00187-00120. <https://doi.org/10.1128/jcm.00187-20>
- Khurshid, Z., Asiri, F. Y. I., & Al Wadaani, H. (2020). Human saliva: Non-invasive fluid for detecting novel coronavirus (2019-nCoV). *International Journal of Environmental Research and Public Health*, 17(7), 2225. <https://doi.org/10.3390/ijerph17072225>
- Kimball, A., Hatfield, K. M., Arons, M., James, A., Taylor, J., Spicer, K., ... Zane, S. (2020). Asymptomatic and presymptomatic SARS-CoV-2 infections in residents of a long-term care skilled nursing facility – King County, Washington, March 2020. *MMWR. Morbidity and Mortality Weekly Report*, 69(13), 377–381. <https://doi.org/10.15585/mmwr.mm6913e1>
- Kirk-Bayley, J., Challacombe, S., Sunkaraneni, S., & Combes, J. (2020). The use of povidone iodine nasal spray and mouthwash during the current COVID-19 pandemic may protect healthcare workers and reduce cross infection. *SSRN Electronic Journal*, <https://doi.org/10.2139/ssrn.3563092>
- Kissler, S. M., Tedijanto, C., Goldstein, E., Grad, Y. H., & Lipsitch, M. (2020). Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. *Science*, eabb5793. <https://doi.org/10.1126/science.abb5793>
- Kluytmans, M., Buiting, A., Bentvelsen, R., van den Bijllaardt, W., van Oudheusden, A., van Rijen, M., ... Kluytmans, J. (2020). SARS-CoV-2 infection in 86 healthcare workers in two Dutch hospitals in March 2020. *medRxiv*, 2020.2003.2023.20041913. <https://doi.org/10.1101/2020.03.23.20041913>
- Kohn, W. G., Collins, A. S., Cleveland, J. L., Harte, J. A., Eklund, K. J., & Malvitz, D. M. (2003). Guidelines for infection control in dental health-care settings-2003. Retrieved from <https://stacks.cdc.gov/view/cdc/6743>
- Kratzel, A., Todt, D., V'kovski, P., Steiner, S., Gultom, M., Thao, T. T. N., ... Pfaender, S. (2020). Inactivation of severe acute respiratory syndrome coronavirus 2 by WHO-recommended hand rub formulations and alcohols. *Emerging Infectious Diseases*, 26(7). <https://doi.org/10.3201/eid2607.200915>
- Kretzer, E. K., & Larson, E. L. (1998). Behavioral interventions to improve infection control practices. *American Journal of Infection Control*, 26(3), 245–253. [https://doi.org/10.1016/S0196-6553\(98\)80008-4](https://doi.org/10.1016/S0196-6553(98)80008-4)
- Lai, C. C., Liu, Y. H., Wang, C.-Y., Wang, Y.-H., Hsueh, S.-C., Yen, M.-Y., ... Hsueh, P.-R. (2020). Asymptomatic carrier state, acute respiratory disease, and pneumonia due to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2): Facts and myths. *Journal of Microbiology, Immunology and Infection*, <https://doi.org/10.1016/j.jmii.2020.02.012>
- Lauer, S. A., Grantz, K. H., Bi, Q., Jones, F. K., Zheng, Q., Meredith, H. R., ... Lessler, J. (2020). The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: Estimation and application. *Annals of Internal Medicine*, 172(9), 577. <https://doi.org/10.7326/M20-0504>
- Leung, N. H. L., Chu, D. K. W., Shiu, E. Y. C., Chan, K.-H., McDevitt, J. J., Hau, B. J. P., ... Cowling, B. J. (2020). Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nature Medicine*, 26(5), 676–680. <http://dx.doi.org/10.1038/s41591-020-0843-2>
- Lindsley, W. G., Noti, J. D., Blachere, F. M., Szalajda, J. V., & Beezhold, D. H. (2014). Efficacy of face shields against cough aerosol droplets from a cough simulator. *Journal of Occupational and Environmental Hygiene*, 11(8), 509–518. <https://doi.org/10.1080/15459624.2013.877591>
- Liu, L., Wei, Q., Alvarez, X., Wang, H., Du, Y., Zhu, H., ... Chen, Z. (2011). Epithelial cells lining salivary gland ducts are early target cells of severe acute respiratory syndrome coronavirus infection in the upper respiratory tracts of rhesus macaques. *Journal of Virology*, 85(8), 4025–4030. <https://doi.org/10.1128/JVI.02292-10>
- Liu, Y., Gayle, A. A., Wilder-Smith, A., & Rocklöv, J. (2020). The reproductive number of COVID-19 is higher compared to SARS coronavirus. *Journal of Travel Medicine*, 27(2), taaa021. <https://doi.org/10.1093/jtm/taaa021>
- Liu, Y., Ning, Z., Chen, Y., Guo, M., Liu, Y., Gali, N. K., ... Lan, K. (2020). Aerodynamic analysis of SARS-CoV-2 in two Wuhan hospitals. *Nature*, <https://doi.org/10.1038/s41586-020-2271-3>
- Liu, Y., Ning, Z., Chen, Y., Guo, M., Liu, Y., Gali, N. K., ... Westerdahl, D. (2020). Aerodynamic characteristics and RNA concentration of SARS-CoV-2 aerosol in Wuhan hospitals during COVID-19 outbreak. *bioRxiv*. <https://doi.org/10.1101/2020.03.08.982637>
- Loftus, R. W., Dexter, F., Parra, M. C., & Brown, J. R. (2020). Importance of oral and nasal decontamination for patients undergoing anesthesia during the COVID-19 era. Publish Ahead of Print. *Anesthesia & Analgesia*. Retrieved from https://journals.lww.com/anesthesia-analgesia/Fulltext/9000/Importance_of_oral_and_nasal_decontamination_for.95698.aspx <https://doi.org/10.1213/ANE.0000000000004854>
- Long, Y., Hu, T., Liu, L., Chen, R., Guo, Q., Yang, L., ... Du, L. (2020). Effectiveness of N95 respirators versus surgical masks against influenza: A systematic review and meta-analysis. *Journal of Evidence-Based Medicine*, n/a (n/a). <https://doi.org/10.1111/jebm.12381>



- Lotfinejad, N., Peters, A., & Pittet, D. (2020). Hand hygiene and the novel coronavirus pandemic: The role of healthcare workers. *Journal of Hospital Infection*, <https://doi.org/10.1016/j.jhin.2020.03.017>.
- Lu, J., Gu, J., Li, K., Xu, C., Su, W., Lai, Z., ... Yang, Z. (2020). COVID-19 outbreak associated with air conditioning in Restaurant, Guangzhou, China, 2020. *Emerging Infectious Diseases*, *26*(7), <https://doi.org/10.3201/eid2607.200764>
- Luo, S.-H., Liu, W., Liu, Z.-J., Zheng, X.-Y., Hong, C.-X., Liu, Z.-R., ... Weng, J.-P. (2020). A confirmed asymptomatic carrier of 2019 novel coronavirus (SARS-CoV-2). *Chinese Medical Journal*, *133*(9), 1123–1125. <https://doi.org/10.1097/cm9.0000000000000798>.
- Lynch, C., Mahida, N., Oppenheim, B., & Gray, J. (2020). Washing our hands of the problem. *Journal of Hospital Infection*, *104*(4), 401–403. <http://dx.doi.org/10.1016/j.jhin.2020.03.010>
- Mallineni, S. K., Innes, N. P., Raggio, D. P., Araujo, M. P., Robertson, M. D., & Jayaraman, J. (2020). Coronavirus disease (COVID-19): Characteristics in children and considerations for dentists providing their care. *International Journal of Paediatric Dentistry*, *30*(3), 245–250. <https://doi.org/10.1111/ipd.12653>
- Marui, V. C., Souto, M. L. S., Rovai, E. S., Romito, G. A., Chambrone, L., & Pannuti, C. M. (2019). Efficacy of preprocedural mouthrinses in the reduction of microorganisms in aerosol: A systematic review. *The Journal of the American Dental Association*, *150*(12), 1015–1026. <https://doi.org/10.1016/j.adaj.2019.06.024>
- Meng, L., Hua, F., & Bian, Z. (2020a). Coronavirus Disease 2019 (COVID-19): Emerging and future challenges for dental and oral medicine. *Journal of Dental Research*, *99*(5), 481–487. <https://doi.org/10.1177/0022034520914246>
- Meng, L., Hua, F., & Bian, Z. (2020b). Response to the Letter to the Editor: How to deal with suspended oral treatment during the COVID-19 epidemic. *Journal of Dental Research*, <https://doi.org/10.1177/0022034520920166>.
- Monaghan, N. P. (2016). Emerging infections – implications for dental care. *British Dental Journal*, *221*(1), 13–15. <https://doi.org/10.1038/sj.bdj.2016.486>.
- Munster, V. J., Koopmans, M., van Doremalen, N., van Riel, D., & de Wit, E. (2020). A novel coronavirus emerging in China—key questions for impact assessment. *New England Journal of Medicine*, *382*(8), 692–694. <https://doi.org/10.1056/NEJMp2000929>
- Narayana, T. V., Mohanty, L., Sreenath, G., & Vidhyadhari, P. (2016). Role of preprocedural rinse and high volume evacuator in reducing bacterial contamination in bioaerosols. *Journal of Oral and Maxillofacial Pathology : JOMFP*, *20*(1), 59–65. <https://doi.org/10.4103/0973-029X.180931>.
- Ngoc, N. M., That, B. T. T., Hong, N. T. T., Dung, N. T. P., Thanh, T. T., Man, D. N. H., ... Thwaites, G. (2020). Duration of viral detection in throat and rectum of a patient with COVID-19. *medRxiv*. <https://doi.org/10.1101/2020.03.07.20032052>
- Nicolaiades, C., Avraam, D., Cueto-Felgueroso, L., González, M. C., & Juanes, R. (2020). Hand-hygiene mitigation strategies against global disease spreading through the air transportation network. *Risk Analysis*, *40*(4), 723–740. <https://doi.org/10.1111/risa.13438>
- Nikitin, N., Petrova, E., Trifonova, E., & Karpova, O. (2014). Influenza virus aerosols in the air and their infectiousness. *Advances in Virology*, *2014*, 1–6. <https://doi.org/10.1155/2014/859090>
- Noti, J. D., Lindsley, W. G., Blachere, F. M., Cao, G., Kashon, M. L., Thewlis, R. E., ... Beezhold, D. H. (2012). Detection of infectious influenza virus in cough aerosols generated in a simulated patient examination room. *Clinical Infectious Diseases*, *54*(11), 1569–1577. <https://doi.org/10.1093/cid/cis237>
- Ong, S. W. X., Tan, Y. K., Chia, P. Y., Lee, T. H., Ng, O. T., Wong, M. S. Y., & Marimuthu, K. (2020). Air, surface environmental, and personal protective equipment contamination by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) From a symptomatic patient. *JAMA*, *323*(16), 1610. <https://doi.org/10.1001/jama.2020.3227>
- Pan, X., Chen, D., Xia, Y., Wu, X., Li, T., Ou, X., ... Liu, J. (2020). Asymptomatic cases in a family cluster with SARS-CoV-2 infection. *The Lancet Infectious Diseases*, *20*(4), 410–411. [https://doi.org/10.1016/S1473-3099\(20\)30114-6](https://doi.org/10.1016/S1473-3099(20)30114-6)
- Peng, X., Xu, X., Li, Y., Cheng, L., Zhou, X., & Ren, B. (2020). Transmission routes of 2019-nCoV and controls in dental practice. *International Journal of Oral Science*, *12*(1), 9. <https://doi.org/10.1038/s41368-020-0075-9>.
- Prati, C., Pelliccioni, G. A., Sambri, V., Chersoni, S., & Gandolfi, M. G. (2020). COVID-19: Its impact on dental schools in Italy, clinical problems in endodontic therapy and general considerations. *International Endodontic Journal*, *53*(5), 723–725. <https://doi.org/10.1111/iej.13291>.
- Rabi, F. A., Al Zoubi, M. S., Kasasbeh, G. A., Salameh, D. M., & Al-Nasser, A. D. (2020). SARS-CoV-2 and coronavirus disease 2019: What we know so far. *Pathogens*, *9*(3), 231. Retrieved from <https://www.mdpi.com/2076-0817/9/3/231> <https://doi.org/10.3390/pathogens9030231>
- Rajgor, D. D., Lee, M. H., Archuleta, S., Bagdasarian, N., & Quek, S. C. (2020). The many estimates of the COVID-19 case fatality rate. *The Lancet Infectious Diseases*. [https://doi.org/10.1016/S1473-3099\(20\)30244-9](https://doi.org/10.1016/S1473-3099(20)30244-9)
- Ran, L., Chen, X., Wang, Y., Wu, W., Zhang, L., & Tan, X. (2020). Risk factors of healthcare workers with corona virus disease 2019: A retrospective cohort study in a designated hospital of Wuhan in China. *Clinical Infectious Diseases*, <https://doi.org/10.1093/cid/ciaa287>
- Rørslett Hardersen, L., Enersen, M., Kristoffersen, A. K., Ørstavik, D., & Sunde, P. T. (2019). Maintenance of the aseptic working field during endodontic treatment. *Acta Odontologica Scandinavica*, *77*(7), 502–507. <https://doi.org/10.1080/00016357.2019.1606935>
- Rothe, C., Schunk, M., Sothmann, P., Bretzel, G., Froeschl, G., Wallrauch, C., ... Hoelscher, M. (2020). Transmission of 2019-nCoV infection from an asymptomatic contact in Germany. *New England Journal of Medicine*, *382*(10), 970–971. <https://doi.org/10.1056/NEJMc2001468>
- Sabino-Silva, R., Jardim, A. C. G., & Siqueira, W. L. (2020). Coronavirus COVID-19 impacts to dentistry and potential salivary diagnosis. *Clinical Oral Investigations*, *24*(4), 1619–1621. <https://doi.org/10.1007/s00784-020-03248-x>
- Samaranayake, L. P., Reid, J., & Evans, D. (1989). The efficacy of rubber dam isolation in reducing atmospheric bacterial contamination. *ASDC Journal of Dentistry for Children*, *56*(6), 442–444. Retrieved from <http://europepmc.org/abstract/MED/2681303>
- Siegel, J. D., Rhinehart, E., Jackson, M., Chiarello, L., & Committee, H. C. I. C. P. A. (2007). 2007 guideline for isolation precautions: Preventing transmission of infectious agents in health care settings. *American Journal of Infection Control*, *35*(10), S65. <https://doi.org/10.1016/j.ajic.2007.10.007>
- Smales, F. C., & Samaranayake, L. P. (2003). Maintaining dental education and specialist dental care during an outbreak of a new coronavirus infection. Part 2: Control of the disease, then elimination. *British Dental Journal*, *195*(12), 679–681. <https://doi.org/10.1038/sj.bdj.4810819>
- Spagnuolo, G., De Vito, D., Rengo, S., & Tatullo, M. (2020). COVID-19 Outbreak: An overview on dentistry. *International Journal of Environmental Research and Public Health*, *17*(6), 2094. Retrieved from <https://www.mdpi.com/1660-4601/17/6/2094> <https://doi.org/10.3390/ijerph17062094>
- Steinle, S., Sleeuwenhoek, A., Mueller, W., Horwell, C. J., Apsley, A., Davis, A., ... Galea, K. S. (2018). The effectiveness of respiratory protection worn by communities to protect from volcanic ash inhalation. Part II: Total inward leakage tests. *International Journal of Hygiene and Environmental Health*, *221*(6), 977–984. <https://doi.org/10.1016/j.ijheh.2018.03.011>.
- Stockwell, R. E., Ballard, E. L., O'Rourke, P., Knibbs, L. D., Morawska, L., & Bell, S. C. (2019). Indoor hospital air and the impact of ventilation on

- bioaerosols: A systematic review. *Journal of Hospital Infection*, 103(2), 175–184. <https://doi.org/10.1016/j.jhin.2019.06.016>.
- Sun, J., He, W. T., Wang, L., Lai, A., Ji, X., Zhai, X., ... Su, S. (2020). COVID-19: Epidemiology, evolution, and cross-disciplinary perspectives. *Trends in Molecular Medicine*, 26(5), 483–495. <https://doi.org/10.1016/j.molmed.2020.02.008>
- The National Institute for Occupational Safety and Health (NIOSH). (2015). Hierarchy of controls. Retrieved from <https://www.cdc.gov/niosh/topics/hierarchy/default.html>
- To, K.-K.-W., Tsang, O.-T.-Y., Yip, C.-C.-Y., Chan, K.-H., Wu, T.-C., Chan, J.-M.-C., ... Yuen, K.-Y. (2020). Consistent detection of 2019 novel coronavirus in saliva. *Clinical Infectious Diseases*, <https://doi.org/10.1093/cid/ciaa149>.
- Tong, Z. D., Tang, A., Li, K. F., Li, P., Wang, H. L., Yi, J. P., ... Yan, J. B. (2020). Potential presymptomatic transmission of SARS-CoV-2, Zhejiang Province, China, 2020. *Emerging Infectious Diseases*, 26(5), 1052–1054. <https://doi.org/10.3201/eid2605.200198>
- van Doremalen, N., Bushmaker, T., Morris, D. H., Holbrook, M. G., Gamble, A., Williamson, B. N., ... Munster, V. J. (2020). Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *New England Journal of Medicine*, 382(16), 1564–1567. <http://dx.doi.org/10.1056/nejmc2004973>
- Verity, R., Okell, L. C., Dorigatti, I., Winskill, P., Whittaker, C., Imai, N., ... Ferguson, N. M. (2020). Estimates of the severity of coronavirus disease 2019: A model-based analysis. *The Lancet Infectious Diseases*. [http://dx.doi.org/10.1016/s1473-3099\(20\)30243-7](http://dx.doi.org/10.1016/s1473-3099(20)30243-7)
- Volgenant, C., & de Soet, J. (2018). Cross-transmission in the dental office: Does this make you ill? *Current Oral Health Reports*, 5(4), 221–228. <https://doi.org/10.1007/s40496-018-0201-3>
- Wax, R. S., & Christian, M. D. (2020). Practical recommendations for critical care and anesthesiology teams caring for novel coronavirus (2019-nCoV) patients. *Canadian Journal of Anesthesia/Journal Canadien D'anesthésie*, 67(5), 568–576. <https://doi.org/10.1007/s12630-020-01591-x>
- WHO (2020). *Coronavirus*. Retrieved from https://www.who.int/health-topics/coronavirus#tab=tab_1
- Xu, H., Zhong, L., Deng, J., Peng, J., Dan, H., Zeng, X., ... Chen, Q. (2020). High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *International Journal of Oral Science*, 12(1), 8. <https://doi.org/10.1038/s41368-020-0074-x>
- Yu, P., Zhu, J., Zhang, Z., & Han, Y. (2020). A familial cluster of infection associated with the 2019 novel coronavirus indicating possible person-to-person transmission during the incubation period. *The Journal of Infectious Diseases*, 221(11), 1757–1761. <https://doi.org/10.1093/infdis/jiaa077>
- Zemouri, C., de Soet, H., Crielaard, W., & Laheij, A. (2017). A scoping review on bio-aerosols in healthcare and the dental environment. *PLoS ONE*, 12(5), e0178007. <https://doi.org/10.1371/journal.pone.0178007>.
- Zemouri, C., Volgenant, C. M. C., Buijs, M. J., Crielaard, W., Rosema, N. A. M., Brandt, B. W., ... de Soet, J. J. (2020). Dental aerosols: Microbial composition and spatial distribution. *Journal of Oral Microbiology*, 12(1), 1762040. <https://doi.org/10.1080/20002297.2020.1762040>
- Zhang, Z., Zhang, L., & Wang, Y. (2020). COVID-19 indirect contact transmission through the oral mucosa must not be ignored. *Journal of Oral Pathology & Medicine*, 49(5), 450–451. <https://doi.org/10.1111/jop.13019>
- Zou, J., Ruan, F., Huang, M., Liang, L., Huang, H., Hong, Z., ... Wu, J. (2020). SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *New England Journal of Medicine*, 382, 970–971. <https://doi.org/10.1056/NEJMc2001737>

How to cite this article: Volgenant CMC, Persoon IF, de Ruijter RAG, de Soet JJ(H). Infection control in dental health care during and after the SARS-CoV-2 outbreak. *Oral Dis*. 2021;27(Suppl. 3):674–683. <https://doi.org/10.1111/odi.13408>