

University of Groningen

Response to "Letter to the Editor Regarding

Brooks, Anne B.; Langer, Jakob; Tervonen, Tommi; Peter Hemmingsen, Mads; Eguchi, Kosei; Bacci, Elizabeth D.

Published in:
Diabetes Technol Ther

DOI:
[10.1007/s13300-020-00900-3](https://doi.org/10.1007/s13300-020-00900-3)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Brooks, A. B., Langer, J., Tervonen, T., Peter Hemmingsen, M., Eguchi, K., & Bacci, E. D. (2020). Response to "Letter to the Editor Regarding: Patient Preferences for Glucagon-like Peptide-1 (GLP-1) Receptor Agonist Treatment of Type 2 Diabetes Mellitus in Japan: A Discrete Choice Experiment". *Diabetes Technol Ther*, 11(10), 2443-2446. <https://doi.org/10.1007/s13300-020-00900-3>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Response to “Letter to the Editor Regarding: Patient Preferences for Glucagon-like Peptide-1 (GLP-1) Receptor Agonist Treatment of Type 2 Diabetes Mellitus in Japan: A Discrete Choice Experiment”

Anne B. Brooks · Jakob Langer · Tommi Tervonen · Mads Peter Hemmingsen ·
Kosei Eguchi · Elizabeth D. Bacci

Received: July 21, 2020
© The Author(s) 2020

Keywords: GLP-1 receptor agonist; Patient preference study; Treatment; Type 2 diabetes

Dear Editor,

We would like to thank you for the opportunity to respond to the issues raised in the letter related to our publication [1] and to provide details of the methodology to address the concerns. In the letter, the author noted concerns about the cardiovascular (CV) outcome risk reduction value used for the semaglutide 0.50 mg profile. The author also requested

clarification and disclosure of the references related to attribute levels.

The author is correct that the CV outcome risk reduction for the semaglutide 0.50 mg profile (26% versus placebo) was based on combined data for the 0.50 and 1.0 mg doses reported in the primary publication of SUSTAIN-6 by Marso et al. [2]. This was in accordance with the primary outcome of the study and to support noninferiority and superiority testing. As presented in the appendix of Marso et al., and noted in the letter, CV risk reduction was not significant for either dose independently (23% [$p = 0.13$] for semaglutide 0.50 mg and 29% [$p = 0.06$] for semaglutide 1.0 mg) [2], which was expected because the study was not powered or intended to assess the doses separately.

Digital Features To view digital features for this article go to <https://doi.org/10.6084/m9.figshare.12728447>.

A. B. Brooks
Patient-Centered Research, Evidera, Bethesda, MD,
USA

J. Langer
Market Access & External Affairs, Novo Nordisk
Pharma Ltd., Tokyo, Japan

T. Tervonen
Patient-Centered Research, Evidera, London, UK

T. Tervonen
Department of Epidemiology, University Medical
Center Groningen, University of Groningen,
Groningen, The Netherlands

M. Peter Hemmingsen
External Valuation, Novo Nordisk A/S, Søborg,
Denmark

K. Eguchi
Medical Affairs, Novo Nordisk Pharma Ltd., Tokyo,
Japan

E. D. Bacci (✉)
Patient-Centered Research, Evidera, Seattle, WA,
USA
e-mail: elizabeth.bacci@evidera.com

Table 1 Attributes, levels, and sources for the discrete choice experiment

Attribute	Level	Represents	Reference
Method of administration	Multi-dose prefilled pen, used with disposable injection needles, with dose adjustment possible	Semaglutide 0.50 mg	Ozempic [®] prescribing information at the time of study conduct
	Single-dose, disposable prefilled pen, with no dose adjustment possible	Dulaglutide 0.75 mg	Trulicity [®] prescribing information at the time of study conduct
HbA1c change	On average, patients achieve a 1.9% reduction in HbA1c level	Semaglutide 0.50 mg	Seino et al. [6]
	On average, patients achieve a 1.6% reduction in HbA1c level	Intermediate level	Not applicable
	On average, patients achieve a 1.4% reduction in HbA1c level	Dulaglutide 0.75 mg	Miyagawa et al. [7]
CV risk reduction	26% reduction of risk in cardiovascular diseases (heart attack, stroke, death due to cardiovascular diseases)	Semaglutide 0.50 mg	Marso et al. [2]
	No data for the benefit or risk in cardiovascular diseases (heart attack, stroke, death due to cardiovascular diseases)	Dulaglutide 0.75 mg	None available
Weight change	On average, patients have a 2.2 kg weight loss	Semaglutide 0.50 mg	Seino et al. [6]
	On average, patients have a 1.1 kg weight loss	Intermediate level	Not applicable
	On average, patients do not have any weight loss	Dulaglutide 0.75 mg	Miyagawa et al. [7]
Common side effects	On average, 1 out of 9 patients will experience transient nausea	Semaglutide 0.50 mg	Seino et al. [6]
	On average, 1 out of 12 patients will experience transient nausea	Intermediate level	Not applicable
	On average, 1 out of 19 patients will experience transient nausea	Dulaglutide 0.75 mg	Miyagawa et al. [7]

By contrast, providing the uncertainty (95% confidence intervals) around the predicted choice probability for the semaglutide 0.50 mg profile versus the dulaglutide 0.75 mg profile would have strengthened our conclusions and might have alleviated the author's concern. Uncertainty around estimates from patient preference studies should be provided to help

interpret results and inform patient-centered benefit–risk assessments [3]. The confidence interval around the predicted choice probability was small (78% [95% confidence interval, 74–82%]), supporting the conclusion that the large majority of participants preferred the semaglutide 0.50 mg profile.

To address the concern of the author about using a 26% CV risk reduction for the semaglutide 0.50 mg profile, we conducted an additional sensitivity analysis for the predicted choice probability. Using a 23% CV risk reduction, the predicted choice probability was 76% (95% confidence interval, 71–80%) in favor of the semaglutide 0.50 mg profile, which is close to the original value and supports the robustness and validity of our original conclusion.

Additional relevant data have been published since the discrete choice experiment was performed. SUSTAIN-7, a head-to-head randomized clinical trial, showed that hemoglobin A1c (HbA1c) and body weight were reduced significantly more with semaglutide 0.50 mg than with dulaglutide 0.75 mg [4]. This was confirmed in a network meta-analysis among patients with type 2 diabetes mellitus in Japan [5]. The REWIND study showed that dulaglutide 1.5 mg reduces cardiovascular risk compared to placebo, although this dosage is still not currently approved in Japan.

Finally, to address the request for clarification and disclosure of the references related to attribute levels, we provide them as Table 1.

Respectfully,
Anne Brooks, BS
Jakob Langer, MS
Tommi Tervonen, PhD
Mads Peter Hemmingsen, MD
Kosei Eguchi, MD, PhD
Elizabeth Dansie Bacci, PhD

ACKNOWLEDGEMENTS

Funding. Novo Nordisk Pharma Ltd. provided the funding for this study and the journal's Rapid Service Fee. Evidera received funding from Novo Nordisk Pharma Ltd. to participate in the study and develop this letter. All authors had full access to all of the data in this study and take complete responsibility for the integrity of the data and accuracy of the data analysis. No Rapid Service Fee was received by the journal for the publication of this article.

Medical Writing Assistance. The authors would like to thank Philip Leventhal, PhD at Evidera for his assistance preparing this letter, which was funded by Novo Nordisk.

Authorship. All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship for this article, take responsibility for the integrity of the work as a whole, and have given their approval for this version to be published.

Disclosures. Jakob Langer is employed by Novo Nordisk Pharma Ltd. Mads Peter Hemmingsen is employed by Novo Nordisk A/S. Kosei Eguchi is employed by Novo Nordisk Pharma Ltd. Anne Brooks is employed by Evidera, which provides consulting and other research services to pharmaceutical, medical device, and related organizations. In this salaried position, Anne Brooks works with a variety of companies and organizations and is precluded from receiving payment or honoraria directly from these organizations for services rendered. Tommi Tervonen is employed by Evidera, which provides consulting and other research services to pharmaceutical, medical device, and related organizations. In this salaried position, Tommi Tervonen works with a variety of companies and organizations, and is precluded from receiving payment or honoraria directly from these organizations for services rendered. Elizabeth Dansie Bacci is employed by Evidera, which provides consulting and other research services to pharmaceutical, medical device, and related organizations. In this salaried position, Elizabeth Dansie Bacci works with a variety of companies and organizations and is precluded from receiving payment or honoraria directly from these organizations for services rendered.

Compliance with Ethics Guidelines. This article is based on previously conducted studies and does not contain any studies with human participants or animals performed by any of the authors.

Peer Review. Please note, contrary to the journal's standard single-blind peer review

process, as a letter this article underwent review by a member of the journal's editorial board.

Open Access. This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License, which permits any non-commercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

1. Brooks A, Langer J, Tervonen T, Hemmingsen MP, Eguchi K, Bacci ED. Patient preferences for GLP-1 receptor agonist treatment of type 2 diabetes mellitus in japan: a discrete choice experiment. *Diabetes Ther.* 2019;10:735–49. <https://doi.org/10.1007/s13300-019-0591-9>.
2. Marso SP, Bain SC, Consoli A, et al. Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med.* 2016;375:1834–44. <https://doi.org/10.1056/NEJMoa1607141>.
3. Mott DJ, Chami N, Tervonen T. Reporting quality of marginal rates of substitution in discrete choice experiments that elicit patient preferences. *Value Health.* 2020. <https://doi.org/10.1016/j.jval.2020.04.1831>.
4. Pratley RE, Aroda VR, Lingvay I, et al. Semaglutide versus dulaglutide once weekly in patients with type 2 diabetes (SUSTAIN 7): a randomised, open-label, phase 3b trial. *Lancet Diabetes Endocrinol.* 2018;6:275–86. [https://doi.org/10.1016/S2213-8587\(18\)30024-X](https://doi.org/10.1016/S2213-8587(18)30024-X).
5. Webb N, Orme M, Witkowski M, Nakanishi R, Langer J. A network meta-analysis comparing semaglutide once-weekly with other GLP-1 receptor agonists in Japanese patients with type 2 diabetes. *Diabetes Ther.* 2018;9:973–86. <https://doi.org/10.1007/s13300-018-0397-1>.
6. Seino Y, Terauchi Y, Osonoi T, et al. Safety and efficacy of semaglutide once weekly vs sitagliptin once daily, both as monotherapy in Japanese people with type 2 diabetes. *Diabetes Obes Metab.* 2018;20:378–88. <https://doi.org/10.1111/dom.13082>.
7. Miyagawa J, Odawara M, Takamura T, Iwamoto N, Takita Y, Imaoka T. Once-weekly glucagon-like peptide-1 receptor agonist dulaglutide is non-inferior to once-daily liraglutide and superior to placebo in Japanese patients with type 2 diabetes: a 26-week randomized phase III study. *Diabetes Obes Metab.* 2015;17:974–83. <https://doi.org/10.1111/dom.12534>.