

University of Groningen

Intestinal stenosis in Crohn's disease shows a generalized upregulation of genes involved in collagen metabolism and recognition that could serve as novel anti-fibrotic drug targets

van Haaften, Wouter Tobias; Blokzijl, Tjasso; Hofker, Hendrik Sijbrand; Olinga, Peter; Dijkstra, Gerard; Bank, Ruud A.; Boersema, Miriam

Published in:
Therapeutic advances in gastroenterology

DOI:
[10.1177/1756284820952578](https://doi.org/10.1177/1756284820952578)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

van Haaften, W. T., Blokzijl, T., Hofker, H. S., Olinga, P., Dijkstra, G., Bank, R. A., & Boersema, M. (2020). Intestinal stenosis in Crohn's disease shows a generalized upregulation of genes involved in collagen metabolism and recognition that could serve as novel anti-fibrotic drug targets. *Therapeutic advances in gastroenterology*, 13. <https://doi.org/10.1177/1756284820952578>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Intestinal stenosis in Crohn's disease shows a generalized upregulation of genes involved in collagen metabolism and recognition that could serve as novel anti-fibrotic drug targets

Wouter Tobias van Haften, Tjasso Blokzijl, Hendrik Sijbrand Hofker, Peter Olinga^{ID}, Gerard Dijkstra, Ruud A. Bank and Miriam Boersema

Abstract

Background and Aims: Crohn's disease (CD) can be complicated by intestinal fibrosis. Pharmacological therapies against intestinal fibrosis are not available. The aim of this study was to determine whether pathways involved in collagen metabolism are upregulated in intestinal fibrosis, and to discuss which drugs might be suitable to inhibit excessive extracellular matrix formation targeting these pathways.

Methods: Human fibrotic and non-fibrotic terminal ileum was obtained from patients with CD undergoing ileocecal resection due to stenosis. Genes involved in collagen metabolism were analyzed using a microfluidic low-density TaqMan array. A literature search was performed to find potential anti-fibrotic drugs that target proteins/enzymes involved in collagen synthesis, its degradation and its recognition.

Results: mRNA expression of collagen type I (*COL1A1*, 0.76 ± 0.28 versus 37.82 ± 49.85 , $p=0.02$) and III (*COL3A1*, 2.01 ± 2.61 versus 68.65 ± 84.07 , $p=0.02$) was increased in fibrotic CD compared with non-fibrotic CD. mRNA expression of proteins involved in both intra- and extracellular post-translational modification of collagens (prolyl- and lysyl hydroxylases, lysyl oxidases, chaperones), collagen-degrading enzymes (MMPs and cathepsin-K), and collagen receptors were upregulated in the fibrosis-affected part. A literature search on the upregulated genes revealed several potential anti-fibrotic drugs.

Conclusion: Expression of genes involved in collagen metabolism in intestinal fibrosis affected terminal ileum of patients with CD reveals a plethora of drug targets. Inhibition of post-translational modification and altering collagen metabolism might attenuate fibrosis formation in the intestine in CD. Which compound has the highest potential depends on a combination anti-fibrotic efficacy and safety, especially since some of the enzymes play key roles in the physiology of collagen.

Keywords: anti-fibrotic drugs, Crohn's disease, intestinal fibrosis

Received: 17 January 2020; revised manuscript accepted: 31 July 2020.

Introduction

Fibrosis in any organ is the result of chronic injury, leading to a disturbed balance in the formation and degradation of an extracellular matrix (ECM) rich in collagen.¹ Crohn's disease (CD) is an inflammatory

bowel disease showing a heterogeneous phenotype. The phenotype, which can change over time, can be classified according to the Montreal classification in non-stricturing/non-penetrating (B1), stricturing (B2), and penetrating disease (B3).²⁻⁶ A stricturing

Ther Adv Gastroenterol

2020, Vol. 13: 1–17

DOI: 10.1177/
1756284820952578

© The Author(s), 2020.
Article reuse guidelines:
[sagepub.com/journals-](https://sagepub.com/journals-permissions)
permissions

Correspondence to:

Peter Olinga
Department of
Pharmaceutical
Technology and
Biopharmacy, University
of Groningen, Ant.
Deusinglaan 1, Groningen,
9713 AV, the Netherlands
p.olinga@rug.nl

Wouter Tobias van Haften
Department of
Gastroenterology and
Hepatology, University
Medical Center Groningen,
University of Groningen,
Groningen, the Netherlands

Department of
Pharmaceutical Technology
and Biopharmacy,
University of Groningen,
Groningen, the Netherlands

Tjasso Blokzijl
Department of Laboratory
Medicine, University of
Groningen, University
Medical Center Groningen,
Groningen, The
Netherlands

Hendrik Sijbrand Hofker
Department of Surgery,
University Medical Center
Groningen, University of
Groningen, Groningen, the
Netherlands

Gerard Dijkstra
Department of
Gastroenterology and
Hepatology, University
Medical Center
Groningen, University of
Groningen, Groningen, the
Netherlands

Ruud A. Bank
Department of Pathology
and Medical Biology,
University Medical Center
Groningen, University of
Groningen, Groningen, the
Netherlands

Miriam Boersema
Department of
Pharmaceutical
Technology and
Biopharmacy, University of
Groningen, Groningen, the
Netherlands

phenotype, which is characterized by intestinal fibrosis, occurs mainly in CD (70%). However, it can also occur in ulcerative colitis (UC; 1.5–11.2%), upon radiation injury, or upon chronic allograft dysfunction after intestinal transplantation.^{7–11} In CD, thickening of the intestinal wall causes symptomatic fibrotic stenosis due to narrowing of the lumen, which requires surgery. In UC, it will lead to thickening and shortening of the colon.¹² The mechanisms involved in transmural intestinal fibrosis may be comparable with those of pathological collagen accumulation in other organs, and drugs tested for fibrosis in other organs might be applicable to intestinal fibrosis as well. So far, despite the tremendous amount of research performed on renal and hepatic fibrosis, pharmacological treatments have become available only for fibrotic conditions such as idiopathic pulmonary fibrosis.^{13,14} No pharmacological therapies against intestinal fibrosis are available so far.

The interstitial matrix of the intestine consists of the fibrillary collagens type I, III and V, and in addition collagen type VI.¹⁵ The non-fibrillar collagen type IV is the main component of the basement membrane, which creates the barrier between the epithelium on the gut luminal side and the lamina propria of the intestine.¹⁶ In addition, intestinal ECM (as ECM of any other organ) consists of fibronectin (FN1) and presumably of elastin, as well as the proteoglycans decorin, biglycan (BGN), and fibromodulin.^{17,18} An increase, especially in the amount of interstitial collagens, is accompanied with thickening and stiffening of the intestinal wall, thereby causing stenosis due to the luminal stricture of the intestine.¹⁹ Net deposited ECM is the result of a complex balance between factors involved in collagen synthesis (including post-transcriptional modification) *versus* collagen degradation. Even though fibrogenesis in the intestine is presumably similar to fibrosis in other organs, the expression of genes involved in collagen homeostasis has not been evaluated before.

Generally, formation of collagen starts with transcription of procollagen mRNA in the nucleus, leading to synthesis on ribosomes of three polypeptide α -chains that are released into the endoplasmic reticulum for post-transcriptional modification.^{20,21} Within the endoplasmic reticulum, certain lysine and proline residues from the α -chains are hydroxylated by lysyl hydroxylases (LHs) and prolyl 3- and 4- hydroxylases (P3H, P4H), respectively. Some of the hydroxylysine residues are subsequently glycosylated by collagen glycosyltransferases. During hydroxylation and

glycosylation, the three procollagen-chains are assembled and further stabilized by formation of intra- and inter-molecular disulfide bonds.²⁰ Triple helix formation requires the aid of chaperones like heat shock protein 47 (HSP47) and FK506 binding protein 10.^{22,23} Procollagens are then transported out of the cell *via* the Golgi apparatus. In the extracellular space, the N- and C-terminal propeptides are enzymatically cleaved off [by enzymes from the “a Disintegrin and Metalloproteinase with Thrombospondin motifs” (ADAMTS, N-terminal) family and by Bone Morphogenetic Protein 1 (BMP1, C-terminal)]. Subsequently, collagens are assembled into fibrils and cross-linking is induced by lysyl oxidases/lysyl oxidase-like (LOX, LOXL) enzymes. Degradation of collagens occurs mostly *via* matrix metalloproteinases (MMPs). MMPs are Zn²⁺-dependent endopeptidases that can degrade a plethora of ECM proteins, including collagen.²⁴ Degradation products of ECM can have chemotactic properties and MMPs are able to activate or degrade several non-ECM substrates like cytokines/chemokines or growth factors. The MMPs thereby play a central role in ECM remodeling as well as in intestinal inflammation. MMPs 1, 2, 3, and 9 activity is elevated in the active mucosal inflammation of both patients with CD and UC, and the balance between MMPs and tissue inhibitors of MMPs (TIMPs) is altered in inflammatory bowel disease (IBD).^{25,26}

Candidate drugs against intestinal fibrosis mostly target pathways of (intestinal) fibrosis such as the mitogen-activated protein kinase (MAPK), Rho-associated protein kinase (ROCK), and transforming growth factor β (TGF- β) pathways.^{27,28} However, genes involved in the assembly of the ECM also comprise targets that can inhibit ECM formation or alter its molecular structure in such a way that ECM will be degraded faster.²⁹ Here, we show an upregulation of mRNA expression of genes involved in collagen metabolism in intestinal fibrosis affected terminal ileum of patients with CD. These results reveal possible drug targets, which are reviewed in this paper.

Methods

Fibrosis affects terminal ileum from patients with CD and from non-fibrotic control patients

Fibrotic and non-fibrotic terminal ileum from patients with CD undergoing ileocecal (re-)resection because of stenosis was obtained. All included

patients with CD had purely stricturing phenotype (Montreal B2, Table 1). The fibrotic/stenotic and the non-fibrosis affected (resection margin) regions were identified macroscopically during surgery. Non-fibrotic non-CD affected tissue was obtained from patients undergoing right-sided hemicolectomy because of an adenocarcinoma (non-cancer affected ileal resection margin, Table 1). All tissue obtained was freshly collected and stored by a laboratory technician in the operating room immediately after resection. Samples were fixed in Tissue-Tek® (O.C.T. Compound, Sakura® Finetek) in the operation room, and frozen in isopentane on dry ice before being stored at -80°C until further use.

Isolation of RNA

To isolate RNA, 10 Tissue-Tek sections (10 μm thick) containing the full thickness (verified by hematoxylin and eosin staining) of the intestinal wall, were cut using a cryostat. Sections were dissolved in TRIzol (Invitrogen, Life Technologies, Carlsbad, CA, USA), after which total RNA was isolated according to the manufacturer's protocol. To avoid genomic DNA contamination, samples were treated with DNase I, Amp Grade (Invitrogen) according to the manufacturer's protocol.

Reverse transcription and TaqMan® gene expression assays

Equal amounts of RNA were reverse transcribed using the Reverse Transcription System (Promega, Madison, WI, USA). Subsequently, complementary DNA was used for quantitative real-time polymerase chain reaction (RT-qPCR) in a microfluidic card-based low-density TaqMan array (Applied Biosystems, Foster City, CA, USA), which enables simultaneous measurement of mRNA expression of 44 genes we selected (Supplemental Table S2). RT-qPCR was performed by loading 100 ng of cDNA per sample using the ViiA™ 7 Real-Time PCR System (Applied Biosystems). The following settings were used: 50°C for 2 min, 95°C for 10 min, and the next two steps were repeated for 50 cycles: 95°C for 12 s and 60°C for 1 min. Threshold cycle numbers >40 were excluded from the analysis. Patients/pairs were removed from the analysis if there was no detectable expression ($\text{Ct} > 40$) in either one of the pairs. Delta-Ct values were calculated using *GAPDH* as a reference gene. Expression in the figures is presented as $2^{-\Delta\text{Ct}}$ on a logarithmic scale.

Table 1. Characteristics of the patients with CD and controls of which terminal ileum was obtained.

	Control (n=4)	CD (n=7)
General		
Gender, % female	2 (50%)	7 (100%)
Age at surgery, years (mean, min-max)	73.1 (69.1–78.2)	33.6 (21.1–54.5)
Disease duration, years, (mean, min-max)	4 (100%)	6.4 (1.8–16.0)
Montreal age at diagnosis [n (%)]		
17–40 years (A2)	NA	6 (85.7%)
>40 years (A3)		1 (14.3%)
Montreal disease behavior [n (%)]		
Stricturing disease (B2)	NA	7 (100%)
Disease location [n (%)]		
Terminal ileum (L1)	NA	4 (57.1%)
Ileocolon (L3)		3 (42.9%)
C-reactive protein before operation [n (%)]		
C-reactive protein > 5 mg/l	NA	1 (14.3%)
C-reactive protein < 5 mg/l		5 (71.4%)
Missing		1 (14.3%)
Clinical disease activity based on HBI before operation [n (%)]		
Disease in remission (HBI < 5)	NA	0 (0%)
Mild disease (HBI 5–7)		0 (0%)
Moderate disease (HBI 8–16)		3 (42.9%)
Severe disease (HBI > 16)		4 (57.1%)
Medication [n (%)]		
Corticosteroids	NA	4 (57.1%)
Thiopurines		4 (57.1%)
Anti-TNF α		1 (14.3%)
Anti-IL12/23		1 (14.3%)
CD, Crohn's disease; HBI, Harvey Bradshaw Index.		

Statistical analysis

Data were statistically analyzed and visualized with graphs using GraphPad Prism software (v6.0). All data was considered to be non-parametric. A Wilcoxon paired signed rank test was used to compare analysis of fibrotic *versus* non-fibrotic ileum from the one patient with CD for gene expression as well as quantification of immunohistochemistry (IHC). A Mann-Whitney *U* test was used to

compare with CD-affected non-fibrotic ileum to non-CD-affected non-fibrotic ileum, and to compare age at surgery. Differences were considered significant at a p value of <0.05 . Median interquartile range (IQR) values are presented in the text and figures.

Literature search

A literature search was performed to find (potential) drugs that target the proteins/enzymes transcribed from genes involved in collagen fibril synthesis and degradation, detected in CD patients with stricturing phenotype by a microfluidic card-based low-density TaqMan array. A comprehensive literature search was conducted to identify relevant drugs. The electronic exploration involved keyword searches in Pubmed. The following search criteria were used (all fields) (“gene name” or “protein name”) and (“inhibitor”, “antagonist” or “agonist”). Targeting drugs tested *in vivo/in silico*, *in vivo* in animals or *in vivo* in humans are listed separately in Table 2.

Ethical considerations

Patients gave written informed consent for anonymous use of patient data and resected parts of human intestine according to the code of conduct for responsible use of surgical left-over material (See: “Code goed gebruik voor gecodeerd lichaamsmateriaal”, Research Code University Medical Center Groningen, <http://www.rug.nl/umcg/research/documents/research-code-info-umcg-nl.pdf>).

Results

Cohort characteristics

This descriptive cohort study included seven patients with fibro-stenotic CD who underwent ileocecal resection and four patients with adenocarcinoma who underwent right-sided hemicolectomy. All patients with CD had a stricturing disease phenotype and had either ileal [$n=4$ (57.1%)] or ileocolonic [$n=3$ (42.9%)] disease. On average, they were 33.6 years old (range 21.1–54.5) and had suffered from CD for 6.4 years (range 1.8–16.0). All patients had clinically active disease before they underwent ileocecal resection [moderate disease $n=3$ (42.9%), severe disease $n=4$ (57.1%)], and they used several different anti-inflammatory drugs before surgery (Table 1). As controls, patients who right-sided hemicolectomy due to adenocarcinoma were included at a

mean age of 73.1 years (range 69.1–78.2). These patients were significantly older than the patients with CD ($p=0.008$). All included patients with CD were female.

Expression of fibrosis markers is increased in macroscopically fibrosis-affected terminal ileum

Using a microfluidic card-based low-density TaqMan array, mRNA expression of a variety of ECM proteins were investigated. mRNA expression of procollagens type I, III, IV, V and IV was increased in fibrotic CD compared with non-fibrotic CD. Especially expression of collagen type I (*COL1A1*, 0.76 ± 0.28 versus 37.82 ± 49.85 , $p=0.02$) and III (*COL3A1*, 2.01 ± 2.61 versus 68.65 ± 84.07 , $p=0.02$) was increased in the fibrosis and the CD affected part compared with non-fibrotic CD (Figure 1A). Upregulation of these procollagens was considered as a positive control for the correct selection of the fibrosis-affected region. mRNA expression of other ECM proteins such as elastin (ELN, 0.01 ± 0.02 versus 0.07 ± 0.10 , $p=0.03$), FN1 (0.53 ± 0.43 versus 2.50 ± 10.52 , $p=0.02$), and BGN (0.1 ± 0.12 versus 2.83 ± 4.46 , $p=0.03$) was also increased in fibrotic CD compared with non-fibrotic CD (Figure 1B). mRNA expression of alpha-smooth muscle actin (generally considered as a marker for myofibroblasts), was also elevated in the fibrosis and CD-affected region compared with non-fibrotic CD (*ACTA2*, 1.03 ± 3.98 versus 16.57 ± 47.31 , $p=0.02$, Figure 1C).

Expression of intra- and extra-cellular modification of collagen fibrils is increased in macroscopically fibrosis-affected terminal ileum

Enzymes involved in intracellular post-translational modification of the collagen fibril, were also upregulated in fibrotic CD compared with non-fibrotic CD. Expression of lysyl hydroxylases 1–3 (*PLOD1*, 0.05 ± 0.06 versus 0.40 ± 0.38 , $p=0.02$; *PLOD2*, 0.23 ± 0.13 versus 0.58 ± 0.69 , $p=0.05$; *PLOD3*, 0.06 ± 0.01 versus 0.29 ± 0.40 , $p=0.02$), prolyl 4-hydroxylases (*P4HA1*, 0.37 ± 0.43 versus 1.04 ± 2.10 , $p=0.02$); *P4HB*, 2.37 ± 3.72 versus 9.48 ± 14.54 , $p=0.02$) and prolyl-3-hydroxylases 1–3 (*P3H1*, 0.022 ± 0.037 versus 0.39 ± 0.54 , $p=0.03$; *P3H2*, 0.05 ± 0.27 versus 0.33 ± 1.77 , $p=0.03$; *P3H3*, 0.03 ± 0.16 versus 0.39 ± 0.91 , $p=0.02$) was increased in the fibrosis- and

Table 2. Possible drug targets in collagen metabolism in intestinal fibrosis.

Gene	Protein	Targeting drugs tested <i>in vitro/in silico</i>	Targeting drugs tested <i>in vivo</i> in animals	Targeting drugs tested <i>in vivo</i> in human
Synthesis				
Procollagens 1-6 including alpha 1-3 helices	Collagen	Pre-transcriptional acting anti-fibrotic drugs	Pre-transcriptional acting anti-fibrotic drugs	Pre-transcriptional acting anti-fibrotic drugs
Intra-cellular post-translational modifications				
Procollagen lysyl hydroxylases (<i>PLOD1-3</i>)	Lysyl hydroxylases 1-3	- Lysyl hydroxylase 1-3 inhibition by Minoxidil ³⁰	- Lysyl hydroxylase 1-3 inhibition by Minoxidil ³¹	- Minoxidil is an FDA-approved anti-hypertensive agent and registered for topical use to treat alopecia
Prolyl 4-Hydroxylases (<i>P4HA1-A3</i>)	Prolyl 4-Hydroxylases	- 1,4 dihydrophenanthroline-4-one-3-carboxylic acid and 8-(N-butyl-N-ethylcarbamoyl)-1,4-dihydrophenanthroline-4-one-3-carboxylic acid ³² - Short-hairpin RNAs (shP4HA2-1 and shP4HA2-2) ³³	- 1,4 dihydrophenanthroline-4-one-3-carboxylic acid and 8-(N-butyl-N-ethylcarbamoyl)-1,4-dihydrophenanthroline-4-one-3-carboxylic acid ³²	- Not known
Prolyl 3-hydroxylases (<i>P3H1, 2</i>)	Prolyl 3-hydroxylases (Leprecans)	- Not known	- Not known	- Not known
Collagen Beta(1-0) Galactosyltransferase 1 (<i>COLGALT1</i>)	Procollagen galactosyltransferase 1	- Carminic acid ³⁴	- Not known	- Not known
Intra-cellular assembly of the triple helix				
Serpin Family H Member 1 (<i>SERP1H1</i>)	Heat shock protein 47	- AK778 and its cleavage product Co1003 ³⁵ - HSP47 small interfering RNA (siRNA) ³⁶ - Four Small Molecule Chemical Inhibitors ³⁷	- HSP47 small interfering RNA (siRNA) ³⁸⁻⁴¹	- Not known
FKBP Prolyl Isomerase 10 (<i>FKBP10</i>)	FK506 Binding Protein 10	- siRNA mediated knockdown ^{42,43} - Peptidyl prolyl isomerase inhibition by tacrolimus ^{22,44}	- Tacrolimus ⁴⁴⁻⁴⁶	- Tacrolimus is used as immunosuppressant for several indications, was not tested as anti-fibrotic drug in human so far ⁴⁷
Extracellular cleavage of propeptides				
A Disintegrin and Metalloproteinase with Thrombospondin motifs (<i>ADAMTS</i>)	A Disintegrin and Metalloproteinase with Thrombospondin motifs	- ADAMTS-2; TIMP-3 ⁴⁸	Not known	Not known

(Continued)

Table 2. (Continued)

Gene	Protein	Targeting drugs tested <i>in vitro/in silico</i>	Targeting drugs tested <i>in vivo</i> in animals	Targeting drugs tested <i>in vivo</i> in human
Bone Morphogenetic Protein 1 (<i>BMP1</i>)	Bone Morphogenetic Protein 1	- α 2-Macroglobulin ⁴⁹ - Acidic dipeptide hydroxamate ⁵⁰	Not known	Not known
Procollagen C-Endopeptidase Enhancer 1-2 (<i>PCOLCE/PCOLCE2</i>)	Procollagen C-Endopeptidase Enhancer	Not known	Not known	Not known
Assembly into collagen fibrils by crosslinking of collagens				
Lysyl oxidases (<i>LOX, LOXL1-4</i>)	Lysyl oxidases	- β -aminopropionitrile (β -APN) ^{51,52} - LOX inhibitory monoclonal antibody ⁵³ - LOXL2 with an inhibitory monoclonal antibody (AB0023) ^{54,55} - LOXL2 [2-chloropyridin-4-yl]methanamine ⁵⁶	- LOX inhibitory monoclonal ab/M64 ^{53,54} - LOXL2 inhibitory monoclonal antibody (AB0023, humanized variant AB0024) ^{54,55} - PXS-S2B, a small-molecule selective LOXL2 inhibitor - PAT-1251, a small-molecule selective LOXL2 inhibitor	- Simtuzumab (anti-LOXL-2 monoclonal antibody) was tested in phase II trials for primary sclerosing cholangitis, NASH induced liver-fibrosis, idiopathic pulmonary fibrosis, Second-Line Treatment of Metastatic KRAS Mutant Colorectal Adenocarcinoma and metastatic pancreatic adenocarcinoma ⁵⁷⁻⁶⁰
Collagen degrading enzymes				
Matrix metalloproteinases (<i>MMP</i>)	Matrix metalloproteinases	Not relevant as this will not work anti-fibrotic		
Tissue inhibitors of Matrix metalloproteinases (<i>TIMP1-4</i>)	Tissue inhibitors of Matrix metalloproteinases	- Several genetic deletion studies performed, no pharmacological inhibitors are known ^{61,62}	- Several genetic deletion studies performed, no pharmacological inhibitors are known ^{61,62}	- Not Known
Cathepsin K (<i>CTSK</i>)	Cathepsin K	Not relevant as this will not work anti-fibrotic		
Collagen receptors				
DDR tyrosine Kinase 1 and 2 (<i>DDR1, 2</i>)	Discoidin Domain Receptor Tyrosine Kinase 1 and 2	- Pyrazolopyrimidine derivatives ⁶³ - DDR1-IN-1 ⁶⁴ - Actinomycin D ⁶⁵ - Monoclonal antibodies against DDR1 ⁶⁶ -Dasatinib/imatinitib/nilotinib ^{67,68}	- Dasatinib, ^{69,70} but is also reported to lung vascular toxicity and predisposes to pulmonary hypertension ⁷¹	- Actinomycin D is clinically used as anti-tumor antibiotic ⁷² - Dasatinib/imatinitib/nilotinib are currently used to treat chronic myeloid leukemia, as well as several fibrotic conditions
Mannose Receptor C Type 2 (<i>MRC2</i>)	Mannose Receptor C Type 2	- Not Known	- Not Known	- Not Known
FDA, United States Food and Drug Administration.				




Figure 1. mRNA expression of procollagens 1–6 (A), of extracellular matrix molecules *BGN*, *DCN*, *ELN*, *FMOD*, *FN1* (B), and of *ACTA2* in fibrotic versus non-fibrotic terminal ileum of patients with CD, and versus non-CD control terminal ileum. Significant differences are depicted as: * $p < 0.05$, ** $p < 0.01$, ns: not significant. Marker levels are presented as median. *ACTA2*, alpha-actin-2; *BGN*, biglycan; CD, Crohn's disease; *DCN*, decorin; *ELN*, elastin; *FMOD*, fibromodulin; *FN1*, fibronectin.

CD-affected region compared with non-fibrotic CD (Figure 2A–C). Expression of *P4HA2* and *P4HA3* was not detectable. Expression of chaperones HSP47 and FK506 binding protein 10 (*SERPINH1*, 0.06 ± 0.02 versus 0.68 ± 0.62 , $p = 0.03$; *FKBP10*, 0.04 ± 0.07 versus 0.51 ± 0.35 , $p = 0.031$) was also increased in fibrotic CD compared with non-fibrotic CD (Figure 2D). mRNA expression of the enzymes related to cleavage of the N- and C- terminal propeptides (*ADAMTS2*, 0.06 ± 0.08 versus 0.27 ± 0.38 , $p = 0.03$; *ADAMTS14*, 0.01 ± 0.01 versus 0.01 ± 0.06 , $p = 0.06$) and bone morphogenetic protein 1 (*BMP1*, 0.07 ± 0.05 versus 0.30 ± 1.26 , $p = 0.031$) was increased in fibrotic CD compared with non-fibrotic CD. Also, the expression of collagen receptors discoidin domain receptor tyrosine kinase 2 (*DDR2*, 0.17 ± 0.09 versus 0.49 ± 1.72 , $p = 0.016$) and of mannose receptor C type 2 (*MRC2*, 0.07 ± 0.06 versus 0.36 ± 0.80 , $p = 0.016$) was increased in the fibrosis- and CD-affected tissue compared with non-fibrotic CD.

Expression of MMPs and their tissue inhibitors is increased in macroscopically fibrosis-affected terminal ileum

Net deposition of collagen depends on the balance between formation and degradation. Collagens are

degraded by matrix-metalloproteinases, which are inhibited by tissue inhibitors of MMPs. *MMP1* (0.02 ± 0.24 versus 1.22 ± 0.88 , $p = 0.031$) and *MMP14* (0.16 ± 0.21 versus 1.51 ± 2.95 , $p = 0.031$) was upregulated in fibrotic CD compared with non-fibrotic CD, as well as *TIMP1* (0.85 ± 3.17 versus 22.78 ± 47.61 , $p = 0.016$).

No differences in expression of the genes described in this study involved in collagen metabolism was observed between non-CD non-fibrotic control tissue and non-fibrotic CD (Figures 1–4 and Supplemental Table S1).

Discussion

To our knowledge, this is the first study examining the expression of genes coding for enzymes involved in the metabolism (such as post-translational modifications) of collagens in intestinal fibrosis in CD. We aimed to reveal a gene signature with potential targets for drugs against intestinal fibrosis based on gene expression of CD-affected fibrotic versus non-fibrotic terminal ileum from the same patient using a microfluidic card-based low-density TaqMan array. A literature search revealed several drugs that interfere with the metabolism of collagens that could be candidates for drug treatment against intestinal




Figure 2. mRNA expression of lysyl hydroxylases 1–3 (*PLOD1–3*) (A), of prolyl 4-hydroxylases (*P4HA1* and *P4HB*) (B), of prolyl 3-hydroxylases (*P3H1*, *P3H2*, *P3H3*) (C), and of heat shock protein 47 (*SERPINH1*) and of FK506 binding protein 10 (*FKBP10*) (D) in fibrotic versus non-fibrotic terminal ileum of patients with CD, and versus non-CD control terminal ileum. Significant differences are depicted as: * $p < 0.05$, ** $p < 0.01$, ns: not significant. Marker levels are presented as median. CD, Crohn's disease.

fibrosis (Table 2). Some of these drugs are already used clinically for other (fibrotic) conditions, whereas some have been tested only either *in vivo* in animals or *in vitro* in cell culture models. Others have been tested only in biochemical models such as binding assays to determine biochemical half-maximal binding concentrations or were identified using screening technologies for small molecule discovery. Outside of the scope of this study, comprehensive reviews and studies are available on pharmacological inhibition of (pre-transcriptional/translational) pathways leading to a lowered production of ECM.^{73,74}

The expression of enzymes involved in the intracellular and extracellular post-translational modifications of collagens (see Table 2) has never been described for intestinal fibrosis in CD. However, expression of several collagens and post-translational modulators of collagens was assessed in colorectal-cancer-associated fibrosis using comparative liquid chromatography with mass spectrometry.⁷⁵ In a study by Afik *et al.*, *PLOD1–3* and *P4HA1* protein expression was upregulated

in colorectal cancer-associated fibrosis compared with more distal non-fibrosis-affected colon tissue, which is in line with our results. This study also reports on increased protein expression of BGN and FN1 in colorectal cancer-associated fibrosis, which is in line with our results.⁷⁵ Upregulation of *PLOD1–3* mRNA was also observed in fibrotic conditions such as idiopathic pulmonary fibrosis.³¹ Furthermore, upregulation of *LOXL-2* has been reported in renal fibrosis and inhibition of *LOXL-2* by several inhibitors in mice *in vivo* successfully reduced the expression of fibrosis markers in several studies.^{76,77} A relative increase in protein expression of collagen type III over type I in fibrostenotic CD compared with inflamed or non-disease affected intestinal tissue reported previously was not observed in this cohort at the mRNA level.^{78,79}

Intracellular post-translational modifications

Pharmacological inhibition of the intracellular post-translational modifications (lysyl hydroxylase, prolyl-3 and -4 hydroxylase and glycosyltransferase,




Figure 3. mRNA expression of *ADAMTS2*, *ADAMTS14* and *BMP1* (A), *PCOLCE2* (B), and LOXs *LOXL1–4* (C) in fibrotic versus non-fibrotic terminal ileum of patients with CD, and versus non-CD control terminal ileum. Significant differences are depicted as: * $p < 0.05$, ** $p < 0.01$, ns: not significant. Marker levels are presented as median. ADAMTS, A disintegrin and metalloproteinase with thrombospondin motifs; BMP1, bone morphogenic protein 1; CD, Crohn's disease; LOX, lysyl oxidase; PCOLCE2, procollagen C-endopeptidase enhancer 2.




Figure 4. mRNA expression of collagen degrading enzymes MMPs *MMP1*, *MMP14*, and *CTSK* (A), and of *TIMP1* (B) in fibrotic versus non-fibrotic terminal ileum of patients with CD, and versus non-CD control terminal ileum. Significant differences are depicted as: * $p < 0.05$, ** $p < 0.01$, ns: not significant. Marker levels are presented as median. CD, Crohn's disease; CTSK, cathepsin K; MMP, matrix metalloproteinase; TIMP1, tissue inhibitor of matrix metalloproteinase 1.

Table 2) could inhibit collagen formation and thereby fibrosis, but these enzymes are pivotal to human physiology. Therefore, the inhibition

should be very enzyme specific and ideally be targeted to the fibrosis-affected area in order to be effective without causing severe side effects. The

importance of these genes is confirmed by the fact that genetic mutations in the genes coding for these enzymes (resulting in aberrant synthesis, degradation and/or modification) can lead to syndromic disorders in musculoskeletal or connective tissues such as osteogenesis imperfecta type VIII (mutation in *P3H1*), Bruck syndrome type 2 (mutation in *PLOD2* resulting in a form of osteogenesis imperfecta), Ehlers-Danlos syndrome type VIA (mutation in *PLOD1*), and severe myopia in the absence of musculoskeletal abnormalities (*P3H2*).²⁹ No mutations are found in genes coding for the P4HA α -subunits, perhaps indicating that loss of PH4 enzyme function leads to premature death of the embryo.²⁹ Pharmacological inhibition of lysyl hydroxylases 1-3 by minoxidil upon stimulation with transforming growth factor beta (TGF- β) was tested *in vitro* in primary human fibroblasts by Zuurmond *et al.* Even though a concentration- and time-dependent reduction in *LHI-3* mRNA was observed, no effect on the total number of pyridinoline cross-links in the collagen matrix was observed, and the authors therefore conclude that minoxidil is unlikely to be anti-fibrotic in these concentrations.³⁰ Shao *et al.* did observe an anti-fibrotic effect of minoxidil in mice in bleomycin-induced pulmonary fibrosis without reporting side effects in these animals.³¹ The side effects of minoxidil when used systemically in humans are, however, severe, which makes further testing unattractive. However, specific inhibition of lysyl hydroxylase-2 (*PLOD-2*) could be attractive as reduction of the pyridinoline cross-links between collagen molecules facilitates easier degradation by endogenous proteinases (i.e., MMPs and cathepsin K). This because cross-linked collagen can be degraded effectively only by MMP-13 and cathepsin K, in contrast to non-crosslinked collagen, which can be degraded by several MMPs.²⁹ Inhibition of another hydroxylase, P4H, was tested by administering two small-molecular inhibitors (both Phenanthrolinones, see Table 2) to rats.³² These compounds were well tolerated in the rat at doses producing sustained inhibition of collagen hydroxylation. Procollagen molecules which are less hydroxylated will accumulate in the endoplasmic reticulum (ER), thereby causing ER-stress, which triggers ER-stress-mediated apoptosis of myofibroblasts.^{80,81} However, P4H inhibition would be effective as an anti-fibrotic target only if the inhibition is constantly maintained, as collagen will be rapidly hydroxylated after P4H activity is restored.³²

Intracellular assembly of the triple helix

Inhibition of the collagen chaperone HSP47 and the FK506 binding protein 10 has promising anti-fibrotic potential. HSP47 expression is upregulated in intestinal fibrosis.^{23,82-84} The anti-fibrotic potential of inhibition of HSP47 is shown by the deletion of *Hsp47* in hepatic stellate cells isolated from Cre-LoxP system *Hsp47* floxed mice, which led to ER stress-mediated apoptosis of the collagen-producing cells.⁸¹ Another study shows that local (submesothelial) delivery of *Hsp47* siRNA conjugated with cationized gelatin microspheres could suppresses peritoneal fibrosis in mice.³⁸ The use of microRNA and small siRNA for several indications *in vivo* in humans has progressed to several phase II and III clinical trials.⁸⁵ The activity of the other collagen chaperone FK506 binding protein 10 (peptidyl prolyl isomerase) can be inhibited by the widely used immunosuppressive tacrolimus. Next to its immunosuppressive properties, tacrolimus can inhibit the chaperone activity of FK506 binding protein 10, and is thereby proposed to have anti-fibrotic properties as well. Results from human embryonic kidney 293-cells and normal human dermal fibroblasts indicate that FK506 binding protein 10 peptidyl prolyl isomerase inhibition activity (which can be inhibited by tacrolimus) is linked to pyridinoline cross-linking by specifically mediating the dimerization of LH2.²² Thereby, tacrolimus not only inhibits the collagen chaperone activity of FK506 binding protein 10, but also decreases the number of pyridinoline cross-links, making the produced collagen more easily degradable by MMPs. Animal studies showed that tacrolimus can prevent alcohol- or carbon tetrachloride (CCl_4)-induced liver fibrosis in rats by inhibiting synthesis of type I collagen polypeptides, without affecting expression of collagen mRNAs.⁴⁵ Results are, however, conflicting since both Patsenker *et al.* (liver fibrosis induced by CCl_4 and bile duct ligation) and Frizell *et al.* (liver fibrosis induced by CCl_4) showed that tacrolimus was not able to inhibit fibrosis formation, but even enhanced fibrogenesis in the liver.^{46,86} The clinical experience with tacrolimus for CD is limited, but remission rates of 44% (range, 7–69%) and response rates of 37% (range, 14–57%) are reported for luminal CD.⁸⁷ No studies were found investigating the incidence of stricturing Crohn's disease in patients receiving tacrolimus *versus* other immunosuppressives. Cohort studies from kidney transplant recipients show that tacrolimus as the

current standard calcineurin inhibitor therapy post-renal transplantation, is not superior in preventing progression to interstitial fibrosis compared with cyclosporin or sirolimus.⁸⁸

Extracellular cleavage of propeptides

Inhibition of the cleavage of C- and N-terminal propeptides could have anti-fibrotic potential as well since it could inhibit the formation of an irreversibly stable collagenous ECM, thereby making it more easily degradable by collagenases that are able to degrade cross-linked collagen.⁴⁸ *In vitro* studies have shown that TIMP-3 can inhibit the procollagen N-proteinase activity of ADAMTS-2, thereby inhibiting procollagen processing in mouse embryonic fibroblasts. Upon stimulation with TIMP-3, reduced amounts of mature $\alpha 1(I)$ chains were observed.⁴⁸ Administration of TIMP-3 *in vivo* in a model for fibrosis has not been performed so far. Inhibition of bone morphogenic protein-1 proteinase activity (BMP-1) was tested only *in vitro* using (modified forms of) $\alpha 2$ -macroglobulin and acidic dipeptide hydroxamate. Also, these compounds have not been tested in *in vivo* models of organ fibrosis. Therapeutic inhibition of procollagen C-endopeptidase enhancer (PCOLCE; an enhancer of BMP-1 activity *in vitro* and *in vivo*), might have the same effect as therapeutic inhibition of BMP-1. In chronic-pressure-overload-induced cardiac fibrosis, PCOLCE2-null hearts demonstrated a decreased collagen content and a lower muscle stiffness compared with wildtype chronic pressure overloaded hearts.⁸⁹

Assembly into collagen fibrils by crosslinking of collagens

The general hypothesis is that targeting extracellular cross-linking by lysyl oxidases [including lysyl oxidase-like (LOXL)] might cause an increase in net degradation of collagen and other ECM molecules, thereby resulting in less fibrosis. Furthermore, it is generally hypothesized and proven in animal studies that inhibition of LOX or LOXL (e.g. anti-LOX or -lysyl oxidase-like 2 antibodies) decreases tumor stiffness and suppresses metastasis.^{53,90} *In vivo* studies on liver fibrosis showed that LOXL2 mediates collagen crosslinking and fibrotic matrix stabilization during liver fibrosis, and independently promotes fibrogenic differentiation of hepatic progenitor cells. By blocking these two

convergent profibrotic pathways, therapeutic LOXL2 inhibition attenuates both parenchymal and biliary fibrosis and promotes fibrosis reversal.⁹¹ However, even though inhibition of LOX(L) was very promising in pre-clinical *in vitro* and *in vivo* studies, several phase II studies testing the effect of the LOXL2 monoclonal antibody simtuzumab did not show an anti-fibrotic or antimetastatic effect.⁵⁷⁻⁶⁰ This study is the first to show upregulation of LOX and LOXL1-4 in CD-associated fibrosis in the terminal ileum. *In vivo* studies using animal models for intestinal fibrosis with LOX(L) knock-out animals or LOX(L) inhibitors, have not yet been performed.

Collagen-degrading enzymes

Based on their function as collagenases, inhibition of MMPs or CTSK is not expected to have anti-fibrotic efficacy. However, mRNA and protein levels of MMPs and CTSK are known to be upregulated in IBD, and it is known that these enzymes have other, for example chemotactic, properties as well.^{24,26} Therefore, inhibition of MMPs or CTSK might have an (indirect) anti-fibrotic effect. Especially MMP-9 is suggested to play a role in intestinal fibrosis and fistulae formation. Inhibition of MMP-9 in a heterotopic transplant model for intestinal fibrosis did reduce collagen content of the intestinal graft after induction of fibrosis by transplantation.⁹² Furthermore, expression of MMP-9 is a marker of mucosal healing, and increased local or serologic expression of MMP-9 is related to penetrating CD.⁹²⁻⁹⁴ Because inhibition of MMP-9 could simultaneously inhibit degradation of ECM and reduce fistulae associated fibrosis, this therapy may be superior for stricturing or penetrating CD compared with current immunosuppressive agents.

The importance of *Mmp-9* in intestinal inflammation in mice was recently questioned.⁹⁵ Very well-controlled studies comparing intestinal (colonic) inflammation induced by dextran sodium sulfate (DSS, both acute and chronic) and 2,4,6-trinitrobenzenesulfonic acid (TNBS) between *Mmp-9* knockout and wildtype mice, did not show a difference in the degree of intestinal inflammation or fibrosis induced. Inhibition of MMP-9 with bioactive peptides did not improve DSS-induced colitis, but the effect of inhibition of MMP-9 on intestinal fibrosis was not tested. De Bruyn *et al.* suggest that upregulation of MMP-9 is a consequence rather than a cause of inflammation of the

colon, and question whether MMP-9 represents a disease target in IBD.⁹⁵ Differences in the pathophysiology of fibrosis between colon and (terminal) ileum, and pathophysiological differences between the models used by de Bruyn *et al.* and Goffin *et al.* allow further testing of MMP-9 inhibitors for stricturing and penetrating CD.^{92,95}

Several MMP inhibitors are being tested in clinical trials as it was hypothesized that metastasis of cancer could be reduced by inhibition of MMP-mediated degradation of tumor-associated fibrosis, thereby reducing cancer progression.⁹⁶ Due to a lack of inhibitory specificity, and insufficient knowledge about the pleiotropic substrates and opposing effects of MMPs (on tumor growth/angiogenesis/modulation of immune response), these trials have all failed.⁹⁶ Especially since ECM remodeling is believed to be part of the pathophysiology of IBD, further testing of inhibitors of (other) MMPs still holds promise.⁹⁷

Whether inhibition of the TIMP-1 would have anti-fibrotic properties by reducing TIMP-mediated inhibition of MMP-activity remains to be studied. In a model of obstructive nephropathy-induced interstitial fibrosis, no amelioration of renal fibrosis was observed in *Timp-1* knockout mice.⁶²

Collagen receptors

Similar to other tyrosine kinase receptors, DDR kinase receptors (DDR-1 and -2) regulate fundamental cellular processes such as adhesion, migration, proliferation, and differentiation. Furthermore, they influence ECM remodeling *via* activation of MMPs.⁹⁸ DDR1 is found mostly in epithelial cells, whereas DDR2 is confined to cells of mesenchymal origin.⁹⁸ Inhibition of DDRs by tyrosine kinase inhibitors (dasatinib/imatinib/inilotinib, which are currently clinically used to treat chronic myeloid leukemia) has anti-fibrotic potential.⁶⁷ A significant decrease in collagen deposition of injured arteries of DDR1-null mice was observed.⁹⁹ Furthermore, type I collagen-dependent upregulation of DDR2 expression in hepatic stellate cells (HSC) establishes a positive feedback loop in activated stellate cells, leading to further proliferation and enhanced invasive activity of HSC.¹⁰⁰

A limitation of this study is that number of patients included is small; therefore, this study was not powered to reveal differences in, for

example, the expression of fibrosis genes depending on the therapy patients were using. However, the validity of the data increases due to the fact that the samples from non-fibrotic CD *versus* fibrotic CD are from the same patient and are therefore paired. Ideally, expression of the detected fibrosis genes should be validated in a replication cohort. Another limitation of this study is that mechanisms by which inhibition of post-translational modification and processing of collagens might attenuate the development of fibrosis in CD patients are only proposed. Mechanistic *in vitro/in vivo* data proving that inhibition of post-translational modification of collagens can inhibit fibrosis formation using one of the proposed mechanisms is not provided and was not the aim of this study. Efficacy for intestinal fibrosis and the underlying mechanisms will be explored in future research using, for example, cell culture of intestinal fibroblasts, precision-cut intestinal slices, or a heterotopic transplant animal model of intestinal fibrosis.^{101–103}

Conclusion

Therapeutic inhibition of fibrosis for patients with CD is not yet possible, but several drugs acting on factors involved in both pre- and post-transcriptional regulation of deposition of collagens and other extracellular matrix molecules are available. Which compound has the highest potential will depend on a combination of safety and anti-fibrotic efficacy. A drug against intestinal fibrosis would ideally be targeted to an enzyme/receptor that is uniquely expressed in a fibro-stenotic intestine in order to minimize systemic side effects. Since fibrosis formation occurs over a long period of time, clinical trials of long duration, a large number of patients, and selecting intestinal fibrosis-relevant endpoints, are warranted.^{73,104} Furthermore, the best route of application of the drug should be determined. This might be intravenous, but could also be oral for example, making use of sustained release by the ColoPulse technology (film coated tablets of targeted delivery in the lower intestinal tract) or topical administration for more distal (radiation-induced) fibrosis.^{105,106} Moreover, sustained release from the staple of a stapled anastomosis might be optional. However, meta-analysis of CD patients post-stricturoplasty showed that (with an overall 5-year surgical recurrence rate of 28%), recurrence occurs mainly at non-stricturoplasty sites in 90% of patients (whereas the site-specific

recurrence rate was 3%).¹⁰⁷ Furthermore, caution should be taken with anti-fibrotic therapy, because disturbance of the balance between collagen formation and degradation may induce a shift towards degradation to such an extent that fistulae or abscesses can occur. The anti-fibrotic capacity of some of the reviewed drugs for fibro-stenotic CD could be further unraveled, and their efficacy can be tested in *in vitro* and *in vivo* models for (CD-associated) intestinal fibrosis. In conclusion, inhibition of post-translational modification of collagens might be suitable to inhibit fibrosis formation in the intestine in CD.

Acknowledgements

We would like to thank the students of the Prometheus Kidney team of the Department of Surgery of the University Medical Center Groningen for collecting the surgical resection material from patients with CD and healthy controls.

Author contributions

WTvH: sample and data collection, analysis of the data, drafting of the manuscript. TB: sample and data collection, HS: sample collection. PO: critically reviewing the manuscript, GD: sample collection, critically reviewing the draft, critically reviewing the manuscript. RAB: data collection, critically reviewing the manuscript. MB: sample and data collection, analysis of the data, critically reviewing the manuscript.

Conflict of interest statement

WTvH has received funding to print his thesis from Ferring b.v., Teva b.v., Tramedico b.v. and Mylan b.v. GD reports outside the submitted work grants from Takeda and Abbvie, Fees for advisory boards from Cosmopharma and Mundipharma, speakers fees from Pfizer, Janssen pharmaceutical and Takeda.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Peter Olinga  <https://orcid.org/0000-0003-4855-8452>

Supplemental material

Supplemental material for this article is available online.

References

1. Piersma B, Bank RA and Boersema M. Signaling in fibrosis: TGF- β , WNT, and YAP/TAZ converge. *Front Med (Lausanne)* 2015; 2: 59.
2. Abraham C and Cho JH. Inflammatory bowel disease. *N Engl J Med* 2009; 361: 2066–2078.
3. Louis E, Collard A, Oger AF, *et al.* Behaviour of Crohn's disease according to the Vienna classification: changing pattern over the course of the disease. *Gut* 2001; 49: 777–782.
4. Smith BRK, Arnott IDR, Drummond HE, *et al.* Disease location, anti-saccharomyces cerevisiae antibody, and NOD2/CARD15 genotype influence the progression of disease behavior in Crohn's disease. *Inflamm Bowel Dis* 2004; 10: 521–528.
5. Louis E, Michel V, Hugot JP, *et al.* Early development of stricturing or penetrating pattern in Crohn's disease is influenced by disease location, number of flares, and smoking but not by NOD2/CARD15 genotype. *Gut* 2003; 52: 552–557.
6. Silverberg MS, Satsangi J, Ahmad T, *et al.* Toward an integrated clinical, molecular and serological classification of inflammatory bowel disease: report of a working party of the 2005 montreal world congress of gastroenterology. *Can J Gastroenterol* 2005; 19(Suppl. A): 5A–36A.
7. Hamama S, Gilbert-Sirieix M, Vozenin M-C, *et al.* Radiation-induced enteropathy: molecular basis of pentoxifylline-vitamin E anti-fibrotic effect involved TGF- β 1 cascade inhibition. *Radiother Oncol* 2012; 105: 305–312.
8. Rieder F, Fiocchi C and Rogler G. Mechanisms, management, and treatment of fibrosis in patients with inflammatory bowel diseases. *Gastroenterology* 2017; 152: 340–350.e6.
9. Fishbein TM, Gondolesi GE and Kaufman SS. Intestinal transplantation for gut failure. *Gastroenterology* 2003; 124: 1615–1628.
10. de Bruyn JR, Meijer SL, Wildenberg ME, *et al.* Development of fibrosis in acute and longstanding ulcerative colitis. *J Crohns Colitis* 2015; 9: 966–972.
11. Baron TH. Benign and malignant colorectal strictures. In: Waye JD, Rex DK and Williams CB (eds) *Colonoscopy: Principles and Practice*. 2nd ed. Chichester, West Sussex, UK: John Wiley & Sons Ltd, 2009, pp. 689–702.
12. Cosnes J, Bourrier A, Nion-Larmurier I, *et al.* Factors affecting outcomes in Crohn's disease over 15 years. *Gut* 2012; 61: 1140–1145.

13. Martinez FJ, Collard HR, Pardo A, *et al.* Idiopathic pulmonary fibrosis. *Nat Rev Dis Primers* 2017; 3: 17074.
14. Horowitz JC and Thannickal VJ. Mechanisms for the resolution of organ fibrosis. *Physiology (Bethesda)* 2019; 34: 43–55.
15. Graham MF, Diegelmann RF, Elson CO, *et al.* Collagen content and types in the intestinal strictures of Crohn's disease. *Gastroenterology* 1988; 94: 257–265.
16. Groulx J-F, Gagné D, Benoit YD, *et al.* Collagen VI is a basement membrane component that regulates epithelial cell-fibronectin interactions. *Matrix Biol* 2011; 30: 195–206.
17. Leeb SN, Vogl D, Grossmann J, *et al.* Autocrine fibronectin-induced migration of human colonic fibroblasts. *Am J Gastroenterol* 2004; 99: 335–340.
18. Allan A, Wyke J, Allan RN, *et al.* Plasma fibronectin in Crohn's disease. *Gut* 1989; 30: 627–633.
19. Johnson LA, Rodansky ES, Sauder KL, *et al.* Matrix stiffness corresponding to strictured bowel induces a fibrogenic response in human colonic fibroblasts. *Inflamm Bowel Dis* 2013; 19: 891–903.
20. Myllyharju J and Kivirikko KI. Collagens, modifying enzymes and their mutations in humans, flies and worms. *Trends Genet* 2004; 20: 33–43.
21. Karsdal MA, Nielsen MJ, Sand JM, *et al.* Extracellular matrix remodeling: the common denominator in connective tissue diseases. Possibilities for evaluation and current understanding of the matrix as more than a passive architecture, but a key player in tissue failure. *Assay Drug Dev Technol* 2013; 11: 70–92.
22. Gjaltema RAF, van der Stoel MM, Boersema M, *et al.* Disentangling mechanisms involved in collagen pyridinoline cross-linking: the immunophilin FKBP65 is critical for dimerization of lysyl hydroxylase 2. *Proc Natl Acad Sci U S A* 2016; 113: 7142–7147.
23. Honzawa Y, Nakase H, Matsumura K, *et al.* IL-17 promotes HSP47 expression and intestinal fibrosis in Crohn's disease. *Gastroenterology* 2011; 140: 493.
24. Ravi A, Garg P and Sitaraman SV. Matrix metalloproteinases in inflammatory bowel disease: boon or a bane? *Inflamm Bowel Dis* 2007; 13: 97–107.
25. Baugh MD, Perry MJ, Hollander AP, *et al.* Matrix metalloproteinase levels are elevated in inflammatory bowel disease. *Gastroenterology* 1999; 117: 814–822.
26. Warnaar N, Hofker HS, Maathuis MHJ, *et al.* Matrix metalloproteinases as profibrotic factors in terminal ileum in Crohn's disease. *Inflamm Bowel Dis* 2006; 12: 863–869.
27. Rockey DC, Bell PD and Hill JA. Fibrosis — a common pathway to organ injury and failure. *N Engl J Med* 2015; 372: 1138–1149.
28. Holvoet T, Devriese S, Castermans K, *et al.* Treatment of intestinal fibrosis in experimental inflammatory bowel disease by the pleiotropic actions of a local rho kinase inhibitor. *Gastroenterology* 2017; 153: 1054–1067.
29. Gjaltema RAF and Bank RA. Molecular insights into prolyl and lysyl hydroxylation of fibrillar collagens in health and disease. *Crit Rev Biochem Mol Biol* 2017; 52: 74–95.
30. Zuurmond A-M, van der Slot-Verhoeven AJ, van Dura EA, *et al.* Minoxidil exerts different inhibitory effects on gene expression of lysyl hydroxylase 1, 2, and 3: implications for collagen cross-linking and treatment of fibrosis. *Matrix Biol* 2005; 24: 261–270.
31. Liu W, Rao S, Fang H, *et al.* Lysyl hydroxylase inhibition by minoxidil blocks collagen deposition and prevents pulmonary fibrosis via TGF- β_1 /Smad3 signaling pathway. *Med Sci Monit* 2018; 24: 8592–8601.
32. Franklin TJ, Morris WP, Edwards PN, *et al.* Inhibition of prolyl 4-hydroxylase in vitro and in vivo by members of a novel series of phenanthrolinones. *Biochem J* 2001; 353: 333–338.
33. Xiong G, Deng L, Zhu J, *et al.* Prolyl-4-hydroxylase α subunit 2 promotes breast cancer progression and metastasis by regulating collagen deposition. *BMC Cancer* 2014; 14: 1.
34. Chang AY and Noble RE. Carminic acid, a non-competitive inhibitor kidney UDP-glucose: galactosylhydroxylysine-collagen glucosyltransferase. *Int J Biochem* 1982; 14: 691–694.
35. Ito S, Ogawa K, Takeuchi K, *et al.* A small-molecule compound inhibits a collagen-specific molecular chaperone and could represent a potential remedy for fibrosis. *J Biol Chem* 2017; 292: 20076–20085.
36. Ruigrok MJR, Xian J-L, Frijlink HW, *et al.* siRNA-mediated protein knockdown in precision-cut lung slices. *Eur J Pharm Biopharm* 2018; 133: 339–348.

37. Thomson CA, Atkinson HM and Ananthanarayanan VS. Identification of small molecule chemical inhibitors of the collagen-specific chaperone Hsp47. *J Med Chem* 2005; 48: 1680–1684.
38. Obata Y, Nishino T, Kushibiki T, *et al.* HSP47 siRNA conjugated with cationized gelatin microspheres suppresses peritoneal fibrosis in mice. *Acta Biomater* 2012; 8: 2688–2696.
39. Xia Z, Abe K, Furusu A, *et al.* Suppression of renal tubulointerstitial fibrosis by small interfering RNA targeting heat shock protein 47. *Am J Nephrol* 2008; 28: 34–46.
40. Otsuka M, Shiratori M, Chiba H, *et al.* Treatment of pulmonary fibrosis with siRNA against a collagen-specific chaperone HSP47 in vitamin A-coupled liposomes. *Exp Lung Res* 2017; 43: 271–282.
41. Kitamura H, Yamamoto S, Nakase H, *et al.* Role of heat shock protein 47 in intestinal fibrosis of experimental colitis. *Biochem Biophys Res Commun* 2011; 404: 599–604.
42. Staab-Weijnitz CA, Fernandez IE, Knüppel L, *et al.* FK506-binding protein 10, a potential novel drug target for idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 2015; 192: 455–467.
43. Liang X, Chai B, Duan R, *et al.* Inhibition of FKBP10 attenuates hypertrophic scarring through suppressing fibroblast activity and extracellular matrix deposition. *J Invest Dermatol* 2017; 137: 2326–2335.
44. Nagano J, Iyonaga K, Kawamura K, *et al.* Use of tacrolimus, a potent antifibrotic agent, in bleomycin-induced lung fibrosis. *Eur Respir J* 2006; 27: 460–469.
45. Manojlovic Z, Blackmon J and Stefanovic B. Tacrolimus (FK506) prevents early stages of ethanol induced hepatic fibrosis by targeting LARP6 dependent mechanism of collagen synthesis. *PLoS One* 2013; 8: e65897.
46. Frizell E, Abraham A, Doolittle M, *et al.* FK506 enhances fibrogenesis in in vitro and in vivo models of liver fibrosis in rats. *Gastroenterology* 1994; 107: 492–498.
47. Scott LJ, McKeage K, Keam SJ, *et al.* Tacrolimus: a further update of its use in the management of organ transplantation. *Drugs* 2003; 63: 1247–1297.
48. Wang W-M, Ge G, Lim NH, *et al.* TIMP-3 inhibits the procollagen N-proteinase ADAMTS-2. *Biochem J* 2006; 398: 515–519.
49. Zhang Y, Ge G and Greenspan DS. Inhibition of bone morphogenetic protein 1 by native and altered forms of α 2-macroglobulin. *J Biol Chem* 2006; 281: 39096–39104.
50. Kallander LS, Washburn D, Hilfiker MA, *et al.* Reverse hydroxamate inhibitors of bone morphogenetic protein 1. *ACS Med Chem Lett* 2018; 9: 736–740.
51. Trackman PC and Kagan HM. Nonpeptidyl amine inhibitors are substrates of lysyl oxidase. *J Biol Chem* 1979; 254: 7831–7836.
52. Tang SS, Trackman PC and Kagan HM. Reaction of aortic lysyl oxidase with beta-aminopropionitrile. *J Biol Chem* 1983; 258: 4331–4338.
53. Erler JT, Bennewith KL, Nicolau M, *et al.* Lysyl oxidase is essential for hypoxia-induced metastasis. *Nature* 2006; 440: 1222–1226.
54. Barry-Hamilton V, Spangler R, Marshall D, *et al.* Allosteric inhibition of lysyl oxidase-like-2 impedes the development of a pathologic microenvironment. *Nat Med* 2010; 16: 1009–1017.
55. Rodriguez HM, Vaysberg M, Mikels A, *et al.* Modulation of lysyl oxidase-like 2 enzymatic activity by an allosteric antibody inhibitor. *J Biol Chem* 2010; 285: 20964–20974.
56. Hutchinson JH, Rowbottom MW, Lonergan D, *et al.* Small molecule lysyl oxidase-like 2 (LOXL2) inhibitors: the identification of an inhibitor selective for LOXL2 over LOX. *ACS Med Chem Lett* 2017; 8: 423–427.
57. Muir AJ, Levy C, Janssen HLA, *et al.* Simtuzumab for primary sclerosing cholangitis: phase 2 study results with insights on the natural history of the disease. *Hepatology* 2019; 69: 684–698.
58. Raghu G, Brown KK, Collard HR, *et al.* Efficacy of simtuzumab versus placebo in patients with idiopathic pulmonary fibrosis: a randomised, double-blind, controlled, phase 2 trial. *Lancet Respir Med* 2017; 5: 22–32.
59. Harrison SA, Abdelmalek MF, Caldwell S, *et al.* Simtuzumab is ineffective for patients with bridging fibrosis or compensated cirrhosis caused by nonalcoholic steatohepatitis. *Gastroenterology* 2018; 155: 1140–1153.
60. Dong H, Benson AB III, Kudrik F, *et al.* A phase II randomized, double-blind, placebo-controlled study of simtuzumab or placebo in combination with gemcitabine for the first-line treatment of pancreatic adenocarcinoma. *Oncologist* 2017; 22: 241–e15.

61. Breynaert C, de Bruyn M, Arijs I, *et al.* Genetic deletion of tissue inhibitor of metalloproteinase-1/TIMP-1 alters inflammation and attenuates fibrosis in dextran sodium sulphate-induced murine models of colitis. *J Crohns Colitis* 2016; 10: 1336–1350.
62. Kim H, Oda T, López-Guisa J, *et al.* TIMP-1 deficiency does not attenuate interstitial fibrosis in obstructive nephropathy. *J Am Soc Nephrol* 2001; 12: 736–748.
63. Gao M, Duan L, Luo J, *et al.* Discovery and optimization of 3-(2-(pyrazolo[1,5-a]pyrimidin-6-yl) ethynyl)benzamides as novel selective and orally bioavailable discoidin domain receptor 1 (DDR1) inhibitors. *J Med Chem* 2013; 56: 3281–3295.
64. Kim H-G, Tan L, Weisberg EL, *et al.* Discovery of a potent and selective DDR1 receptor tyrosine kinase inhibitor. *ACS Chem Biol* 2013; 8: 2145–2150.
65. Siddiqui K, Kim GW, Lee DH, *et al.* Actinomycin D identified as an inhibitor of discoidin domain receptor 2 interaction with collagen through an insect cell based screening of a drug compound library. *Biol Pharm Bull* 2009; 32: 136–141.
66. Carafoli F, Mayer MC, Shiraishi K, *et al.* Structure of the discoidin domain receptor 1 extracellular region bound to an inhibitory Fab fragment reveals features important for signaling. *Structure* 2012; 20: 688–697.
67. Day E, Waters B, Spiegel K, *et al.* Inhibition of collagen-induced discoidin domain receptor 1 and 2 activation by imatinib, nilotinib and dasatinib. *Eur J Pharmacol* 2008; 599: 44–53.
68. Bantscheff M, Eberhard D, Abraham Y, *et al.* Quantitative chemical proteomics reveals mechanisms of action of clinical ABL kinase inhibitors. *Nat Biotechnol* 2007; 25: 1035–1044.
69. Yilmaz O, Oztay F and Kayalar O. Dasatinib attenuated bleomycin-induced pulmonary fibrosis in mice. *Growth Factors* 2015; 33: 366–375.
70. Mohammadalipour A, Karimi J, Khodadadi I, *et al.* Dasatinib prevent hepatic fibrosis induced by carbon tetrachloride (CCl₄) via anti-inflammatory and antioxidant mechanism. *Immunopharmacol Immunotoxicol* 2017; 39: 19–27.
71. Guignabert C, Phan C, Seferian A, *et al.* Dasatinib induces lung vascular toxicity and predisposes to pulmonary hypertension. *J Clin Invest* 2016; 126: 3207–3218.
72. Hill CR, Cole M, Errington J, *et al.* Characterisation of the clinical pharmacokinetics of actinomycin D and the influence of ABCB1 pharmacogenetic variation on actinomycin D disposition in children with cancer. *Clin Pharmacokinet* 2014; 53: 741–751.
73. Bettenworth D and Rieder F. Medical therapy of stricturing Crohn's disease: what the gut can learn from other organs - a systematic review. *Fibrogenesis Tissue Repair* 2014; 7: 5.
74. Bettenworth D and Rieder F. Reversibility of stricturing Crohn's disease—fact or fiction? *Inflamm Bowel Dis* 2016; 22: 241–247.
75. Afik R, Zigmond E, Vugman M, *et al.* Tumor macrophages are pivotal constructors of tumor collagenous matrix. *J Exp Med* 2016; 213: 2315–2331.
76. Stangenberg S, Saad S, Schilter HC, *et al.* Lysyl oxidase-like 2 inhibition ameliorates glomerulosclerosis and albuminuria in diabetic nephropathy. *Sci Rep* 2018; 8: 1–10.
77. Cosgrove D, Dufek B, Meehan DT, *et al.* Lysyl oxidase like-2 contributes to renal fibrosis in Col4α3/Alport mice. *Kidney Int* 2018; 94: 303–314.
78. Lawrance IC, Maxwell L and Doe W. Inflammation location, but not type, determines the increase in TGF-β1 and IGF-1 expression and collagen deposition in IBD intestine. *Inflamm Bowel Dis* 2001; 7: 16–26.
79. Wegrowski J, Lafuma C, Lefaix JL, *et al.* Modification of collagen and noncollagenous proteins in radiation-induced muscular fibrosis. *Exp Mol Pathol* 1988; 48: 273–285.
80. Xu C, Bailly-Maitre B and Reed JC. Endoplasmic reticulum stress: cell life and death decisions. *J Clin Invest* 2005; 115: 2656–2664.
81. Kawasaki K, Ushioda R, Ito S, *et al.* Deletion of the collagen-specific molecular chaperone Hsp47 causes endoplasmic reticulum stress-mediated apoptosis of hepatic stellate cells. *J Biol Chem* 2015; 290: 3639–3646.
82. Nakase H, Honzawa Y and Chiba T. Heat shock protein 47 is a new candidate molecule as anti-fibrotic treatment of Crohn's disease. *Aliment Pharmacol Ther* 2010; 31: 926–927; author reply 927–928.
83. Honzawa Y, Nakase H, Takeda Y, *et al.* Heat shock protein 47 can be a new target molecule for intestinal fibrosis related to inflammatory bowel disease. *Inflamm Bowel Dis* 2010; 16: 2004–2006.

84. Honzawa Y, Nakase H, Shiokawa M, *et al.* Involvement of interleukin-17A-induced expression of heat shock protein 47 in intestinal fibrosis in Crohn's disease. *Gut* 2014; 63: 1902–1912.
85. Chakraborty C, Sharma AR, Sharma G, *et al.* Therapeutic miRNA and siRNA: moving from bench to clinic as next generation medicine. *Mol Ther Nucleic Acids* 2017; 8: 132–143.
86. Patsenker E, Schneider V, Ledermann M, *et al.* Potent antifibrotic activity of mTOR inhibitors sirolimus and everolimus but not of cyclosporine A and tacrolimus in experimental liver fibrosis. *J Hepatol* 2011; 55: 388–398.
87. McSharry K, Dalzell AM, Leiper K, *et al.* Systematic review: the role of tacrolimus in the management of Crohn's disease. *Aliment Pharmacol Ther* 2011; 34: 1282–1294.
88. Oberbauer R. Progression of interstitial fibrosis in kidney transplantation. *Clin J Am Soc Nephrol* 2016; 11: 2110–2112.
89. Baicu CF, Zhang Y, Van Laer AO, *et al.* Effects of the absence of procollagen C-endopeptidase enhancer-2 on myocardial collagen accumulation in chronic pressure overload. *Am J Physiol Heart Circ Physiol* 2012; 303: H234–H240.
90. Levental KR, Yu H, Kass L, *et al.* Matrix crosslinking forces tumor progression by enhancing integrin signaling. *Cell* 2009; 139: 891–906.
91. Ikenaga N, Peng Z-W, Vaid KA, *et al.* Selective targeting of lysyl oxidase-like 2 (LOXL2) suppresses hepatic fibrosis progression and accelerates its reversal. *Gut* 2017; 66: 1697–1708.
92. Goffin L, Fagagnini S, Vicari A, *et al.* Anti-MMP-9 antibody: a promising therapeutic strategy for treatment of inflammatory bowel disease complications with fibrosis. *Inflamm Bowel Dis* 2016; 22: 2041–2057.
93. van Haaften WT, Mortensen JH, Karsdal MA, *et al.* Misbalance in type III collagen formation/degradation as a novel serological biomarker for penetrating (Montreal B3) Crohn's disease. *Aliment Pharmacol Ther* 2017; 46: 26–39.
94. de Bruyn M, Arijis I, De Hertogh G, *et al.* Serum neutrophil gelatinase B-associated lipocalin and matrix metalloproteinase-9 complex as a surrogate marker for mucosal healing in patients with Crohn's disease. *J Crohns Colitis* 2015; 9: 1079–1087.
95. de Bruyn M, Breynaert C, Arijis I, *et al.* Inhibition of gelatinase B/MMP-9 does not attenuate colitis in murine models of inflammatory bowel disease. *Nat Commun* 2017; 8: 1–15.
96. Vandenbroucke RE and Libert C. Is there new hope for therapeutic matrix metalloproteinase inhibition? *Nat Rev Drug Discov* 2014; 13: 904–927.
97. Shimshoni E, Yablecovitch D, Baram L, *et al.* ECM remodelling in IBD: innocent bystander or partner in crime? The emerging role of extracellular molecular events in sustaining intestinal inflammation. *Gut* 2015; 64: 367–372.
98. Leitinger B. Transmembrane collagen receptors. *Annu Rev Cell Dev Biol* 2011; 27: 265–290.
99. Hou G, Vogel W and Bendeck MP. The discoidin domain receptor tyrosine kinase DDR1 in arterial wound repair. *J Clin Invest* 2001; 107: 727–735.
100. Olaso E, Ikeda K, Eng FJ, *et al.* DDR2 receptor promotes MMP-2-mediated proliferation and invasion by hepatic stellate cells. *J Clin Invest* 2001; 108: 1369–1378.
101. Meier JK-H, Scharl M, Miller SN, *et al.* Specific differences in migratory function of myofibroblasts isolated from Crohn's disease fistulae and strictures. *Inflamm Bowel Dis* 2011; 17: 202–212.
102. Pham BT, van Haaften WT, Oosterhuis D, *et al.* Precision-cut rat, mouse and human intestinal slices can be used as a novel model for intestinal fibrosis in inflammatory bowel disease. *Physiol Rep* 2015; 3: e12323.
103. Hausmann M, Rechsteiner T, Caj M, *et al.* A new heterotopic transplant animal model of intestinal fibrosis. *Inflamm Bowel Dis* 2013; 19: 2302–2314.
104. Vermeire S, Sands BE, Bonovas S, *et al.* Identification of endpoints for development of antifibrosis drugs for treatment of Crohn's disease. *Gastroenterology* 2018; 155: 76–87.
105. Maurer JM, Schellekens RCA, Van Rieke HM, *et al.* Gastrointestinal pH and transit time profiling in healthy volunteers using the IntelliCap system confirms Ileo-Colonic release of ColoPulse tablets. *PLoS One* 2015; 10: e0129076.
106. van Dieren JM, van Bodegraven AA, Kuipers EJ, *et al.* Local application of tacrolimus in distal colitis: feasible and safe. *Inflamm Bowel Dis* 2009; 15: 193–198.
107. Yamamoto T, Fazio VW and Tekkis PP. Safety and efficacy of strictureplasty for Crohn's disease: a systematic review and meta-analysis. *Dis Colon Rectum* 2007; 50: 1968–1986.