

University of Groningen

Biosynthesis of lanthionine-constrained agonists of G protein-coupled receptors

Moll, Gert N; Kuipers, Anneke; Rink, Rick; Bosma, Tjibbe; de Vries, Louwe; Namsolleck, Pawel

Published in:
Biochemical Society Transactions

DOI:
[10.1042/BST20200427](https://doi.org/10.1042/BST20200427)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Moll, G. N., Kuipers, A., Rink, R., Bosma, T., de Vries, L., & Namsolleck, P. (2020). Biosynthesis of lanthionine-constrained agonists of G protein-coupled receptors. *Biochemical Society Transactions*, 48(5), 2195-2203. <https://doi.org/10.1042/BST20200427>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Review Article

Biosynthesis of lanthionine-constrained agonists of G protein-coupled receptors

 Gert N. Moll^{1,2}, Anneke Kuipers¹, Rick Rink¹, Tjibbe Bosma¹, Louwe de Vries¹ and Pawel Namsolleck¹

¹Lanthio Pharma, a MorphoSys AG Company, 9727 DL Groningen, The Netherlands; ²Department of Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, 9747 AG Groningen, The Netherlands

Correspondence: Gert N. Moll (moll@lanthiopharma.com) or (g.n.moll@rug.nl)



The conformation with which natural agonistic peptides interact with G protein-coupled receptor(s) (GPCR(s)) partly results from intramolecular interactions such as hydrogen bridges or is induced by ligand–receptor interactions. The conformational freedom of a peptide can be constrained by intramolecular cross-links. Conformational constraints enhance the receptor specificity, may lead to biased activity and confer proteolytic resistance to peptidic GPCR agonists. Chemical synthesis allows to introduce a variety of cross-links into a peptide and is suitable for bulk production of relatively simple lead peptides. Lanthionines are thioether bridged alanines of which the two alanines can be introduced at different distances in chosen positions in a peptide. Thioether bridges are much more stable than disulfide bridges. Biosynthesis of lanthionine-constrained peptides exploiting engineered Gram-positive or Gram-negative bacteria that contain lanthionine-introducing enzymes constitutes a convenient method for discovery of lanthionine-stabilized GPCR agonists. The presence of an N-terminal leader peptide enables dehydratases to dehydrate serines and threonines in the peptide of interest after which a cyclase can couple the formed dehydroamino acids to cysteines forming (methyl)lanthionines. The leader peptide also guides the export of the formed lanthionine-containing precursor peptide out of Gram-positive bacteria via a lanthipeptide transporter. An engineered cleavage site in the C-terminus of the leader peptide allows to cleave off the leader peptide yielding the modified peptide of interest. Lanthipeptide GPCR agonists are an emerging class of therapeutics of which a few examples have demonstrated high efficacy in animal models of a variety of diseases. One lanthipeptide GPCR agonist has successfully passed clinical Phase Ia.

Introduction

Today nearly 50 G protein-coupled receptor (GPCR) peptide drugs have been approved, and the pipeline is filled with an increasing number [1]. Many therapeutic peptides on the market are agonists of GPCRs [1,2]. GPCRs contain seven transmembrane segments and extracellular and intracellular loops. GPCR activity can be modulated by ligands that bind to their extracellular or transmembrane parts [3,4]. GPCRs can be activated by a variety of extracellular stimuli, among which interaction with peptides. GPCRs and their selectivity are very susceptible to allosteric modulation [5,6]. Extracellular GPCR stimulation activates G proteins that initiate intracellular signal transduction cascades. In addition, other GPCR-linked pathways, independent of G proteins, can be activated as well [7].

Structural stability of peptides can be imposed by hydrophobic or electrostatic intramolecular interactions and/or cross-links and may lead to increased GPCR-binding affinity by decreasing the entropic cost to adopt the GPCR-binding conformation. In the case of GPCR agonists, a conformational constraint imposed on a peptide, may induce a structure which is suitable or required for functional binding to a GPCR.

Received: 19 July 2020

Revised: 16 September 2020

Accepted: 17 September 2020

Version of Record published:
14 October 2020

Importantly, lanthionine-induced cyclization of a peptide may enhance the receptor specificity [8] or even the receptor subtype specificity [9]. A lanthionine somatostatin showed a similar high affinity for rat somatostatin receptor 5 (rSSTR5) compared with somatostatin [1–14] and sandostatin. However, it exhibited about 50 times weaker binding affinity for mouse somatostatin receptor 2b (mSSTR2b) than sandostatin [8]. Lanthionine enkephalin analogs with D-Ala in position 2 bind to both μ - and δ -opioid receptors and showed picomolar potencies [9]. The receptor specificity of a cyclized ligand has an additional impact, when the ligand up-regulates the GPCR level as reported for the angiotensin II type 2 receptor [10]. Furthermore, in specific cases, cyclized peptides can exert biased receptor agonism. Biased agonism by lanthi-apelins via the APJ receptor has been observed as witnessed by relatively stronger stimulation of the G protein pathway compared with the arrestin pathway [11]. This can translate to improved efficacy in the clinic by selectively stimulating therapeutic actions but avoiding activating detrimental β -arrestin-dependent pathways [12]. Selected lanthi-apelins, which more strongly stimulated the G protein pathway than the β -arrestin pathway did neither cause vasodilation nor increase heart rate in rats (Lanthio Pharma unpublished data, Figure 1).

In addition to the primary importance of conformational constraints of receptor agonists for their interaction with the receptor, several secondary advantages may occur such as the enhanced capacity to pass membranes and resistance to breakdown by peptidases and increased physical stability. Resistance to breakdown by peptidases in organ homogenates has been demonstrated amongst others for lanthionine-stabilized angiotensin-(1-7) [13,14] and for lanthionine-stabilized analogs of gonadotropin-releasing hormone I [15]. The capacity to pass membranes and the property of proteolytic resistance may lead to increased bioavailability after oral and




Figure 1. Lanthipeptide biased agonism via APJ receptor.

Cartoon of hypothetical stimulation of the APJ receptor by a strictly biased agonist that stimulates G protein-dependent pathways but not β -arrestin dependent pathways. APJ apelin receptor; AT₁R angiotensin II type 1 receptor; HNE1 sodium/hydrogen exchanger 1; NCX sodium–calcium exchanger; AC adenylyl cyclase; PKA protein kinase A; MEK 1/2 dual specificity mitogen-activated protein kinase kinase 1/2; PKC ϵ protein kinase C epsilon type; MLCK myosin light-chain kinase; MLC myosin light-chain; PLC phospholipase C; IP₃ inositol trisphosphate; SR sarcoplasmic reticulum; AQP2 aquaporin 2; BP blood pressure.

pulmonary delivery as observed for lanthionine-stabilized angiotensin-(1-7) [14], less frequent administration, lower doses and prolonged shelf life. These features apply also to conformationally constrained antagonists/blockers [16], which are outside the scope of this minireview which focusses on agonistic peptides. Some naturally occurring peptide GPCR agonists are constrained by disulfide bridges, e.g. vasopressin, oxytocin, cortistatin, urotensin II, endothelin I, somatostatin, adrenomedullin and calcitonin.

GPCR peptide agonists have been cyclized synthetically by a variety of cross-links, such as beta lactam [17], lysinoalanine [18], disulfide [12], artificial lipidated crosslinkers [19] and lanthionines [9]. Known lanthionine rings in lanthipeptide GPCR agonists range from i, i+3 (two amino acids between the lanthionine) to i, i+6 (five amino acids between the lanthionine) [8,9,11,13,20,21]. A thioether bridge is shorter than a disulfide bridge and a lysinoalanine. The exact structures resulting from the introduction of a lanthionine depend on the amino acid sequence and the position of the lanthionine and the resulting ring-size. Bacterial production of lanthionine-constrained agonistic peptides is a convenient method in the discovery of drug candidates. Obviously, the introduction of a lanthionine, either by insertion into the existing amino acid sequence or by replacing existing amino acids with (half of) a lanthionine, may strongly modulate the agonistic activity of the peptide, ranging from completely loss of activity to enhanced activity. Screening of lanthionine-containing variants will be required to select pharmacologically valuable peptides.

In many therapeutic peptides, the C-terminus is amidated which precludes carboxypeptidase action and may contribute to optimal receptor interaction. Peptides can be enzymatically amidated by peptidyl-glycine alpha-amidating monooxygenase or carboxypeptidase Y, which is likely compatible with bacterial production of lanthionine-containing peptides. Following bacterial production chemical and enzymatic amidation has been demonstrated for gonadotropin-releasing hormone I [15].

As broadly applicable, highly convenient methods have been developed for the bacterial synthesis of lanthionine-stabilized peptides, we here provide an overview on their emerging application in the discovery of lanthionine-constrained GPCR agonists.

Biosynthesis of lanthipeptides

The word lanthionine stems from the latin word *lana* which means wool, in which lanthionines were first observed, and thiol (sulfhydryl). Lanthionines are thioether bridged amino acids such as alanine-S-alanine, aminobutyric acid-S-alanine or alanine-S-aminobutyric acid. Depending on the distances between the thioether bridged residues, smaller or larger ring structures are formed in the peptide. Lanthionine-containing peptides, so-called lanthipeptides [22] are naturally produced for instance by several Gram-positive bacteria [23].

Lanthipeptides are synthesized as precursor peptides that contain an N-terminal leader peptide and a modifiable core peptide. This leader peptide enables the interaction with modification enzymes that can dehydrate serines and threonines in the core peptide and couple the formed dehydroamino acids to cysteines thus forming lanthionines and methylanthionines. The cyclization is stereospecific: either D,L (methyl)lanthionines or L,D (methyl)lanthionines are formed. The enzymatic cyclization is also regiospecific. This means that a dehydroamino acid in a certain position is coupled to only one cysteine position yielding only one ring position structure and, in case of more than one lanthionine in one peptide, only one ring pattern. In *Lactococcus lactis*, the leader peptide allows the export of the modified precursor peptide out of the cell. These leader peptides and the corresponding modification and transporter enzymes have been subject of a detailed review [24]. For some of these enzyme systems, a membrane-bound enzyme complex has been described which is composed of the modification enzymes and a dedicated transporter. The leader peptide is cleaved off either intracellularly by a bifunctional transporter/leaderpeptidase or extracellularly by a leader peptidase or non-dedicated peptidases.

The best-known lanthipeptide is the pentacyclic lantibiotic nisin [25] which is produced by *L. lactis*. The nisin precursor peptide is dehydrated by the dehydratase NisB [26,27], after which a cyclase NisC [28] couples dehydroamino acids to cysteines followed by export by the transporter NisT and removal of the leader peptide by the extracellular leader peptidase NisP [29].

Exploitation of bacteria for the biosynthesis of engineered lanthipeptides

Following the elucidation of the precursor sequence of the lanthipeptide gallidermin, as early as 1988, it was recognized that the introduction of (methyl)lanthionines into therapeutic peptides, such as agonistic peptide hormones might constitute a valuable opportunity to discover specific and stable lanthionine-containing therapeutics [30].

However, this concept was entirely dependent on the yet unknown substrate specificity of the complex of lanthipeptide modification and transport enzymes. In 2004, the first GPCR agonist peptide, an angiotensin-(1-7) variant fused to the nisin leader, which was totally unrelated to nisin, was modified by the nisin dehydratase and exported out of *L. lactis* by the nisin transporter NisT [31]. Subsequent studies demonstrated that the dehydratase NisB and the nisin cyclase NisC could modify a broad range of peptides unrelated to nisin, and that also the transporter could accommodate the export of a variety of fusion peptides of the nisin leader peptide and a peptide of interest [32–35]. In the C-terminal part of the nisin leader peptide, general cleavage sites could be engineered to allow convenient clean removal of the leader peptide from the produced peptide of interest [29]. By engineering a signal sequence at the N-terminus of the leader peptide constructs peptides could also be exported out of *L. lactis* via the SEC secretion system provided they had limited bulkiness [36,37]. Excellent *in vitro* studies by Wilfred van der Donk and co-workers furthermore demonstrated that a bifunctional lanthionine-introducing LanM enzyme, which performs both the dehydration as well as the cyclization step, also has a broad substrate specificity [38].

Several types of lanthionine-introducing enzymes were demonstrated to be functional in *Escherichia coli* [39,40] allowing intracellular modification and subsequent harvesting after disruption of the cells. The use of either Gram-positive or Gram-negative bacteria would thus constitute a cheap and effective method for producing a variety of rationally designed lanthipeptides to subsequently pursue the discovery of GPCR agonists with interesting pharmacokinetics and -dynamics.

Substrate specificity of lanthionine-introducing enzymes

An excellent, comprehensive and thorough review focuses on lanthionine-introducing enzymes, their structure and mechanism of action [23]. Here, we summarize some of the progress made with respect to engineering novel lanthipeptides mainly by using bacteria containing the nisin modification enzymes. While the attainment of strict rules with respect to the substrate specificities of lanthionine-introducing enzymes has been challenging, some practical guidelines for the design of modifiable peptides have been reached for some lanthipeptide biosynthesis systems [32]. The substrate specificity of lanthionine-introducing enzymes was first investigated via an *in silico* study on all available lanthipeptide structures and subsequently experimentally validated with the nisin modification enzymes [32,34]. Directly flanking residues of serines and threonines clearly had an impact on the extent of dehydration by NisB (Figure 2). Hydrophobic amino acids, especially when present on both




Figure 2. Biosynthesis of a lanthipeptide.

A threonine in the core peptide of interest preceded by a cleavable leader peptide (Lp) is dehydrated by a LanB dehydratase forming a dehydrobutyryne (Dhb) after which a LanC cyclase couples the formed dehydrobutyryne to a cysteine forming a D,L-methylanthionine (dAbu-S-Ala). Directly flanking amino acids (light green) of the Thr and of the Cys (dark green) may affect the extent of dehydration and cyclization [32,34,35]. The enzymes LanB and LanC can also introduce a lanthionine via LanB-catalyzed dehydration of a serine yielding dehydroalanine and subsequent LanC-catalyzed coupling of the dehydroalanine to a cysteine. However, dehydroalanines are much more reactive than dehydrobutyrynes and can spontaneously react with cysteines without stereospecificity.

sides of the Ser/Thr favored dehydration whereas hydrophilic amino acids did not. Hydrophilic amino acids, especially Arg, Asp, Glu and Gly, simultaneously present at both sides of the Ser/Thr, prevented dehydration. The requirement of hydrophobic amino acids directly flanking dehydratable serines/threonines might be caused by the negative charge of glutamyl tRNA which acts as cofactor in the dehydration process [23]. The influence on dehydration of the combination of one hydrophobic and one hydrophilic amino acid as directly flanking residues of Ser/Thr differed from case to case, and often led to partial dehydration. Thr was generally dehydrated to a larger extent than Ser. Ser/Thr at the very C-terminus escaped dehydration possibly as a result of the hydrophilicity of the carboxyl group. The presence of an already present lanthionine ring might in specific sequence contexts hamper the dehydration of Ser/Thr by a LanM enzyme [41]. Dehydration at a position as remote from the leader as 46 positions could still be dehydrated [34], but it is not known yet where the upper limit is. Interestingly, an elegant high-throughput screening system allowed to obtain mutant NisB enzymes with adapted substrate specificity with respect to the amino acids that directly flank Ser/Thr [42].

The substrate specificity of NisC was first investigated with model peptides. Dehydroalanine appeared to be very reactive leading to spontaneous lanthionine formation in bacterially produced peptides even in the absence of NisC as evidenced by comparing cells expressing NisBTC with cells with only NisBT [35]. In the case of spontaneous cyclization in which highly reactive dehydroalanines readily couple to cysteines, more than one isomer can be formed. To be complete it should also be mentioned that in specific sequences spontaneous cyclization between dehydroalanines and cysteines can be stereospecific as well. In these cases, the sequence imposes the stereochemistry. For NisC-catalyzed cyclization, it appears complex to generate guidelines for the design of peptides that are well cyclized or escape cyclization. Bulky residues in small ring structures appear to be unfavorable [35]. Glycines appear to be favorable and the impact of prolines is hard to predict. For cyclization, negatively charged amino acids as N-terminally directly flanking residues of a cysteine seem to be favored over positively charged amino acids.

The stereospecificity of NisC-cyclized GPCR agonist, 4,7 lanthionine-angiotensin-(1-7) was investigated using the Ni₂B method, which opens the ring structure while retaining the D or L conformation [43]. Using reference peptide with combinations of D-Ala/L-Ala in positions 4 and 7, the NisC-cyclized cAng-(1-7) isomer turned out to be D,L [R. Rink unpublished data]. Using the same method it was demonstrated that also 4,7 NisC-cyclized LHRH had the D,L conformation [15]. This is in accordance with the D,L stereospecificity of all the lanthionine and methylanthionine rings in nisin. Designed, NisC-cyclized intertwined lanthionine structures of different sizes have been synthesized [35].

The substrate specificity of the nisin transporter is not precisely known yet. Generally, positively charged peptides [44] seem to be better exported than negatively charged peptides in correspondence with the cationicity of the natural substrate, nisin. The mechanism of export and the involvement of NisT in the observed export of a large cell-wall spanning protein out of *L. lactis* may require detailed mechanistic analyses [45]. Generally, the efficiency of NisT-mediated export of peptides that are longer than nisin decreases significantly with increasing peptide length.

The membrane-bound leader peptidase NisP appears to be highly specific for lanthionine-containing precursor nisin [31]. In contrast, an engineered water-soluble truncated NisP was able to cleave off the leader peptide from a variety of substrate peptides [46]. Apart from the nisin biosynthetic machinery, many other lanthipeptide biosynthesis systems exist. Interestingly, the single bifunctional enzyme ProcM naturally introduces lanthionine rings in 29 different substrate peptides [47]. Clearly, this enzyme is a valuable tool in the design and synthesis of lanthionine-containing GPCR agonists. Recently, a powerful high-throughput screening method of NisB mutants with adapted substrate specificity has been demonstrated, thus opening the way to analogously obtain many tailor-made modification enzymes [42].

Lanthipeptide GPCR agonists

Lanthipeptide GPCR agonists are an emerging class of peptide therapeutics. A number of them is listed in Table 1. Most data are available on 4,7 lanthionine-constrained angiotensin-(1-7) which has shown high efficacy in many animal disease models such as acute respiratory distress syndrome [49,50], acute lung injury [20], pulmonary arterial hypertension [51], myocardial infarct [52], diabetic nephropathy [54], type 1 and 2 diabetes [55] and cerebral stroke [56]. It is completely resistant to breakdown by ACE [13] and pulmonary delivery is feasible [14]. Its *ex vivo* vasodilatory action was inhibited by two peptides, D-Pro7-Ang-(1-7) and D-Ala7-Ang-(1-7), that are considered antagonists of the MAS1 receptor [13].

Table 1. Lanthipeptide GPCR agonists

Lanthipeptide GPCR agonists	References
Lanthionine-enkephalins	[9,48]
Lanthionine-somatostatin	[8]
Cyclic angiotensin-(1-7)	[13,14,20,49–56]
Cyclic apelin	[11]
Cyclic galanins	[21]
LP2	[20,57]

Lanthionine naturally occurs in lanthionine-ketimine present in the human brain [58]; thioethers are ubiquitously present in the human body in methionines. The lanthipeptide AT₂R agonist LP2 [20,57] has been tested in clinical Phase I where it showed at the tested doses safety and favorable pharmacokinetics. Also complex lanthipeptides, other than receptor agonists, like the lantibiotics NBV302 ([59], BioCentury|August 13, 2012) and duramycin [60], have been successfully tested in clinical safety trials. Duramycin interacts with phosphatidylethanolamine and it can uncouple mitochondrial respiration in isolated mitochondria [61]. Duramycin activates an alternative chloride channel in cystic fibrosis nasal and airway epithelia. By this, duramycin bypasses the transmembrane conductance regulator, a chloride channel, which is dysfunctional in cystic fibrosis. Under therapeutic conditions after administration via inhalation, duramycin plasma concentration stays below 0.5 nM [62]. Until now no data directly point at any intrinsic toxicity of lanthionines.

Next to potentially high value as therapeutics lanthipeptide, GPCR agonists may have value as stable research tools with higher GPCR specificity than the corresponding natural linear peptides. Natural peptides such as those derived from angiotensin I are *in vivo* rapidly degraded, making it difficult to assign their precise effects. In this respect, stable analogs may allow more reliable conclusions as to which peptide acts via which receptor leading to which effects. High GPCR specificity of lanthionine-constrained agonists may also prove useful in the elucidation of GPCR heterodimerization which occurs for instance for MAS1 and AT₂ receptors [63]. While biosynthesis methods are very convenient in the discovery process of therapeutic candidates, upscaled chemical production methods [64–67] are in development and base-assisted sulfur extrusion has already successfully been applied for bulk production of GMP material of the clinically tested LP2.

Perspectives

- The target specificity, stability and safety of lanthipeptide GPCR agonists make them an important class of candidate therapeutics.
- Currently, it is clear that several biosynthetic systems have relaxed substrate specificity which allows their exploitation in lanthipeptide drug discovery. In addition, the feasibility of high-throughput screening for tailor-made lanthionine-introducing enzymes with adapted substrate specificity for rational design and biosynthesis of lanthipeptide GPCR agonists has recently been demonstrated.
- Future developments comprise the discovery and clinical development of multiple lanthipeptide GPCR agonists as therapeutics for treating human diseases.

Competing Interest

All authors are employees of LanthioPep B.V., which organization is the owner of patents on the biosynthesis of lanthipeptides and specific lanthipeptide GPCR agonists. G.N.M. is the director of LanthioPep B.V.

Open Access

Open access for this article was enabled by the participation of the University of Groningen in an all-inclusive Read & Publish pilot with Portland Press and the Biochemical Society.

Author Contributions

G.N.M. contributed to the overall outline of the manuscript; R.R. with unpublished data, A.K., R.R. and T.B. to the paragraphs on biosynthesis, R.R. to the chemical synthesis, A.K., L.d.V. and P.N. to GPCRs; P.N. made Figure 1.

Abbreviations

GPCRs, G protein-coupled receptors; mSSTR2b, mouse somatostatin receptor 2b; rSSTR5, rat somatostatin receptor 5.

References

- 1 Davenport, A.P., Scully, C.C.G., de Graaf, C., Brown, A.J.H. and Maguire, J.J. (2020) Advances in therapeutic peptides targeting G protein-coupled receptors. *Nat. Rev. Drug Discov.* **19**, 389–413 <https://doi.org/10.1038/s41573-020-0062-z>
- 2 Reichert, J., Pechon, P., Tartar, A., and Dunn, M.K. (2010) Development Trends for Peptide Therapeutics. Peptide Therapeutics Foundation. San Diego, California 92121
- 3 Wang, W., Qiao, Y. and Li, Z. (2018) New insights into modes of GPCR activation. *Trends Pharmacol. Sci.* **39**, 367–386 <https://doi.org/10.1016/j.tips.2018.01.001>
- 4 Weis, W.I. and Kobilka, B.K. (2018) The molecular basis of G protein-coupled receptor activation. *Annu. Rev. Biochem.* **87**, 897–919 <https://doi.org/10.1146/annurev-biochem-060614-033910>
- 5 van der Westhuizen, E.T., Valant, C., Sexton, P.M. and Christopoulos, A. (2015) Endogenous allosteric modulators of G protein-coupled receptors. *J. Pharmacol. Exp. Ther.* **353**, 246–260 <https://doi.org/10.1124/jpet.114.221606>
- 6 Thal, D.M., Glukhova, A., Sexton, P.M. and Christopoulos, A. (2018) Structural insights into G-protein-coupled receptor allostery. *Nature* **559**, 45–53 <https://doi.org/10.1038/s41586-018-0259-z>
- 7 Walther, C. and Ferguson, S.S. (2015) Minireview: role of intracellular scaffolding proteins in the regulation of endocrine g protein-coupled receptor signaling. *Mol. Endocrinol.* **29**, 814–830 <https://doi.org/10.1210/me.2015-1091>
- 8 Ösabay, G., Prokai, L., Kim, H.S., Medzhradszky, K.F., Coy, D.H., Liapakis, G. et al. (1997) Lanthionine-somatostatin analogs: synthesis, characterization, biological activity, and enzymatic stability studies. *J. Med. Chem.* **40**, 2241–2251 <https://doi.org/10.1021/jm960850i>
- 9 Rew, Y., Malkmus, S., Svensson, C., Yaksh, T.L., Chung, N.N., Schiller, P.W. et al. (2002) Synthesis and biological activities of cyclic lanthionine enkephalin analogues: delta-opioid receptor selective ligands. *J. Med. Chem.* **45**, 3746–3754 <https://doi.org/10.1021/jm020108k>
- 10 Toedebusch, R., Belenchia, A. and Pulakat, L. (2018) Cell-specific protective signaling induced by the novel AT₂R-agonist NP-6A4 on human endothelial and smooth muscle cells. *Front. Pharmacol.* **9**, 928 <https://doi.org/10.3389/fphar.2018.00928>
- 11 Rink, R. (2019) Cyclic apelin analogs. Patent EP3122764 (B1)
- 12 Brame, A.L., Maguire, J.J., Yang, P., Dyson, A., Torella, R., Cheriyan, J. et al. (2015) Design, characterization, and first-in-human study of the vascular actions of a novel biased apelin receptor agonist. *Hypertension* **65**, 834–840 <https://doi.org/10.1161/HYPERTENSIONAHA.114.05099>
- 13 Kluskens, L.D., Nelemans, S.A., Rink, R., de Vries, L., Meter-Arkema, A., Wang, Y. et al. (2009) Angiotensin-(1-7) with thioether bridge: an angiotensin-converting enzyme-resistant, potent angiotensin-(1-7) analog. *J. Pharmacol. Exp. Ther.* **328**, 849–854 <https://doi.org/10.1124/jpet.108.146431>
- 14 de Vries, L., Reitzema-Klein, C.E., Meter-Arkema, A., van Dam, A., Rink, R., Moll, G.N. et al. (2010) Oral and pulmonary delivery of thioether-bridged angiotensin-(1-7). *Peptides* **31**, 893–898 <https://doi.org/10.1016/j.peptides.2010.02.015>
- 15 Rink, R., Arkema-Meter, A., Baudoin, I., Post, E., Kuipers, A., Nelemans, S.A. et al. (2010) To protect peptide pharmaceuticals against peptidases. *J. Pharmacol. Toxicol. Methods* **61**, 210–218 <https://doi.org/10.1016/j.vascn.2010.02.010>
- 16 Bosma, T., Rink, R., Moosmeier, M.A. and Moll, G.N. (2019) Genetically encoded libraries of constrained peptides. *ChemBioChem* **20**, 1754–1758 <https://doi.org/10.1002/cbic.201900031>
- 17 Murage, E.N., Gao, G., Bisello, A. and Ahn, J.M. (2010) Development of potent glucagon-like peptide-1 agonists with high enzyme stability via introduction of multiple lactam bridges. *J. Med. Chem.* **53**, 6412–6420 <https://doi.org/10.1021/jm100602m>
- 18 Kuipers, A., de Vries, L., de Vries, M.P., Rink, R., Bosma, T. and Moll, G.N. (2017) Semi-microbiological synthesis of an active lysinoalanine-bridged analog of glucagon-like-peptide-1. *Peptides* **91**, 33–39 <https://doi.org/10.1016/j.peptides.2017.03.004>
- 19 Yang, P.Y., Zou, H., Amso, Z., Lee, C., Huang, D. and Woods, A.K. (2020) New generation oxyntomodulin peptides with improved pharmacokinetic profiles exhibit weight reducing and anti-steatotic properties in mice. *Bioconjug. Chem.* **31**, 1167–1176 <https://doi.org/10.1021/acs.bioconjugchem.0c00093>
- 20 Wagenaar, G.T., Laghmani el, H., Fidder, M., Sengers, R.M., de Visser, Y.P., de Vries, L. et al. (2013) Agonists of MAS oncogene and angiotensin II type 2 receptors attenuate cardio-pulmonary disease in rats with neonatal hyperoxia-induced lung injury. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **305**, L341–L351 <https://doi.org/10.1152/ajplung.00360.2012>
- 21 Kuipers, A. (2019) Cyclic galanin-analogs and uses thereof. Patent US 10501514, B2
- 22 Arnison, P.G., Bibb, M.J., Bierbaum, G., Bowers, A.A., Bugni, T.S., Bulaj, G. et al. (2013) Ribosomally synthesized and post-translationally modified peptide natural products: overview and recommendations for a universal nomenclature. *Nat. Prod. Rep.* **30**, 108–160 <https://doi.org/10.1039/C2NP20085F>

- 23 Repka, L.M., Chekan, J.R., Nair, S.K. and van der Donk, W.A. (2017) Mechanistic understanding of lanthipeptide biosynthetic enzymes. *Chem. Rev.* **117**, 5457–5520 <https://doi.org/10.1021/acs.chemrev.6b00591>
- 24 Plat, A., Kuipers, A., Rink, R. and Moll, G.N. (2013) Mechanistic aspects of lanthipeptide leaders. *Curr. Protein Pept. Sci.* **14**, 85–96 <https://doi.org/10.2174/1389203711314020001>
- 25 Lubelski, J., Rink, R., Khusainov, R., Moll, G.N. and Kuipers, O.P. (2008) Biosynthesis, immunity, regulation, mode of action and engineering of the model lantibiotic nisin. *Cell. Mol. Life Sci.* **65**, 455–476 <https://doi.org/10.1007/s00018-007-7171-2>
- 26 Garg, N., Salazar-Ocampo, L.M. and van der Donk, W.A. (2013) In vitro activity of the nisin dehydratase NisB. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 7258–7263 <https://doi.org/10.1073/pnas.1222488110>
- 27 Ortega, M.A., Hao, Y., Zhang, Q., Walker, M.C., van der Donk, W.A. and Nair, S.K. (2015) Structure and mechanism of the tRNA-dependent lantibiotic dehydratase NisB. *Nature* **517**, 509–512 <https://doi.org/10.1038/nature13888>
- 28 Li, B., Yu, J.P., Brunzelle, J.S., Moll, G.N., van der Donk, W.A. and Nair, S.K. (2006) Structure and mechanism of the lantibiotic cyclase involved in nisin biosynthesis. *Science* **311**, 1464–1467 <https://doi.org/10.1126/science.1121422>
- 29 Plat, A., Kluskens, L.D., Kuipers, A., Rink, R. and Moll, G.N. (2011) Requirements of the engineered leader peptide of nisin for inducing modification, export, and cleavage. *Appl. Environ. Microbiol.* **77**, 604–611 <https://doi.org/10.1128/AEM.01503-10>
- 30 Schnell, N., Entian, K.D., Schneider, U., Götz, F., Zähler, H., Kellner, R. et al. (1988) Prepeptide sequence of epidermin, a ribosomally synthesized antibiotic with four sulphide-rings. *Nature* **333**, 276–278 <https://doi.org/10.1038/333276a0>
- 31 Kuipers, A., de Boef, E., Rink, R., Fekken, S., Kluskens, L.D., Driessen, A.J. et al. (2004) NisT, the transporter of the lantibiotic nisin, can transport fully modified, dehydrated, and unmodified prenisin and fusions of the leader peptide with non-lantibiotic peptides. *J. Biol. Chem.* **279**, 22176–22182 <https://doi.org/10.1074/jbc.M312789200>
- 32 Rink, R., Kuipers, A., de Boef, E., Leenhouts, K.J., Driessen, A.J.M., Moll, G.N. et al. (2005) Lantibiotic structures as guidelines for the design of peptides that can be modified by lantibiotic enzymes. *Biochemistry* **44**, 8873–8882 <https://doi.org/10.1021/bi050081h>
- 33 Kluskens, L.D., Kuipers, A., Rink, R., de Boef, E., Fekken, S., Driessen, A.J. et al. (2005) Post-translational modification of therapeutic peptides by NisB, the dehydratase of the lantibiotic nisin. *Biochemistry* **44**, 12827–12834 <https://doi.org/10.1021/bi050805p>
- 34 Rink, R., Wierenga, J., Kuipers, A., Kluskens, L.D., Driessen, A.J.M., Kuipers, O.P. et al. (2007a) Production of dehydroamino acid-containing peptides by *Lactococcus lactis*. *Appl. Environ. Microbiol.* **73**, 1792–1796 <https://doi.org/10.1128/AEM.02350-06>
- 35 Rink, R., Kluskens, L.D., Kuipers, A., Driessen, A.J., Kuipers, O.P. and Moll, G.N. (2007c) Nisc, the cyclase of the lantibiotic nisin, can catalyze cyclization of designed nonlantibiotic peptides. *Biochemistry* **46**, 13179–13189 <https://doi.org/10.1021/bi700106z>
- 36 Kuipers, A., Wierenga, J., Rink, R., Kluskens, L.D., Driessen, A.J., Kuipers, O.P. et al. (2006) Sec-mediated transport of posttranslationally dehydrated peptides in *Lactococcus lactis*. *Appl. Environ. Microbiol.* **72**, 7626–7633 <https://doi.org/10.1128/AEM.01802-06>
- 37 Kuipers, A., Rink, R. and Moll, G.N. (2009) Translocation of a thioether-bridged azurin peptide fragment via the sec pathway in *Lactococcus lactis*. *Appl. Environ. Microbiol.* **75**, 3800–3802 <https://doi.org/10.1128/AEM.00341-09>
- 38 Levengood, M.R., Kerwood, C.C., Chatterjee, C. and van der Donk, W.A. (2009) Investigation of the substrate specificity of lactacin 481 synthetase by using nonproteinogenic amino acids. *Chembiochem* **10**, 911–919 <https://doi.org/10.1002/cbic.200800752>
- 39 Shi, Y., Yang, X., Garg, N. and van der Donk, W.A. (2011) Production of lantipeptides in *Escherichia coli*. *J. Am. Chem. Soc.* **133**, 2338–2341 <https://doi.org/10.1021/ja109044r>
- 40 Nagao, J., Harada, Y., Shioya, K., Aso, Y., Zendo, T., Nakayama, J. et al. (2005) Lanthionine introduction into nukacin ISK-1 prepeptide by co-expression with modification enzyme NukM in *Escherichia coli*. *Biochem. Biophys. Res. Commun.* **336**, 507–513 <https://doi.org/10.1016/j.bbrc.2005.08.125>
- 41 Kuipers, A., Meijer-Wierenga, J., Rink, R., Kluskens, L.D. and Moll, G.N. (2008) Mechanistic dissection of the enzyme complexes involved in biosynthesis of lactacin 3147 and nisin. *Appl. Environ. Microbiol.* **74**, 6591–6597 <https://doi.org/10.1128/AEM.01334-08>
- 42 Zhao, X., Cebrian, R., Fu, Y., Rink, R., Bosma, T., Moll, G.N. et al. (2020) High-throughput screening for substrate specificity-adapted mutants of the nisin dehydratase NisB. *ACS Synth. Biol.* **9**, 1468–1478 <https://doi.org/10.1021/acssynbio.0c00130>
- 43 Martin, N.I., Sprules, T., Carpenter, M.R., Cotter, P.D., Hill, C., Ross, R.P. et al. (2004) Structural characterization of lactacin 3147, a two-peptide lantibiotic with synergistic activity. *Biochemistry* **43**, 3049–3056 <https://doi.org/10.1021/bi0362065>
- 44 Plat, A., Kuipers, A., de Lange, J.G., Moll, G.N. and Rink, R. (2011) Activity and export of engineered nisin-(1-22) analogs. *Polymers (Basel)* **3**, 1282–1296 <https://doi.org/10.3390/polym3031282>
- 45 Bosma, T., Kuipers, A., Bulten, E., de Vries, L., Rink, R. and Moll, G.N. (2011) Bacterial display and screening of posttranslationally thioether-stabilized peptides. *Appl. Environ. Microbiol.* **77**, 6794–6801 <https://doi.org/10.1128/AEM.05550-11>
- 46 Lagedroste, M., Smits, S.H.J. and Schmitt, L. (2017) Substrate specificity of the secreted nisin leader peptidase NisP. *Biochemistry* **56**, 4005–4014 <https://doi.org/10.1021/acs.biochem.7b00524>
- 47 Tang, W. and van der Donk, W.A. (2012) Structural characterization of four prochlorosins: a novel class of lantipeptides produced by planktonic marine cyanobacteria. *Biochemistry* **51**, 4271–4279 <https://doi.org/10.1021/bi300255s>
- 48 Levengood, M.R. and van der Donk, W.A. (2008) Use of lantibiotic synthetases for the preparation of bioactive constrained peptides. *Bioorg. Med. Chem. Lett.* **18**, 3025–3028 <https://doi.org/10.1016/j.bmcl.2008.01.062>
- 49 Wösten-van Asperen, R.M., Lutter, R., Specht, P.A., Moll, G.N., van Woensel, J.B., van der Loos, C.M. et al. (2011) Acute respiratory distress syndrome leads to reduced ratio of ACE/ACE2 activities and is prevented by angiotensin-(1-7) or an angiotensin II receptor antagonist. *J. Pathol.* **225**, 618–627 <https://doi.org/10.1002/path.2987>
- 50 Namsolleck, P. and Moll, G.N. (2020) Does activation of the protective renin-angiotensin system have therapeutic potential in COVID-19? *Mol. Med.* **26**, 80 <https://doi.org/10.1186/s10020-020-00211-0>
- 51 Breilting, S., Krauszman, A., Parihar, R., Walther, T., Friedberg, M.K. and Kuebler, W.M. (2015) Dose-dependent, therapeutic potential of angiotensin-(1-7) for the treatment of pulmonary arterial hypertension. *Pulm. Circ.* **5**, 649–657 <https://doi.org/10.1086/683696>
- 52 Durik, M., van Veghel, R., Kuipers, A., Rink, R., Haas-Jimoh Akanbi, M., Moll, G. et al. (2012) The effect of the thioether-bridged, stabilized angiotensin-(1-7) analogue cyclic ang-(1-7) on cardiac remodeling and endothelial function in rats with myocardial infarction. *Int. J. Hypertens.* **2012**, 536426 <https://doi.org/10.1155/2012/536426>

- 53 Pessôa, S., Becher, B., Van Veghel, P.M., De Vries, R., Tempel, R., Sneep, D. (2015) Effect of a stable angiotensin-(1-7) analogue on progenitor cell recruitment and cardiovascular function post myocardial infarction. *J. Am. Heart Assoc.* **4**, e001510 <https://doi.org/10.1161/JAHA.114.001510>
- 54 Cassis, P., Locatelli, M., Corna, D., Villa, S., Rottoli, D., Cerullo, D. et al. (2019) Addition of cyclic angiotensin-(1-7) to angiotensin-converting enzyme inhibitor therapy has a positive add-on effect in experimental diabetic nephropathy. *Kidney Int.* **96**, 906–917 <https://doi.org/10.1016/j.kint.2019.04.024>
- 55 Kuipers, A., Moll, G.N., Wagner, E. and Franklin, R. (2019) Efficacy of lanthionine-stabilized angiotensin-(1-7) in type 1 and type 2 diabetes mouse models. *Peptides* **112**, 78–84 <https://doi.org/10.1016/j.peptides.2018.10.015>
- 56 Kuipers, A., Moll, G.N., Levy, A., Krakovsky, M. and Franklin, R. (2020) Cyclic angiotensin-(1-7) contributes to rehabilitation of animal performance in a rat model of cerebral stroke. *Peptides* **123**, 170193 <https://doi.org/10.1016/j.peptides.2019.170193>
- 57 Wagenaar, G.T., Sengers, R.M., El Laghmani, H., Chen, X., Lindeboom, M.P., Roks, A.J. et al. (2014) Angiotensin II type 2 receptor ligand PD123319 attenuates hyperoxia-induced lung and heart injury at a low dose in newborn rats. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **307**, L261–L272 <https://doi.org/10.1152/ajplung.00345.2013>
- 58 Fontana, M., Brunori, A., Costa, M. and Antonucci, A. (1997) Detection of cystathione ketimine and lanthionine ketimine in human brain. *Neurochem. Res.* **22**, 821–824 <https://doi.org/10.1023/A:1022083809994>
- 59 Sandiford, S.K. (2019) Current developments in lantibiotic discovery for treating *Clostridium difficile* infection. *Expert Opin. Drug Discov.* **14**, 71–79 <https://doi.org/10.1080/17460441.2019.1549032>
- 60 Zeitlin, P.L., Boyle, M.P., Guggino, W.B. and Molina, L. (2004) A phase I trial of intranasal Moli1901 for cystic fibrosis. *Chest* **125**, 143–149 <https://doi.org/10.1378/chest.125.1.14>
- 61 Sokolove, P.M., Kester, M.B. and Westphal, P.A. (1991) Duramycin effects on the structure and function of heart mitochondria. II. Energy conversion reactions. *Arch. Biochem. Biophys.* **287**, 180–185 [https://doi.org/10.1016/0003-9861\(91\)90404-7](https://doi.org/10.1016/0003-9861(91)90404-7)
- 62 Zebedin, E., Koenig, X., Radenkovic, M., Pankevych, H., Todt, H., Freissmuth, M. et al. (2008) Effects of duramycin on cardiac voltage-gated ion channels. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **377**, 87–100 <https://doi.org/10.1007/s00210-007-0248-5>
- 63 Leonhardt, J., Villela, D.C., Teichmann, A., Münter, L.M., Mayer, M.C. and Mardah, M. (2017) Evidence for heterodimerization and functional interaction of the angiotensin type 2 receptor and the receptor MAS. *Hypertension* **69**, 1128–1135 <https://doi.org/10.1161/HYPERTENSIONAHA.116.08814>
- 64 Galande, A.K., Trent, J.O. and Spatola, A.F. (2003) Understanding base-assisted desulfurization using a variety of disulfide-bridged peptides. *Biopolymers* **71**, 534–551 <https://doi.org/10.1002/bjip.10532>
- 65 Knerr, P.J. and van der Donk, W.A. (2012) Chemical synthesis and biological activity of analogues of the lantibiotic epilancin 15X. *J. Am. Chem. Soc.* **134**, 7648–7651 <https://doi.org/10.1021/ja302435y>
- 66 Gieselmann, M.D., Zhu, Y., Zhou, H., Galonic, D. and van der Donk, W.A. (2002) Selenocysteine derivatives for chemoselective ligations. *ChemBiochem* **3**, 709–716 [https://doi.org/10.1002/1439-7633\(20020802\)3:8<709::AID-CBIC709>3.0.CO;2-8](https://doi.org/10.1002/1439-7633(20020802)3:8<709::AID-CBIC709>3.0.CO;2-8)
- 67 Ongey, E.L. and Neubauer, P. (2016) Lanthipeptides: chemical synthesis versus in vivo biosynthesis as tools for pharmaceutical production. *Microb. Cell. Fact.* **15**, 97 <https://doi.org/10.1186/s12934-016-0502-y>