

University of Groningen

Odontoblast markers and dentine reactions in carious primary molars with and without hypomineralised enamel defects

Shah, Janita; Manton, David J.; McCullough, Michael J.; Rajan, Sadna

Published in:
International Journal of Paediatric Dentistry

DOI:
[10.1111/ipd.12750](https://doi.org/10.1111/ipd.12750)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Shah, J., Manton, D. J., McCullough, M. J., & Rajan, S. (2021). Odontoblast markers and dentine reactions in carious primary molars with and without hypomineralised enamel defects. *International Journal of Paediatric Dentistry*, 31(4), 451-458. <https://doi.org/10.1111/ipd.12750>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Odontoblast markers and dentine reactions in carious primary molars with and without hypomineralised enamel defects

Janita Shah^{1,4}  | David J. Manton^{1,2}  | Michael J. McCullough³  | Sadna Rajan¹ 

¹Paediatric Dentistry, Melbourne Dental School, The University of Melbourne, Melbourne, Vic., Australia

²Centrum voor Tandheelkunde en Mondzorgkunde, Universitair Medisch Centrum Groningen, Groningen, The Netherlands

³Oral Anatomy, Medicine and Surgery, Melbourne Dental School, The University of Melbourne, Melbourne, Vic., Australia

⁴Oral Health Services, Health Care Agency, Mahé, Republic of Seychelles

Correspondence

Sadna Rajan, Melbourne Dental School, The University of Melbourne, 720 Swanston St, Carlton Vic. 3053, Australia.
Email: sadna.v@unimelb.edu.au

Abstract

Background: Wnt/ β -Catenin signalling and DMP1 have key roles in tertiary dentinogenesis.

Aim: To compare the relationship between remaining dentine thickness (RDT), tertiary dentine thickness (TDT), β -catenin and dentine matrix protein 1 (DMP1) in carious second primary molar teeth with normal (SPM) and hypomineralised enamel (HSPM).

Design: Extracted carious SPM and HSPM were fixed, sectioned (5 μ m) and stained with haematoxylin and eosin or with indirect immunofluorescence for β -catenin and DMP1. Image analysis was performed to determine RDT, TDT, β -catenin and DMP1 intensity in the odontoblast layer and dentine-pulp complex.

Results: Carious SPM ($n = 11$; mean RDT = 1536.1 μ m) and HSPM ($n = 12$; mean RDT = 1179.9 μ m) had mean TDT 248.6 μ m and 518.1 μ m, respectively ($P = .02$). There were no significant differences in intensity values in the odontoblast layer and dentine-pulp complex for β -catenin and DMP1 for both groups.

Conclusion: There was no observable variation in Wnt/ β -catenin and DMP1 expression between HSPM and SPM despite a statistically significant twofold increased TDT in HSPM compared with SPM that had similar RDT. Thus, the observed increased TDT in HSPM is more likely due to an earlier onset of repair processes rather than an amplified response to caries.

KEYWORDS

DMP1, hypomineralised enamel, odontoblasts, primary teeth, Wnt/ β -catenin

1 | INTRODUCTION

Odontoblasts are the first pulpal cells encountered by dentine-invading pathogens and their released products. They defend the pulp against bacterial infection by releasing antibacterial agents (nitric oxide, β -defensins and liposaccharide-binding proteins) and secreting pro-inflammatory cytokines and chemokines.¹ Their ability to produce tertiary dentine however remains their most important response to carious lesion progression. A mild pulpal inflammatory response, combined with the leaching of signalling molecules

and growth factors from the dentine, can stimulate tertiary dentine formation.²

Carious lesion progression involves bacteria and their by-products, breaking down dentinal tubule walls and releasing signalling molecules, such as the Small-Integrin-Binding Ligand N-linked Glycoproteins (SIBLINGs)³ and matrix metalloproteinases,¹ and growth factors such as tumour growth factor- β super family, insulin-like growth factor-1 and 2, fibroblast growth factor-2 and several angiogenic growth factors.² Dentine matrix protein 1 (DMP1), a member of the SIBLINGs family,³ is a highly

phosphorylated extracellular matrix protein that has been detected in inflamed permanent pulps, both within the sub-odontoblast cell layer at the periphery of the pulp and within the core of the pulp.⁴ DMP1 plays a key role in regulation of odontoblast differentiation and dentine mineralisation⁵ and has been identified as a marker for odontoblast differentiation⁶⁻⁹ and maturity.^{7,10,11} The presence of DMP1 in inflamed pulps and absence in normal pulps indicate that DMP1 also plays an important role in the inflammatory process,⁴ increasing the release of pro-inflammatory mediators (interleukin-6 and interleukin-8) from dental pulp fibroblasts, which are responsible for recruiting inflammatory cells such as neutrophils to the site of injury. Therefore, DMP1 may be considered as a marker for assessing the pro-inflammatory activity of pulpal fibroblasts and angiogenesis.⁴

In initial, slowly progressing carious lesions, signalling molecules interact with post-mitotic odontoblasts to upregulate their formation of reactionary dentine. Further lesion progression may result in necrosis of odontoblasts. The interaction of signalling molecules with dental pulp stem cells (DPSCs) can trigger their differentiation to odontoblast-like cells that produce reparative dentine.² Due to the cyclic nature of carious lesion progression, tertiary dentine is usually a combination of reactionary and reparative dentine.¹² The interaction of signalling molecules with odontoblasts and DPSCs can activate several signalling transduction pathways involved in tertiary dentinogenesis. These include p38 mitogen-activated protein kinase, transforming growth factor/small mother against decapentaplegic, mitogen-activated protein kinase/extracellular signal-regulated kinase, c-Jun N-terminal kinase and wingless/ β -catenin (Wnt/ β -catenin) pathways.¹³

The Wnt/ β -catenin signalling pathway plays an important role in tertiary dentinogenesis by promoting the proliferation and differentiation of DPSCs to odontoblast-like cells.¹⁴⁻¹⁷ This is thought to occur via the activation of the Runt-related transcription factor 2.¹⁵ The protein β -catenin is the main component of the Wnt/ β -catenin pathway and is believed to be the bottleneck through which signals pass.¹⁵ Odontoblasts are Wnt responsive¹⁴ and the Wnt/ β -catenin pathway activates both post-mitotic and odontoblast-like cells.¹⁸ Therefore β -catenin, which indicates activation of the Wnt/ β -catenin pathway, can be considered a marker for assessing tertiary dentinogenesis.¹³⁻¹⁵ Wnt/ β -catenin signalling also stimulates angiogenesis via the induction of angiogenic regulators such as interleukin-8.¹⁹ The upregulation of angiogenic signalling during pulpal inflammation contributes considerably to the pathogenesis associated with survival and differentiation of DPSCs into mature odontoblast-like cells.^{20,21} Therefore, angiogenic activity can be assessed through β -catenin markers that indicate activation of the Wnt/ β -catenin signalling pathway.²⁰

Why this paper is important to paediatric dentists

- There appears to be no differences in the intensity of the molecular processes involved in the stimulation of Wnt/ β -catenin and DMP1 between HSPM and SPM.
- With similar remaining dentine thickness, HSPM produce twofolds tertiary dentine compared to SPM.
- HSPM may mount an earlier response to caries compared to SPM. This has an implication on clinical management of HSPM.

There is limited research related to the role of Wnt/ β -catenin signalling and DMP1 in tertiary dentine formation in carious primary teeth with both normal and hypomineralised enamel. Therefore, the aim of this study is to compare the relationship between remaining dentine thickness (RDT), tertiary dentine thickness (TDT), and β -catenin and DMP1 biomarkers for tertiary dentine in carious second primary molar teeth with normal (SPM) and hypomineralised enamel (HSPM).

2 | MATERIALS AND METHODS

Ethical approval for the study was obtained from the University of Melbourne Human Ethics Research Committee (HREC number 1 749 762.1). Parental consent and child assent were obtained prior to tooth collection. A convenience sample of 23 extracted carious second primary molar teeth (SPM $n = 11$, HSPM $n = 12$) were collected from 23 healthy children (mean age 5.5 ± 1.6 y, range 3-7.5 y), requiring dental extractions under general anaesthesia at the Royal Dental Hospital Melbourne between April and December 2018. Teeth with clinical or radiographical signs of pulp pathosis were excluded. Collected teeth were deidentified prior to laboratory processing. The percentage of total root length that had undergone any resorption was calculated using Kramer and Ireland's normative data.²²

Immediately following extraction, teeth were sectioned at the root apices and placed in fixative (10% neutral buffered formalin) for 24 h at room temperature prior to decalcification in 20% ethylene diamine tetra-acetic acid solution at pH 7.4. Decalcified teeth were cut longitudinally through the carious lesion. Tooth halves were embedded in paraffin wax and sectioned at 5 μ m with a microtome through the deepest part of the carious lesion. Two sections per tooth were stained with either haematoxylin and eosin ($n = 46$) or stained using

indirect immunofluorescence for β -catenin ($n = 46$) and DMP1 ($n = 46$).

Prior to immunostaining, slides were immersed in boiling antigen retrieval solution (10 mM sodium citrate) and quenched (0.25% potassium permanganate) for 20 min to reduce autofluorescence. Slides were then incubated in phosphate-buffered saline and triton containing 10% goat serum for 30 min at room temperature. Following this, sections were labelled with either monoclonal antibody to β -catenin (rabbit anti- β -catenin E247, dilution 1:100, Abcam, Cambridge, UK (ab32572)) or polyclonal antibody to DMP1 (rabbit anti-DMP1, dilution 1:300, Abcam, Cambridge, UK (ab103203)) for 1 h at room temperature. Slides were subsequently incubated with polyclonal fluorescent secondary antibodies (goat anti-rabbit IgG H&L Alexa Fluor 488 pre-adsorbed, dilution 1:500, Abcam, Cambridge, UK (ab150081)) for 1 h at room temperature. Nuclear counterstaining (Hoechst 33 342, Abcam, Cambridge, UK (ab228551)) was performed for 10 min prior to cover slipping. No positive labelling was seen in any of the controls.

Haematoxylin and eosin-stained slides were viewed in Brightfield Standard Mode with the Nikon Eclipse TE2000-U microscope (Nikon Corporation, Japan) at magnifications of $\times 1.5$ and $\times 20$. Remaining dentine thickness (RDT/ μm), the shortest distance between the cavity base and pulp-dentine border, and tertiary dentine thickness (TDT/ μm), the longest distance between the cavity base and calcio-traumatic line, were obtained using the measurement tool in the Fiji (Fiji Is Just ImageJ, open source project hosted on GitHub, <https://fiji.sc/>) analysis software. Immuno-stained slides were viewed in Wide-Field Fluorescent Mode with the Nikon Eclipse TE2000-U microscope at X20 magnification. A TG-2A long pass filter cube was used to view Alexa Fluor 488 fluorescence, and a DAPI Bandpass filter cube was used to view the Hoechst 33 342 fluorescence. 16-bit multi-channel colour images of pulp sections subjacent to the cavity base were captured using a high sensitivity Nikon Ds-Fi1 colour camera (Nikon Corporation, Japan) with standardised exposure settings (333ms, Gamma = 0, Offset = 0). The colour images were converted to 16-bit grey images in Fiji, and subsequently, the mean grey value (MGV) of β -catenin and DMP1 within the odontoblast layer and the dentine-pulp complex were obtained using the measurement tool in Fiji.

A single examiner performed all the measurements and intra-examiner reproducibility was assessed by repeating measurements for RDT, TDT, β -catenin and DMP1 intensities in the odontoblast layer and dentine-pulp complex on 22% of the samples 1 week after the initial measurement.

The data obtained had normal distribution and therefore parametric tests were used. Pearson's correlation was used to test for statistically significant correlations between RDT, TDT and intensities of β -catenin and DMP1. The t test for independent samples determined whether there were any

statistically significant differences between the two groups: SPM and HSPM. The statistical significance level was set at $P < .05$.

3 | RESULTS

SPM (occlusal caries: $n = 9$; proximal caries: $n = 2$) were collected from children (4 boys, 7 girls) with mean age 5.6 ± 1.4 y, range 3-7.5 y. The degree of root resorption ranged between 7.4% and 63.2% (mean $27.5 \pm 16.7\%$) and RDT (remaining dentine thickness) ranged between 880.9 μm and 2165.7 μm . HSPM (occlusal caries: $n = 9$; proximal caries: $n = 3$) were collected from children (8 boys, 4 girls) with mean age 5.4 ± 1.8 y, range 3-7.5 y. The degree of root resorption ranged between 9.4% and 57.0% (mean $28.7 \pm 16.7\%$) and RDT ranged between 631.6 μm and 1957.2 μm .

Tertiary dentine was observed in nine of eleven SPM (mean RDT 1457.1 ± 459.0 μm ; Figure 1A-C). The two SPM without tertiary dentine had RDT of 2051.6 μm and 1731.9 μm . Tertiary dentine was observed in all HSPM ($n = 12$) (Figure 2A-C). TDT ranged between 0 μm and 579.8 μm in SPM, and between 150.6 μm and 1235.2 μm in HSPM.

DMP1 was detected within the peripheral odontoblast layer and the mineralised tertiary dentine subjacent to the base of the cavity, as well as deeper within the core of the pulp tissue (Figure 1D) in all SPM teeth. DMP1 was detected in supra- and intranuclear locations within the odontoblast-like cells (Figure 1F). The intensity of DMP1 ranged between 1290.5 MGv and 4457.8 MGv in the odontoblast layer, and between 123.7 MGv and 1164.1 MGv in the dentine-pulp complex of SPM. β -Catenin was detected within the peripheral odontoblast layer subjacent to the base of the cavity in all SPM teeth (Figure 1G). The accumulation was mainly localised within the nuclei of the odontoblast-like cells and DPCs (Figure 1I). The intensity of β -catenin ranged between 649.4 MGv and 2640.2 MGv in the odontoblast layer, and between 1920.0 MGv and 12 413.0 MGv in the dentine-pulp complex of SPM.

Similarly, in all HSPM teeth, DMP1 was detected within the peripheral odontoblast layer and the mineralised tertiary dentine subjacent to the base of the cavity, as well as deeper within the core of the pulp tissue (Figure 2D). DMP1 was also detected in supra- and intranuclear locations within the odontoblast-like cells (Figure 2F). With HSPM, the intensity of DMP1 ranged between 2138.6 MGv and 5821.6 MGv in the odontoblast layer, and between 231.6 MGv and 811.7 MGv in the dentine-pulp complex. β -Catenin was detected within the peripheral odontoblast layer subjacent to the base of the cavity in all HSPM samples (Figure 2G). The accumulation was mainly localised within the nuclei of the odontoblast-like cells and DPCs (Figure 2I). β -Catenin was




FIGURE 1 A, Occlusal view of a second primary molar with an occlusal cavitated lesion B, $\times 1.5$ Haematoxylin and eosin image C, $\times 20$ Haematoxylin and eosin image D, DMP1 (green); localisation of DMP1 in peripheral odontoblast layer (white arrow), and deeper within pulp (gold arrow) E, Nuclear staining with Hoechst 33 342 (blue) F, Merge of D & E; intranuclear (yellow arrow) and supranuclear (brown arrow) localisation of DMP1 G, β -catenin (red); localisation of β -catenin in peripheral odontoblast layer (white arrow) H, Nuclear staining with Hoechst 33 342 (blue) I, Merge of G & H; intranuclear localisation of β -catenin (yellow arrow). Abbreviations: TDT, tertiary dentine thickness; RDT, remaining dentine thickness; CT, calcitriatic line; TD, tertiary dentine; Od, odontoblast-like cells; DMP1, dentine matrix protein 1; B-catenin, β -catenin

also noted in the endothelium lining of nearby pulpal blood vessels (Figure 3B). With HSPM, the intensity of β -catenin ranged between 501.1 MGv and 3220.0 MGv in the odontoblast layer, and between 1675.0 MGv and 14 389.0 MGv in the dentine-pulp complex. All measurements were within the 95% limits of agreement showing high intra-examiner reproducibility.

A moderately negative correlation was noted between RDT and TDT in SPM (Pearson's coefficient = -0.494 , $P = .061$) and in HSPM (Pearson's coefficient = -0.461 , $P = .066$). There was a significant difference between the mean TDT in SPM ($248.6 \pm 205.8 \mu\text{m}$) and HSPM ($518.1 \pm 282.0 \mu\text{m}$) with a 108.4% (approximately twofold) increase in the mean TDT in HSPM (Table 1). There were no statistically significant correlations between RDT and the intensity of β -catenin and DMP-1 in the odontoblast layer and dentine-pulp

complex in both SPM and HSPM teeth. There was no statistically significant difference in the mean RDT in SPM and HSPM (Table 1). Although the mean intensity of β -catenin and DMP1 in the odontoblast layer and dentine-pulp complex were greater in HSPM compared to SPM, this was not statistically significant (Table 1).

4 | DISCUSSION

Tertiary dentine is deposited subjacent to the base of the carious lesion at pulp-dentine interface and its thickness is proportional to the duration and intensity of the stimulus.¹² As lesion progression becomes closer to pulp tissue, injury to pulp triggers the formation of tertiary dentine. Increased inflammatory changes in pulpal innervation, vascularity and immune cell




FIGURE 2 A, Occlusal view of a hypomineralised second primary molar with a disto-occlusal cavitated lesion B, $\times 1.5$ Haematoxylin and eosin image C, $\times 20$ Haematoxylin and eosin image D, DMP1 (green); localisation of DMP1 in peripheral odontoblast layer (white arrow), and deeper within pulp (gold arrow) E, Nuclear staining with Hoechst 33 342 (blue) F, Merge of D & E; intranuclear (yellow arrow) and supranuclear (brown arrow) localisation of DMP1 G, β -catenin (red); localisation of β -catenin in peripheral odontoblast layer (white arrow) H, Nuclear staining with Hoechst 33 342 (blue), I, Merge of G & H; intranuclear localisation of β -catenin (yellow arrow). Abbreviations: TDT, tertiary dentine thickness; RDT, remaining dentine thickness; CT, calciotraumatic line; TD, tertiary dentine; Od, odontoblast-like cells; DMP1, dentine matrix protein 1; B-catenin, β -catenin

accumulation,²³ as well as an increase in pulpal TRPV1 expression and vascular flow²⁴ have been reported in hypomineralised teeth compared to teeth with normal enamel. It has been suggested that sensitive intact hypomineralised first permanent molars have an underlying pulpal inflammation triggered by an ingress of biofilm-derived acids, bacteria,²⁵ and their by-products into the dentinal tubules via the porosities in the hypomineralised enamel²³ leading to a cascade of events resulting in earlier tertiary dentine formation.^{25,26} In this study, we found no statistically significant difference in the mean RDT in carious SPM and HSPM, but a statistically significant twofold increase in mean TDT in carious HSPM when compared to carious SPM suggests an amplified or an earlier onset of dentinal repair processes in the pulps of HSPM.

Pulpal injury and inflammation activate the Wnt/ β -catenin signalling pathway which plays an important role in tertiary

dentogenesis. Implantation of an exogenous liposome-reconstituted form of Wnt3A protein into mice molar pulps activated Wnt/ β -catenin signalling, promoted differentiation of odontoblast-like cells and reduced apoptosis, culminating in a superior repair response.¹⁴ Similarly, direct pulp capping of mice molars with glycogen synthase kinase 3 (GSK-3) inhibitors, Wnt/ β -catenin signalling agonists, resulted in greater reparative dentine formation at pulp exposure sites when compared with plain collagen sponge or mineral trioxide aggregate.²⁷⁻²⁹ The reparative dentine was found to be close to native dentine compositions at defect sizes translatable to small human lesions.²⁹ Animal immunofluorescence studies performed on dental pulps have demonstrated the translocation of β -catenin into nuclei of the odontoblast-like cells coating the reparative dentine.¹⁵⁻¹⁷ The distribution of β -catenin within the peripheral odontoblast layer was noted to be




FIGURE 3 Presence of β -catenin in cells lining blood vessels in the pulp of a carious hypomineralised second primary molar A, Haematoxylin and eosin staining x20 magnification B, immunofluorescence (red) detection of β -catenin within peripheral odontoblast layer and endothelial lining of blood vessels C, Nuclear staining with Hoechst 33 342 (blue) (D) β -catenin and nuclear staining merged. Abbreviations: Od, odontoblast-like cells; bv, blood vessels; B-catenin, β -catenin

TABLE 1 Difference between mean RDT, TDT and intensity of β -catenin and DMP1 in HSPM and SPM teeth

	HSPM ^a		SPM ^b		(t test) P
	Mean	SD	Mean	SD	
RDT (μ m)	1179.9	471.7	1536.1	443.5	.08
TDT (μ m)	518.1	282.0	248.6	205.8	.02*
β -catenin (Od) (MGV)	1931.4	928.5	1425.7	726.7	.16
β -catenin (DPC) (MGV)	8129.4	3912.0	6670.1	3085.2	.34
DMP1 (Od) (MGV)	3360.1	1012.6	2865.6	897.4	.23
DMP1 (DPC) (MGV)	541.4	187.6	495.1	296.5	.66

Abbreviations: DMP1, dentine matrix protein 1; DPC, dentine-pulp complex; HSPM, hypomineralised second primary molar; MGV, mean grey value (unit of measurement of intensity); RDT, remaining dentine thickness; SD, standard deviation, P, two-tail significance; SPM, second primary molar; TDT, tertiary dentine thickness, Od, odontoblast-like cells; μ m, micrometre.

^an = 12.

^bn = 11.

* $P < .05$.

broader and deeper at the exposure site or base of the cavity, subjacent to the reparative dentine.¹⁵⁻¹⁷ The findings of this study are in agreement with the results of the animal immunofluorescence studies. The presence of β -catenin in the peripheral odontoblast layer of pulps without tertiary dentine formation suggest that the Wnt/ β -catenin signalling pathway had been activated in the dental pulp cells activating stem cell

differentiation to form odontoblast-like cells. An interesting finding was the presence of β -catenin in the endothelial lining of nearby blood vessels. Angiogenic effects of the Wnt/ β -catenin pathway have been reported,¹⁹⁻²¹ and the findings of this study suggest that the β -catenin in the endothelial cells was promoting endothelial cell proliferation and angiogenesis via the Wnt/ β -catenin signalling pathway.

DMP1 is another important biomarker for odontoblast differentiation that has been investigated in this study. Direct pulp capping of rat maxillary first molars with recombinant DMP1-impregnated collagen matrix,⁶ and with Delta like 1 homolog, a DMP1 agonist,³⁰ stimulated the differentiation of mouse stem cells to odontoblast-like cells resulting in reparative dentine deposition. Immunohistochemical studies have localised DMP1 within the nuclei of undifferentiated DPSCs and odontoblast-like cells at pulp exposure sites in rat⁷ and human primary⁹ molars. DMP1 has a dual role in tertiary dentinogenesis.⁴ Within the nucleus, DMP1 regulates specific genes that control odontoblast-like differentiation.^{7,8} Within the supranuclear (Golgi) region of secretory odontoblast-like cells and the mineralising dentine matrix,⁷⁻¹⁰ DMP1 is involved in dentine mineralisation. The findings of this study indicate that DMP1 has a dual role in tertiary dentinogenesis. In addition to the peripheral odontoblast layer, DMP-1 has also been localised deeper within the pulp tissue of inflamed permanent pulps with normal enamel where it is thought to play an important role in inflammation by stimulating the release of interleukin-6 and interleukin-8 from pulp fibroblasts, which are responsible for recruiting inflammatory cells such as neutrophils to the site of injury.⁴ To the best of our knowledge, this is the first study to localise DMP1 in inflamed pulps of human primary molars with normal and

hypomineralised enamel, and the findings are in agreement with a previous study.¹¹ Therefore, in addition to its active involvement in the dentinal repair process in carious HSPM and SPM, DMP1 may also be participating in the development of inflammatory changes in the pulp by stimulating pro-inflammatory activity of pulp fibroblasts.

Research investigating clinical applications of pulp healing and repair mechanisms in primary teeth is still in its infancy and is currently limited to animal studies and histological outcomes. Due to the limited sample size and the findings being histological observations rather than experimental findings, it is difficult to make definitive remarks on the clinical implications of the results of this study. Nevertheless, the results have provided further insights into the molecular processes involved in tertiary dentinogenesis in carious HSPM and SPM. A statistically significant two-fold increase in mean TDT in carious HSPM when compared to carious SPM suggests an amplified or an earlier onset of repair processes in the dentine-pulp complex of HSPM. The absence of a statistically significant difference between mean intensity of β -catenin and DMP1 in the odontoblast layer and dentine-pulp complex in carious HSPM and SPM however suggests that the intensity of molecular processes involved in tertiary dentinogenesis via the stimulation of Wnt/ β -catenin and DMP1 in HSPM and SPM are similar, and the increased TDT in HSPM pulps is more likely due to an earlier onset of repair processes rather than an amplified response to caries. An extension of the present project, with a larger sample size to observe interactions between the many variables, would be valuable and may identify further significant relationships that may have been lost to type II error.

5 | CONCLUSION

Similar Wnt/ β -catenin and DMP1 activities were observed in the peripheral odontoblast layer and the dentine-pulp complex between HSPM and SPM. The pulps of HSPM appear to respond earlier to carious lesions, via activation of Wnt/ β -catenin and DMP1 signalling pathways, to produce a twofold increased TDT compared to SPM with similar RDT.

ACKNOWLEDGEMENTS

We would like to thank the various teams from The University of Melbourne for their technical support: Mr Paul McMillan and Ellie Cho from the Biological Optical Microscopy Platform, Advanced Microscopy Facility; Ms Su Toulson, Ms Deanne Catmull and Prof. Rodrigo Mariño from Melbourne Dental School; and Ms Laura Leone from the School of Biomedical Sciences.

CONFLICT OF INTEREST

All authors deny any conflicts of interest in this study.

AUTHOR CONTRIBUTIONS

SR and JS conceived idea. JS, DJM, MJM and SR designed the project. JS collected data; JS and SR contributed to analysis. JS and SR lead the writing. JS, DJM, MJM and SR approved the final manuscript.

ORCID

Janita Shah  <https://orcid.org/0000-0002-1390-2455>

David J. Manton  <https://orcid.org/0000-0002-4570-0620>

Michael J. McCullough  <https://orcid.org/0000-0001-7829-4684>

[org/0000-0001-7829-4684](https://orcid.org/0000-0001-7829-4684)

Sadna Rajan  <https://orcid.org/0000-0002-4576-4894>

REFERENCES

1. Farges J-C, Alliot-licht B, Renard E, et al. Dental pulp defence and repair mechanisms in dental caries. *Mediat Inflamm.* 2015;2015:230251.
2. Cooper PR, Takahashi Y, Graham LW, Simon S, Imazato S, Smith AJ. Inflammation–regeneration interplay in the dentine–pulp complex. *J Dent.* 2010;38(9):687-697.
3. Fisher LW, Fedarko NS. Six genes expressed in bones and teeth encode the current members of the SIBLING family of proteins. *Connect Tissue Res.* 2003;44(suppl 1):33-40.
4. Abd-Elmeguid A, Yu D, Kline WL, Moqbel R, Vliagoftis H. Dentin Matrix Protein-1 Activates Dental Pulp Fibroblasts. *J Endod.* 2012;38:75-80.
5. Narayanan K. Dual functional roles of dentine matrix protein 1 - Implications in biomineralisation and gene transcription by activation of intracellular Ca²⁺ store. *J Biol Chem.* 2003;278(19):17500.
6. Almushayt A, Narayanan K, Zaki AE, George A. Dentine matrix protein 1 induces cytodifferentiation of dental pulp stem cells into odontoblasts. *Gene Ther.* 2005;13(7):611-620.
7. Shigetani Y, Yoshida K, Kuratate M, et al. Temporospatial localization of dentine matrix protein 1 following direct pulp capping with calcium hydroxide in rat molars. *Int Endod J.* 2015;48(6):573-581.
8. Aguiar MC, Arana-Chavez VE. Immunocytochemical detection of dentine matrix protein 1 in experimentally induced reactionary and reparative dentine in rat incisors. *Arch Oral Biol.* 2010;55(3):210-214.
9. Lourenco Neto N, Marques NC, Fernandes AP, et al. Immunolocalisation of dentine matrix protein-1 in human primary teeth treated with different pulp capping materials. *J Biomed Mater Res B.* 2016;104(1):165-169.
10. Massa LF, Ramachandran A, George A, Arana-Chavez VE. Developmental appearance of dentine matrix protein 1 during the early dentinogenesis in rat molars as identified by high-resolution immunocytochemistry. *Histochem Cell Biol.* 2005;124(3):197-205.
11. Simon S, Smith AJ, Lumley PJ, et al. Molecular characterisation of young and mature odontoblasts. *Bone.* 2009;45(4):693-703.
12. Smith AJ, Cassidy N, Hb P, Bègue-Kirn C, Ruch VJ, Lesot H. Reactionary dentinogenesis. *Int J Dev Biol.* 1995;39:273-280.
13. da Rosa WLO, Piva E, da Silva AF. Disclosing the physiology of pulp tissue for vital pulp therapy. *Int Endod J.* 2018;51(8):829-846.
14. Hunter D, Bardet C, Mouraret S, et al. Wnt Acts as a Prosurvival Signal to Enhance Dentine Regeneration. *J Bone Miner Res.* 2015;30(7):1150-1159.
15. Han N, Zheng Y, Li R, et al. Beta-catenin enhances odontoblastic differentiation of dental pulp cells through activation of Runx2. *PLoS One.* 2014;9(2):e88890.

16. Yoshioka S, Takahashi Y, Abe M, et al. Activation of the Wnt/ β -catenin pathway and tissue inhibitor of metalloprotease 1 during tertiary dentinogenesis. *J Biochem.* 2013;153(1):43-50.
17. Babb R, Chandrasekaran D, Carvalho Moreno Neves V, Sharpe PT. Axin2-expressing cells differentiate into reparative odontoblasts via autocrine Wnt/ β -catenin signaling in response to tooth damage. *Sci Rep.* 2017;7(1):3102.
18. Han XL, Liu M, Voisey A, et al. Post-natal effect of overexpressed DKK1 on mandibular molar formation. *J Dent Res.* 2011;90(11):1312-1317. <https://doi.org/10.1177/0022034511421926>
19. Masckauchan N, Shawber C, Funahashi Y, Li C-M, Kitajewski J. Wnt/ β -Catenin Signaling Induces Proliferation, Survival and Interleukin-8 in Human Endothelial Cells. *Angiogenesis.* 2005;8:43-51.
20. Matsushita K, Motani R, Sakuta T, et al. The role of vascular endothelial growth factor in human dental pulp cells: induction of chemotaxis, proliferation, and differentiation and activation of the AP-1-dependent signaling pathway. *J Dent Res.* 2000;79(8):1596-1603.
21. Tran-Hung L, Laurent P, Camps J, About I. Quantification of angiogenic growth factors released by human dental cells after injury. *Arch Oral Biol.* 2008;53(1):9-13.
22. Kramer W, Ireland R. Measurements of the primary teeth. *J Dentistry Child.* 1959;26:252-261.
23. Rodd H, Boissonade F, Day P. Pulpal status of hypomineralised permanent molars. *Paediatr Dent.* 2007;29(6):514-520.
24. Rodd H, Christmas C, Day P, Boissonade MF. Pulpal expression of TRPV1 in molar incisor hypomineralisation. *Eur Arch Paediatr Dent.* 2008;8:184-188.
25. Fagrell TG, Lingstrom P, Olsson S, Steiniger F, Noren JG. Bacterial invasion of dentinal tubules beneath apparently intact but hypomineralized enamel in molar teeth with molar incisor hypomineralisation. *Int J Paediatr Dent.* 2008;18(5):333-340.
26. Heijs S, Dietz W, Norén J, Blanksma N, Jälevik B. Morphology and chemical composition of dentin in permanent first molars with the diagnose MIH. *Swed Dent J.* 2007;31:155-164.
27. Neves VCM, Babb R, Chandrasekaran D, Sharpe PT. Promotion of natural tooth repair by small molecule GSK3 antagonists. *Article. Sci Rep.* 2017;7:39654.
28. Babb RC, Chandrasekaran D, Zaugg LK, Sharpe PT. A Mouse Model to Study Reparative Dentinogenesis. *Methods Mol Biol.* 2019;1922:111-119. https://doi.org/10.1007/978-1-4939-9012-2_11
29. Zaugg LK, Banu A, Walther AR, et al. Translation Approach for Dentine Regeneration Using GSK-3 Antagonists. *J Dent Res.* 2020;99(5):544-551.
30. Walker JV, Zhuang H, Singer D, et al. Transit amplifying cells coordinate mouse incisor mesenchymal stem cell activation. *Nat Commun.* 2019;10(1):3596.

How to cite this article: Shah J, Manton DJ, McCullough MJ, Rajan S. Odontoblast markers and dentine reactions in carious primary molars with and without hypomineralised enamel defects. *Int J Paediatr Dent.* 2020;00:1–8. <https://doi.org/10.1111/ipd.12750>