

University of Groningen

Extraction and Quantitative Analysis of Goat Milk Oligosaccharides

Chatziioannou, Anastasia Chrysovalantou; Benjamins, Eric; Pellis, Linette; Haandrikman, Alfred; Dijkhuizen, Lubbert; van Leeuwen, Sander S.

Published in:
Journal of Agricultural and Food Chemistry

DOI:
[10.1021/acs.jafc.1c00499](https://doi.org/10.1021/acs.jafc.1c00499)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2021

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Chatziioannou, A. C., Benjamins, E., Pellis, L., Haandrikman, A., Dijkhuizen, L., & van Leeuwen, S. S. (2021). Extraction and Quantitative Analysis of Goat Milk Oligosaccharides: Composition, Variation, Associations, and 2'-FL Variability. *Journal of Agricultural and Food Chemistry*, 69(28), 7851-7862. <https://doi.org/10.1021/acs.jafc.1c00499>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Extraction and Quantitative Analysis of Goat Milk Oligosaccharides: Composition, Variation, Associations, and 2'-FL Variability

Anastasia Chrysovalantou Chatziioannou, Eric Benjamins, Linette Pellis, Alfred Haandrikman, Lubbert Dijkhuizen, and Sander S. van Leeuwen*

Cite This: *J. Agric. Food Chem.* 2021, 69, 7851–7862

Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Human milk oligosaccharides (hMOS) are associated with health benefits for newborns. We studied the composition of goat MOS (gMOS) from colostrum up to the 9th month of lactation to conceive an overview of the structures present and their fate. Potential correlations with factors such as age, parity, and lifetime milk production were examined. An effective method for gMOS extraction and ultra-high-performance liquid chromatography coupled to fluorescence detection (UPLC-FLD) analysis was established, following 2-aminobenzamide gMOS labeling. Considerable biological variability was highlighted among the 12 quantified gMOS and the 9 non-quantified structures in the individual milk samples. Most characteristic, 2'-fucosyllactose was present in 73.7% of the milk samples analyzed, suggesting the possibility of a secretor/non-secretor goat genotype, similar to humans. Contributing factors to the observed biological variability were goat age, parity, lifetime milk production, and the kids' sex. The results significantly contribute to the current understanding of (variations in) gMOS composition.

KEYWORDS: goat milk, UPLC-FLD, oligosaccharides, 2'-fucosyllactose, secretor

INTRODUCTION

One of the most abundant classes of macronutrients found in human milk are carbohydrates, mainly lactose. Among carbohydrates, the free milk oligosaccharides, referring to glycans that are not bound to glycoproteins, are suggested to have significant biological functions. Human milk oligosaccharides (hMOS) are present in relatively high concentrations in human milk (5–20 g/L).¹ The diversity of the structures of these hMOS is high. It is mentioned that in human milk, 162 hMOS have been structurally identified and more than 240 compositional varieties have been annotated.² These are all based on five monosaccharide units: galactose (Gal), glucose (Glc), *N*-acetylglucosamine (GlcNAc), fucose (Fuc), and *N*-acetylneuraminic acid (Neu5Ac). The typical length of these hMOS is between three and seven monosaccharide units. Composition of hMOS differs between individual mothers due to genetics, diet, and epigenetics, and within the mothers over the course of lactation.^{3–6} However, most variance is yet unexplained.

Recent studies have unraveled the importance of hMOS, providing a long list of functionalities.^{7,8} Besides the hMOS “prebiotic” character toward certain beneficial bacteria (e.g., *Bifidobacterium longum* subsp. *infantis*), it is reported that they inhibit pathogen adhesion to epithelial cells⁹ and have immune modulation effects.¹⁰ Protection of the neonate from certain diseases (e.g., necrotizing enterocolitis) has been reported in animal and human studies,^{11,12} while possible enhanced offspring brain development has been studied in animal studies.^{13,14} Potential protective effects of hMOS toward the breastfeeding mothers have been suggested, either by influencing the microbiota of the mammary gland or by contributing to its epithelial cell responses.⁸

The high structural diversity of hMOS has not been reported for MOS of domesticated animals (non-human mammals).^{15,16}

In all studies of domesticated animals, less variety of structures is reported together with an overall lower amount of MOS. In hMOS, a high percentage of the compounds found are neutral and mostly (~70%) fucosylated,¹⁷ while this is not the case for the MOS of domesticated dairy animals, which are mostly sialylated structures.¹⁷


Undoubtedly, human milk is the preferable source of oligosaccharides for neonates. However, not all mothers can or want to breastfeed. These mothers depend on infant formula, where bovine milk is the most chosen option for infant formula production. Several studies have reported on bovine MOS composition, reviewed recently.¹⁸ The differences between bovine milk and human milk in terms of MOS content and structural diversity have been highlighted by those studies.^{8,15} Goat milk has higher levels of MOS compared to other non-human mammals and a richer variety of structures.¹⁹ Goat milk-based infant formula is becoming increasingly popular. However, goat milk oligosaccharides (gMOS) studies are limited, especially those providing quantitative data. Some of those providing quantitative information are based on rather small sample sizes.^{20–22} Other gMOS studies cover a selection of predefined targets resulting in underestimation of the total gMOS content.²³ Different analytical instruments and MOS

Received: February 1, 2021

Revised: May 25, 2021

Accepted: June 25, 2021

Published: July 7, 2021






Figure 1. Milk sample collection schedule per goat during lactation. The collection was performed as follows: during the first 3 days (three milk collections per day); during days 4–10 (one milk collection per day); every week (one milk collection per week for the rest of the first month); and every month (one milk collection per month).

extraction methods may impact the precision of the quantitation. Finally, some gMOS studies focus only on specific structures (i.e., 3', 6'-sialyllactose and disialyllactose²³), not facilitating the understanding of the complete gMOS "fingerprint".²⁴

In this study, a high-throughput method for the extraction of MOS from goat milk samples and their analysis with ultra-high-performance liquid chromatography coupled to fluorescence detection (UPLC-FLD) was established. In total, gMOS composition was assessed for 57 Dutch Saanen goats for 9 months over the course of lactation from day 1 to mature milk, aiming to provide average concentrations of the identified structures. The 57 goats of the study were divided over two independent farms, following different feeding regimes, providing the chance to evaluate feeding as a factor in gMOS composition. Moreover, associations between oligosaccharide composition and factors, such as age, parity, diet, volume of milk produced, and milk characteristics, were studied.

EXPERIMENTAL SECTION

Materials. The standards of 3'-galactosyllactose (3'-GL), 4'-galactosyllactose (4'-GL), 6'-galactosyllactose (6'-GL), 6'-sialyl-N-acetyllactosamine (6'-SLN), and lacto-N-neotetraose (LNnT) of minimum 95% purity were purchased from Carbosynth (Berkshire, U.K.). 2'-Fucosyllactose (2'-FL), 3-fucosyllactose (3-FL), 3'-sialyllactose (3'-SL), and 6'-sialyllactose (6'-SL) of >95% purity were from IsoSep AB (Tullinge, Sweden).

Ultrapure water (18.2 MΩ) was generated by a Milli-Q water system (Millipore Merck).

Methanol (MeOH) and acetonitrile (MeCN) (HPLC gradient grade) were from Boom BV (Meppel, the Netherlands). Anhydrous acetic acid (glacial, 100%) was from Merck (Darmstadt, Germany).

2-Methylpyridine borane complex (95%), 2-aminobenzamide (2-AB, ≥98%), dimethyl sulfoxide (DMSO, ≥99.5%), and cellulose microcrystalline powder were from Sigma-Aldrich (Steinheim, Germany).

Trifluoroacetic acid (TFA, 99%) was purchased by Acros Organics (New Jersey).

The 150 mg/4 mL porous graphitic carbon (PGC) solid-phase extraction (SPE) cartridges (Extract-Clean SPE Carbo) for the removal of excessive amounts of lactose were purchased from Aurora Borealis Control BV (Schoonebeek, the Netherlands). The 96-well plate filters used for the removal of the excessive labeling reagent and the final purification of the oligosaccharides before injection in the analytical system were the AcroPrep 96-well 0.2 μm GHP from Pall Corporation (Puerto Rico). The Amicon Ultra-15 spin-filters with a 10 kDa cutoff were from Millipore.

Goat Milk Samples. Goat milk was obtained from two Dutch goat farms between February 2018 and January 2019. Goats were individually milked using a bucket milking system. Full milkings were

homogenized, and samples of ~10 mL were collected from 57 Dutch Saanen (Melkgeit) goats. The samples were stored at −20 °C in the laboratory until collection and then stored at −80 °C in the laboratory until further analysis. The kidding and, hence, the offset of sample collection fell within a 44-day span for all goats of farm #1 (25 February to 11 April 2018) and within 14 days for farm #2 (26 February to 13 March 2018). The two farms followed different feeding regimes, farm #1 using homogenized (pelletized) feed at libitum and farm #2 using diverse feeding based on crop excess in the local area, supplemented by pelletized feed. The sample collection was performed at multiple time points per lactation period (Figure 1). For colostrum milk, three samples per day were collected in the first 3 days of lactation (day 1 (D1), day 2 (D2), and day 3 (D3)). For the next 7 days, one sample was collected per day, and after this, one sample per week till the end of the first month (day 31 (D31)) of lactation. The sample collection was continued for the next 8 months, collecting one sample per month. A total of 28 samples per goat were collected from the 57 goats of the two farms. Moreover, tank (pooled) mature milk of all farms providing milk to Ausnutria BV and pooled colostrum milk from one farm was used as reference milk during the optimization of the extraction method and as a quality control (QC) sample during the analysis of the samples of the individual goats.

Optimization of Oligosaccharide Extraction and Lactose Removal. Optimization experiments for the efficient extraction of gMOS were executed with the pooled sample of colostrum and the pooled sample of mature goat milk (reference milk). First, direct labeling to 20 μL of reference goat milk was performed, following the Austin et al. approach for human milk.²⁵ Moreover, direct centrifugation (21 130g for 30 min) of reference goat milk and reference goat colostrum was attempted, prior to the labeling of the defatted extracts. The same tests were repeated after dilution of the above-mentioned samples with Milli-Q water in a 1:1 ratio.

Additional experiments were performed at larger scale to obtain an enriched gMOS profile. For this purpose, precipitation of the caseins by lowering the pH to 4.6 was attempted. For this test, a volume of 10 mL of reference mature goat milk was employed, diluted 1:1 with Milli-Q water. The pH was reduced to 4.6 by gradual addition of 1 M HCl. The samples were centrifuged (4000g for 30 min at 4 °C), and the supernatant was collected and applied to spin-filters with a 10 kDa cutoff. Subsequently, the decaseinated fraction was further processed to gMOS analysis as described below.

Another larger-scale method aimed to "salt out" the proteins by the slow addition of ammonium sulfate. To 10 mL of reference goat milk diluted 1:1 with Milli-Q water, 10 g of ammonium sulfate was added. The samples were centrifuged (4000g for 30 min at 4 °C), and the supernatant was collected. The deproteinated sample was processed as described below to analyze gMOS.

Since the above experiments highlighted the problem of lactose interference with the obtained gMOS profiles, further tests were performed to achieve sufficient and selective removal of lactose. Extraction of gMOS with acetone and methanol at a sample-to-solvent ratio of 1:2 was applied prior to porous graphitic carbon (PGC) solid-phase extraction (SPE). Also, direct application of undiluted milk (1

Table 1. Oligosaccharides Identified and Quantified in the 57 Goat Milk Samples of the Study and Their Concentrations (mg L⁻¹) at Three Time Points: Colostrum of the First and Second Days and Mature Milk of the 31st Day of Lactation^a

Abbreviation	Name	hMOS	Structure	Mass detected (Da)	Retention time (min)	Day 1 1st Colostrum sample			Day 2 4th Colostrum sample			Day 31		
						Mean conc. mg/L	Min	Max	Mean conc. mg/L	Min	Max	Mean conc. mg/L	Min	Max
2'-FL	2'-Fucosyl-lactose	Y		609.2	13.1	41.41	2.14	268.95	**** 62.92	0.58	236.30	5.12	0.35	69.51
3'-GL	3'-Galactosyl-lactose	Y		625.2	19.1	147.94	62.67	300.60	**** 51.96	12.63	123.82	31.69	13.62	74.40
6'-GL	6'-Galactosyl-lactose	Y		625.2	20	65.67	5.27	170.67	**** 13.84	0.75	71.95	8.05	1.03	36.31
3'-SL	3'-Sialyl-lactose	Y		754.3	21.3	198.92	28.91	429.48	**** 72.53	2.48	242.18	43.62	0.42	98.00
6'-SLN	6'-Sialyl-N-acetyllactosamine	N		795.3	23.0	107.2	3.1	281.8	**** 27.32	0.48	81.03	7.36	0.05	15.54
3'-NGL	3'-N-glycolyl-neuraminyllactose	N		770.3	24.4	57.71	6.8	171.47	41.88	3.69	142.25	45.32	6.03	163.68
6'-SL	6'-Sialyl-lactose	Y		754.3	24.6	284.41	30.11	674.87	**** 87.93	6.41	220.37	41.47	6.69	187.34
NGLN	N-glycolyl-neuraminyllactosamine	N		811.3	25.6	48.29	2.39	188.44	**** 19.88	0.65	109.20	2.98	0.37	9.34
6'-NGL	6'-N-glycolyl-neuraminyllactose	N		770.3	27.4	11.93	3.02	29.55	*** 45.89	9.43	150.55	30.29	5.84	82.78
DSL	Disialyl-lactose	N		1045.4	28.0	66.1	7.0	209.7	*** 3.39	0.26	11.38	2.19	0.14	7.30
SHL	Sialyl-galactosyl-lactose	N		916.3	28.3	47.98	14.77	130.63	**** 3.67	0.10	11.58	2.46	0.08	7.60
NADHL	N-acetylglucosaminyl-dihexosyl-lactose	N		990.4	33.1	25.47	4.03	67.67	**** 6.90	0.48	17.07	0.69	0.38	1.03
Total OS						1102.95			438.11			221.25		
Total Acidic OS						822.46 (75%)			302.49 (69%)			175.70 (79%)		
Total Neutral OS						280.49 (25%)			135.62 (31%)			45.55 (21%)		
Total Fucosylated OS						41.41 (4%)			62.92 (14%)			5.12 (2%)		
Total Sialylated OS						822.46 (75%)			302.49 (69%)			175.70 (79%)		

^aThe symbols used in the graphical presentation of structures correspond to Glc: blue circle filled, Gal: yellow circle filled, Fuc: red triangle filled, GlcNAc: blue square filled, Neu5Ac: pink diamond filled, Neu5Gc: turquoise diamond filled. The column hMOS indicates the presence (Y) or absence (N) of the structure in human milk studies. The stars indicate the result of the Wilcoxon paired *t*-test performed between D2 and D31 (***p* < 0.001, ****p* < 0.0001). The data presented are for 57 goats, except for 2'-FL, which was detected in 42 goats.

mL) to PGC columns was investigated. For the PGC-SPE filtration of the goat milk samples, the protocol of Robinson et al.²⁶ was applied. The SPE cartridges were initially conditioned by adding two times 1 mL of 80% acetonitrile (MeCN) in Milli-Q water containing 0.1% trifluoroacetic acid (TFA), followed by an equilibration step using three times 1 mL of Milli-Q water with 0.1% TFA. After their reconstitution in 1 mL of Milli-Q water, the samples were applied to SPE cartridges. A series of different washing steps were tested; Milli-Q/0.1% TFA, 2% MeCN/0.1% TFA, 4% MeCN/0.1% TFA, and 6% MeCN/0.1% TFA. Finally, the purified gMOS was eluted by two times with 0.5 mL of 50% MeCN/0.1% TFA under all tested conditions. MeCN was removed from the samples under reduced pressure (Speedvac Concentrator, Thermo Fisher Scientific) at 45 °C, and the samples were freeze-dried for optimal labeling results.

Labeling of Oligosaccharides. After removal of excess lactose, the extracted and freeze-dried gMOS were labeled with the fluorophore 2-aminobenzamide (0.7 M, 2-AB) and catalyzed with 1 M 2-picoloneborane according to Ruhaak et al.,²⁷ by a reductive amination reaction. This labeling mix was prepared freshly in 7:3 dimethyl sulfoxide (DMSO)/glacial acetic acid solution; 50 μL of labeling mix was added per sample, vortexed, centrifuged, and incubated for 2 h at 65 °C.

Excessive amounts of free labeling reagent and reducing agent (2-picoloneborane) were removed by filtering in a 96-well plate format with microcrystalline cellulose, following a previously reported method with small modifications.²⁸ A 200 mg/mL microcrystalline cellulose suspension in water was prepared, and 200 μL was applied to each well of the 0.2 μm GHP hydrophilic polypropylene filter. After washing each well with 200 μL (repeated twice), they were equilibrated. Four different concentrations were tested for the equilibration/washing of the wells after application of the samples. The concentrations tested were 100% MeCN, 90:10 MeCN/Milli-Q, 80:20 MeCN/Milli-Q, and 50:50 MeCN/Milli-Q. The equilibration was performed using 3 × 200 μL of the mixture mentioned above and washing with 4 × 200 μL of the same mixture. Finally, the elution was performed with 400 μL of Milli-Q water.

Analysis of Oligosaccharides. The labeled and purified oligosaccharides were analyzed using an Ultimate 3000 UHPLC system (Thermo Fisher Scientific) coupled to an FP-920 fluorescence detector (Jasco, Inc.). The detector was set for excitation and emission at 330 and 420 nm, respectively.

The chromatographic separation was conducted on an Acquity UPLC Glycan BEH Amide column (2.1 mm × 100 mm, 130 Å, 1.7 μm) and an Acquity UPLC Glycan BEH Amide VanGuard pre-column (2.1 mm × 5 mm, 130 Å, 1.7 μm) both from Waters. The column was maintained at 40 °C. The 10 times diluted sample (2 μL) was injected in the system, under gradient elution with a quaternary solvent system. The solvent system used consisted of acetonitrile (solvent A), 250 mM formic acid in 10% acetonitrile in Milli-Q water (pH 3.0) adjusted with ammonia (solvent B), and 10% acetonitrile in Milli-Q water (solvent C). The method used for the analysis was based on a gradient where solvent B was constant at 5%. The elution was performed with a gradient of 40 min moving from 5 to 27% solvent C. The gradient was followed by a cleaning step with 20% solvent B and 20% solvent C for 5 min and a subsequent equilibration step to the initial conditions of the analysis for 12 min, resulting in a total analysis time of 57 min. The flow applied for optimal chromatographic separation was 0.5 mL/min.

Quantification was performed using maltopentaose as an internal standard (IS) and general calibrant, based on the study of Austin et al.²⁵ This compound was selected, instead of the laminaritrinose in Austin's study, as it was eluting in an area of the chromatogram free of oligosaccharide peaks for both mature and colostrum milk samples.

Commercial standards (2'-FL, 3'-GL, 6'-GL, 3'-SL, 6'-SL, and 6'-SLN) were used for identification of gMOS. For the identification of peaks not co-eluting with the analytical standards, the same chromatographic system and fluorescence detector were coupled inline with a time-of-flight mass spectrometer (MaXis Plus, Bruker). The MS was operated in positive-ion mode, scanning between *m/z* 300 and 2000, with collision energy 7 eV, collision RF 80 Vpp, and transfer time 100 μs.

Quality control (QC) samples were utilized to monitor the stability of the analytical system and the consistency of the sample preparation.

For this purpose, reference mature goat milk samples were treated in the same way as the individual goat's milk samples, and were inserted in every analytical run every 10–15 samples.^{29,30}

Data Analysis. For the statistical analysis of the two farms, pairwise comparisons were assessed via the Mann–Whitney and Welch's test, depending on the outcome of the normality test. To check the normality, both D'Agostino and Pearson test and Shapiro–Wilk test were taken into consideration.

The descriptive parameters tested for potential comparisons with the gMOS levels were the age of the goats, milk production per day, lifetime milk production, % fat, % protein, number of pregnancies (parity), pseudocyesis, and the outcome of pregnancy in terms of sex of the goat kid. Pearson and Spearman correlation tests were employed for the factors described by quantitative data, while Mann–Whitney and Welch's tests were employed for the outcome of pregnancy factor. Pearson or Spearman test and Mann–Whitney or Welch's tests were selected based on both D'Agostino–Pearson and Shapiro–Wilk normality tests.

RESULTS AND DISCUSSION

Experimental Optimization. Initial experiments, using direct labeling of the goat milk²⁵ resulted in poor oligosaccharide




Figure 2. Overview of the relative intensities of the gMOS identified and quantified in the present study in D2 (colostrum) and D31 time points (mature). The oligosaccharides are presented per characteristic epitope and sorted by their relative abundance in a clockwise direction. Structures presented with an offset from the pie chart and highlighted with bold font in the legend indicate the common structures among hMOS and gMOS.

profiles both in reference mature milk and in the pooled colostrum samples. These were dominated by a lactose peak and almost no detectable oligosaccharides. Goat milk and cow milk differ in general composition from human milk. For example, the level of total protein in goat milk is almost 3.5 times higher than in human milk.³¹ Also, the lactose:MOS ratio in goat milk varies between 150 and 180, while in human milk, it is 8.5–13.6.¹⁹ The difference in lactose ratio may result in a very poor chromatogram. The higher protein levels may interfere with the labeling reaction. Moreover, the system setup used in this study does not have a switching valve between the guard and main columns to wash out excess 2-AB and lactose.²⁵ A second experiment with centrifugal defatting prior to labeling was performed, giving similar results, indicating that the fat levels were not the main cause of the poor chromatograms. Attempts to centrifuge the goat milk reference and goat colostrum milk samples, with or without dilution, led to troublesome supernatant collection, due to the formation of a fatty layer on top of the supernatant-liquid phase. Collection of supernatants was even more laborious during the processing of colostrum samples, slowing down the procedure and hindering the high-throughput character of the application.

Two larger-scale experiments were performed, involving precipitation of casein by acidification and salting out of proteins with ammonium sulfate. These approaches resulted in high-quality chromatograms clearly showing oligosaccharide peaks. However, these labor-intensive approaches would be prohibitive toward high-throughput sample handling, which was one of the requirements of this study.

Acetone extraction of oligosaccharides led to the formation of a pellet, and the extraction of gMOS structures in the supernatant remained poor. A possible explanation is that the pellet formed also caused precipitation of the carbohydrate structures, not allowing their exchange with the solvent. Methanol extraction resulted in a precipitate of the fat/protein content of milk, and the supernatant extract yielded a chromatogram rich in oligosaccharides. Direct application of 1 mL of milk to the PGC filters was not applicable, as the filters became clogged.

In view of the above results, extraction of oligosaccharides with methanol was chosen as the first step to remove most of the fat/protein content of the milk. After thawing overnight at 4 °C, 120 μ L of milk was mixed with 240 μ L of methanol. The samples were vortexed thoroughly and centrifuged for 15 min at 21 130g at 4 °C. The clear supernatant was collected, and the samples were dried using a Speedvac evaporator.

For the subsequent removal of the excessive amount of lactose interfering with the oligosaccharide analysis, filtration through PGC-SPE filters was applied. After initial conditioning, equilibration, and application of the samples on the SPE filters (see the [Experimental section](#)), different washing steps were tested. Optimal removal of lactose was achieved by washing the column with 4% MeCN/0.1% TFA; higher amounts of MeCN did not result in improved lactose removal and/or oligosaccharide recovery. Elution and drying of the samples resulted in extraction of purified gMOS.

After labeling of the oligosaccharides, excess of free label was removed by filtration in 96-well plate format with microcrystalline cellulose. Four different ratios of the MeCN/Milli-Q mixture for the equilibration and washing of the samples were examined. The results showed that 100% MeCN was not sufficient to remove excess label, yielding a broad peak in the first minutes of the chromatogram and overloading the detector. The ratio of 80:20 MeCN/Milli-Q and 50:50 MeCN/Milli-Q resulted in a loss of oligosaccharides during the washing steps and, hence, in non-reproducible oligosaccharide profiles. Optimal conditions were achieved when equilibrating and washing with 90:10 MeCN/Milli-Q that resulted in maximal removal of the labeling mixture without detectable loss of oligosaccharides. Repeat analysis of the reference sample with the final method showed high reproducibility in results.

Samples of different time points of different goats were analyzed in different batches of 60 to 65 samples. To monitor the performance of the analytical system and the repeatability of the extraction/labeling applied, two different controls were included in the analysis. Through the inclusion of quality control (QC) samples during the sample preparation of each batch, and their repetitive injection during the analytical run, both the batch-to-batch extraction efficiency and the analytical system stability were verified. The coefficient of variation (CV) for all of the structures detected in the QC samples ([Table 1](#)) did not exceed 15% within each batch and between different batches of analysis. The second control was the inclusion of an internal standard (IS) during sample preparation. The CV of maltopentaose (internal standard) did not exceed 10% within and between




Figure 3. Representative HPLC-FLD chromatograms of milk sample analysis of two goats (#14 and #18) at two time points (colostrum of the 2nd day and mature milk of the 31st day of lactation) and of the reference tank milk provided by Ausnutria BV. The identified structures are presented using the corresponding symbol nomenclature (see Table 1).




Figure 4. Indicative trend for the 3'-NGL (left) and 3'-SL (right) levels in the period between the 2nd month (M2) and the 9th month (M9) of lactation. Each box plot extends from 25th to 75th percentiles, indicating the median value.

Table 2. Descriptive Characteristics of the Goats of the Study

variable	farm #1 (<i>n</i> = 27)	farm #2 (<i>n</i> = 30)
age (years, mean ± SD)	4 ± 3	3 ± 3
production (milk, kg/day, mean ± SD)	3.5 ± 0.6	3.1 ± 0.5
lifetime production (milk, kg, mean ± SD)	4180 ± 2911	3028 ± 2708
% fat (mean ± SD)	4.0 ± 0.7	4.2 ± 0.4
% protein (mean ± SD)	3.4 ± 0.4	3.3 ± 0.2
parity (mean ± SD)	3 ± 2	2 ± 1
pseudocyesis (<i>n</i> , %)		7 (23%)
kids' sex (female <i>n</i> , %)		11 (37%)
(male <i>n</i> , %)		11 (37%)
(female and male <i>n</i> , %)		8 (27%)
feeding regime	pelletized feed	feed based on seasonal availability

batches. The final method is suitable for large sample sets and provides reproducible results.

Oligosaccharide Composition and Variation. Milk samples from the 57 goats participating in the study, early and later during the lactation period (colostrum and mature milk),

were analyzed. From preliminary analysis performed with 22 goats for all time points until the second month of lactation, it was highlighted that the oligosaccharide profiles followed a similar pattern for all 22 goats; in the first days of lactation (colostrum), large variations occurred in the levels of oligosaccharides, followed by a gradual decrease over time. After the first few days, most structures were decreased to a minimum level (Figure S1). For the rest of the time points up until month 9, three goats from each farm were analyzed, showing that the profiles remained the same until the last sample point (9 months of lactation). Samples of all 57 goats were analyzed for selected time points, reflecting the colostrum sample profile (day 1—1st colostrum sample; day 2—4th colostrum sample) and the mature milk (day 31). Table 1 summarizes the 12 gMOS structures identified and quantified in the samples of all 57 goats and their mean concentrations in the selected colostrum and mature time points.

Peaks annotated as 2'-FL, 3'-GL, 6'-GL, 3'-SL, 6'-SL, and 6'-SLN were identified based on comparison of the retention times with commercial standards and verified by in-spiking with standards and UPLC-TOF-FLD mass spectrometric analysis.³² Peaks eluting at 24.4 and 27.4 min were identified as two isomers of *N*-glycolyl-neuraminyl-lactose (3'- and 6'-NGL, respec-




Figure 5. Comparisons of goat milk MOS structures per farm and time point (D2 colostrum and D31 mature milk). The error bars indicate the minimum and maximum values and the stars the outcome of the pairwise comparison with the Mann–Whitney and Welch’s test, based on the normality test performed. The *p*-values are presented with stars: * <0.05 , ** <0.01 , *** <0.001 , and **** <0.0001 .

tively), based on the mass *m/z* 770.3 determined by the UPLC-TOF-FLD analysis. The peak at 25.6 min was characterized as *N*-glycolyl-neuraminyl-lactosamine (NGLN, *m/z* 811.3). Disialyllactose (DSL, *m/z* 1045.4) and sialyl-galactosyllactose

(SHL, *m/z* 916.3) were assigned to the peaks at 28 and 28.3 min respectively, while *N*-acetyl-glycosaminyl-dihexosyl-lactose (NADHL, *m/z* 990.4) was assigned to the peak at 33.1 min (Table 1). Based on previous observations of this structure and

Table 3. Spearman and Pearson Correlation Test Outcome of the MOS Studied for the Factors Age, Parity, and Lifetime Milk Production^a

Abbreviation	Name	r parameter	Age		Parity		Lifetime milk production	
			D2	D31	D2	D31	D2	D31
2'-FL	2'-Fucosyl-lactose	Spearman r	0.1034	0.1665	0.0874	0.0500	0.1887	0.2237
		95% CI	-0.2284 to 0.4137	-0.1812 to 0.4771	-0.2437 to 0.4003	-0.2924 to 0.3811	-0.1444 to 0.4833	-0.1231 to 0.5218
		R ²						
3'-GL	3'-Galactosyl-lactose	p value	0.5310	0.3319	0.5967	0.7720	0.2500	0.1897
		Spearman r	-0.0856	0.4170	-0.1200	0.1874	-0.1579	0.4429
		95% CI	-0.3714 to 0.2150	0.1282 to 0.6406	-0.3981 to 0.1783	-0.1211 to 0.4629	-0.4329 to 0.1440	0.1593 to 0.6589
6'-GL	6'-Galactosyl-lactose	R ²						
		p value	0.5674	0.0049	0.4167	0.2176	0.2891	0.0026
		Spearman r	0.0510	0.5583	-0.6686	0.3920	0.0575	0.5502
3'-SL	3'-Sialyl-lactose	95% CI	-0.2479 to 0.3410	0.3051 to 0.7377	-0.3520 to 0.2297	0.1025 to 0.6203	-0.2418 to 0.3468	0.2945 to 0.7323
		R ²						
		p value	0.7337	<0.0001	0.6516	0.0077	0.7012	0.0001
SLN	Sialyl-N-acetyl-lactosamine	r	S -0.2323	P -0.2291	S -0.2422	P -0.1197	S -0.1920	P -0.3008
		95% CI	-0.4936 to 0.0675	-0.4925 to 0.0727	-0.4989 to 0.0537	-0.3992 to 0.1801	-0.4610 to 0.1094	-0.5487 to -0.0043
		R ²						
3'-NGL	3'-N-glycolyl-neuraminyl-lactose	p value	0.1162	0.1346	0.0972	0.4333	0.1961	0.0472
		Spearman r	-0.2604	-0.3433	-0.2927	-0.3052	-0.1839	-0.3366
		95% CI	-0.5159 to 0.0377	-0.5870 to -0.0427	-0.5387 to -0.0001	-0.5575 to -0.0039	-0.4544 to 0.1177	-0.5819 to -0.0351
6'-SL	6'-Sialyl-lactose	R ²						
		p value	0.0771	0.0225	0.0435	0.0415	0.2160	0.0255
		Spearman r	0.6971	0.4482	0.6705	0.5924	0.6055	0.4064
NGLN	N-glycolyl-neuraminyl-lactosamine	95% CI	0.5060 to 0.8229	0.1657 to 0.6627	0.4706 to 0.8049	0.3540 to 0.7585	0.3779 to 0.7641	0.1157 to 0.6330
		R ²						
		p value	<0.0001	0.0023	<0.0001	<0.0001	<0.0001	0.0062
6'-NGL	6'-N-glycolyl-neuraminyl-lactose	r	P -0.2713	S -0.1944	P -0.2463	S -0.2307	P -0.3045	S -0.2119
		95% CI	-0.5181 to 0.0172	-0.4715 to 0.1177	-0.4958 to 0.0407	-0.4977 to 0.0763	-0.5441 to -0.0190	-0.4856 to 0.0996
		R ²						
DSL	Disialyl-lactose	p value	0.0651	0.2062	0.0915	0.1274	0.0374	0.1673
		Spearman r	0.5370	0.2644	0.4445	0.3080	0.5223	0.2474
		95% CI	0.2873 to 0.7183	-0.0443 to 0.5270	0.1752 to 0.6519	0.0069 to 0.5578	0.2686 to 0.7083	-0.0625 to 0.5137
SHL	Sialyl-galactosyl-lactose	R ²						
		p value	<0.0001	0.0829	0.0015	0.0396	0.0002	0.1055
		Spearman r	0.5146	0.4094	0.5485	0.5018	0.4591	0.3699
NADHL	N-acetyl-glucosaminyl-dihexosyl-lactose	95% CI	0.2587 to 0.7030	0.1192 to 0.6352	0.3054 to 0.7245	0.2358 to 0.6978	0.1897 to 0.6643	0.0730 to 0.6065
		R ²						
		p value	0.0002	0.0058	<0.0001	0.0004	0.0012	0.0135
SHL	Sialyl-galactosyl-lactose	Spearman r	-0.2483	-0.2274	-0.2450	-0.1778	-0.2376	-0.2295
		95% CI	-0.5145 to 0.0614	-0.5070 to 0.0957	-0.5091 to 0.6117	-0.4643 to 0.1425	-0.5061 to 0.0725	-0.5087 to 0.0935
		R ²						
SHL	Sialyl-galactosyl-lactose	p value	0.1041	0.1529	0.1047	0.2601	0.1204	0.1490
		Spearman r	-0.3197	-0.2669	-0.3663	-0.2950	-0.2644	-0.2272
		95% CI	-0.5618 to -0.0270	-0.5317 to 0.0455	-0.5947 to -0.0831	-0.5505 to 0.0111	-0.5191 to 0.0334	-0.5007 to 0.0876
NADHL	N-acetyl-glucosaminyl-dihexosyl-lactose	R ²						
		p value	0.0285	0.0836	0.0105	0.0519	0.0725	0.1429
		Spearman r	-0.0458	-0.7071	0.0135	-0.7071	-0.0346	-0.4000
NADHL	N-acetyl-glucosaminyl-dihexosyl-lactose	95% CI	-0.3364 to 0.2528		-0.2797 to 0.3043		-0.3264 to 0.2633	
		R ²						
		p value	0.7597	0.4000	0.9276	0.4000	0.8175	0.5167

^aThe correlations were assessed in both colostrum (D2) and mature milk (D31). The MOS/factor highlighted with green are those for which significant correlations were found based on the *p* value (<0.05). Pearson or Spearman test was selected based on both D'Agostino–Pearson and Shapiro–Wilk normality tests. S and P indications provide the data deriving from Spearman and Pearson test, respectively.

the relative abundances reported, identification as NADHL appears more likely than other structures fitting the observed mass, e.g., α -galactosyl-lacto-N-neotetraose.^{20,21} Moreover, in some of the samples analyzed, a peak was identified at 24.7 min as lacto-N-neotetraose (LNnT) based on the detected mass (m/z 828.3) and by comparing with the retention time of the commercial standard. However, due to the very low abundance of the peak and its elution immediately after the high abundance peak of 6'-SL, it was not possible to quantify LNnT. It should be underlined that the 3-FL commercial standard was washed out during the PGC-SPE filtration step. This effect of the PGC filtration on 3-FL has been reported previously.^{33,34} Hence, information about the 3-FL structure cannot be provided for the analyzed samples. Figure 2 provides an overview of the structures identified in the current study and their relative abundances in colostrum (D2) and mature milk (D31).

It is worth noting that the peaks indicated in Table 1 are the main peaks found both in colostrum and mature milk and correspond to the broadest panel of quantified structures in goat milk. However, in the colostrum of the first 3 days, eight

additional peaks were detected, six of which represented <2% of the total oligosaccharide content. Two of those eight unidentified peaks (~18.5 min) were observed in higher levels in the colostrum samples, but not in the mature time points. The TOF-MS analysis indicated galactosyllactose-like structures (m/z 625.2). The retention times of those peaks were compared with the 4'-galactosyllactose analytical standard, although this structure has not been reported previously in goat milk. Based on the retention time, the peaks of interest do not correspond to 4'-GL. A potential candidate for such peaks observed only in the colostrum samples is the 3'- α -galactosyllactose, reported previously in goat milk^{24,35} and, in a similar analytical system, eluting slightly earlier than 3'-GL,¹⁷ which is the case in our analysis also.

Previous studies have mainly reported relative abundances. In such a study, Albrecht et al. provided the broadest set of identified gMOS up to date and their relative abundances.¹⁷ All structures reported in that study with a relative abundance higher than 1% were quantified in the present study, as well as some of the lower detected structures (i.e., 2'-FL). It is




Figure 6. Boxplots of the gMOS highlighted as significantly different based on the outcome of the pregnancy in colostrum (D2) and mature milk (D31). The stars indicate the significance based on the *p* value of the Welch's or the Mann–Whitney test performed, depending on the outcome of the normality test (D'Agostino–Pearson and Shapiro–Wilk tests) (**p* value < 0.05, ***p* value < 0.01).

important to highlight that the authors used only pooled mature goat milk from a non-explicitly mentioned number of goats ($n \geq 3$, undefined breeds), and this could explain the differences in the levels of certain compounds when comparing with our findings. Concerning the extra structures identified in Albrecht's study, the identification of structures took place after pre-fractionation into neutral and acidic oligosaccharide pools.¹⁷ The method suggested in our study provides a more high-throughput analysis of the most abundant gMOS structures.

Significant differences in the profiles of oligosaccharides were observed between individual goats, in agreement with variation between Murciano-Granadina goats reported previously.²⁰ Four representative chromatograms of the study, belonging to milk samples of two goats of the same farm, following the same feeding regime, in two different time points colostrum of D2/4th collected sample and mature milk of (D31), are presented in Figure 3. Characteristic variations deriving from the time point of lactation (differences between D2 vs D31 for each goat) and biological factors (differences between the two goats) were observed. Figure 3 also shows the chromatogram of the tank milk (Ausnutria BV), which was used as a quality control sample during all performed analyses.

From colostrum (D1) to mature milk of the 9th month, almost all observed structures decreased in concentration in the milk samples for all of the studied goats. The only exception to this trend was the 3'-NGL structure, which remained high during later time points of lactation (D31, Figure 3, bottom chromatogram). Analysis of additional data points of later samples (up to the 9th month of lactation (M9)) showed that the levels of 3'-NGL even increased at later time points. The decrease in concentration of the identified gMOS is observed in

the data presented in Table 1, while an indicative trend of 3'-NGL over the course of lactation is given Figure 4. In Figure 4, the trend of 3'-SL is also given for comparison purposes. Furthermore, it should be noted that the concentrations change very strongly the first days of lactation, but the difference between day 17 and day 31 post-partum for all goats was already small (Figure S2). After day 31, the month 2 sample showed no significant change with day 31 for 22 goats tested. A full analysis of all samples collected for six goats also showed no significant change across the 9 months; although observation of the means of all oligosaccharides suggested a downward trend, statistical significance was not observed, with the exception of 3'-NGL, which showed a continuous upward trend with a significance in change between day 31 and month 9, but not between day 2 and day 31.

A noteworthy decrease in the concentration of most structures was observed, also between the colostrum samples (D1 and D2, Table 1). An exception to this trend is 2'-FL, which for some of the goats kept on increasing during D1 after parturition, reaching a peak during D2 and markedly decreasing during the next days of lactation. A very significant decrease in levels of sialylated structures has been reported previously for bovine colostrum.³⁶ The decrease rates for the structures reported by Nakamura et al. are similar to the decrease rates we observed in our study of the gMOS for the first 24 h (3'-SL 63.5% decrease, 6'-SL 69.1% decrease, and 6'-SLN 74.5% decrease).

It is worth mentioning that the variability perceived in the first post-partum sample was very broad, adding to the biological variation and complexity observed also in later time points of lactation. The extremely high levels of oligosaccharides in the

first post-partum sample could be related to providing the goat lambs with key components contributing to their healthy development.

Previous studies on goat and human MOS [20–22] also reported an overall decrease in the levels of MOS throughout lactation. The increase of 3'-NGL, the only structure showing an average increase in later time points (Figure 4), may be associated with biological functions in the later stages of lactation. This observation contrasts with previous studies reporting a decline in the levels of both NGL isomers over time.²² The different trend that 3'-NGL follows over the lactation period was observed in our study already after 31 days. The difference with the above-mentioned study could be the fact that the time points studied by Martin-Ortiz et al. were up to 40 days, while in our case, the samples expanded up to 9 months after the delivery. Further investigation of the biosynthesis and the potential biological functionality of 3'-NGL is required. To conclude, the presented data indicate that most gMOS decrease in concentration during the first weeks of lactation, after which the concentrations become stable, with the exception of 3'-NGL (Figure S2).

The most characteristic example of the differences observed in the levels of gMOS between the goats is that of 2'-FL. More specifically, 2'-FL was detected only in 73.7% of the goats studied. Looking at each farm, 21 out of the 27 goats (77.8%) from farm #1 and 21 out of the 30 goats (70%) from farm #2 produce 2'-FL. When 2'-FL was absent in the colostrum samples, it also did not appear in samples of later time points. In Table 1, the mean, min, and max values presented for 2'-FL correspond only to those goat samples in which this structure could be detected (73.7%). These data present a similar phenotype division as observed for human secretor vs non-secretor individuals,^{8,33} which is the result of disabling mutations in the FUT2 gene. So far, there have been no studies showing such a phenotype division in non-human mammals.

Among the "secretor" goats, there is a 2'-FL concentration range of 0.6–236.3 mg/L in the D2 (4th collected) samples. Within this group of secretors, two subgroups were observed; at D2, 30 goats were found with 0.6–69.9 mg/L and 12 goats with 97.8–236.3 mg/L (5 goats between 97.8 and 157.1 mg/L, and 7 goats between 184.2 and 236.3 mg/L). At the D31 time point, the 2'-FL concentrations were less diverse, ranging from 0.4 to 14.9 mg/L, with only one goat having a level of 69.5 mg/L. A similar division in high and low 2'-FL producers has been observed in human milk.^{37,38} The absence of 2'-FL in milk samples of some goats may be caused by very low levels of this oligosaccharide in samples from all time points. However, the oligosaccharide enrichment in the PGC-SPE filtration step was a strategy used to improve the detection of oligosaccharides even when present in very low levels. Austin et al., in a similar analysis type, reported a LOD level for 2'-FL of 3.9 mg/L.²⁵ Future studies focusing on genetic analysis could enlighten the genetic variance related with the very low levels/absence of 2'-FL in milk from some goats. It can be concluded that goat milk contains interesting levels of milk oligosaccharides, particularly in the first few days of lactation. It should be noted, however, that the concentrations are significantly lower than in human milk, which remains the best standard for infant nutrition.

Comparison of Oligosaccharide Levels between Farms. The differences in the levels of gMOS between the two farms were examined. Relevant data known for the goats and their milk characteristics included in this study are summarized in Table 2. The comparisons of the identified gMOS per farm at

the D2 and D31 time points are presented in Figure 5. Going from colostrum (D2) to mature milk (D31), a significant reduction in the levels of gMOS is clear for all of the structures in milk samples from both farms, except 3'-NGL (Figures 3 and 5). The pairwise comparisons of the two farms in colostrum (D2) and mature milk (D31) highlighted that the 6'-SLN, DSL, and SHL structure concentrations were significantly different (corresponding *p*-values were *p* < 0.005 at both time points) between farms. It is noteworthy that DSL and SHL are two of the least abundant structures detected and identified in the present study. The concentrations of two other structures, 6'-SL and NGLN, were significantly different between the two farms only at D31 with *p*-values <0.05 and <0.01, respectively.

The two farms were following different feeding regimes, which may play a role in the observed differences. Other factors, however, also may lead to the different levels of these structures, such as environmental or genetic ones. In human milk studies, the effect of the diet has been investigated with some studies reporting an interrelationship between hMOS levels and the diet of the mothers.³⁹ It should be emphasized that the sample collection from both goat farms was performed during the same period of the year with only a few days difference from each other. Therefore, any differences derived from seasonal variations can most likely be excluded. The differences observed between the farms suggest that a feeding intervention study in goats may yield interesting insights into influences that diet can have on gMOS composition in goat milk. However, further differences that may exist between different farms also should be taken into consideration, such as age of goats, parity, and milk yield. To draw more solid conclusions, controlled intervention studies with standardized conditions should be considered. The current study provides several leads for interesting factors to study in such a controlled experiment.

Correlations between gMOS and Available Goat Metadata. With the available metadata (Table 2), and the quantitative data collected from the gMOS milk fingerprinting, different correlations were assessed. Three parameters were found correlating with gMOS concentrations, namely, goat age, parity, and lifetime milk production. Table 3 provides information on the gMOS that were correlated with the above-mentioned factors. All three factors mainly correlated with the concentrations of two structures (3'-NGL and 6'-NGL), both in the colostrum samples (D2 time point of lactation) and in the mature milk (D31). These three factors are correlated with the structure 6'-GL only at D31 and with NGLN only at D2. The mechanisms involved remain to be studied. The average daily milk production, the average % fat and % protein content, and the appearance of pseudocytosis did not show any correlation with the gMOS fingerprints determined in the present study.

It should be emphasized that goat age, parity, and lifetime milk production are highly related to each other (*p* value < 0.0001 for age vs parity and age vs lifetime production comparisons), and it is not possible to conclude which has the most significant effect. Goat age is very strongly correlated with lifetime milk production and with the number of previous pregnancies of the goats (Figure S3). In the case of protein-bound glycosylation in human serum, it has been shown that age has an influence on the structural diversity of glycans.⁴⁰ Therefore, it seems logical that age may be the dominant factor in the correlation with gMOS levels, but, to our knowledge, this is the first time that this is reported for goats. Age, also, has been associated with the levels of other hMOS structures (fucosylacto-*N*-hexaose

(FLNH) and of difucosyllacto-*N*-hexaose (DFLNH)).²⁵ McGuire et al. studied also parity in humans,⁵ but no significant differences were reported in hMOS levels. Recently, another group reported that parity was associated independently with some hMOS levels.⁴¹ The same was concluded by Claps et al. who underlined an association of goat parity with the structures 3'-SL, 6'-SL, and DSL.²³ In our study, none of these structures correlated with parity.

For the lamb sex, pairwise comparison was performed among three groups (mothers of one female, mothers of one male, and mothers of both male and female lambs). Among the mothers that had lambs of both sexes, two goats had triplets of two daughters and one son and six goats had twins of one daughter and one son. The varied outcomes, female-only, male-only, or mixed-gender lambs, are distributed equally across the age ranges of the mother goats. Correlations were corrected for the age/parity effects mentioned earlier. For these comparisons, significant differences were highlighted for 3'-SL, NADHL, and DSL at the D2 time point and for 3'-SL, NGLN, and SHL at the D31 time point (Figure 6). Goat mothers of female lambs had significantly higher 3'-SL levels than mothers of both gender lambs at D2 (mean values, 96.2 and 32.6 mg/L, respectively; Mann–Whitney test, $p = 0.016$) and at D31 (66.3 and 25.1 mg/L, respectively; Welch's test, $p = 0.002$). The same structure was found in significantly higher levels in the comparison of mothers with male and both gender lambs at D31 (52.9 and 25.1 mg/L, respectively; Welch's test, $p = 0.025$).

At D2, mothers of female lambs had higher NADHL levels compared to those of male lambs (10.2 and 8.3 mg/L, respectively; Welch's test, $p = 0.023$) and compared to both gender lambs (10.2 and 4.5 mg/L, respectively; Welch's test, $p = 0.003$). The milk of mothers of male lambs had significantly lower levels of DSL compared to both gender lambs at D2 (3.7 and 7.0 mg/L, respectively; Welch's test, $p = 0.017$). At D31, the milk of mothers giving birth to females contained lower levels of NGLN compared to mothers with male lambs (2.2 and 3.9 mg/L, respectively; Welch's test, $p = 0.012$) and to both genders' pregnancy outcome (2.2 and 4.6 mg/L, respectively; Welch's test, $p = 0.024$). Comparing the pregnancy outcomes of female and both genders, SHL was also found significant at D31 (3.2 and 1.9 mg/L, respectively; Welch's test, $p = 0.013$).

A recent hMOS study also correlated the sex of the neonates with the concentration of specific structures, with a focus on 2'-FL, LNT, LNnT, and other neutral core hMOS.⁴² Our study showed that this correlation also may be found in the milk of goats, but in this case, mainly acidic gMOS are involved. It should be noted that, in contrast to human milk, in milk of goats and other domesticated animals, the acidic structures are predominant (Table 1).

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jafc.1c00499>.

HPLC-FLD chromatograms of gMOS (Figure S1); PCA analysis of gMOS data (Figure S2); and correlation graphs depicting the factors of age with parity and lifetime milk production (Figure S3) (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Sander S. van Leeuwen – Sector Human Nutrition and Health, Department of Laboratory Medicine, University Medical Center Groningen (UMCG), 9713 GZ Groningen, The Netherlands; orcid.org/0000-0003-0017-2841; Email: s.s.van.leeuwen@umcg.nl

Authors

Anastasia Chrysovalantou Chatziioannou – CarbExplore Research BV, 9747 AN Groningen, The Netherlands; orcid.org/0000-0002-1973-7542

Eric Benjamins – Ausnutria BV, 8025 BM Zwolle, The Netherlands

Linette Pellis – Ausnutria BV, 8025 BM Zwolle, The Netherlands

Alfred Haandrikman – Ausnutria BV, 8025 BM Zwolle, The Netherlands

Lubbert Dijkhuizen – CarbExplore Research BV, 9747 AN Groningen, The Netherlands

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.jafc.1c00499>

Funding

This project received financial support from Ausnutria BV.

Notes

The authors declare the following competing financial interest(s): E.B., L.P., and A.H. are employed by Ausnutria BV; A.C.C. and L.D. are employed by CarbExplore Research BV, which has received financial support from Ausnutria BV, at least partly used to also fund the work of S.S.v.L.

■ ACKNOWLEDGMENTS

Special thanks go to the two farms that participated in this study, providing all milk samples and the related information of the studied goat subjects, essential for a better evaluation of the results obtained. The authors also thank Dr Hjalmar Permentier and the personnel of the Interfaculty Mass Spectrometry Center, Groningen, the Netherlands, for assistance with the UPLC-FLD-TOF-MS system and the provided measurement time.

■ REFERENCES

- (1) Thurl, S.; Munzert, M.; Boehm, G.; Matthews, C.; Stahl, B. Systematic Review of the Concentrations of Oligosaccharides in Human Milk. *Nutr. Rev.* **2017**, *75*, 920–933.
- (2) Urashima, T.; Hirabayashi, J.; Sato, S.; Kobata, A. Human Milk Oligosaccharides as Essential Tools for Basic and Application Studies on Galectins. *Trends Glycosci. Glycotechnol.* **2018**, *30*, SE51–SE65.
- (3) Samuel, T. M.; Binia, A.; de Castro, C. A.; Thakkar, S. K.; Billeaud, C.; Agosti, M.; Al-Jashi, I.; Costeira, M. J.; Marchini, G.; Martínez-Costa, C.; Picaud, J.-C.; Stiris, T.; Stoicescu, S.-M.; Vanpeé, M.; Domellöf, M.; Austin, S.; Sprenger, N. Impact of Maternal Characteristics on Human Milk Oligosaccharide Composition over the First 4 Months of Lactation in a Cohort of Healthy European Mothers. *Sci. Rep.* **2019**, *9*, No. 11767.
- (4) Ma, L.; McJarow, P.; Jan Mohamed, H. J. B.; Liu, X.; Welman, A.; Fong, B. Y. Lactational Changes in the Human Milk Oligosaccharide Concentration in Chinese and Malaysian Mothers' Milk. *Int. Dairy J.* **2018**, *87*, 1–10.
- (5) McGuire, M. K.; Meehan, C. L.; McGuire, M. A.; Williams, J. E.; Foster, J.; Sellen, D. W.; Kamau-Mbutia, E. W.; Kamundia, E. W.; Mbugua, S.; Moore, S. E.; Prentice, A. M.; Kvist, L. J.; Otoo, G. E.; Brooker, S. L.; Price, W. J.; Shafii, B.; Placek, C.; Lackey, K. A.; Robertson, B.; Manzano, S.; Ruíz, L.; Rodríguez, J. M.; Pareja, R. G.;

Bode, L. What's Normal? Oligosaccharide Concentrations and Profiles in Milk Produced by Healthy Women Vary Geographically. *Am. J. Clin. Nutr.* **2017**, *105*, 1086–1100.

(6) Davis, J. C. C.; Lewis, Z. T.; Krishnan, S.; Bernstein, R. M.; Moore, S. E.; Prentice, A. M.; Mills, D. A.; Lebrilla, C. B.; Zivkovic, A. M. Growth and Morbidity of Gambian Infants Are Influenced by Maternal Milk Oligosaccharides and Infant Gut Microbiota. *Sci. Rep.* **2017**, *7*, No. 40466.

(7) Triantis, V.; Bode, L.; van Neerven, R. J. J. Immunological Effects of Human Milk Oligosaccharides. *Front. Pediatr.* **2018**, *6*, No. 190.

(8) Bode, L. Human Milk Oligosaccharides: Every Baby Needs a Sugar Mama. *Glycobiology* **2012**, *22*, 1147–1162.

(9) Manthey, C. F.; Autran, C. A.; Eckmann, L.; Bode, L. Human Milk Oligosaccharides Protect against Enteropathogenic Escherichia Coli Attachment in Vitro and EPEC Colonization in Suckling Mice. *J. Pediatr. Gastroenterol. Nutr.* **2014**, *58*, 165–168.

(10) Bode, L.; Rudloff, S.; Kunz, C.; Strobel, S.; Klein, N. Human Milk Oligosaccharides Reduce Platelet-Neutrophil Complex Formation Leading to a Decrease in Neutrophil β 2 Integrin Expression. *J. Leukocyte Biol.* **2004**, *76*, 820–826.

(11) Jantscher-Krenn, E.; Zherebtsov, M.; Nissan, C.; Goth, K.; Guner, Y. S.; Naidu, N.; Choudhury, B.; Grishin, A. V.; Ford, H. R.; Bode, L. The Human Milk Oligosaccharide Disialyllacto-N-Tetraose Prevents Necrotising Enterocolitis in Neonatal Rats. *Gut* **2012**, *61*, 1417–1425.

(12) Autran, C. A.; Kellman, B. P.; Kim, J. H.; Asztalos, E.; Blood, A. B.; Spence, E. C. H.; Patel, A. L.; Hou, J.; Lewis, N. E.; Bode, L. Human Milk Oligosaccharide Composition Predicts Risk of Necrotising Enterocolitis in Preterm Infants. *Gut* **2018**, *67*, 1064–1070.

(13) Vázquez, E.; Barranco, A.; Ramírez, M.; Gruart, A.; Delgado-García, J. M.; Martínez-Lara, E.; Blanco, S.; Martín, M. J.; Castany, E.; Buck, R.; Prieto, P.; Rueda, R. Effects of a Human Milk Oligosaccharide, 2'-Fucosyllactose, on Hippocampal Long-Term Potentiation and Learning Capabilities in Rodents. *J. Nutr. Biochem.* **2015**, *26*, 455–465.

(14) Jacobi, S. K.; Yatsunenko, T.; Li, D.; Dasgupta, S.; Yu, R. K.; Berg, B. M.; Chichlowski, M.; Odle, J. Dietary Isomers of Sialyllactose Increase Ganglioside Sialic Acid Concentrations in the Corpus Callosum and Cerebellum and Modulate the Colonic Microbiota of Formula-Fed Piglets. *J. Nutr.* **2016**, *146*, 200–208.

(15) Urashima, T.; Taufik, E.; Fukuda, K.; Asakuma, S. Recent Advances in Studies on Milk Oligosaccharides of Cows and Other Domestic Farm Animals. *Biosci., Biotechnol., Biochem.* **2013**, *77*, 455–466.

(16) Urashima, T.; Saito, T.; Nakamura, T.; Messer, M. Oligosaccharides of Milk and Colostrum in Non-Human Mammals. *Glycoconjugate J.* **2001**, *18*, 357–371.

(17) Albrecht, S.; Lane, J. A.; Mariño, K.; Al Busadah, K. A.; Carrington, S. D.; Hickey, R. M.; Rudd, P. M. A Comparative Study of Free Oligosaccharides in the Milk of Domestic Animals. *Br. J. Nutr.* **2014**, *111*, 1313–1328.

(18) Robinson, R. C. Structures and Metabolic Properties of Bovine Milk Oligosaccharides and Their Potential in the Development of Novel Therapeutics. *Front. Nutr.* **2019**, *6*, No. 1147.

(19) Martínez-Ferez, A.; Rudloff, S.; Guadix, A.; Henkel, C. A.; Pohlentz, G.; Boza, J. J.; Guadix, E. M.; Kunz, C. Goats' Milk as a Natural Source of Lactose-Derived Oligosaccharides: Isolation by Membrane Technology. *Int. Dairy J.* **2006**, *16*, 173–181.

(20) Martín-Ortiz, A.; Salcedo, J.; Barile, D.; Bunyatratchata, A.; Moreno, F. J.; Martín-García, I.; Clemente, A.; Sanz, M. L.; Ruiz-Matute, A. I. Characterization of Goat Colostrum Oligosaccharides by Nano-Liquid Chromatography on Chip Quadrupole Time-of-Flight Mass Spectrometry and Hydrophilic Interaction Liquid Chromatography-Quadrupole Mass Spectrometry. *J. Chromatogr. A* **2016**, *1428*, 143–153.

(21) Meyrand, M.; Dallas, D. C.; Caillat, H.; Bouvier, F.; Martin, P.; Barile, D. Comparison of Milk Oligosaccharides between Goats with and without the Genetic Ability to Synthesize As1-Casein. *Small Rumin. Res.* **2013**, *113*, 411–420.

(22) Martín-Ortiz, A.; Barile, D.; Salcedo, J.; Moreno, F. J.; Clemente, A.; Ruiz-Matute, A. I.; Sanz, M. L. Changes in Caprine Milk Oligosaccharides at Different Lactation Stages Analyzed by High Performance Liquid Chromatography Coupled to Mass Spectrometry. *J. Agric. Food Chem.* **2017**, *65*, 3523–3531.

(23) Claps, S.; Di Napoli, M. A.; Caputo, A. R.; Rufrano, D.; Sepe, L.; Di Trana, A. Factor Affecting the 3' Sialyllactose, 6' Sialyllactose and Disialyllactose Content in Caprine Colostrum and Milk: Breed and Parity. *Small Rumin. Res.* **2016**, *134*, 8–13.

(24) van Leeuwen, S. S.; te Poele, E. M.; Chatziioannou, A. C.; Benjamins, E.; Haandrikman, A.; Dijkhuizen, L. Goat Milk Oligosaccharides: Their Diversity, Quantity, and Functional Properties in Comparison to Human Milk Oligosaccharides. *J. Agric. Food Chem.* **2020**, *68*, No. 13469.

(25) Austin, S.; Béné, T. Quantitative Determination of Non-Lactose Milk Oligosaccharides. *Anal. Chim. Acta* **2018**, *1010*, 86–96.

(26) Robinson, R. C.; Colet, E.; Tian, T.; Poulsen, N. A.; Barile, D. An Improved Method for the Purification of Milk Oligosaccharides by Graphitised Carbon-Solid Phase Extraction. *Int. Dairy J.* **2018**, *80*, 62–68.

(27) Ruhaak, L. R.; Steenvoorden, E.; Koeleman, C. A. M.; Deelder, A. M.; Wuhler, M. 2-Picoline-Borane: A Non-Toxic Reducing Agent for Oligosaccharide Labeling by Reductive Amination. *Proteomics* **2010**, *10*, 2330–2336.

(28) Ruhaak, L. R.; Huhn, C.; Waterreus, W.-J.; de Boer, A. R.; Neustüss, C.; Hokke, C. H.; Deelder, A. M.; Wuhler, M. Hydrophilic Interaction Chromatography-Based High-Throughput Sample Preparation Method for N-Glycan Analysis from Total Human Plasma Glycoproteins. *Anal. Chem.* **2008**, *80*, 6119–6126.

(29) Gika, H. G.; Theodoridis, G. A.; Wingate, J. E.; Wilson, I. D. Within-Day Reproducibility of an HPLC–MS-Based Method for Metabonomic Analysis: Application to Human Urine. *J. Proteome Res.* **2007**, *6*, 3291–3303.

(30) Virgiliou, C.; Valianou, L.; Witting, M.; Moritz, F.; Fotakis, C.; Zoumpoulakis, P.; Chatziioannou, A. C.; Lazaros, L.; Makrydimas, G.; Chatzimeletiou, K.; Raikos, N.; Theodoridis, G. A. Metabolic Profile of Human Coelomic Fluid. *Bioanalysis* **2017**, *9*, 37–51.

(31) Verduci, E.; D'Elia, S.; Cerrato, L.; Comberiat, P.; Calvani, M.; Palazzo, S.; Martelli, A.; Landi, M.; Trikamjee, T.; Peroni, D. G. Cow's Milk Substitutes for Children: Nutritional Aspects of Milk from Different Mammalian Species, Special Formula and Plant-Based Beverages. *Nutrients* **2019**, *11*, No. 1739.

(32) Remoroza, C. A.; Liang, Y.; Mak, T. D.; Mirokhin, Y.; Sheetlin, S. L.; Yang, X.; San Andres, J. V.; Power, M. L.; Stein, S. E. Increasing the Coverage of a Mass Spectral Library of Milk Oligosaccharides Using a Hybrid-Search-Based Bootstrapping Method and Milks from a Wide Variety of Mammals. *Anal. Chem.* **2020**, *92*, 10316–10326.

(33) van Leeuwen, S. S. Challenges and Pitfalls in Human Milk Oligosaccharide Analysis. *Nutrients* **2019**, *11*, No. 2684.

(34) Xu, G.; Davis, J. C.; Goonatilake, E.; Smilowitz, J. T.; German, J. B.; Lebrilla, C. B. Absolute Quantitation of Human Milk Oligosaccharides Reveals Phenotypic Variations during Lactation. *J. Nutr.* **2017**, *147*, 117–124.

(35) Urashima, T.; Bubb, W. A.; Messer, M.; Tsuji, Y.; Taneda, Y. Studies of the Neutral Trisaccharides of Goat (Capra Hircus) Colostrum and of the One- and Two-Dimensional ¹H and ¹³C NMR Spectra of 6'-N-Acetylglucosaminylactose. *Carbohydr. Res.* **1994**, *262*, 173–184.

(36) Nakamura, T.; Kawase, H.; Kimura, K.; Watanabe, Y.; Ohtani, M.; Arai, I.; Urashima, T. Concentrations of Sialyloligosaccharides in Bovine Colostrum and Milk during the Parturition and Early Lactation. *J. Dairy Sci.* **2003**, *86*, 1315–1320.

(37) Nakhla, T.; Fu, D.; Zopf, D.; Brodsky, N. L.; Hurt, H. Neutral Oligosaccharide Content of Preterm Human Milk. *Br. J. Nutr.* **1999**, *82*, 361–367.

(38) Elwakiel, M.; Hageman, J. A.; Wang, W.; Szeto, I. M.; van Goudoever, J. B.; Hettinga, K. A.; Schols, H. A. Human Milk Oligosaccharides in Colostrum and Mature Milk of Chinese Mothers:

Lewis Positive Secretor Subgroups. *J. Agric Food Chem.* **2018**, *66*, 7036–7043.

(39) Qiao, Y.; Feng, J.; Yang, J.; Gu, G. The Relationship between Dietary Vitamin A Intake and the Levels of Sialic Acid in the Breast Milk of Lactating Women. *J. Nutr. Sci. Vitaminol.* **2013**, *59*, 347–351.

(40) Ruhaak, L. R.; Uh, H.-W.; Beekman, M.; Hokke, C. H.; Westendorp, R. G. J.; Houwing-Duistermaat, J.; Wuhler, M.; Deelder, A. M.; Slagboom, P. E. Plasma Protein N-Glycan Profiles Are Associated with Calendar Age, Familial Longevity and Health. *J. Proteome Res.* **2011**, *10*, 1667–1674.

(41) Azad, M. B.; Robertson, B.; Atakora, F.; Becker, A. B.; Subbarao, P.; Moraes, T. J.; Mandhane, P. J.; Turvey, S. E.; Lefebvre, D. L.; Sears, M. R.; Bode, L. Human Milk Oligosaccharide Concentrations Are Associated with Multiple Fixed and Modifiable Maternal Characteristics, Environmental Factors, and Feeding Practices. *J. Nutr.* **2018**, *148*, 1733–1742.

(42) M Tonon, K.; B de Morais, M.; F V Abrão, A. C.; Miranda, A.; B Morais, T. Maternal and Infant Factors Associated with Human Milk Oligosaccharides Concentrations According to Secretor and Lewis Phenotypes. *Nutrients* **2019**, *11*, No. 1358.