

## University of Groningen

### Low-frequency noise

Erdelyi, Kristina; Fuermaier, Anselm; Koerts, Janneke; Tucha, Lara; Tucha, Oliver

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Publication date:*  
2021

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Erdelyi, K., Fuermaier, A., Koerts, J., Tucha, L., & Tucha, O. (2021). *Low-frequency noise: Experiences from individuals reporting LFN complaints*. Poster session presented at 13th ICBEN congress on noise as a public health problem.

#### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Low-Frequency Noise: Experiences from individuals reporting LFN complaints

Kristina Erdélyi<sup>1</sup>, Anselm Fuermaier<sup>1</sup>, Janneke Koerts<sup>1</sup>, Lara Tucha<sup>2</sup>, Oliver Tucha<sup>1,2,3</sup>

<sup>1</sup> Department of Clinical and Developmental Neuropsychology, University of Groningen, Groningen, The Netherlands, <sup>2</sup> Department of Psychiatry and Psychotherapy, University Medical Center Rostock, Rostock, Germany, <sup>3</sup> Department of Psychology, Maynooth University, National University of Ireland, Maynooth, Ireland

## Low-Frequency Noise (LFN)

Low-Frequency Noise (LFN) refers to sound waves around/below the hearing threshold (below 125 Hz).<sup>1</sup>

Sources of LFN can be natural (e.g.: sea waves, wind) but are mostly man-made (e.g.: traffic, turbines, pumps).<sup>1</sup>

Sources and complaints of LFN are rising. An estimated 2% of the Dutch population experience severe annoyance from LFN.<sup>1,2,3</sup>

## Self-reported complaints of individuals reporting LFN

Physical: Headache, dizziness, palpitation, neck/ back/ aural pain<sup>2,4,5</sup>

Psychological: Sleeping difficulties, fatigue, annoyance, stress, feelings of depression, frustration<sup>2,4,5</sup>

Cognition: Difficulties in concentration, cognitively demanding tasks, executive functions<sup>2,4,5,6</sup>

Well-being & daily functioning: Lower quality of life, job loss, incapacity to work, relationship or family problems, housing issues<sup>2,4</sup>

## LFN research is urgently needed

LFN is still a hardly recognized and investigated environmental stressor.

The source of reported sounds is often not found (e.g.: by inaccurate measurement methods or complaints originating from a possibly non-LFN related cause).<sup>2</sup>

Previous research often used small and specific groups, unstructured observations or subjective reports, short-term laboratory LFN-exposures, or applied different definitions of LFN. Research objectifying subjective complaints does not allow for conclusions yet.

## Individual factors and LFN

Knowledge on individual factors associated with noise reactions in this group is scarce.<sup>5</sup> Current research indicates:

- 1) Sociodemographic factors: LFN-perceptions are most commonly reported by individuals between 50-70 years and in 2/3 of cases by females.<sup>2,4,7</sup>
- 2) Personality factors: Introverts seem more sensitive to and more annoyed by LFN and perform worse on cognitive tasks during LFN exposure despite being faster compared to extraverts.<sup>8,9,10</sup> However, firm conclusions regarding this or other aspects of personality cannot be made so far.

## Type of LFN perception

There seems to be substantial heterogeneity regarding the reported type of LFN perceptions. Some general tendencies seem to be:<sup>2,4,7</sup>

- 1) Location: mainly indoors, especially at home.
- 2) Time: mostly all the time or at night only, rarely at daytime only.
- 3) Type: by hearing (often a humming, sometimes a buzzing, throbbing, or engine-like sound). But also feelings of pressure and vibrations are reported.

The considerable individual differences in LFN perceptions necessitate further investigation and the examination of possible subgroups.

## The current research of the University of Groningen

Within a systematic, large-scale research project, a present study investigates

- 1) the demographic and personal characteristics of individuals reporting LFN complaints
- 2) the specific LFN perceptions (including location, time, type, circumstances)
- 3) reported physical, psychological, and social health-related symptoms and restrictions in daily living

in comparison to individuals who are not perceiving LFN.



## References

1. Rijksinstituut voor Volksgezondheid en Milieu. (2020). Laagfrequentgeluid [Fact sheet]. Retrieved from <https://www.rivm.nl/sites/default/files/2020-09/Factsheet%20laagfrequent%20geluid.pdf>
2. Erasmus, M., & van der Plas, D., (2020). Jaarrapportage 2019 Laagfrequentgeluid. Retrieved from <https://www.laagfrequentgeluid.nl/download/jaarrapportage-scroll-2019.pdf>
3. van Kamp, I., Breugelmans, O., R., P., van Poll, H., F., P., M., Baliatsas, C., van Kempen, E., E., M., M. (2018). Meldingen over en hinder van Laagfrequent Geluid of het horen van een bromtoon in Nederland: Inventarisatie. Retrieved from <https://www.rivm.nl/bibliotheek/rapporten/2018-0119.pdf>
4. Leventhall, G., Pelmear, P., & Benton, S. (2003). A review of Published research on Low Frequency Noise and its effects: UK Department for Environment, Food and Rural Affairs. <https://doi.org/EPG1/2/50>
5. Baliatsas, C., van Kamp, I., van Poll, R., & Yzermans, J. (2016). Health effects from low-frequency noise and infrasound in the general population: is it time to listen? a systematic review of observational studies. *Science of the Total Environment*, 557-558, 163-169.
6. Erdélyi K., Fuermaier, A. B. M., Koerts, J., Tucha, L., & Tucha, O. (2019). Laagfrequent geluid als omgevingsstressor in relatie tot neuropsychologische beperkingen. *Psychopraxis Neuropraxis*, 23(3), 48-52. <https://doi.org/10.1007/s12474-019-00221-7>
7. Vasudevan, R. N., and Gordon, C. G. (1977): Experimental study of annoyance due to low frequency environmental noise. *Applied Acoustics* 10(1), 57-69.
8. Abbasi, M., Tokhi, M. O., Falahati, M., Yazdanirad, S., Ghaljahi, M., Etemadinezhad, S., & Jaffari, T. P. R. (2020). Effect of personality traits on sensitivity, annoyance and loudness perception of low- and high-frequency noise. *Journal of Low Frequency Noise, Vibration and Active Control*, 146134842094581, 146134842094581-146134842094581. <https://doi.org/10.1177/1461348420945818>
9. Alimohammadi, I., Sandrock, S., & Gohari, M. R. (2013). The effects of low frequency noise on mental performance and annoyance. *Environmental Monitoring and Assessment*, 185(8), 7043-7051.
10. Babamiri, M., Moatamedzadeh, M., Golmohammadi, R., Derakhshan, J., & Farhadian, M. (2017). Investigating the role of personality type in the effects of low frequency noise on students' cognitive performances. *Journal of Occupational Hygiene Engineering* 4(2):59-66. <https://doi.org/10.21859/johe.4.2.59>