

University of Groningen

Geometric analysis of nonlinear differential-algebraic equations via nonlinear control theory

Chen, Yahao; Respondek, Witold

Published in:
Journal of Differential Equations

DOI:
[10.1016/j.jde.2022.01.008](https://doi.org/10.1016/j.jde.2022.01.008)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2022

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Chen, Y., & Respondek, W. (2022). Geometric analysis of nonlinear differential-algebraic equations via nonlinear control theory. *Journal of Differential Equations*, 314, 161-200.
<https://doi.org/10.1016/j.jde.2022.01.008>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Geometric analysis of nonlinear differential-algebraic equations via nonlinear control theory

Yahao Chen ^{a,*}, Witold Respondek ^b

^a *Bernoulli Institute, University of Groningen, the Netherlands*

^b *Normandie Université, INSA-Rouen, LMI, France*

Received 2 May 2021; revised 8 December 2021; accepted 6 January 2022

Available online 19 January 2022

Abstract

For nonlinear differential-algebraic equations (DAEs), we define two kinds of equivalences, namely, the external and internal equivalence. Roughly speaking, the word “external” means that we consider a DAE (locally) everywhere and “internal” means that we consider the DAE on its (locally) maximal invariant submanifold (i.e., where its solutions exist) only. First, we revise the geometric reduction method in DAEs solution theory and formulate an implementable algorithm to realize that method. Then a procedure called explicitation with driving variables is proposed to connect nonlinear DAEs with nonlinear control systems and we show that the driving variables of an explicitation system can be reduced under some involutivity conditions. Finally, due to the explicitation, we use some notions from nonlinear control theory to derive two nonlinear generalizations of the Weierstrass form.

© 2022 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

MSC: 34A09; 93B17; 93B27; 93B10; 93C10

Keywords: Geometric control methods; Nonlinear differential-algebraic equations; Explicitation; Internal equivalence; Invariant submanifolds; Nonlinear Weierstrass form

* Corresponding author.


E-mail addresses: yahao.chen@rug.nl (Y. Chen), witold.respondek@insa-rouen.fr (W. Respondek).

1. Introduction

Consider a nonlinear differential-algebraic equation (DAE) of the form

$$\Xi : E(x)\dot{x} = F(x), \tag{1}$$

where $x \in X$ is a vector of the generalized states and X is an open subset of \mathbb{R}^n (or an n -dimensional manifold).



The maps $E : TX \rightarrow \mathbb{R}^l$ and $F : X \rightarrow \mathbb{R}^l$ (see the above diagram, where $\pi : TX \rightarrow X$ is the canonical projection from the tangent bundle TX onto X) are smooth and the word “smooth” will mean throughout this paper C^∞ -smooth. We will denote a DAE of the form (1) by $\Xi_{l,n} = (E, F)$ or, simply, Ξ . Equation (1) is affine with respect to the velocity \dot{x} , so sometimes it is called a quasi-linear DAE (see e.g., [1,2]) and can be considered as an affine Pfaffian system since the rows E^i of E are actually differential 1-forms on X (for linear Pfaffian systems, see e.g. [3]), so E is an \mathbb{R}^l -valued differential 1-form on X . A semi-explicit DAE is of the form

$$\Xi^{SE} : \begin{cases} \dot{x}_1 = F_1(x_1, x_2), \\ 0 = F_2(x_1, x_2), \end{cases} \tag{2}$$

where $x_1 \in X_1$ is a vector of state variables and $x_2 \in X_2$ is a vector of algebraic or free variables (since there are no differential equations for x_2) with X_1 and X_2 being open subsets of \mathbb{R}^q and \mathbb{R}^{n-q} , respectively (or q - and $(n - q)$ -dimensional manifolds, respectively), the maps $F_1 : X_1 \times X_2 \rightarrow TX_1$ and $F_2 : X_1 \times X_2 \rightarrow \mathbb{R}^{l-q}$ are smooth. A linear DAE of the form

$$\Delta : E\dot{x} = Hx \tag{3}$$

will be denoted by $\Delta_{l,n} = (E, H)$ or, simply, Δ , where $E \in \mathbb{R}^{l \times n}$ and $H \in \mathbb{R}^{l \times n}$. Both the semi-explicit DAE Ξ^{SE} and the linear DAE Δ can be seen as special cases of DAE Ξ . The motivation of studying DAEs is their frequent presence in modeling of practical systems as electrical circuits [2,4], chemical processes [5,6], mechanical systems [7–9], etc.

There are three main results of this paper. The first result (section 2) concerns analyzing a DAE (locally) everywhere (i.e., externally) or considering the restriction of the DAE to a submanifold (i.e., internally), which corresponds to the external equivalence (see Definition 2.10) and the internal equivalence (see Definition 2.17), respectively. The difference between the two equivalences will be illustrated by their relations with the solutions. In order to analyze the existence of solutions, we use a concept called *locally maximal invariant submanifold* (see Definition 2.2), which is a submanifold where the solutions of a DAE exist and can be constructed via a geometric reduction method shown in section 2.1. Note that the geometric reduction method is not new in the theory of nonlinear DAEs, see e.g., [1,2,10–12] and the recent papers [13–15]. In the

present paper, we will show a practical implementation of this method via an algorithm summarized in section 2.2. In our recent publication [16], the geometric reduction method was applied to DAE control systems, i.e., (1) with additional input variables u . Some results in section 2 can be seen as a special case, i.e., when the inputs u are absent, of the corresponding results in [16], we will address both the connections and differences between section 2 and [16] in Remark 2.22 below.

Note that considering only the restriction of a DAE means that we only care about where and how the solutions of that DAE evolve. However, when a nominal point is not on the maximal invariant submanifold (which is common for practical systems, since an initial point could be anywhere), there are no solutions passing through that point but we still want to steer the solutions to the submanifold and thus we must follow the rules indicated by the “external” form of the DAE, thus considering DAEs everywhere is also important, see [17], where we use external equivalence to study jump solutions of nonlinear DAEs. Hence both the internal and the external analysis play crucial roles for DAEs and one needs to make a suitable choice among them depending on whether the purpose is to study C^1 -solutions of DAEs evolving on locally maximal invariant submanifolds only or to study discontinuous solutions of DAEs starting from inadmissible initial points.

The second result of this paper (section 3) is a nonlinear counterpart of the results of [18], in which we have shown that one can associate a class of linear control systems to any linear DAE (by the procedure of explicitation for linear DAEs). In the present paper, to any nonlinear DAE, by introducing extra variables (called driving variables), we can attach a class of nonlinear control systems. Moreover, we show that the driving variables in this explicitation procedure can be fully reduced under some involutivity conditions which explains when a DAE Ξ is ex-equivalent to a semi-explicit DAE Ξ^{SE} .

It is well-known (see e.g., [19], [20]) that any linear DAE Δ of the form (3) is ex-equivalent (via linear transformations) to the Kronecker canonical form **KCF**. In particular, if Δ is *regular*, i.e., the matrices E and H are square ($l = n$) and $|sE - H| \neq 0, \forall s \in \mathbb{C}$, then Δ is ex-equivalent (also via linear transformations) to the Weierstrass form **WF** [21] (see (19) below). The studies on normal forms and canonical forms of DAEs can be found in [19,21–24] for the linear case and in [15,25,26] for the nonlinear case. The last result of this paper (section 4) is to use such concepts as zero dynamics, relative degree and invariant distributions of the nonlinear control theory [27,28] to derive nonlinear generalizations of the **WF**. In the linear case, canonical forms as the **KCF** and the **WF** are closely related to a geometric concept named the Wong sequences [29] (see Remark 2.6 below). In [23], relations between the **WF** and the Wong sequences have been built and in [24], the importance of the Wong sequences for the geometric analysis of linear DAEs is reconfirmed. In the present paper, we propose generalizations of the Wong sequences for nonlinear DAEs and show their importance in analyzing structure properties.

The paper is organized as follows. In section 2.1, we discuss the existence of solutions of DAEs by revising the geometric reduction method. The latter method is implemented via a recursive algorithm in section 2.2. In section 2.3, we compare the notions of external equivalence and internal equivalence. The major contributions in section 2 are Proposition 2.8, which summarizes the results of geometric reduction algorithm, and Theorem 2.20, which gives characterizations for the uniqueness of DAE solutions via two novel notions: internal equivalence and internal regularity. In section 3.1, we propose the explicitation (with driving variables) procedure to connect nonlinear DAEs to nonlinear control systems. In section 3.2, we show when a nonlinear DAE is externally equivalent to a semi-explicit one and how this problem is related to the explicitation procedure. The most important result in section 3 is Theorem 3.6, which relates equivalences of

DAEs and those of their explicitations, this result will be used throughout the paper for applying nonlinear control theory to nonlinear DAEs. Two nonlinear generalizations of the Weierstrass form are given in Theorem 4.1 and Theorem 4.5 in section 4, respectively. The form in Theorem 4.1 is derived via the zero dynamics algorithm and that in Theorem 4.5 is related to relative degree and invariant distributions in nonlinear control theory. Section 5 contains the conclusions and the perspectives of the paper. The proofs of the main results are put into Appendix.

The following notations will be used throughout the paper. We use $\mathbb{R}^{n \times m}$ to denote the set of real valued matrices with n rows and m columns, $GL(n, \mathbb{R})$ to denote the group of invertible matrices of $\mathbb{R}^{n \times n}$ and I_n to denote the $n \times n$ -identity matrix. For a linear map L , we denote by $\text{rank } L$, $\text{ker } L$ and $\text{Im } L$, the rank, the kernel and the image of L , respectively. Denote by $T_x M$ the tangent space of a submanifold M of \mathbb{R}^n at $x \in M$ and by \mathcal{C}^k the class of functions which are k -times differentiable with continues k -th derivative. For a smooth map $f : X \rightarrow \mathbb{R}$, we denote its differential by $df = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i = [\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n}]$ and for a vector-valued map $f : X \rightarrow \mathbb{R}^m$,

where $f = [f_1, \dots, f_m]^T$, we denote its differential by $Df = \begin{bmatrix} df_1 \\ \vdots \\ df_m \end{bmatrix}$. For two column vectors $v_1 \in \mathbb{R}^m$ and $v_2 \in \mathbb{R}^n$, we write $(v_1, v_2) = [v_1^T, v_2^T]^T \in \mathbb{R}^{m+n}$.

2. Solutions and equivalences of nonlinear DAEs

2.1. The geometric reduction method revisited

In this section, we revise the geometric reduction method in the DAEs solution theory, other formulations of this method can be consulted in section 3.4 of [2], Chapter IV of [1] and [13] for DAEs and [14,16] for DAE control systems. We start from the definition of a solution for a DAE.

Definition 2.1. A solution of a DAE $\Xi_{l,n} = (E, F)$ is a \mathcal{C}^1 -curve $x : I \rightarrow X$ defined on an open interval I such that for all $t \in I$, the curve $x(\cdot)$ satisfies $E(x(t))\dot{x}(t) = F(x(t))$.

Throughout this paper, we will be interested only in solutions of Ξ that are at least \mathcal{C}^1 . A given point x_0 is called *consistent* (or *admissible*) if there exists at least one solution $x(\cdot)$ of Ξ satisfying $x(t_0) = x_0$ (i.e., $E(x_0)\dot{x}(t_0) = F(x_0)$) for a certain $t_0 \in I$; we will denote by S_c the *consistency set*, i.e., the set of all consistent points.

Definition 2.2 (*invariant and locally invariant submanifolds*). Consider a DAE $\Xi_{l,n} = (E, F)$ defined on X . A smooth connected embedded submanifold M of X is called *invariant* if for any point $x_0 \in M$, there exists a solution $x : I \rightarrow X$ of Ξ such that $x(t_0) = x_0$ for a certain $t_0 \in I$ and $x(t) \in M$ for all $t \in I$. Given a point $x_p \in X$, we will say that a submanifold M containing x_p is *locally invariant* (around x_p) if there exists an open neighborhood $U \subseteq X$ of x_p such that $M \cap U$ is invariant.

Proposition 2.3. Consider a DAE $\Xi_{l,n} = (E, F)$ and fix a point x_p . Let M be a smooth connected embedded submanifold containing x_p . If M is a locally invariant submanifold around x_p , then $F(x) \in E(x)T_x M$ for all $x \in M$ around x_p . Conversely, assume that there exists an open neighborhood U of x_p such that, at all $x \in M \cap U$, we have $F(x) \in E(x)T_x M$ and, additionally, $\dim E(x)T_x M = \text{const.}$, then M is a locally invariant submanifold.

The proof is given in Appendix A.

Remark 2.4. Note that the assumption that $\dim E(x)T_x M = \text{const.}$ of Proposition 2.3 is not a necessary condition to conclude that M is an invariant submanifold, but it excludes singular points of DAEs and helps to view a DAE as an ordinary differential equation (ODE) defined on the invariant submanifold. Take the following DAE for an example:

$$\Xi_{1,1} : x\dot{x} = x^2,$$

where $x \in X = \mathbb{R}$. Let $M = X$, then clearly, $F(x) = x^2 \in x \cdot T_x X$, at any $x \in M = \mathbb{R}$. We have $\dim E(x)T_x M$ equals 1 for $x \neq 0$ and is 0 for $x = 0$, so $\dim E(x)T_x M \neq \text{const.}$, for all $x \in M$. Nevertheless, for any $x_0 \in M = \mathbb{R}$, there exists a unique solution $x(t)$ satisfying $x(0) = x_0$, namely, $x(t) = e^t x_0$. Therefore $M = \mathbb{R}$ is an invariant submanifold.

A locally invariant submanifold M^* (around x_p) is called locally *maximal*, if there exists a neighborhood U of x_p such that for any other locally invariant submanifold M , we have $M \cap U \subseteq M^* \cap U$. The *geometric reduction method* for DAEs is the following recursive procedure, which can be used to construct locally maximal invariant submanifold M^* .

Definition 2.5 (*geometric reduction method*). Consider a DAE $\Xi_{l,n} = (E, F)$, fix a point $x_p \in X$ and let U_0 be an open connected subset of X containing x_p . Set $M_0 = X, M_0^c = U_0$. Suppose that there exist an open neighborhood U_{k-1} of x_p and a sequence of smooth connected embedded submanifolds $M_{k-1}^c \subsetneq \cdots \subsetneq M_0^c$ of U_{k-1} for a certain $k \geq 1$, has been constructed. Define recursively

$$M_k := \{x \in M_{k-1}^c : F(x) \in E(x)T_x M_{k-1}^c\}. \tag{4}$$

Then either $x_p \notin M_k$ or $x_p \in M_k$, and in the latter case, assume that there exists a neighborhood U_k of x_p such that $M_k^c = M_k \cap U_k$ is a smooth embedded submanifold (which can always be assumed connected by taking U_k sufficiently small).

Remark 2.6. For a linear DAE $\Delta = (E, H)$ of the form (3), define a sequence of subspaces (one of the Wong sequences [29]) by

$$\mathcal{V}_0 = \mathbb{R}^n, \quad \mathcal{V}_k = H^{-1}E\mathcal{V}_{k-1}, \quad k \geq 1.$$

If we apply the iterative construction of M_k by (4) to the DAE Δ , we get $M_k^c = \mathcal{V}_k, \forall k \geq 0$. Thus the sequence of submanifolds M_k can be seen as a nonlinear generalization of the sequence \mathcal{V}_k .

The following proposition shows that the geometric reduction method above can be used to construct locally maximal invariant submanifold M^* and to deduce that the consistency set S_c , on which the solutions exist, coincides locally with M^* .

Proposition 2.7. *In the geometric reduction method of Definition 2.5, there always exists $k^* \leq n$ such that either k^* is the smallest integer for which $x_p \notin M_{k^*+1}$ or k^* is the smallest integer such that $x_p \in M_{k^*+1}^c$ and $M_{k^*+1}^c \cap U_{k^*+1} = M_{k^*}^c \cap U_{k^*+1}$. In the latter case, we assume that $\dim E(x)T_x M_{k^*+1}^c = \text{const.}$ in a neighborhood $U^* \subseteq U_{k^*+1}$ of x_p in X and then*

- (i) x_p is consistent and $M^* = M_{k^*+1}^c$ is a locally maximal invariant submanifold around x_p .
- (ii) M^* coincides locally with the consistency set S_c , i.e., $M^* \cap U = S_c \cap U^*$ (take a smaller U^* if necessary).

We omit the proof of Proposition 2.7 because it can be seen as a special case (i.e., the control input u is absent) of Proposition 2 of [16] for constructing locally controlled invariant submanifolds of DAE control systems, the reader can consult the proof therein.

2.2. An algorithm for the geometric reduction method

Now we present an algorithm which implements in practice the geometric reduction method. Note that the results of Proposition 2.8 and Theorem 4.1 below will be based on the algorithm.

In what follows, we use the algorithm to show that any DAE Ξ has isomorphic solutions with an “internal” DAE Ξ^* defined on its locally maximal invariant submanifold M^* , which is a practical application (via the algorithm) of Proposition 2.7. In the statement of Proposition 2.8 below, we refer to the submanifold $M^* = M_{k^*+1}^c$, the neighborhood $U^* = U_{k^*+1}^*$, the coordinates $(z^*, \bar{z}_1, \dots, \bar{z}_{k^*})$ on U^* , and the DAE $\Xi_{r^*,n^*}^* = (E^*, F^*)$ defined on M^* by the algorithm, where $E^* = E_{k^*+1} : M^* \rightarrow \mathbb{R}^{r^* \times n^*}$, $F^* = F_{k^*+1} : M^* \rightarrow \mathbb{R}^{r^*}$, $n^* = n_{k^*} = n_{k^*+1}$, $r^* = r_{k^*+1}$ come from Step $k^* + 1$ of the algorithm.

Proposition 2.8 (isomorphic solutions). Consider a DAE $\Xi_{l,n} = (E, F)$ and fix a point $x_p \in X$. Suppose for each Step k ($1 \leq k \leq k^* + 1$) of the algorithm that there exists a neighborhood $U_k \subseteq U_{k-1} \subseteq X$ of x_p such that

Assumption 1: $\text{rank } \tilde{E}_k(z_{k-1}) = \text{const.} = r_k, \forall z_{k-1} \in W_k = U_k \cap M_{k-1}^c$;

Assumption 2: $x_p \in M_k$ and $\text{rank } D\tilde{F}_k^2(z_{k-1}) = \text{const.} = n_{k-1} - n_k$ for $z_{k-1} \in M_k \cap U_k$.

Then M_k^c , for $k = 0, \dots, k^* + 1$, given by (4), are smooth connected embedded submanifolds and $\dim E(x)T_x M^* = \text{const.}$ for all $x \in M^* \cap U^*$. Thus by Proposition 2.7, $x_p \in M^*$ is a consistent point and M^* is a locally maximal invariant submanifold around x_p , given by $M^* = \{x \mid \bar{z}_1(x) = 0, \dots, \bar{z}_{k^*}(x) = 0\}$. Furthermore, for the DAE $\Xi_{r^*,n^*}^* = (E^*, F^*)$, given on M^* by

$$\Xi^* : E^*(z^*)\dot{z}^* = F^*(z^*), \tag{6}$$

where $z^* = z_{k^*+1} = z_{k^*}$ are local coordinates on M^* , we have $\text{rank } E^*(z^*) = r^*, \forall z^* \in M^*$, i.e., $E^*(z^*)$ is of full row rank.

Moreover, the DAE Ξ^* has isomorphic solutions with $\Xi_{l,n}$, i.e., there exists a local diffeomorphism $\Psi : U^* \rightarrow \Psi(U^*)$, $\Psi(x) = \hat{z} = (z^*, \bar{z}) = (z^*, \bar{z}_1, \dots, \bar{z}_{k^*})$, transforming the set of all solutions of $\Xi_{l,n}$ on U^* into that of $\hat{\Xi}_{\hat{l},\hat{n}} = (\hat{E}, \hat{F})$ on $\Psi(U^*)$, where $\hat{l} = r^* + (n - n^*)$, $\hat{n} = n$, given by

$$\hat{\Xi} : \begin{cases} \hat{E}^*(z^*)\dot{\hat{z}}^* = \hat{F}^*(z^*), \\ \bar{z}_1 = 0, \dots, \bar{z}_{k^*} = 0. \end{cases} \tag{7}$$

We omit the proof of Proposition 2.8 because it can be derived from that of Theorem 2(i) in [16].

Algorithm Geometric reduction algorithm for nonlinear DAEs.

Initialization: Consider $\Xi_{l,n} = (E, F)$, fix $x_p \in X$ and let $U_0 \subseteq X$ be an open connected subset containing x_p . Set $z_0 = x$, $E_0(z_0) = E(x)$, $F_0(z_0) = F(x)$, $M_0^c = U_0$, $r_0 = l$, $n_0 = n$, and $\Xi_0 = (E_0, F_0)$. Below all sets U_k are open in X and W_k are open in M_{k-1}^c .

Step k : Suppose that we have defined at Step $k - 1$: an open neighborhood $U_{k-1} \subseteq X$ of x_p , a smooth embedded connected submanifold M_{k-1}^c of U_{k-1} and a DAE $\Xi_{k-1} = (E_{k-1}, F_{k-1})$ given by smooth matrix-valued maps

$$E_{k-1} : M_{k-1}^c \rightarrow \mathbb{R}^{r_{k-1} \times n_{k-1}}, \quad F_{k-1} : M_{k-1}^c \rightarrow \mathbb{R}^{r_{k-1}},$$

whose arguments are denoted $z_{k-1} \in M_{k-1}^c$.

1: Rename the maps as $\tilde{E}_k = E_{k-1}$, $\tilde{F}_k = F_{k-1}$ and define $\tilde{\Xi}_k := (\tilde{E}_k, \tilde{F}_k)$.

Assumption 1: There exists an open neighborhood $U_k \subseteq U_{k-1} \subseteq X$ of x_p such that $\text{rank } \tilde{E}_k(z_{k-1}) = \text{const.} = r_k, \forall z_{k-1} \in W_k = U_k \cap M_{k-1}^c$.

2: Find a smooth map $Q_k : W_k \rightarrow GL(r_{k-1}, \mathbb{R})$, such that \tilde{E}_k^1 of $Q_k \tilde{E}_k = \begin{bmatrix} \tilde{E}_k^1 \\ 0 \end{bmatrix}$ is of full row rank and denote $Q_k \tilde{F}_k = \begin{bmatrix} \tilde{F}_k^1 \\ \tilde{F}_k^2 \end{bmatrix}$, where $\tilde{E}_k^1 : W_k \rightarrow \mathbb{R}^{r_k \times n_{k-1}}$, $\tilde{F}_k^2 : W_k \rightarrow \mathbb{R}^{r_{k-1} - r_k}$ (so all the matrices depend on z_{k-1}); such a map Q_k exists by Dolezal’s theorem [30], see also [31].

3: Following (4), define $M_k = \{z_{k-1} \in W_k \mid \tilde{F}_k^2(z_{k-1}) = 0\}$.

Assumption 2: $x_p \in M_k$ and $\text{rank } D\tilde{F}_k^2(z_{k-1}) = \text{const.} = n_{k-1} - n_k$ for $z_{k-1} \in M_k \cap U_k$.

4: By Assumption 2, $M_k \cap U_k$ is a smooth embedded submanifold and by taking again a smaller U_k , we may assume that $M_k^c = M_k \cap U_k$ is connected and choose new coordinates $(z_k, \bar{z}_k) = \psi_k(z_{k-1})$ on W_k , where $\bar{z}_k = \bar{\varphi}_k(z_{k-1}) = (\bar{\varphi}_k^1(z_{k-1}), \dots, \bar{\varphi}_k^{n_{k-1} - n_k}(z_{k-1}))$, with $d\bar{\varphi}_k^1(z_{k-1}), \dots, d\bar{\varphi}_k^{n_{k-1} - n_k}(z_{k-1})$ being all independent rows of $D\tilde{F}_k^2(z_{k-1})$, and $z_k = \varphi_k(z_{k-1}) = (\varphi_k^1(z_{k-1}), \dots, \varphi_k^{n_k}(z_{k-1}))$ are any complementary coordinates such that $\psi_k = (\varphi_k, \bar{\varphi}_k)$ is a local diffeomorphism.

5: Set $\hat{E}_k = Q_k \tilde{E}_k \left(\frac{\partial \bar{\varphi}_k}{\partial z_{k-1}} \right)^{-1}$, $\hat{F}_k = Q_k \tilde{F}_k$. By Definition 2.10, $\tilde{\Xi}_k \stackrel{ex}{\sim} \hat{\Xi}_k = (\hat{E}_k, \hat{F}_k)$ via Q_k and ψ_k , where

$$\hat{\Xi}_k : \begin{bmatrix} \hat{E}_k^1(z_k, \bar{z}_k) & \bar{E}_k^1(z_k, \bar{z}_k) \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{z}_k \\ \dot{\bar{z}}_k \end{bmatrix} = \begin{bmatrix} \hat{F}_k^1(z_k, \bar{z}_k) \\ \hat{F}_k^2(z_k, \bar{z}_k) \end{bmatrix} \tag{5}$$

with $\hat{E}_k^1 : W_k \rightarrow \mathbb{R}^{r_k \times n_k}$, $\hat{F}_k^1 \circ \psi_k = \tilde{F}_k^1$, $\hat{F}_k^2 \circ \psi_k = \tilde{F}_k^2$ and $[\hat{E}_k^1 \circ \psi_k \quad \bar{E}_k^1 \circ \psi_k] = \tilde{E}_k^1 \left(\frac{\partial \psi_k}{\partial z_{k-1}} \right)^{-1}$.

6: Set $\bar{z}_k = 0$ to define the following reduced and restricted DAE on $M_k^c = \{z_{k-1} \in W_k \mid \bar{z}_k = 0\}$ by

$$\Xi_k : E_k(z_k) \dot{z}_k = F_k(z_k),$$

where $E_k(z_k) = \hat{E}_k^1(z_k, 0)$, $F_k(z_k) = \hat{F}_k^1(z_k, 0)$ are matrix-valued maps and $E_k : M_k^c \rightarrow \mathbb{R}^{r_k \times n_k}$, $F_k : M_k^c \rightarrow \mathbb{R}^{r_k}$.

Repeat: Step k for $k = 1, 2, 3, \dots$, until $n_{k+1} = n_k$, and set $k^* = k$.

Result: Set $n^* = n_{k^*} = n_{k^*+1}$, $r^* = r_{k^*+1}$, $M^* = M_{k^*+1}^c$, $U^* = U_{k^*+1}$, $z^* = z_{k^*+1} = z_{k^*}$ and $\Xi^* = (E^*, F^*)$ with $E^* = E_{k^*+1}$, $F^* = F_{k^*+1}$.

Remark 2.9. (i) The geometric reduction algorithm is a constructive application of Proposition 2.7 but with more assumptions. **Assumption 1** is made to produce the full row rank matrices \tilde{E}_k^1 and the zero-level set $M_k = \{z_{k-1} \in W_k \mid \tilde{F}_k^2(z_{k-1}) = 0\}$. **Assumption 2** assures that $M_k \cap U_k$ is a smooth embedded submanifold and makes it possible to use the components of \tilde{F}_k^2 , with

linearly independent differentials, as a part of new local coordinates. Note that these two assumptions are mild constant rank assumptions made in a neighborhood U_k of the point x_p . In some cases (see Example 2.23 below), the neighborhood U_k can be the whole generalized state space $U_k = X$. While in other cases, we may need to take a smaller neighborhood $U_k \subsetneq U_{k-1}$ such that both assumptions hold on U_k for each Step k , which may result in obtaining a small neighborhood $U^* = U_{k^*+1} \subseteq U_k$ at Step $k^* + 1$.

(ii) The integers r_k, n_k of the geometric reduction algorithm, satisfy, for each $k \geq 1$,

$$\begin{cases} l = r_0 \geq r_1 \geq \dots \geq r_k \geq \dots \geq 0, & n = n_0 \geq n_1 \geq \dots \geq n_k \geq \dots \geq 0, \\ n_{k-1} \geq r_k, & r_{k-1} - r_k \geq n_{k-1} - n_k. \end{cases}$$

2.3. External equivalence, internal equivalence and internal regularity

Two linear DAEs $E\dot{x} = Hx$ and $\tilde{E}\dot{\tilde{x}} = \tilde{H}\tilde{x}$ are called externally equivalent [18] or strictly equivalent [20], if there exist constant invertible matrices Q and P such that $QEP^{-1} = \tilde{E}$ and $QHP^{-1} = \tilde{H}$. Analogously, we define the external equivalence of two nonlinear DAEs as follows.

Definition 2.10 (external equivalence). Two DAEs $\Xi_{l,n} = (E, F)$ and $\tilde{\Xi}_{l,n} = (\tilde{E}, \tilde{F})$ defined on X and \tilde{X} , respectively, are called externally equivalent, shortly ex-equivalent, if there exist a diffeomorphism $\psi : X \rightarrow \tilde{X}$ and a smooth map $Q : X \rightarrow GL(l, \mathbb{R})$ such that

$$\psi^*\tilde{E} = QE \quad \text{and} \quad \psi^*\tilde{F} = QF,$$

where $\psi^*\tilde{E}$ and $\psi^*\tilde{F}$ denote the pull-back [3] of the \mathbb{R}^l -valued differential 1-form \tilde{E} on \tilde{X} and \mathbb{R}^l -valued function \tilde{F} (0-form) on \tilde{X} , respectively, that is,

$$\tilde{E}(\psi(x)) = Q(x)E(x) \left(\frac{\partial \psi(x)}{\partial x} \right)^{-1} \quad \text{and} \quad \tilde{F}(\psi(x)) = Q(x)F(x). \tag{8}$$

The ex-equivalence of two DAEs will be denoted by $\Xi \overset{ex}{\sim} \tilde{\Xi}$. If $\psi : U \rightarrow \tilde{U}$ is a local diffeomorphism between neighborhoods U of x_p and \tilde{U} of \tilde{x}_p , and $Q(x)$ is defined on U , we will speak about local ex-equivalence.

Note that the map Q and the diffeomorphism ψ above should be smooth maps in order to guarantee the smoothness of \tilde{E} and \tilde{F} for the DAE $\tilde{\Xi}$. The following observation relates ex-equivalence with solutions.

Remark 2.11. The ex-equivalence preserves trajectories, i.e., for two DAEs $\Xi \overset{ex}{\sim} \tilde{\Xi}$, if a \mathcal{C}^1 -curve $x(\cdot)$ is a solution of Ξ passing through $x_0 = x(t_0)$, then $\tilde{x} = \psi \circ x$ is a solution of $\tilde{\Xi}$ passing through $\tilde{x}_0 = \psi(x_0)$; but even if we can smoothly conjugate all trajectories of two DAEs, they are not necessarily ex-equivalent. For example, consider $\Xi_1 = (E_1, F_1)$ and $\Xi_2 = (E_2, F_2)$, where $E_1(x) = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $F_1(x) = \begin{bmatrix} x_3^2 \\ x_1 \\ x_2 \end{bmatrix}$, $E_2(x) = \begin{bmatrix} 0 & x_1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$, $F_2(x) = \begin{bmatrix} x_3^2 \\ x_1 \\ x_2 \end{bmatrix}$. Then for both DAEs Ξ_1 and Ξ_2 , the maximal invariant submanifold is $M^* = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_2 = x_3 = 0\}$ and for any $(x_{10}, x_{20}, x_{30}) = (0, 0, x_{30}) \in M^*$, the unique solution of both systems is $x_1(t) = x_2(t) = 0$,

$x_3(t) = \frac{x_{30}}{1-x_{30}t}$. Nevertheless, the DAEs are not ex-equivalent since the distribution $\ker E_1$ is involutive but the distribution $\ker E_2$ is not (clearly, the ex-equivalence of two DAEs preserves the involutivity of $\ker E_1$ and $\ker E_2$ since if $\Xi_1 \stackrel{ex}{\sim} \Xi_2$, via Q and ψ , then $\ker E_2 = \frac{\partial \psi}{\partial x} \ker E_1$).

The results of Proposition 2.8 above show clearly the reason behind Remark 2.11: if we assume two DAEs Ξ and $\tilde{\Xi}$ to have corresponding solutions, this assumption only gives the information that the two internal DAE Ξ^* and $\tilde{\Xi}^*$, which have isomorphic solutions with Ξ and $\tilde{\Xi}$, respectively, are ex-equivalent when restricted to M^* and \tilde{M}^* , respectively, i.e., via a diffeomorphism between the submanifolds M^* and \tilde{M}^* and an invertible map Q defined on the invariant submanifold M^* . We do not know, however, whether the diffeomorphism and the map Q can be extended outside the submanifold M^* . In fact, outside the manifolds M^* and \tilde{M}^* , the two DAEs may have completely different behaviors or even different size of system matrices. This analysis gives a motivation to introduce the concept of *internal equivalence* of two DAEs (see the formal Definition 2.17), which is defined by the ex-equivalence of two internal DAEs. In Proposition 2.8, the internal DAE Ξ^* is defined with the help of the geometric reduction algorithm. Now we introduce two notions: local restriction and full row rank reduction, which can be used to define the internal DAE Ξ^* of a DAE Ξ (which we call the reduction of local M^* -restriction of Ξ , see Proposition 2.16) without going through the algorithm when the invariant submanifold M^* is a priori given. The local restriction of a DAE to a submanifold N (invariant or not) is defined as follows.

Definition 2.12 (local restriction). Consider a DAE $\Xi_{l,n} = (E, F)$ and a smooth connected embedded submanifold $N \subseteq X$ containing a point x_p . Let $\psi(x) = z = (z_1, z_2)$ be local coordinates on a neighborhood U of x_p such that $N \cap U = \{z_2 = 0\}$ and z_1 are thus coordinates on $N \cap U$. The restriction of Ξ to $N \cap U$, called local N -restriction of Ξ and denoted $\Xi|_N$, is

$$\Xi|_N : \tilde{E}(z_1, 0) \begin{bmatrix} \dot{z}_1 \\ 0 \end{bmatrix} = \tilde{F}(z_1, 0), \tag{9}$$

where $\tilde{E} \circ \psi = E \left(\frac{\partial \psi}{\partial x} \right)^{-1}$, $\tilde{F} \circ \psi = F$.

For any DAE $\Xi_{l,n} = (E, F)$, there may exist some redundant equations (in particular, some trivial algebraic equations $0 = 0$ and some dependent equations). In the linear case, we have defined the full rank reduction of a linear DAE (see Definition 6.4 of [18]). We now generalize this notion of reduction to nonlinear DAEs to get rid of their redundant equations.

Definition 2.13 (reduction). For a DAE $\Xi_{l,n} = (E, F)$, assume $\text{rank } E(x) = \text{const.} = q$. Then there exists a smooth map $Q : X \rightarrow GL(l, \mathbb{R})$ such that E_1 of $QE = \begin{bmatrix} E_1 \\ 0 \end{bmatrix}$ is of full row rank q , denote $QF = \begin{bmatrix} F_1 \\ F_2 \end{bmatrix}$. Assume that $\text{rank } DF_2(x) = \text{const.} = \hat{l} - q \leq l - q$. Then the full row rank reduction, shortly reduction, of Ξ , denoted by Ξ^{red} , is the DAE

$$\Xi^{red} : \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix} \dot{x} = \begin{bmatrix} F_1(x) \\ \hat{F}_2(x) \end{bmatrix},$$

where $\hat{F}_2 : X \rightarrow \mathbb{R}^{\hat{l}-q}$ with $D\hat{F}_2$ being all independent rows of DF_2 .

Remark 2.14. Note that the existence of smooth map Q in Definition 2.13 is guaranteed by Dolezal’s theorem [30] and also its generalization [31] under the constant rank assumption of E . Since the choice of $Q(x)$ is not unique, the reduction of Ξ is not unique either. Nevertheless, since $Q(x)$ preserves the solutions, each reduction Ξ^{red} has the same solutions as the original DAE Ξ .

For a locally invariant submanifold M , we consider the local M -restriction $\Xi|_M$ of Ξ , and then we construct a reduction of $\Xi|_M$ and denote it by $\Xi|_M^{red}$. Notice that the order matters: to construct $\Xi|_M^{red}$, we first restrict and then reduce while reducing first and then restricting will not give $\Xi|_M^{red}$ but another DAE $\Xi^{red}|_M$, which may have redundant equations as seen from the following example.

Example 2.15. Consider the following nonlinear DAE $\Xi : \begin{bmatrix} 1 & 1 \\ x & 0 \\ 0 & 0 \\ e^y & e^y \end{bmatrix} \begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} = \begin{bmatrix} x^2 \\ x^3 \\ xy \\ e^y x^2 \end{bmatrix}$ defined on $X = \mathbb{R}^2$. Fix a point $(x_p, y_p) = (1, 0)$, then it is clear that $M^* = \{(x, y) \in \mathbb{R}^2 : x > 0, y = 0\}$ is a locally maximal invariant submanifold around x_p . Set $\psi(x, y) = (z_1, z_2) = (x, y)$ as coordinates on X . Then the M^* -restriction of Ξ , by Definition 2.12, is $\Xi|_{M^*} : \begin{bmatrix} 1 \\ z_1 \\ 0 \\ 1 \end{bmatrix} \dot{z}_1 = \begin{bmatrix} z_1^2 \\ z_1^3 \\ 0 \\ z_1^2 \end{bmatrix}$ and the reduction of $\Xi|_{M^*}$ is $\Xi|_{M^*}^{red} : q(z_1)\dot{z}_1 = q(z_1)z_1^2$, where $q(z_1)$ can be any non-zero function (illustrating that the reduction is not unique). On the other hand, $\Xi^{red}|_{M^*}$ is $\begin{bmatrix} 1 \\ z_1 \\ 0 \end{bmatrix} \dot{z}_1 = \begin{bmatrix} z_1^2 \\ z_1^3 \\ 0 \end{bmatrix}$, and clearly, has redundant equations.

Proposition 2.16. Consider a DAE $\Xi_{l,n} = (E, F)$ and fix a point x_p . Let M be an \bar{n} -dimensional locally invariant submanifold of Ξ around x_p . Assume that $\dim E(x)T_x M = \text{const.} = \bar{r}$ for all $x \in M$ around x_p . Then any reduction $\Xi|_M^{red}$ of the local M -restriction of Ξ is a DAE of the form (1) and the dimensions related to $\Xi|_M^{red}$ are \bar{r} and \bar{n} , i.e., $\Xi|_M^{red} = \bar{\Xi}_{\bar{r},\bar{n}}$. Moreover, the matrix \bar{E} of $\bar{\Xi}_{\bar{r},\bar{n}} = (\bar{E}, \bar{F})$ is of full row rank \bar{r} .

Proof. We skip the proof since we have already constructed $\Xi|_M^{red}$ for M being an invariant submanifold, see (29) in the proof of Proposition 2.3; it is clear that $\bar{E} = [\bar{E}_1^1 \ \bar{E}_1^2]$, $\bar{F} = \bar{F}_1$ and $\text{rank } \bar{E} = \bar{r}$. \square

The definition of the internal equivalence of two DAEs is given as follows.

Definition 2.17. (internal equivalence) Consider two DAEs $\Xi = (E, F)$ and $\tilde{\Xi} = (\tilde{E}, \tilde{F})$, and fix two points $x_p \in X$ and $\tilde{x}_p \in \tilde{X}$. Let M^* and \tilde{M}^* be two locally maximal invariant submanifolds of Ξ and $\tilde{\Xi}$, around x_p and \tilde{x}_p , respectively. Assume that $\dim E(x)T_x M^* = \text{const.}$ for $x \in M^*$ around x_p and $\dim \tilde{E}(\tilde{x})T_{\tilde{x}} \tilde{M}^* = \text{const.}$ for $\tilde{x} \in \tilde{M}^*$ around \tilde{x}_p . Then, Ξ and $\tilde{\Xi}$ are called locally internally equivalent, shortly in-equivalent, if $\Xi|_{M^*}^{red}$ and $\tilde{\Xi}|_{\tilde{M}^*}^{red}$ are ex-equivalent, locally around x_p and \tilde{x}_p , respectively. Denote the in-equivalence of two DAEs by $\Xi \stackrel{in}{\sim} \tilde{\Xi}$.

Remark 2.18. Note that under the assumption that $\dim E(x)T_x M^*$ and $\dim \tilde{E}(\tilde{x})T_{\tilde{x}} \tilde{M}^*$ are constant, by Proposition 2.16 applied to M^* , we have $\Xi|_{M^*}^{red} = \Xi_{r^*, n^*}^*$ and $\tilde{\Xi}|_{M^*}^{red} = \tilde{\Xi}_{\tilde{r}^*, \tilde{n}^*}^*$, where $r^* = \dim E(x)T_x M^*$, $n^* = \dim M^*$ and $\tilde{r}^* = \dim \tilde{E}(\tilde{x})T_{\tilde{x}} \tilde{M}^*$, $\tilde{n}^* = \dim \tilde{M}^*$. The dimensions l and n , related to Ξ , and \tilde{l} and \tilde{n} related to $\tilde{\Xi}$ are not required to be the same. However, if Ξ and $\tilde{\Xi}$ are in-equivalent, then by definition, $\Xi|_{M^*}^{red} = \Xi_{r^*, n^*}^*$ and $\tilde{\Xi}|_{M^*}^{red} = \tilde{\Xi}_{\tilde{r}^*, \tilde{n}^*}^*$ are locally ex-equivalent and thus the dimensions related to them have to be the same, i.e., $r^* = \tilde{r}^*$ and $n^* = \tilde{n}^*$ (and $l^* = r^* = \tilde{r}^* = \tilde{l}^*$ since all reductions of Ξ and $\tilde{\Xi}$ are of full row rank).

Now we will study the uniqueness of solutions of DAEs with the help of the notion of internal equivalence (some other results of uniqueness of DAE solutions can be consulted in e.g., [11, 12]). We will say that a solution $x : I \rightarrow M^*$ of a DAE Ξ satisfying $x(t_0) = x_0$, where $t_0 \in I$ and $x_0 \in M^*$, is maximal if for any solution $\tilde{x} : \tilde{I} \rightarrow M^*$ such that $t_0 \in \tilde{I}$, $\tilde{x}(t_0) = x_0$ and $x(t) = \tilde{x}(t)$, $\forall t \in I \cap \tilde{I}$, we have $\tilde{I} \subseteq I$.

Definition 2.19. (internal regularity) Consider a DAE $\Xi_{l,n} = (E, F)$ and let M^* be a locally maximal invariant submanifold around a point $x_p \in M^*$. Then Ξ is called locally *internally regular* (around x_p) if there exists a neighborhood $U \subseteq X$ of x_p such that for any point $x_0 \in M^* \cap U$, there exists only one maximal solution $x : I \rightarrow M^* \cap U$ satisfying $x(t_0) = x_0$ for a certain $t_0 \in I$.

Theorem 2.20. Consider a DAE $\Xi_{l,n} = (E, F)$ and let M^* be an n^* -dimensional locally maximal invariant submanifold around a point $x_p \in M^*$. Assume that $\dim E(x)T_x M^* = \text{const.} = r^*$ for all $x \in M^*$ around x_p . Then the following conditions are equivalent:

- (i) Ξ is internally regular around x_p ;
- (ii) $\dim M^* = \dim E(x)T_x M^*$, i.e., $n^* = r^*$, for all $x \in M^*$ around x_p ;
- (iii) Ξ is locally internally equivalent to

$$\dot{z}^* = f^*(z^*), \tag{10}$$

for $z^* \in M^* \cap U$, where U is a neighborhood of x_p and f^* is a smooth vector field on $M^* \cap U$.

The proof is given in Appendix A.

Remark 2.21. Theorem 2.20 is a nonlinear generalization of the results on the internal regularity of linear DAEs in [18] (see also [32], where the internal regularity is called autonomy). As stated in Theorem 6.11 of [18], a linear DAE $\Delta = (E, H)$, given by (3), is internally regular if and only if the maximal invariant subspace \mathcal{M}^* of Δ (i.e., the largest subspace such that $H\mathcal{M}^* \subseteq E\mathcal{M}^*$) satisfies $\dim \mathcal{M}^* = \dim E\mathcal{M}^*$. A nonlinear counterpart of the last condition is (ii) of Theorem 2.20 and thus M^* is a natural nonlinear generalization of \mathcal{M}^* . Observe that M^* is the limit of M_k as \mathcal{V}^* is the limit of \mathcal{V}_k , defined in Remark 2.6. Moreover, we have shown in [18] that the maximal invariant subspace $\mathcal{M}^* = \mathcal{V}^*$, where \mathcal{V}^* coincides with the limit of the Wong sequence \mathcal{V}_k defined in Remark 2.6.

For a DAE control system of the form $E(x)\dot{x} = F(x) + G(x)u$, the internal regularization problem, i.e., to find a feedback law $u = \gamma(x)$ such that the DAE $E(x)\dot{x} = F(x) + G(x)\gamma(x)$ is

internally regular, was discussed in [16], via the geometric reduction method. The connections and differences between section 2 and [16] are summarized in the following remark.

Remark 2.22. Note that the algorithm in section 2.2, Propositions 2.7 and 2.8, Remark 2.9 of the present paper are the special cases of, respectively, Algorithm 1, Proposition 2, Theorem 2(i) and Remark 6 in [16] with inputs u been absent. There are, however, essential differences between results of section 2 and those of [16]. Firstly, the main contribution of section 2 is not to discuss geometric reduction method but is to analyze the differences between external equivalence and internal equivalence for DAEs, the notions of restriction $\Xi|_M$ and reduction $\Xi|_M^{red}$ (essential for internal equivalence and formalized in Definitions 2.12 and 2.13, respectively) are novel, while the internal feedback equivalence of DAE control systems was not discussed in [16]. Secondly, Proposition 2.3 is different from the special case of Proposition 1 in [16] because the latter is a necessary and sufficient condition under a suitable constant rank assumption, while, as shown in Proposition 2.3, such constant rank assumption is actually not required for proving one direction, i.e., for proving that if M is locally invariant, then $F(x) \in E(x)T_x M$ for all x around M . Thirdly, Assumption 1 of algorithm, that is, $\text{rank } \tilde{E}_k(\cdot) = \text{const.}$, is replaced by $\text{rank} [\tilde{E}_k(\cdot), \tilde{G}_k(\cdot)]$ in algorithm of [16]. It means that in the latter some singularities can be compensated by the control u while algorithm of section 2.2 detects intrinsic singularities of the pair (E, F) . At last, more detailed explanations on the assumptions of geometric reduction method, such as the requirements $x_p \in M_k$, the connectivity of M_k in Definition 2.5, the necessity of using U_{k^*+1} rather than U_{k^*} in Proposition 2.7, can be consulted in Remark 1(iii) and (iv) of [16].

Example 2.23. Consider a DAE $\Xi_{6,6} = (E, F)$ with the generalized state $x = (x_1, x_2, x_3, x_4, x_5, x_6) \in X$, where $X = \{x \in \mathbb{R}^6 : x_1 \neq x_6, x_6 > 0\}$,

$$\begin{bmatrix} -\ln x_6 & x_6(x_3+x_5) & \frac{x_1 x_5 \ln x_6}{x_1-x_6} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1-\frac{x_1}{x_6} & 0 \\ 0 & 0 & 0 & 0 & x_5 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ \dot{x}_4 \\ \dot{x}_5 \\ \dot{x}_6 \end{bmatrix} = \begin{bmatrix} (x_1-x_6)(x_3+x_5)-(x_2x_6-x_6^2-x_1) \ln x_6 \\ x_5-x_2+x_6 \\ (1-\frac{x_1}{x_6})(x_6^2-x_6x_2+x_4) \\ x_6+x_5(x_6^2-x_6x_2+x_4) \\ \frac{x_1}{x_6} \\ x_3+x_5 \end{bmatrix}. \quad (11)$$

We consider Ξ around a point $x_p = (0, 1, 0, 0, 0, 1)$ and apply to Ξ the algorithm given in section 2.2.

Step 1: We have $\text{rank } E(x) = r_1 = 4$ on $U_1 = X$. Since E is already in the desired form, set $Q_1 = I_6$ to get

$$M_1 = \{x \in X : Q_1 F(x) \in \text{Im } Q_1 E(x)\} = \left\{x \in X : \frac{x_1}{x_6} = 0, x_3 + x_5 = 0\right\}.$$

It is clear that $x_p \in M_1$ and $M_1^c = M_1 \cap U_1 = M_1$ is a locally smooth connected embedded submanifold and $n_1 = \dim M_1^c = 4$. Then choose new coordinates $\bar{z}_1 = (\bar{x}_1, \bar{x}_3) = (\frac{x_1}{x_6}, x_3 + x_5)$ and keep the remaining coordinates $z_1 = (x_2, x_4, x_5, x_6)$ unchanged. The system in new coordinates, denoted $\hat{\Xi}_1$, takes the form

$$\hat{\Xi}_1 : \begin{bmatrix} x_5 \bar{x}_3 & 0 & \frac{-\bar{x}_1 x_5 \ln x_6}{\bar{x}_1 - 1} & \frac{-\bar{x}_1 \ln x_6}{x_6} & -x_6 \ln x_6 & \frac{\bar{x}_1 x_5 \ln x_6}{\bar{x}_1 - 1} \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\bar{x}_1 & 0 & 0 & 1 \\ 0 & 0 & x_5 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{x}_2 \\ \dot{x}_4 \\ \dot{x}_5 \\ \dot{x}_6 \\ \dot{\bar{x}}_1 \\ \dot{\bar{x}}_3 \end{bmatrix} = \begin{bmatrix} \bar{x}_3 x_6 (\bar{x}_1 - 1) - x_6 \ln x_6 (x_2 - x_6 - \bar{x}_1) \\ x_5 - x_2 + x_6 \\ (1 - \bar{x}_1)(x_6^2 - x_6 x_2 + x_4) \\ x_6 + x_5 (x_6^2 - x_6 x_2 + x_4) \\ \bar{x}_1 \\ \bar{x}_3 \end{bmatrix}.$$

By setting $\bar{z}_1 = (\bar{x}_1, \bar{x}_3) = 0$, we get the reduction of M_1^c -restriction of $\hat{\Xi}_1$ (see Definition 2.12 and 2.13) as

$$\Xi_1 = \hat{\Xi}_1|_{M_1^c}^{red} : \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & x_5 & -1 \end{bmatrix} \begin{bmatrix} \dot{x}_2 \\ \dot{x}_4 \\ \dot{x}_5 \\ \dot{x}_6 \end{bmatrix} = \begin{bmatrix} -x_6 \ln x_6 (x_2 - x_6) \\ x_5 - x_2 + x_6 \\ (x_6^2 - x_6 x_2 + x_4) \\ x_6 + x_5 (x_6^2 - x_6 x_2 + x_4) \end{bmatrix}.$$

Step 2: Consider the DAE $\Xi_1 = (E_1, F_1)$. We have $\dim E(x)T_x M_1^c = \text{rank } E_1(z_1) = r_2 = 2$ around x_p (on $W_2 = M_1^c \cap U_2 = M_1^c$, where $U_2 = U_1 = X$). Set $Q_1 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$ and define M_2 by

$$M_2 = \{z_1 : Q_1 F_1(\bar{z}_1) \in \text{Im } Q_1 E_1(z_1)\} = \{z_1 : x_2 - x_6 = 0, x_6^2 - x_6 x_2 + x_4 = 0\}.$$

It is clear that $x_p \in M_2$, $M_2^c = M_2 \cap U_2 = M_2$ and $n_2 = \dim M_2^c = 2$. Then choose new coordinates $\bar{z}_2 = (\bar{x}_2, \bar{x}_4) = (x_2 - x_6, x_6^2 - x_6 x_2 + x_4)$ and keep the remaining coordinates $z_2 = (x_5, x_6)$ unchanged. For the system in new coordinates, denoted $\hat{\Xi}_2$, by a similar procedure as in Step 1, we can define the reduction of M_2^c -restriction of $\hat{\Xi}_2$ as

$$\Xi_2 = \hat{\Xi}_2|_{M_2^c}^{red} : \begin{bmatrix} 0 & 0 \\ x_5 & -1 \end{bmatrix} \begin{bmatrix} \dot{x}_5 \\ \dot{x}_6 \end{bmatrix} = \begin{bmatrix} x_5 \\ x_6 \end{bmatrix}.$$

Step 3: For $\Xi_2 = (E_2, F_2)$, we have $\dim E(x)T_x M_2^c = \text{rank } E_2(z_2) = r_2 = 1$ in $W_3 = M_2^c$. By definition, $M_3^c = M_3 = \{z_2 : x_5 = 0\}$. It can be observed that $\dim M_3^c = n_3 = 1$ and by a similar construction as at former steps, we have

$$\Xi_3 = \bar{\Xi}_2|_{M_3^c}^{red} : -\dot{x}_6 = x_6.$$

Step 4: We have $M_4^c = M_3^c$ ($\dim M_4^c = n_4 = n_3 = 1$) and $\dim E(x)T_x M_4^c = r_4 = 1$, thus $k^* = 3$ and the algorithm stops at Step $k^* + 1 = 4$. Therefore, by Proposition 2.8,

$$\begin{aligned} M^* &= M_4^c = \left\{ x \in \mathbb{R}^6 : x_1 = x_3 = x_4 = x_5 = 0, x_2 = x_6, x_6 > 0 \right\} \\ &= \left\{ x \in \mathbb{R}^6 : \bar{x}_1 = \dots = \bar{x}_5 = 0, \bar{x}_6 > 0 \right\} \end{aligned}$$

is locally maximal invariant and $x_p \in M^*$ is a consistent point. Moreover, since $x_6(t) = e^{-t} x_{60}$ is the unique maximal solution of $\Xi^* = \Xi_3$ passing through $x_0 \in M^*$, we have that $x(t) = \Psi^{-1}(x_6(t), 0, 0, 0, 0, 0) = (0, e^{-t} x_{60}, 0, 0, 0, e^{-t} x_{60})$ is the unique maximal solution of Ξ passing through $x_0 = \Psi^{-1}(x_{60}, 0, 0, 0, 0, 0) \in M^*$, where $\Psi(x) = (x_6, \frac{x_1}{x_6}, x_2 - x_6, x_3 + x_5, x_6^2 - x_2 x_6 + x_4, x_5)$ is a local diffeomorphism (actually, $z^* = x_6$). Hence the DAE Ξ is internally

regular around x_p by definition, which illustrates the results of Theorem 2.20 since $\dim M^* = n_4 = \dim E(x)T_x M^* = r_4 = 1$, and Ξ is in-equivalent to the ODE: $\dot{x}_6 = -x_6$.

3. Analysis of nonlinear DAEs via explicitation

3.1. Explicitation with driving variables of nonlinear DAEs

The explicitation (with driving variables) of a DAE Ξ is the following procedure.

- For a DAE $\Xi_{l,n} = (E, F)$, assume that $\text{rank } E(x) = \text{const.} = q$ in a neighborhood $U \subseteq X$ of a point $x_p \in X$. Then by Dolezal’s theorem [30] (see also [31]), there exists a smooth map $Q : U \rightarrow GL(l, \mathbb{R})$ such that $Q(x)E(x) = \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix}$, where $E_1 : U \rightarrow \mathbb{R}^{q \times n}$, and $\text{rank } E_1(x) = q$. Thus Ξ is, locally on U , ex-equivalent via $Q(x)$ to

$$\begin{cases} E_1(x)\dot{x} = F_1(x), \\ 0 = F_2(x), \end{cases} \tag{12}$$

where $Q(x)F(x) = \begin{bmatrix} F_1(x) \\ F_2(x) \end{bmatrix}$, and where $F_1 : U \rightarrow \mathbb{R}^q, F_2 : U \rightarrow \mathbb{R}^{l-q}$.

- The matrix $E_1(x)$ is of full row rank q , choose its right inverse $E_1^\dagger(x)$, i.e., $E_1 E_1^\dagger = I_q$ and set $f(x) = E_1^\dagger(x)F_1(x)$. The collection of all \dot{x} satisfying $E_1(x)\dot{x} = F_1(x)$ of (12) is given by the differential inclusion:

$$\dot{x} \in f(x) + \ker E_1(x) = f(x) + \ker E(x), \tag{13}$$

where $f(x)$ is a smooth vector field on X .

- Since $\ker E(x)$ is a distribution of constant rank $n - q$, choose locally $m = n - q$ independent vector fields g_1, \dots, g_m on X such that $\ker E(x) = \text{span}\{g_1, \dots, g_m\}(x)$. Then by introducing *driving variables* $v_i, i = 1, \dots, m$, we parametrize the affine distribution $f(x) + \ker E_1(x)$ and thus all solutions of (13) are given by all solutions (corresponding to all controls $v_i(t) \in \mathbb{R}$) of

$$\dot{x} = f(x) + \sum_{i=1}^m g_i(x)v_i. \tag{14}$$

- Form a matrix $g(x) = [g_1(x), \dots, g_m(x)]$. Then, we rewrite equation (14) as $\dot{x} = f(x) + g(x)v$, where $v = (v_1, \dots, v_m)$, and set $h(x) = F_2(x)$. We claim, see Proposition 3.5 below, that all solutions of DAE (12) (and thus of the original DAE Ξ) are in one-to-one correspondence with all solutions (corresponding to all C^0 -controls $v(t)$) of

$$\begin{cases} \dot{x} = f(x) + g(x)v, \\ 0 = h(x). \end{cases} \tag{15}$$

- To (15), we attach the control system $\Sigma = \Sigma_{n,m,p} = (f, g, h)$, given by

$$\Sigma : \begin{cases} \dot{x} = f(x) + g(x)v, \\ y = h(x), \end{cases} \tag{16}$$

where $n = \dim x$, $m = \dim v$, $p = \dim y$. Clearly, $m = n - q$ and $p = l - q$ (we will use these dimensional relations in the following discussion). In the above way, we attach a control system Σ to a DAE Ξ (actually, a class of control systems, see Proposition 3.2 below).

Definition 3.1. (explicitation with driving variables) Given a DAE $\Xi_{l,n} = (E, F)$, fix a point $x_p \in X$ and assume that $\text{rank } E(x) = \text{const.}$ locally around x_p . Then, by a (Q, v) -explicitation we will call any control system $\Sigma = \Sigma_{n,m,p} = (f, g, h)$ given by (16) with

$$f(x) = E_1^\dagger F_1(x), \quad \text{Im } g(x) = \ker E(x), \quad h(x) = F_2(x),$$

where $QE(x) = \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix}$, $QF(x) = \begin{bmatrix} F_1(x) \\ F_2(x) \end{bmatrix}$. The class of all (Q, v) -explicitations will be called shortly the explicitation class. If a particular control system Σ belongs to the explicitation class of Ξ , we will write $\Sigma \in \mathbf{Expl}(\Xi)$.

Notice that a given Ξ has many (Q, v) -explicitations since the construction of $\Sigma \in \mathbf{Expl}(\Xi)$ is not unique: there is a freedom in choosing $Q(x)$, $E_1^\dagger(x)$, and $g(x)$. As a consequence of this non-uniqueness of construction, the explicitation Σ of Ξ is a system defined up to a *feedback transformation*, an *output multiplication* and a *generalized output injection* (or, equivalently, a class of systems).

Proposition 3.2. Assume that a control system $\Sigma_{n,m,p} = (f, g, h)$ is a (Q, v) -explicitation of a DAE $\Xi_{l,n} = (E, F)$ corresponding to a choice of invertible matrix $Q(x)$, right inverse $E_1^\dagger(x)$, and matrix $g(x)$. Then a control system $\tilde{\Sigma}_{n,m,p} = (\tilde{f}, \tilde{g}, \tilde{h})$ is a (\tilde{Q}, \tilde{v}) -explicitation of $\Xi_{l,n}$ corresponding to a choice of invertible matrix $\tilde{Q}(x)$, right inverse $\tilde{E}_1^\dagger(x)$, and matrix $\tilde{g}(x)$ if and only if Σ and $\tilde{\Sigma}$ are equivalent via a v -feedback transformation of the form $v = \alpha(x) + \beta(x)\tilde{v}$, a generalized output injection $\gamma(x)y = \gamma(x)h(x)$ and an output multiplication $\tilde{y} = \eta(x)y$, which map

$$f \mapsto \tilde{f} = f + \gamma h + g\alpha, \quad g \mapsto \tilde{g} = g\beta, \quad h \mapsto \tilde{h} = \eta h, \tag{17}$$

where α, β and η are smooth matrix-valued functions of appropriate sizes, $\gamma = (\gamma_1, \dots, \gamma_p)$ is a p -tuple of smooth vector fields on X , and β and η are invertible.

The proof is given in Appendix B. Since the explicitation of a DAE is a class of control systems, we will propose now an equivalence relation for control systems. An equivalence of two nonlinear control systems is usually defined by state coordinates transformations and feedback transformations (e.g. see [27,28]), and sometimes output coordinates transformations [33]. In the present paper, we define a more general system equivalence of two control systems as follows.

Definition 3.3. (system equivalence) Consider two control systems $\Sigma_{n,m,p} = (f, g, h)$ and $\tilde{\Sigma}_{n,m,p} = (\tilde{f}, \tilde{g}, \tilde{h})$ defined on X and \tilde{X} , respectively. The systems Σ and $\tilde{\Sigma}$ are called system equivalent, or shortly sys-equivalent, denoted by $\Sigma \overset{\text{sys}}{\sim} \tilde{\Sigma}$, if there exist a diffeomorphism $\psi : X \rightarrow \tilde{X}$, matrix-valued functions $\alpha : X \rightarrow \mathbb{R}^m$, $\beta : X \rightarrow GL(m, \mathbb{R})$, $\eta : X \rightarrow GL(p, \mathbb{R})$, and a p -tuple of vector fields $\gamma = (\gamma_1, \dots, \gamma_p)$ on X such that

$$\tilde{f} \circ \psi = \frac{\partial \psi}{\partial x} (f + \gamma h + g\alpha), \quad \tilde{g} \circ \psi = \frac{\partial \psi}{\partial x} g\beta, \quad \tilde{h} \circ \psi = \eta h.$$

If $\psi : U \rightarrow \tilde{U}$ is a local diffeomorphism between neighborhoods U of x_p and \tilde{U} of \tilde{x}_p , and $\alpha, \beta, \gamma, \eta$ are defined locally on U , we will speak about local sys-equivalence.

Remark 3.4. The above defined sys-equivalence of two nonlinear control systems generalizes the Morse equivalence of two linear control systems (see [18,34]).

The following proposition shows that solutions of any DAE are in a one-to-one correspondence with solutions of its (Q, v) -explicitation.

Proposition 3.5. Consider a DAE $\Xi_{l,n} = (E, F)$ and let a control system $\Sigma_{n,m,p} = (f, g, h)$ be a (Q, v) -explicitation of Ξ , i.e., $\Sigma \in \mathbf{Expl}(\Xi)$. Then a C^1 -curve $x(\cdot)$ is a solution of Ξ if and only if there exists $v(\cdot) \in C^0$ such that $(x(\cdot), v(\cdot))$ is a solution of Σ respecting the output constraints $y = 0$, i.e., a solution of (15).

The proof is given in Appendix B. The following theorem is a fundamental result of the present paper, which shows that sys-equivalence for explicitation systems (control systems) is a true counterpart of the ex-equivalence for DAEs.

Theorem 3.6. Consider two DAEs $\Xi_{l,n} = (E, F)$ and $\tilde{\Xi}_{l,n} = (\tilde{E}, \tilde{F})$. Assume that $\text{rank } E(x)$ and $\text{rank } \tilde{E}(\tilde{x})$ are constant around two points x_p and \tilde{x}_p , respectively. Then for any two control systems $\Sigma_{n,m,p} = (f, g, h) \in \mathbf{Expl}(\Xi)$ and $\tilde{\Sigma}_{n,m,p} = (\tilde{f}, \tilde{g}, \tilde{h}) \in \mathbf{Expl}(\tilde{\Xi})$, we have that locally $\Xi \stackrel{\text{ex}}{\sim} \tilde{\Xi}$ if and only if $\Sigma \stackrel{\text{sys}}{\sim} \tilde{\Sigma}$.




Fig. 1. Ex-equivalence of DAEs and sys-equivalence of control systems.

The proof is given in Appendix B. We use Fig. 1 to illustrate the results of Theorem 3.6. In order to show how the explicitation can be useful in the DAEs theory, we discuss below how the analysis of DAEs of sections 2.1 and 2.3 is related to the notion of zero dynamics of nonlinear control theory. For a nonlinear control system $\Sigma_{n,m,p} = (f, g, h)$ and a nominal point x_p , assume $h(x_p) = 0$. Recall its zero dynamics algorithm [27,28].

Step 1: set $N_1 = h^{-1}(0)$. Step k ($k > 1$): assume for some neighborhood $U_{k-1} \subseteq X$ of x_p , $N_{k-1}^c = N_{k-1} \cap U_{k-1}$ is a smooth embedded and connected submanifold such that $x_p \in N_{k-1}^c$. Set

$$N_k = \{x \in N_{k-1}^c : f(x) \in T_x N_{k-1}^c + \text{span}\{g_1(x), \dots, g_m(x)\}\}. \tag{18}$$

For a control system $\Sigma = (f, g, h)$, a smooth embedded connected submanifold N containing a point x_p is called *output zeroing* if (i) $h(x) = 0, \forall x \in N$; (ii) N is locally controlled invariant around x_p (i.e., $\exists u : N \rightarrow \mathbb{R}^m$ and a neighborhood U_p of x_p such that $f(x) + g(x)u(x) \in T_x N, \forall x \in N \cap U_p$). An output zeroing submanifold N^* is locally maximal if for some neighborhood U of x_p , any other output zeroing submanifold N' satisfies $N' \cap U \subseteq N^* \cap U$.

Remark 3.7. (i) It is shown in [27] that N_k is invariant under feedback transformations. Then consider a control system $\tilde{\Sigma} = (\tilde{f}, \tilde{g}, \tilde{h})$, given by applying a *generalized output injection* and an *output multiplication* to Σ , i.e., $\tilde{f} = f + \gamma h, \tilde{g} = g, \tilde{h} = \eta h$, where $\gamma = (\gamma_1, \dots, \gamma_p), \gamma_i : X \rightarrow TX$ and $\eta : X \rightarrow GL(p, \mathbb{R})$. By $\tilde{N}_0 = \tilde{h}^{-1}(0) = h^{-1}(0)$ (since $\eta(x)$ is invertible) and for

$$\begin{aligned} \tilde{N}_k &= \left\{ x \in \tilde{N}_{k-1}^c : f(x) + \gamma h(x) \in T_x \tilde{N}_{k-1}^c + \text{span}\{\tilde{g}_1, \dots, \tilde{g}_m\}(x) \right\} \\ &= \left\{ x \in \tilde{N}_{k-1}^c : f(x) + 0 \in T_x \tilde{N}_{k-1}^c + \text{span}\{g_1, \dots, g_m\}(x) \right\}, \end{aligned}$$

we have $\tilde{N}_k = N_k$ for $k \geq 0$, which means that N_k of the zero dynamics algorithm is invariant under *generalized output injection* and *output multiplication*.

(ii) The sequence of submanifolds N_k^c of the zero dynamics algorithm is well-defined for the class $\mathbf{Expl}(\Xi)$, i.e., does not depend on the choice of $\Sigma \in \mathbf{Expl}(\Xi)$. Indeed, since by Proposition 3.2 any two systems $\Sigma, \Sigma' \in \mathbf{Expl}(\Xi)$ are equivalent via a v -feedback, a generalized output injection, and an output multiplication, then by the argument in item (i) above, we have $\tilde{N}_k = N_k$.

Proposition 3.8. Consider a DAE $\Xi_{l,n} = (E, F)$ satisfying $\text{rank } E(x) = q = \text{const.}$ around a point x_p and a control system $\Sigma = (f, g, h) \in \mathbf{Expl}(\Xi)$. Denote $\mathcal{G}(x) = \text{span}\{g_1, \dots, g_m\}(x)$, where the vector fields $g_i, 1 \leq i \leq m$, are the columns of g . The following conditions

- (A1) for Ξ , the submanifold M_k^c of the geometric reduction method of section 2.1 is smooth, embedded, connected and $\dim E(x)T_x M_{k^*}^c = \text{const.}$ for all $x \in M_{k^*}^c$ around x_p ,
- (A2) for Σ , the submanifold N_k^c of the zero dynamics algorithm above is smooth, embedded, connected and $\dim \mathcal{G}(x) \cap T_x N_{k^*}^c = \text{const.}$ for all $x \in N_{k^*}^c$ around x_p (see Proposition 6.1.1 in [27]),

are equivalent for each $k \geq 1$. Assume that either (A1) or (A2) holds, then the maximal invariant submanifold $M^* = M_{k^*}^c$ of Ξ coincides with the maximal output zeroing submanifold $N^* = N_{k^*}^c$ of Σ . Moreover, Ξ is internally regular (around x_p) if and only if $\mathcal{G}(x_p) \cap T_{x_p} N^* = 0$ (equation (6.4) of [27]).

The proof is given in Appendix B.

Remark 3.9. By Proposition 3.8, if there exists a unique $u = u(x)$ that renders N^* output zeroing and locally maximal control invariant for a control system $\Sigma \in \mathbf{Expl}(\Xi)$, then the original DAE Ξ is internally regular. Since the zero dynamics do not depend on the choice of explicitation, the internal regularity of Ξ corresponds to the fact that the zero output constraint $y(t) = 0$ of any control system $\Sigma \in \mathbf{Expl}(\Xi)$ can be achieved by a unique control $u(t)$ or, equivalently, the zero dynamics of Σ is a unique vector field on N^* .

The explicitation can be also used to characterize solutions of DAEs which are not necessarily internally regular, that is, the restricted DAE Ξ^* , given by (6), has non-unique maximal solutions (recall that Ξ^* has isomorphic solutions with the original DAE Ξ by Proposition 2.8). We now apply the explicitation method to Ξ^* to have the following result.

Proposition 3.10. *Consider a DAE $\Xi = (E, F)$ and fix a point $x_p \in X$. Assume that the locally maximal invariant submanifold M^* around x_p exists and can be constructed via the algorithm in section 2.2. Then the reduction of local M^* -restriction of Ξ , denoted by $\Xi|_{M^*}^{red}$, coincides with the DAE $\Xi^* : E^*(z^*)\dot{z}^* = F^*(z^*)$ of Proposition 2.8 with $E^*(z^*)$ being of full row rank r^* . We have*

- (i) *A curve $z^* : I \rightarrow M^*$ is a solution of Ξ^* if and only if it is an integral curve of the affine distribution $\mathcal{A}(z^*) = f^*(z^*) + \ker E(z^*)$, i.e., $\dot{z}^*(\cdot) \in \mathcal{A}(z^*(\cdot))$, where $f^* = (E^*)^\dagger F^*$.*
- (ii) *C^1 -solutions of Ξ^* are in one-to-one correspondence with those of any (Q, v) -explicitation $\Sigma^* \in \mathbf{Expl}(\Xi^*)$ of the form*

$$\Sigma^* : z^* = f^*(z^*) + g^*(z^*)v,$$

which is a control system without outputs, where $\text{Im } g^ = \ker E$, $g^* = (g_1^*, \dots, g_m^*)$ and $v = (v_1, \dots, v_m)$, and $v(t) \in C^0$.*

- (iii) *If $\ker E = \ker E^*$ is involutive, then Ξ^* is ex-equivalent (that is, the original DAE Ξ is in-equivalent) to a semi-explicit DAE of the form*

$$\dot{z}_1^* = F_1(z_1^*, z_2^*),$$

which can be seen as a control system that is not affine with respect to the control z_2^ .*

Proof. We omit the proof since item (i) is clear, and items (ii) and (iii) can be easily deduced by applying, respectively, the results of Proposition 3.5 and that of Theorem 3.13 (see below) to Ξ^* . \square

3.2. Driving variable reducing and semi-explicit DAEs

Now we will show by an example that sometimes we can reduce some of driving variables of a (Q, v) -explicitation.

Example 3.11. Consider a DAE $\Xi = (E, F)$, given by

$$\begin{bmatrix} \sin x_3 & -\cos x_3 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \end{bmatrix} = \begin{bmatrix} F_1(x) \\ x_1^2 + x_2^2 - 1 \end{bmatrix},$$

where $F_1 : X \rightarrow \mathbb{R}$ is smooth. By $\text{rank } E(x) = 1$, the explicitation class $\mathbf{Expl}(\Xi)$ is not empty. A control system $\Sigma \in \mathbf{Expl}(\Xi)$ is:

$$\Sigma : \begin{cases} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \end{bmatrix} = \begin{bmatrix} \sin x_3 \\ -\cos x_3 \\ 0 \end{bmatrix} F_1(x) + \begin{bmatrix} 0 & \cos x_3 \\ 0 & -\sin x_3 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}, \\ y = x_1^2 + x_2^2 - 1, \end{cases}$$

where $[\sin x_3 \ -\cos x_3 \ 0]^T$ is a right inverse of $E_1(x) = [\sin x_3 \ -\cos x_3 \ 0]$. Now consider the last equation in the dynamics of Σ , which is $\dot{x}_3 = v_1$. Observe that v_1 acts on \dot{x}_3 only, which implies that v_1 is decoupled from the other part of the dynamics. Thus, we may get rid of v_1 and regard x_3 as a new control. Thus the dynamics of Σ become:

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \sin x_3 F_1(x) \\ -\cos x_3 F_1(x) \end{bmatrix} + \begin{bmatrix} \cos x_3 \\ -\sin x_3 \end{bmatrix} v_2,$$

where x_1 and x_2 are new states, x_3 and v_2 are the new control inputs. By rectifying the vector field $g_2 = \cos x_3 \frac{\partial}{\partial x_1} - \sin x_3 \frac{\partial}{\partial x_2}$, we can reduce v_2 in a similar way. We are, however, not able to reduce v_1 and v_2 simultaneously.

Before giving the main result of this subsection, we formally define what we mean by “reducing” variables of a control system Σ .

Definition 3.12 (*driving variable reduction*). For a control system $\Sigma_{n,m,p} = (f, g, h)$, let \mathcal{G}^{red} be an involutive sub-distribution of constant rank k of the distribution $\mathcal{G} = \text{span}\{g_1, \dots, g_m\}$. There exists a feedback transformation and a coordinates change such that, locally, $\mathcal{G}^{red} = \text{span}\left\{\frac{\partial}{\partial x_1^1}, \dots, \frac{\partial}{\partial x_2^k}\right\}$ and Σ takes the form

$$\begin{cases} \dot{x}_1 = f_1(x_1, x_2) + \sum_{i=1}^{m-k} g_1^i(x_1, x_2)v_1^i, \\ \dot{x}_2 = v_2, \\ y = h(x_1, x_2), \end{cases}$$

where $v_2 = (v_2^1, \dots, v_2^k)$. We will say that Σ can be \mathcal{G}^{red} -reduced to the following control system

$$\begin{cases} \dot{x}_1 = f_1(x_1, x_2) + \sum_{i=1}^{m-k} g_1^i(x_1, x_2)v_1^i, \\ y = h(x_1, x_2), \end{cases}$$

where x_2 is a new control and the reduced state x_1 is of dimension $n - k$. We say that Σ can be fully reduced if $\mathcal{G}^{red} = \mathcal{G}$.

Now we connect reducing of control systems with semi-explicit DAEs.

Theorem 3.13. For a DAE $\Xi_{l,n} = (E, F)$, the following statements are equivalent around a point $x_p \in X$:

- (i) $\text{rank } E(x) = \text{const.}$ and the distribution $\ker E(x)$ is involutive.
- (ii) Ξ is locally ex-equivalent to a semi-explicit DAE $\Xi^{SE} : \begin{cases} \dot{x}_1 = F_1(x_1, x_2), \\ 0 = F_2(x_1, x_2). \end{cases}$
- (iii) Any control system $\Sigma = (f, g, h) \in \mathbf{Expl}(\Xi)$ can be fully reduced.

The proof is given in Appendix B.

Remark 3.14. (i) Observe that if Ξ is ex-equivalent to Ξ^{SE} , then by rewriting $x_2 = w$ and choosing the output $y = F_2(x_1, w)$, we get the following control system Σ^w with an input w ,

$$\Sigma^w : \begin{cases} \dot{x}_1 = F_1(x_1, w), \\ y = F_2(x_1, w). \end{cases}$$

The above system Σ^w has the same number of variables as Ξ . Thus Σ^w is an *explicitation without driving variables* of Ξ . So there are two kinds of explicitation for nonlinear DAEs, namely, explicitation with, or without, driving variables (the latter is possible if and only if $\ker E$ is involutive).

(ii) A linear DAE $\Delta = (E, H)$, given by (3), has always two kinds of explicitations, since the rank of E is always constant and the distribution $\mathcal{G} = \ker E$ is always involutive. The relations and differences of the two explicitations for linear DAEs are discussed in [35] and Chapter 3 of [36] (note that the explicitation without driving variables for linear DAEs is called the (Q, P) -explicitation there).

4. Nonlinear generalizations of the Weierstrass form

In this subsection, we will use the explicitation (with driving variables) procedure to transform an internally regular DAE $\Xi_{l,n} = (E, F)$ with $l = n$, into normal forms under the external equivalence. A linear regular DAE is always ex-equivalent (via linear transformations) to the Weierstrass form **WF** [21], given by

$$\mathbf{WF} : \begin{bmatrix} N & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} \dot{z} \\ \dot{z}^* \end{bmatrix} = \begin{bmatrix} I & 0 \\ 0 & A \end{bmatrix} \begin{bmatrix} z \\ z^* \end{bmatrix}, \tag{19}$$

where $N = \text{diag}(N_1, \dots, N_m)$, with $N_i, i = 1, \dots, m$ being nilpotent matrices of index ρ_i , i.e., $N_i^j \neq 0$ for all $j = 1, \dots, \rho_i - 1$ and $N_i^{\rho_i} = 0$. The following theorem generalizes that result and shows that any internally regular nonlinear DAE (under the assumption that some ranks are constant) is always ex-equivalent to a nonlinear Weierstrass form **NWF1** (see (20) below). Note that $\bar{\phi}_k$ in the algorithm in section 2.2, defined on $W_k \subseteq M_k^c$, can be considered as maps on $U_0 \subseteq X$ by taking $\bar{\Phi}_k = \bar{\phi}_k \circ \varphi_{k-1} \circ \dots \circ \varphi_1(x)$. Then for $k \geq 1$, set $H_k = [\bar{\Phi}_1 \ \dots \ \bar{\Phi}_k]^T$ and H_0 is empty. **Assumption 2** of the algorithm of section 2.2 says that $\text{rank } D\bar{F}_k^2(z_{k-1}) = \text{const.}$ for $z_{k-1} \in M_k \cap U_k$. In (A2) below, we replace it by a stronger rank assumption on a neighborhood $U \subseteq X$ of x_p .

Theorem 4.1. Consider a DAE $\Xi_{l,n} = (E, F)$, assume that $\text{rank } E(x) = \text{const.} = q$ around a point x_p . Also assume in the geometric reduction algorithm in section 2.2 that

- (A1) $\dim E(x)T_x M_k^c = \text{const.}$ for $x \in M_k^c$ around $x_p, 1 \leq k \leq k^*$;
- (A2) $\text{rank} \begin{bmatrix} D^{H_{k-1}} \\ D\bar{F}_k^2 \end{bmatrix} (x) = \text{const.}$ for $1 \leq k \leq k^*$ (H_0 is absent) and for all x around x_p ;
- (A3) $l = n$ and $\dim M^* = \dim E(x)T_x M^*,$ i.e., $r^* = n^*,$ for all $x \in M^*$ around x_p .

Then Ξ is internally regular and there exists a neighborhood U of x_p such that Ξ is locally on U ex-equivalent to the DAE (20), represented by the nonlinear Weierstrass form

$$\mathbf{NWF1} : \left[\begin{array}{cccc|c} N_{\rho_1} & 0 & \cdots & 0 & \\ 0 & N_{\rho_2} & \ddots & \vdots & \\ \vdots & \ddots & \ddots & 0 & \\ 0 & \cdots & 0 & N_{\rho_m} & \\ \hline & G(z, z^*) & & & I \end{array} \right] \begin{bmatrix} \dot{z}_1 \\ \dot{z}_2 \\ \vdots \\ \dot{z}_m \\ \dot{z}^* \end{bmatrix} = \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_m \\ f^*(z, z^*) \end{bmatrix} + \begin{bmatrix} a_1 + b_1 \dot{z}^\rho \\ a_2 + b_2 \dot{z}^\rho \\ \vdots \\ a_m + b_m \dot{z}^\rho \\ 0 \end{bmatrix}, \quad (20)$$

where $z_i = (z_i^1, \dots, z_i^{\rho_i})$, $z = (z_1, \dots, z_m)$, and (z, z^*) are new coordinates, and $\dot{z}^\rho = (\dot{z}_1^{\rho_1}, \dot{z}_2^{\rho_2}, \dots, \dot{z}_m^{\rho_m})$, with $m = n - q$. The indices ρ_i , $1 \leq i \leq m$, satisfy $\rho_1 \leq \rho_2 \leq \dots \leq \rho_m$.

More specifically, for $1 \leq i \leq m$, the $\rho_i \times \rho_i$ nilpotent matrices N_{ρ_i} and the ρ_i -dimensional vector-valued functions $a_i + b_i \dot{z}^\rho$ are of the following form

$$N_{\rho_i} = \begin{bmatrix} 0 & & & & \\ 1 & 0 & & & \\ & \ddots & \ddots & & \\ & & & 1 & 0 \end{bmatrix}, \quad a_i + b_i \dot{z}^\rho = \begin{bmatrix} 0 \\ a_i^1 + \sum_{s=1}^m b_{i,s}^1 \dot{z}_s^{\rho_s} \\ \vdots \\ a_i^{\rho_i-1} + \sum_{s=1}^m b_{i,s}^{\rho_i-1} \dot{z}_s^{\rho_s} \end{bmatrix},$$

where the functions $a_i^k, b_{i,s}^k$ satisfy $a_i^k|_{M_k^c} = b_{i,s}^k|_{M_k^c} = 0$, for $1 \leq k \leq \rho_i - 1$.

The proof of Theorem 4.1 is given in Appendix C. This proof is closely related to the zero dynamics algorithm for nonlinear control systems shown in [27] and the construction procedure of the above normal form is not difficult but quite tedious, so in order to avoid reproducing the zero dynamics algorithm, we will use some results directly from [27] with small modifications.

Remark 4.2. (i) Assumption (A1) of Theorem 4.1 is equivalent to **Assumption 1** of the geometric reduction algorithm in section 2.2. By Theorem 2.20, we know that (A3) of Theorem 4.1 implies that Ξ is internally regular around x_p .

(ii) A component-wise expression of the above **NWF1** is

$$\mathbf{NWF1} : \begin{cases} 0 = z_i^1, & 1 \leq i \leq m, \\ \dot{z}_i^k = z_i^{k+1} + a_i^k + \sum_{s=1}^m b_{i,s}^k \dot{z}_s^{\rho_s}, & 1 \leq k \leq \rho_i - 1, \\ \dot{z}^* = f^* - G\dot{z}, \end{cases}$$

where $a_i^k, b_{i,s}^k, f^*$ and G depend on (z, z^*) .

(iii) The submanifolds $M_k^c, k \geq 1$, of the algorithm are given by

$$M_k^c = \left\{ (z, z^*) : z_i^j = 0, 1 \leq i \leq m, 1 \leq j \leq k \right\},$$

and the maximal invariant submanifold M^* is given by

$$M^* = \{(z, z^*) : z_i^j = 0, 1 \leq i \leq m, 1 \leq j \leq \rho_i\}.$$

Therefore, an equivalent condition for $a_i^k|_{M_k^c} = b_{i,s}^k|_{M_k^c} = 0$ is that $a_i^k, b_{i,s}^k \in \mathbf{I}^k$, where \mathbf{I}^k is the ideal generated by $z_i^j, 1 \leq i \leq m, 1 \leq j \leq k$ in the ring of smooth functions of z_b^a and z_c^* .

(iv) We see that all maximal solutions $(z(\cdot), z^*(\cdot))$ are unique and of the form $(0, z^*(\cdot))$, where $z^*(\cdot)$ are maximal solutions of the ODE $\dot{z}^* = f^*(0, z^*)$ on M^* , which agrees with the result of Theorem 2.20(iii).

Example 4.3 (continuation of Example 2.23). Consider the DAE $\Xi_{6,6} = (E, F)$ of (11) around the point $x_p = (0, 1, 0, 0, 0, 1)$. A control system $\Sigma_{6,2,2} \in \mathbf{Expl}(\Xi)$ is

$$\Sigma : \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ \dot{x}_4 \\ \dot{x}_5 \\ \dot{x}_6 \end{bmatrix} = \begin{bmatrix} x_6(x_2-x_6)-x_1 \\ \frac{x_1}{x_6}-1 \\ 0 \\ x_5-x_2+x_6 \\ x_4-x_6(x_2-x_6) \\ -x_6 \end{bmatrix} + \begin{bmatrix} x_6(x_3+x_5) & \frac{x_1x_5}{x_6} \\ \ln x_6 & 0 \\ 0 & \frac{x_1}{x_6}-1 \\ 0 & 0 \\ 0 & 1 \\ 0 & x_5 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix}, \quad \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} \frac{x_1}{x_6} \\ x_3+x_5 \end{bmatrix}.$$

It can be observed from Example 2.23 that the assumptions (A1)-(A3) of Theorem 4.1 are satisfied. Now via the following local change of coordinates defined on $U = X = \{x \in X : x_6 > 0, x_1 \neq x_6\}$:

$$z_1^1 = \frac{x_1}{x_6}, \quad z_1^2 = x_2 - x_6, \quad z_1^3 = x_3 + x_5, \quad z_2^2 = x_4 - x_2x_6 + x_6^2, \quad z_2^3 = x_5, \quad z^* = x_6,$$

we can bring Σ into the system Σ' below, which is of the zero dynamics form (40),

$$\Sigma' : \begin{cases} y_1 = z_1^1 \\ \dot{z}_1^1 = z_2^1 + z_2^1 v_1 \\ \dot{z}_1^2 = z_1^1 + \ln z^* \cdot v_1 - z_2^3 v_2 \\ y_2 = z_2^1 \\ \dot{z}_2^1 = z_2^2 + z_1^1 v_2 \\ \dot{z}_2^2 = z_2^3 + z^*(z_1^1 + \ln z^* \cdot v_1 - z_2^3 v_2) - z_1^2 z_2^3 v_2 \\ \dot{z}_2^3 = z_2^2 + v_2 \\ \dot{z}^* = -z^* + z_2^3 v_2, \end{cases} \Rightarrow \Sigma'' : \begin{cases} y_1 = z_1^1 \\ \dot{z}_1^1 = z_2^1 - \frac{z_1^1 z_2^1}{\ln z^*} + \frac{z_2^1}{\ln z^*} \tilde{v}_1 + \frac{z_2^1 z_2^3}{\ln z^*} \tilde{v}_2 \\ \dot{z}_1^2 = \tilde{v}_1 \\ y_2 = z_2^1 \\ \dot{z}_2^1 = z_2^2 - z_1^1 z_2^2 + z_1^1 \tilde{v}_2 \\ \dot{z}_2^2 = z_2^3 + z^* \tilde{v}_1 + z_1^2 z_2^2 z_2^3 - z_1^2 z_2^3 \tilde{v}_2 \\ \dot{z}_2^3 = \tilde{v}_2 \\ \dot{z}^* = -z^* - z_2^2 z_2^3 + z_2^3 \tilde{v}_2, \end{cases}$$

where the feedback transformation

$$\begin{bmatrix} \tilde{v}_1 \\ \tilde{v}_2 \end{bmatrix} = \begin{bmatrix} \frac{x_1}{x_6} \\ x_4-x_2x_6+x_6^2 \end{bmatrix} + \begin{bmatrix} \ln x_6 & -x_5 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix},$$

brings the system Σ' into the system Σ'' above. In order to eliminate $z^* \tilde{v}_1$ in $\dot{z}_2^2 = z_2^3 + z^* \tilde{v}_1 + z_1^2 z_2^2 z_2^3 - z_1^2 z_2^3 \tilde{v}_2$ of Σ'' , we define the change of coordinates

$$\tilde{z}_1^1 = z_1^1, \quad \tilde{z}_1^2 = z_1^2, \quad \tilde{z}_2^1 = z_2^1 - z^* z_1^1, \quad \tilde{z}_2^2 = z_2^2 - z^* z_1^2, \quad \tilde{z}_2^3 = z_2^3,$$

and the output multiplication $\begin{bmatrix} \tilde{y}_1 \\ \tilde{y}_2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ z^* & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$. Then the system Σ'' becomes

$$\tilde{\Sigma} : \begin{cases} \tilde{y}_1 = \tilde{z}_1^1 \\ \dot{\tilde{z}}_1^1 = \tilde{z}_1^2 - \frac{\tilde{z}_1^1(\tilde{z}_2^1 + \tilde{z}_1^1 z^*)}{\ln z^*} + \frac{(\tilde{z}_2^1 + \tilde{z}_1^1 z^*)}{\ln z^*} \tilde{v}_1 + \frac{(\tilde{z}_2^1 + \tilde{z}_1^1 z^*) \tilde{z}_2^3}{\ln z^*} \tilde{v}_2 \\ \dot{\tilde{z}}_1^2 = \tilde{v}_1 \\ \tilde{y}_2 = \tilde{z}_2^1 \\ \dot{\tilde{z}}_2^1 = \tilde{z}_2^2 + \tilde{z}_1^1(\tilde{z}_2^2 + \tilde{z}_1^2 z^*)(\tilde{z}_2^3 - 1) + \frac{\tilde{z}_1^1 \tilde{z}_2^2 z^*}{\ln z^*} - \frac{(\tilde{z}_2^1 + \tilde{z}_1^1 z^*) z^*}{\ln z^*} \tilde{v}_1 - (\tilde{z}_1^1 \tilde{z}_2^3 + \frac{(\tilde{z}_2^1 + \tilde{z}_1^1 z^*) \tilde{z}_2^3 z^*}{\ln z^*}) \tilde{v}_2 \\ \dot{\tilde{z}}_2^2 = \tilde{z}_2^3 + \tilde{z}_1^2 z^* \\ \dot{\tilde{z}}_2^3 = \tilde{v}_2 \\ \dot{z}^* = -z^* - \tilde{z}_2^3(\tilde{z}_2^2 + \tilde{z}_1^2 z^*) + \tilde{z}_2^3 \tilde{v}_2. \end{cases}$$

Now we drop all the tildes in the system $\tilde{\Sigma}$ for ease of notation. By setting $y_1 = y_2 = 0$, replacing $v_1 = \dot{z}_1^2$, $v_2 = \dot{z}_2^3$, and deleting the equations $\dot{z}_1^2 = v_1$, $\dot{z}_2^3 = v_2$, we get the following DAE Ξ from $\tilde{\Sigma}$,

$$\Xi : \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \dot{z}_1^1 \\ \dot{z}_1^2 \\ \dot{z}_2^1 \\ \dot{z}_2^2 \\ \dot{z}_2^3 \\ \dot{z}^* \end{bmatrix} = \begin{bmatrix} z_1^1 \\ z_1^2 \\ z_2^1 \\ z_2^2 \\ z_2^3 \\ -z^* - z_2^3(z_2^2 + z_1^2 z^*) \end{bmatrix} + \begin{bmatrix} 0 \\ a_1^1 + b_{11}^1 \dot{z}_1^2 + b_{12}^1 \dot{z}_2^3 \\ 0 \\ a_2^1 + b_{21}^1 \dot{z}_1^2 + b_{22}^1 \dot{z}_2^3 \\ a_2^2 \\ 0 \end{bmatrix}, \tag{21}$$

where $a_1^1 = -\frac{z_1^1(z_2^1 + z_1^1 z^*)}{\ln z^*}$, $b_{11}^1 = \frac{(z_2^1 + z_1^1 z^*)}{\ln z^*}$, $b_{12}^1 = \frac{(z_2^1 + z_1^1 z^*) z_2^3}{\ln z^*}$, $a_2^1 = z_1^1(z_2^2 + z_1^2 z^*)(z_2^3 - 1) + \frac{z_1^1 z_2^2 z^*}{\ln z^*}$, $b_{21}^1 = -\frac{(z_2^1 + z_1^1 z^*) z^*}{\ln z^*}$, $b_{22}^1 = z_1^1 z_2^3 + \frac{(z_2^1 + z_1^1 z^*) z_2^3 z^*}{\ln z^*}$, $a_2^2 = z_1^2 z^*$. It is clear that $\tilde{\Sigma} \in \mathbf{Expl}(\tilde{\Xi})$, thus we have $\Xi \stackrel{ex}{\sim} \tilde{\Xi}$ since $\Sigma \in \mathbf{Expl}(\Xi)$ and $\Sigma \stackrel{sys}{\sim} \tilde{\Sigma}$ (see Theorem 3.6). The above DAE $\tilde{\Xi}$ is in the **NWF1** of (20) and the sequence of submanifolds M_k^c of the geometric reduction algorithm can be expressed as $M_1^c = \{(z, z^*) : z_1^1 = z_1^2 = 0\}$, $M_2^c = \{(z, z^*) \in M_1^c : z_2^1 = z_2^2 = 0\}$ and

$$M^* = M_3^c = \{(z, z^*) \in M_2^c : z_2^3 = 0\}.$$

The functions $a_1^1, b_{11}^1, b_{12}^1, a_2^1, b_{21}^1, b_{22}^1 \in \mathbf{I}^1$ vanish on M_1^c , and the function $a_2^2 \in \mathbf{I}^2$ vanishes on M_2^c .

The form **NWF1** of Theorem 4.1 is related to the zero dynamics of nonlinear control systems. In the remaining part of this section, we will use the notions of (vector) relative degree and invariant distributions of nonlinear control theory to study when a DAE Ξ is ex-equivalent to a simpler form

$$\mathbf{NWF2} : \begin{bmatrix} N & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} \dot{z} \\ \dot{z}^* \end{bmatrix} = \begin{bmatrix} z \\ f^*(z^*) \end{bmatrix}, \tag{22}$$

where $N = \text{diag}(N_1, \dots, N_m)$, with $N_i \in \mathbb{R}^{\rho_i \times \rho_i}$, $i = 1, \dots, m$, being nilpotent matrices of index ρ_i . The **NWF2** is a perfect nonlinear counterpart of the linear **WF** because the nonlinear terms

G, a_i and b_j of **NWF1** are absent in **NWF2** and f^* depends on z^* -variables only. We now recall the definitions of (vector) relative degree and (conditional) invariant distributions for nonlinear control systems.

Definition 4.4 (relative degree [27]). A square control system $\Sigma_{n,m,m} = (f, g, h)$ has a (vector) relative degree $\rho = (\rho_1, \dots, \rho_m)$ at a point x_p if (i) $L_{g_j} L_f^k h_i(x) = 0$ for all $1 \leq j \leq m, k < \rho_i - 1$, for all $1 \leq i \leq m$, and for all x in a neighborhood of x_p ; (ii) the $m \times m$ decoupling matrix $D(x) = (L_{g_j} L_f^{\rho_i - 1} h_i(x))$, $1 \leq i, j \leq m$, is invertible around x_p .

For a nonlinear control system $\Sigma_{n,m,p} = (f, g, h)$, define a sequence of distributions S_i by

$$\begin{cases} S_1 & := \mathcal{G} = \text{span}\{g_1, \dots, g_m\}, \\ S_{i+1} & := S_i + [f, S_i \cap \ker dh] + [\mathcal{G}, S_i \cap \ker dh], \\ S^* & := \sum_{i \geq 1} S_i, \end{cases} \tag{23}$$

where $[f, v]$ stands for the Lie bracket of vector fields f and v , and $[f, V] = \{[f, v] : v \in V\}$.

Theorem 4.5. For a nonlinear DAE $\Xi_{n,n} = (E, F)$ (i.e., $l = n$), assume that $\text{rank } E(x) = \text{const.}$ around a point $x_p \in X$. Then Ξ is locally ex-equivalent to the **NWF2**, given by (22), around x_p if and only if there exists a control system $\Sigma = \Sigma_{n,m,m} = (f, g, h) \in \mathbf{Expl}(\Xi)$ such that

- (i) the system Σ has a well-defined relative degree $\rho = (\rho_1, \dots, \rho_m)$ at $x = x_p$;
- (ii) the distributions S_i of Σ , defined by (23), are involutive for all $1 \leq i \leq n - 1$.

We omit the proof the Theorem 4.5 since it is a consequence of Theorem 3.6 and some results from nonlinear control theory, see Remark 4.6(i) below.

Remark 4.6. (i) Note that, under conditions (i) and (ii) of Theorem 4.5, using the results of [33], we can transform the system Σ into the following form (called the input-output special form in [33]) via suitable coordinates transformations and feedback transformations,

$$\begin{cases} \dot{z}^* = \bar{f}^*(z^*, y), \\ \dot{z}_i^j = z_i^{j+1}, \quad 1 \leq j \leq \rho_i - 1, \quad 1 \leq i \leq m, \\ \dot{z}_i^{\rho_i} = v_i, \\ y_i = z_i. \end{cases} \tag{24}$$

Rewrite $\bar{f}^*(z^*, y) = \bar{f}^*(z^*, 0) + \gamma(z^*, y)y$ for some smooth function γ , then we can always get rid of the y -variables in $\bar{f}^*(z^*, y)$ by an output injection $\bar{f}^* \mapsto \bar{f}^* - \gamma y = f^*$, where $f^* = f^*(z^*)$. Thus the system Σ is always sys-equivalent to the system $\tilde{\Sigma}$ below

$$\Sigma \stackrel{\text{sys}}{\sim} \tilde{\Sigma} : \begin{cases} \dot{z}^* = f^*(z^*), \\ \dot{z}_i^j = z_i^{j+1}, \quad 1 \leq j \leq \rho_i - 1, \\ \dot{z}_i^{\rho_i} = v_i, \quad 1 \leq i \leq m, \\ y_i = z_i. \end{cases} \xleftrightarrow{\text{Thm. 3.6}} \Xi \stackrel{\text{ex}}{\sim} \tilde{\Xi} : \begin{cases} \dot{z}^* = f^*(z^*), \\ 0 = z_i, \quad 1 \leq i \leq m, \\ \dot{z}_i^j = z_i^{j+1}, \quad 1 \leq j \leq \rho_i - 1. \end{cases}$$

So by Theorem 3.6, the DAE Ξ is ex-equivalent to $\tilde{\Xi}$ represented in the **NWF2** since $\tilde{\Sigma} \in \mathbf{Expl}(\tilde{\Xi})$.

(ii) The linear counterparts of the distributions S_i , given by (23), for linear control systems of the form $\Lambda : \begin{cases} \dot{x} = Ax + Bv \\ y = Cv \end{cases}$ is $\mathcal{W}_1 = \text{Im } B$, $\mathcal{W}_{i+1} = A(\mathcal{W}_i \cap \ker C) + \text{Im } B$, and are called the conditional invariant subspaces. We have shown in [35] that for a linear DAE $\Delta = (E, H)$, if a control system $\Lambda \in \mathbf{Expl}(\Delta)$, then for all $i \geq 1$, the subspaces \mathcal{W}_i coincide with the Wong sequences \mathcal{W}_i of Δ , given by $\mathcal{W}_1 = \ker E$, $\mathcal{W}_{i+1} = E^{-1}H\mathcal{W}_i$. Therefore, the sequences of distributions S_i can be seen as a nonlinear generalization of the Wong sequence subspaces \mathcal{W}_i .

(iii) Although conditions (i) and (ii) of Theorem 4.5 are necessary and sufficient for Ξ being locally ex-equivalent to **NWF2**, it is, in general, not easy to check them because the relative degree and the involutivity of distributions S_i are *not* invariant under output multiplications and output injections (the two properties are invariant under coordinates changes and feedback). From Proposition 3.2, we know that a control system $\Sigma \in \mathbf{Expl}(\Xi)$ is defined up to a feedback transformation, an output multiplication and a generalized output injection. So it is possible that for one system in $\mathbf{Expl}(\Xi)$, conditions (i) and (ii) hold while for another explicitation system the two conditions (or one of them) are not satisfied. The problem of finding easily checkable conditions for a DAE being ex-equivalent to the **NWF2** remains open and, in view of the above analysis, is challenging.

Example 4.7. Consider a classical pendulum system shown in Fig. 2, which is a hanging rigid wire with a ball attached to its end (see also [7,12,13]).




Fig. 2. A hanging wire with a ball attached at the end.

In this system, m is the mass of the ball, τ is the tension force of the wire, θ is the angle between the wire and the y -axis. The system can be modeled by Newton’s law:

$$m\ddot{x} = -\tau \sin \theta = -\frac{\tau x}{l}, \quad m\ddot{y} = -\tau \cos \theta - mg = -\frac{\tau y}{l} - mg, \tag{25}$$

under the following constraint

$$x^2 + y^2 = l^2. \tag{26}$$

Denote $x_1 = x$, $x_2 = \dot{x}$, $x_3 = y$, $x_4 = \dot{y}$, $x_5 = \frac{\tau}{ml}$, then (25) and (26) result in a nonlinear DAE $\Xi_{5,5} = (E, F)$, given by

$$\Xi : \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ \dot{x}_4 \\ \dot{x}_5 \end{bmatrix} = \begin{bmatrix} x_2 \\ -x_5x_1 \\ x_4 \\ -x_5x_3-g \\ x_1^2+x_3^2-l^2 \end{bmatrix}. \tag{27}$$

A control system $\Sigma_{5,1,1} = (f, g, h) \in \mathbf{Expl}(\Xi)$ can be chosen as

$$\Sigma : \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \\ \dot{x}_4 \\ \dot{x}_5 \end{bmatrix} = \begin{bmatrix} x_2 \\ -x_5x_1 \\ x_4 \\ -x_5x_3-g \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} v, \quad y = x_1^2 + x_3^2 - l^2.$$

We consider a nominal point x_p , given by

$$x_p = (x_{1p}, x_{2p}, x_{3p}, x_{4p}, x_{5p}) = (0, 0, l, 0, -\frac{g}{l}).$$

Clearly, the relative degree of Σ is $\rho = 3$ at x_p because $L_g h(x) = L_g L_f h(x) \equiv 0$ and $L_g L_f^2 h(x_p) \neq 0$. By (23), we have

$$S_1 = \text{span}\{g\}, \quad S_2 = \text{span}\{g, ad_f g\}, \quad S_3 = \text{span}\{g, ad_f g, ad_f^2 g\}, \quad S^* = S_4 = S_3,$$

where $g = \frac{\partial}{\partial x_5}$, $ad_f g = -x_1 \frac{\partial}{\partial x_2} - x_3 \frac{\partial}{\partial x_4}$, $ad_f^2 g = x_1 \frac{\partial}{\partial x_1} - x_2 \frac{\partial}{\partial x_2} + x_3 \frac{\partial}{\partial x_3} - x_4 \frac{\partial}{\partial x_4}$. It can be deduced that the distributions $S_i, i = 1, 2, 3$, are all involutive around x_p . Thus conditions (i) and (ii) of Theorem 4.5 are satisfied. Then via the following local coordinates transformation

$$\begin{aligned} \tilde{x}_1 &= -x_1/x_3, & \tilde{x}_2 &= x_1x_4 - x_2x_3, & \tilde{x}_3 &= x_1^2 + x_3^2 - l^2, \\ \tilde{x}_4 &= x_1x_2 + x_3x_4, & \tilde{x}_5 &= -x_5(x_1^2 + x_3^2) + x_2^2 + x_4^2 - gx_3, \end{aligned} \tag{28}$$

and a suitable feedback transformation, the system Σ can be transformed into its input-output special form (see (24)), given by

$$\begin{bmatrix} \dot{\tilde{x}}_1 \\ \dot{\tilde{x}}_2 \\ \dot{\tilde{x}}_3 \\ \dot{\tilde{x}}_4 \\ \dot{\tilde{x}}_5 \end{bmatrix} = \begin{bmatrix} \frac{\tilde{x}_2(1+\tilde{x}_1^2)}{l^2+\tilde{x}_3} \\ -g\tilde{x}_1 \left(\frac{l^2+\tilde{x}_3}{1+\tilde{x}_1^2} \right)^{1/2} \\ \tilde{x}_4 \\ \tilde{x}_5 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix} \tilde{v}, \quad y = \tilde{x}_3.$$

Notice that we can always write $\frac{\tilde{x}_2(1+\tilde{x}_1^2)}{l^2+\tilde{x}_3} = \frac{\tilde{x}_2(1+\tilde{x}_1^2)}{l^2} + \tilde{x}_3 a(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$ and $-g\tilde{x}_1 \left(\frac{l^2+\tilde{x}_3}{1+\tilde{x}_1^2} \right)^{1/2} = -\frac{gl\tilde{x}_1}{\sqrt{1+\tilde{x}_1^2}} + \tilde{x}_3 b(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$ for some smooth functions a and b . After using an output injection to get rid of the terms $\tilde{x}_3 a(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$ and $\tilde{x}_3 b(\tilde{x}_1, \tilde{x}_2, \tilde{x}_3)$, we can see that Σ is locally sys-equivalent to the system $\tilde{\Sigma}$ below.

$$\Sigma^{sys} \sim \tilde{\Sigma} : \begin{cases} \dot{\tilde{x}}_1 = \frac{\tilde{x}_2(1+\tilde{x}_1^2)}{l^2}, \\ \dot{\tilde{x}}_2 = -\frac{gl\tilde{x}_1}{\sqrt{1+\tilde{x}_1^2}}, \\ \dot{\tilde{x}}_3 = \tilde{x}_4, \\ \dot{\tilde{x}}_4 = \tilde{x}_5, \\ \dot{\tilde{x}}_5 = \tilde{v}, \\ y = \tilde{x}_3. \end{cases} \xleftrightarrow{\text{Thm. 3.6}} \Xi^{ex} \sim \tilde{\Xi} : \begin{cases} \dot{\tilde{x}}_1 = \frac{\tilde{x}_2(1+\tilde{x}_1^2)}{l^2}, \\ \dot{\tilde{x}}_2 = -\frac{gl\tilde{x}_1}{\sqrt{1+\tilde{x}_1^2}}, \\ \dot{\tilde{x}}_3 = \tilde{x}_4, \\ \dot{\tilde{x}}_4 = \tilde{x}_5, \\ 0 = \tilde{x}_3. \end{cases}$$

Hence the original DAE Ξ is locally ex-equivalent to $\tilde{\Xi}$, represented in the **NWF2**, via $Q =$

$$\begin{bmatrix} 1 & 0 & 0 & 0 & -a \\ 0 & 1 & 0 & 0 & -b \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix} \text{ and the coordinates transformations given by (28).}$$

5. Conclusions and perspectives

In this paper, we first revise the geometric reduction method for the existence of solutions of nonlinear DAEs, then we define the notions of internal and external equivalence and their differences are discussed by analyzing their relations with solutions. We show that the internal regularity (existence and uniqueness of solutions) of a DAE is equivalent to the fact that the DAE is internally equivalent to an ODE (without free variables) on its maximal invariant submanifold. A procedure named explicitation with driving variables is proposed to connect nonlinear DAEs with nonlinear control systems. We show that the external equivalence for two DAEs is the same as the system equivalence for their explicitation systems. Moreover, we show that Ξ is externally equivalent to a semi-explicit DAE if and only if the distribution defined by $\ker E(x)$ is of constant rank and involutive. If so, the driving variables of a control system $\Sigma \in \mathbf{Expl}(\Xi)$ can be fully reduced. Finally, two nonlinear generalizations of the Weierstrass form **WF** are proposed based on the explicitation method and the notions of nonlinear control theory, such as zero dynamics, relative degree and invariant distributions.

Several results of the paper can be used for further studies on nonlinear DAE systems. The geometric reduction algorithm and internal equivalence can be used for the stability analysis since DAEs has isomorphic solutions with an ODE (with free variables) by Propositions 2.8 and 3.10, the stability of the ODE clearly indicates that of the original DAE. The explicitation method shown in section 3, in particular, the results of Theorem 3.6, are fundamental tools for applying nonlinear geometric control theory to solve problems like exact linearization, disturbance decoupling, controllability and observability analysis for DAEs systems (possibly with extra controls or disturbances). Finally, as the Weierstrass form is useful for linear DAEs, the two nonlinear Weierstrass form can be applied to study e.g., index analysis, jump solutions, impulse-freeness, of nonlinear DAEs.

Acknowledgment

The first author is currently supported by Vidi-grant 639.032.733. The authors would like to thank the reviewers whose suggestions and remarks have improved the final presentation of the paper.

Appendix A. Proofs of Proposition 2.3 and Theorem 2.20

Proof of Proposition 2.3. Suppose that M is a locally invariant submanifold around x_p . By Definition 2.2, there exists a neighborhood U of x_p such that for any point $x_0 \in M \cap U$, there exists a solution $x : I \rightarrow M \cap U$ satisfying $x(t_0) = x_0$ for a certain $t_0 \in I$. Then we have $F(x(t)) = E(x(t))\dot{x}(t) \in E(x(t))T_{x(t)}M, \forall t \in I$. It follows that $F(x_0) \in E(x_0)T_{x_0}M$ by taking $t = t_0$. Hence $F(x) \in E(x)T_xM$ for all $x \in M \cap U$.

Conversely, suppose that $\dim E(x)T_xM = \text{const.} = \bar{r}$ and $F(x) \in E(x)T_xM$ locally for all $x \in M \cap U$. Notice that M is a smooth connected embedded submanifold, thus there exists a smaller neighborhood U_1 of x_p and local coordinates $\psi(x) = z = (z_1, z_2)$ on U_1 such that $M \cap U_1 = \{z_2 = 0\}$, where z_1 are any complementary coordinates, with $\dim z_1 = \bar{n}, \dim z_2 = n - \bar{n}$ and $\bar{n} = \dim M$. In the local z -coordinates, the DAE Ξ has the following form

$$E(x) \left(\frac{\partial \psi(x)}{\partial x} \right)^{-1} \left(\frac{\partial \psi(x)}{\partial x} \right) \dot{x} = F(x) \Rightarrow \begin{bmatrix} \tilde{E}_1(z) & \tilde{E}_2(z) \end{bmatrix} \begin{bmatrix} \dot{z}_1 \\ \dot{z}_2 \end{bmatrix} = \tilde{F}(z),$$

where $\tilde{E}_1 : U_1 \rightarrow \mathbb{R}^{l \times \bar{n}}, \tilde{E}_2 : U_1 \rightarrow \mathbb{R}^{l \times (n - \bar{n})}, [\tilde{E}_1 \circ \psi \quad \tilde{E}_2 \circ \psi] = E \left(\frac{\partial \psi}{\partial x} \right)^{-1}$ and $\tilde{F} \circ \psi = F$. By setting $z_2 = 0$, we consider the following DAE defined locally on M (denoted by $\Xi|_M$ and called the local M -restriction of Ξ , see Definition 2.12):

$$\Xi|_M : \tilde{E}_1(z_1, 0)\dot{z}_1 = \tilde{F}(z_1, 0).$$

Then by $\dim E(x)T_xM = \text{const.} = \bar{r}$ for all $x \in M$ around x_p , there exists a neighborhood $U_2 \subseteq U_1$ of x_p such that $\text{rank } \tilde{E}_1(z_1, 0) = \bar{r}, \forall z_1 \in M \cap U_2$. So by Dolezal’s theorem, see also [31], there exists a smooth map [30] $Q : M \cap U_2 \rightarrow GL(l, \mathbb{R})$ such that $\tilde{E}_1(z_1)$ of $Q(z_1)\tilde{E}_1(z_1, 0) = \begin{bmatrix} \tilde{E}_1(z_1) \\ 0 \end{bmatrix}$ is of full row rank \bar{r} . Rewrite $\tilde{E}_1(z_1)\dot{z}_1 = [\tilde{E}_1^1(z_1) \quad \tilde{E}_1^2(z_1)] \begin{bmatrix} \dot{z}_1^1 \\ \dot{z}_1^2 \end{bmatrix}$, where $z_1 = (z_1^1, z_1^2), \tilde{E}_1^1 : M \cap U_2 \rightarrow \mathbb{R}^{\bar{r} \times \bar{r}}$ and $\tilde{E}_1^2 : M \cap U_2 \rightarrow \mathbb{R}^{\bar{r} \times (n - \bar{r})}$ and denote $Q(z_1)\tilde{F}(z_1, 0) = \begin{bmatrix} \tilde{F}_1(z_1) \\ \tilde{F}_2(z_1) \end{bmatrix}$. Without loss of generality, we assume that $\tilde{E}_1^1(z_1)$ is invertible (if not, we permute the components of z_1 such that the first \bar{r} columns of $\tilde{E}_1(z_1)$ are independent). Now by the assumption that $F(x) \in E(x)T_xM$ for all $x \in M$ around x_p , there exists a neighborhood $U_3 \subseteq U_2$ such that $\tilde{F}(z) \in \tilde{E}(z)T_zM$ for all $z \in M \cap U_3$, i.e.,

$$\begin{bmatrix} \tilde{F}_1(z_1) \\ \tilde{F}_2(z_1) \end{bmatrix} \in \text{Im} \begin{bmatrix} \tilde{E}_1^1(z_1) & \tilde{E}_1^2(z_1) \\ 0 & 0 \end{bmatrix}.$$

It follows that $\tilde{F}_2(z_1) \equiv 0$ for all $z_1 \in M \cap U_3$. Then consider the following DAE (which is actually a reduction of $\Xi|_M$, denoted by Ξ_M^{red} , see Definition 2.13)

$$\Xi_M^{\text{red}} : [\tilde{E}_1^1(z_1) \quad \tilde{E}_1^2(z_1)] \begin{bmatrix} \dot{z}_1^1 \\ \dot{z}_1^2 \end{bmatrix} = \tilde{F}_1(z_1). \tag{29}$$

Note that a \mathcal{C}^1 -curve $z_1 : I \rightarrow M \cap U_3$ is a solution of (29) passing through $z_{10} = (z_{10}^1, z_{10}^2)$ if and only if $x(\cdot) = \psi^{-1}(z_1(\cdot), 0)$ is a solution of Ξ passing through $x_0 = \psi^{-1}(z_{10}, 0)$. Observe

that for any initial point $z_{10} \in M \cap U_3$, there always exists a solution $z_1(\cdot)$ of (29) such that $z_1(t_0) = z_{10}$ for a certain $t_0 \in I$ and $z_1(t) \in M \cap U_3, \forall t \in I$. Indeed, rewrite DAE (29) as the following ODE (recall that $\bar{E}_1^1(z_1)$ is invertible):

$$\dot{z}_1^1 = \left(\bar{E}_1^1(z_1)\right)^{-1} \left(\bar{F}_1(z_1) - \bar{E}_1^2(z_1)\dot{z}_1^2\right). \tag{30}$$

It is always possible to parameterize solutions $z_1(\cdot) = (z_1^1(\cdot), z_1^2(\cdot))$ of (30) as follows. Denote $\dot{z}_1^2 = v, f(z_1) = (E_1^1)^{-1}\bar{F}_1(z_1)$ and $g(z_1) = (E_1^1)^{-1}E_1^2(z_1)$, then (30) can be expressed as

$$\begin{cases} \dot{z}_1^1 = f(z_1) + g(z_1)v, \\ \dot{z}_1^2 = v, \end{cases} \tag{31}$$

(called a (Q, v) -explicitation of (29), see Definition 3.1), and for any solution $(z_1(\cdot), v(\cdot))$ of (31), with $v \in C^0$, the curve $z_1(\cdot)$ is a C^1 -solution of (29) satisfying $z_1(t_0) = z_{10}$ (see Proposition 3.5). It follows that for any point $x_0 = \psi^{-1}(z_{10}, 0) \in M \cap U_3$, there always exists a solution $x(\cdot) = \psi^{-1}(z_1(\cdot), 0)$ of Ξ such that $x(t_0) = x_0$ for a certain $t_0 \in I$ and that $x(t) \in M \cap U_3$ for all $t \in I$, so M is a locally invariant submanifold of Ξ around x_p by definition. \square

Proof of Theorem 2.20. Since M^* is locally invariant around x_p , via a similar construction to that shown in the proof of Proposition 2.3, we can get a DAE $\Xi|_{M^*}^{red}$ of the form (29) (if the maximal invariant submanifold M^* is constructed via the algorithm in section 2.2, then $\Xi|_{M^*}^{red}$ coincides with the DAE Ξ^* of (6) from the results of that algorithm). Note that $\Xi|_{M^*}^{red}$ can be seen as an ODE possibly with free variables (see (30) and (31), where z_1^2 are free variables), and that $\Xi|_{M^*}^{red}$ has isomorphic solutions with Ξ (see Proposition 2.8). Thus Ξ is internally regular around x_p , i.e., there exists only one maximal solution passing through any $x_0 \in M^*$ around x_p if and only if no free variables are present in $\Xi^* = \Xi|_{M^*}^{red}$, i.e., $[\bar{E}_1^1, \bar{E}_1^2]$ of (29) is invertible or, equivalently, $n^* = \dim M^* = \dim E(x)T_x M^* = r^*$ for all $x \in M^*$ around x_p (i.e., E^* of (6) is invertible). Moreover, it is clear that $[\bar{E}_1^1, \bar{E}_1^2]$ is invertible if and only if $\Xi|_{M^*}^{red}$ of (29) (or Ξ^* , given by (6)) is ex-equivalent to an ODE (10) without free variables, where $f^* = [\bar{E}_1^1, \bar{E}_1^2]^{-1}\bar{F}_1$ (or $f^* = (E^*)^{-1}F^*$), that is, Ξ is internally equivalent to (10) around x_p . \square

Appendix B. Proofs of Proposition 3.2, Proposition 3.5, Theorem 3.6, Proposition 3.8 and Theorem 3.13

Proof of Proposition 3.2. *If.* Throughout the proof below, we may drop the argument x for the maps $f(x), g(x), h(x), \dots$, for ease of notation. Suppose that Σ and $\tilde{\Sigma}$ are equivalent via transformations given by (17). First, $\text{Im } \tilde{g} = \text{Im } g\beta = \ker E_1 = \ker E$ implies that \tilde{g} is another choice such that $\text{Im } \tilde{g} = \ker E$. Moreover, we have

$$\tilde{\Sigma} : \begin{cases} \dot{x} = \tilde{f} + \tilde{g}\tilde{v} = f + g\alpha + \gamma h + g\beta v = E_1^\dagger F_1 + g\alpha + \gamma F_2 + g\beta v, \\ \tilde{y} = \tilde{h} = \eta h. \end{cases}$$

Pre-multiplying the differential part $\dot{x} = E_1^\dagger F_1 + g\alpha + \gamma F_2 + g\beta v$ of $\tilde{\Sigma}$ by E_1 , we get (note that $\text{Im } g = \ker E_1$)

$$\begin{cases} E_1 \dot{x} = F_1 + E_1 \gamma F_2, \\ \tilde{y} = \eta h. \end{cases}$$

Thus $\tilde{\Sigma}$ is an (I, \tilde{v}) -explicitation of the following DAE:

$$\begin{bmatrix} E_1 \\ 0 \end{bmatrix} \dot{x} = \begin{bmatrix} F_1 + E_1 \gamma F_2 \\ \eta F_2 \end{bmatrix}.$$

Since the above DAE can be obtained from Ξ via $\tilde{Q} = Q'Q$, where $Q' = \begin{bmatrix} I_q & E_1 \gamma \\ 0 & \eta \end{bmatrix}$, it proves that $\tilde{\Sigma}$ is a (\tilde{Q}, \tilde{v}) -explicitation of Ξ corresponding to the choice of invertible matrix $\tilde{Q} = Q'Q$. Finally, by $E_1 \tilde{f} = F_1 + E_1 \gamma F_2$, we get $\tilde{f} = \tilde{E}_1^\dagger (F_1 + \gamma F_2)$ for the above choice of right inverse \tilde{E}_1^\dagger of E_1 .

Only if. Suppose that $\tilde{\Sigma} \in \mathbf{Expl}(\Xi)$ via \tilde{Q} , \tilde{E}_1^\dagger and \tilde{g} . First, by $\text{Im } \tilde{g} = \ker E = \text{Im } g$, there exists an invertible matrix β such that $\tilde{g} = g\beta$. Moreover, since E_1^\dagger is a right inverse of E_1 if and only if any solution \dot{x} of $E_1 \dot{x} = w$ is given by $E_1^\dagger w$, we have $E_1 E_1^\dagger F_1 = F_1$ and $E_1 \tilde{E}_1^\dagger F_1 = F_1$. It follows that $E_1 (\tilde{E}_1^\dagger - E_1^\dagger) F_1 = 0$, so $(\tilde{E}_1^\dagger - E_1^\dagger) F_1 \in \ker E_1$. Since $\ker E_1 = \text{Im } g$, it follows that $(\tilde{E}_1^\dagger - E_1^\dagger) F_1 = g\alpha$ for a suitable α . Furthermore, since Q is such that E_1 of $QE = \begin{bmatrix} E_1 \\ 0 \end{bmatrix}$ is of full row rank, any other \tilde{Q} , such that \tilde{E}_1 of $\tilde{Q}E = \begin{bmatrix} \tilde{E}_1 \\ 0 \end{bmatrix}$ is of full row rank, must be of the form $\tilde{Q} = Q'Q$, where $Q' = \begin{bmatrix} Q_1 & Q_2 \\ 0 & Q_4 \end{bmatrix}$. Thus via \tilde{Q} , Ξ is ex-equivalent to

$$Q' \begin{bmatrix} E_1 \\ 0 \end{bmatrix} \dot{x} = Q' \begin{bmatrix} F_1 \\ F_2 \end{bmatrix} \Rightarrow \begin{bmatrix} Q_1 E_1 \\ 0 \end{bmatrix} \dot{x} = \begin{bmatrix} Q_1 F_1 + Q_2 F_2 \\ Q_4 F_2 \end{bmatrix}.$$

The equation on the right-hand side of the above can be expressed (using \tilde{E}_1^\dagger and \tilde{g}) as:

$$\begin{cases} \dot{x} = \tilde{E}_1^\dagger F_1 + \tilde{E}_1^\dagger Q_1^{-1} Q_2 F_2 + \tilde{g} v = E_1^\dagger F_1 + g\alpha + E_1^\dagger Q_1^{-1} Q_2 h + g\beta \tilde{v}, \\ 0 = Q_4 F_2 = Q_4 h. \end{cases}$$

Thus the explicitation of Ξ via \tilde{Q} , \tilde{E}_1^\dagger and \tilde{g} is

$$\tilde{\Sigma}: \begin{cases} \dot{x} = E_1^\dagger F_1 + g\alpha + \gamma h + g\beta \tilde{v} = f + \gamma h + g(\alpha + \beta \tilde{v}) = \tilde{f} + \tilde{g} \tilde{v}, \\ \tilde{y} = \eta h = \tilde{h}, \end{cases}$$

where $\gamma = E_1^\dagger Q_1^{-1} Q_2$, $\eta = Q_4$. Therefore, we can see that Σ and $\tilde{\Sigma}$ are equivalent via the transformations of the form (17). \square

Proof of Proposition 3.5. Consider the DAE (12) of the (Q, v) -explicitation procedure. Since Q -transformations preserve solutions of Ξ , system (12) resulting from a Q -transformation of Ξ has the same solutions as Ξ . Thus we need to prove that (12) and (15) have corresponding solutions for any choices of E_1^\dagger and g . Moreover, the second equation $0 = F_2(x)$ of (12) coincides with $0 = h(x)$ of (15). So we only need to prove that $x(t) \in \mathcal{C}^1$ is a solution of $E_1(x)\dot{x} = F_1(x)$ if and only if there exists $v(t) \in \mathcal{C}^0$ such that $(x(t), v(t))$ is a solution of $\dot{x} = f(x) + g(x)v$

independently of the choice of E_1^\dagger , defining $f(x) = E_1^\dagger(x)F_1(x)$, and of the choice of g satisfying $\text{Im } g(x) = \ker E_1(x)$.

If. Suppose that $(x(t), v(t))$ is a solution of $\dot{x} = f(x) + g(x)v$. Then we have $\dot{x}(t) = f(x(t)) + g(x(t))v(t)$. Pre-multiplying the latter equation by $E_1(x(t))$, we get that

$$E_1(x(t))\dot{x}(t) = E_1(x(t))f(x(t)) = E_1(x(t))E_1^\dagger(x(t))F_1(x(t)) = F_1(x(t)),$$

which proves that $x(t)$ is a solution of $E_1(x)\dot{x} = F_1(x)$.

Only if. Suppose that $x(t)$ is a solution of $E_1(x)\dot{x} = F_1(x)$. Rewrite $E_1(x)\dot{x}$ as $\begin{bmatrix} E_1^1(x) & E_1^2(x) \end{bmatrix} \times \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix}$, where $E_1^1 : U \rightarrow \mathbb{R}^{q \times q}$ is smooth and $x = (x_1, x_2)$. Then, by taking a smaller neighborhood U , if necessary, we assume that $E_1^1(x)$ is invertible locally around x_p (if not, we permute the components of x such that the first q columns of $E_1(x)$ are independent). Thus a choice of right inverse of E_1 is $E_1^\dagger = \begin{bmatrix} (E_1^1)^{-1} \\ 0 \end{bmatrix}$. So the maps f and g can be defined as $f := E_1^\dagger F_1 = \begin{bmatrix} (E_1^1)^{-1} F_1 \\ 0 \end{bmatrix}$, $g := \begin{bmatrix} -(E_1^1)^{-1} E_2 \\ I_m \end{bmatrix}$. Set $v(t) = \dot{x}_2(t)$, then $v \in C^0$ and it is clear that if $x(t) = ((x_1(t), x_2(t)))$ is a solution of $E_1(x)\dot{x} = F_1(x)$, then $(x(t), v(t))$ solves $\dot{x} = f(x) + g(x)v$ since

$$\begin{bmatrix} E_1^1(x(t)) & E_1^2(x(t)) \end{bmatrix} \begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} = F_1(x(t)) \Rightarrow \dot{x}_1(t) = (E_1^1)^{-1} F_1(x(t)) - (E_1^1)^{-1} E_1^2(x(t))\dot{x}_2(t).$$

Notice that if we choose another right inverse \tilde{E}_1^\dagger of E_1 and another matrix \tilde{g} such that $\text{Im } \tilde{g} = \ker E_1$, then by Proposition 3.2, we have

$$\dot{x} = \tilde{f}(x) + \tilde{g}(x)\tilde{v} \Leftrightarrow \dot{x} = f(x) + g(x)(\alpha(x) + \beta(x)v).$$

We thus conclude that there exists $\tilde{v}(t) = \alpha(x(t)) + \beta(x(t))v(t) = \alpha(x(t)) + \beta(x(t))\dot{x}_2(t)$ such that $(x(t), \tilde{v}(t))$ solves $\dot{x} = \tilde{f}(x) + \tilde{g}(x)\tilde{v}$. Therefore, Ξ has corresponding solutions with any (Q, v) -explicitation Σ independently of the choice of Q, E_1^\dagger and g . \square

Proof of Theorem 3.6. By the assumptions that $\text{rank } E(x) = \text{const.} = q$ and $\text{rank } \tilde{E}(\tilde{x}) = \text{const.} = \tilde{q}$ around x_p and \tilde{x}_p , respectively, we have that Ξ and $\tilde{\Xi}$ are locally ex-equivalent to

$$\Xi' : \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix} \dot{x} = \begin{bmatrix} F_1(x) \\ F_2(x) \end{bmatrix} \quad \text{and} \quad \tilde{\Xi}' : \begin{bmatrix} \tilde{E}_1(\tilde{x}) \\ 0 \end{bmatrix} \dot{\tilde{x}} = \begin{bmatrix} \tilde{F}_1(\tilde{x}) \\ \tilde{F}_2(\tilde{x}) \end{bmatrix},$$

respectively, where $E_1(x)$ and $\tilde{E}_1(\tilde{x})$ are full row rank matrices and their ranks are q and \tilde{q} , respectively. By Definition 3.1, we have

$$\begin{aligned} f(x) &= E_1^\dagger(x)F_1(x), \quad \text{Im } g(x) = \ker E_1(x), \quad h(x) = F_2(x), \\ \tilde{f}(\tilde{x}) &= \tilde{E}_1^\dagger(\tilde{x})\tilde{F}_1(\tilde{x}), \quad \text{Im } \tilde{g}(\tilde{x}) = \ker \tilde{E}_1(\tilde{x}), \quad \tilde{h}(\tilde{x}) = \tilde{F}_2(\tilde{x}). \end{aligned} \tag{32}$$

Note that explicitation systems are defined up to a feedback, an output multiplication and a generalized output injection. Any two control systems belonging to $\mathbf{Expl}(\Xi)$ are sys-equivalent to each other and so are any two control systems belonging to $\mathbf{Expl}(\tilde{\Xi})$. Thus the choice of an explicitation system makes no difference for the proof of sys-equivalence. Without loss of

generality, we will use $f(x)$, $g(x)$, $h(x)$ and $\tilde{f}(x)$, $\tilde{g}(x)$, $\tilde{h}(x)$ given in (32) for the remaining part of this proof.

If. Suppose $\Sigma \overset{sys}{\sim} \tilde{\Sigma}$ in a neighborhood U of x_p . By Definition 3.3, there exists a diffeomorphism $\tilde{x} = \psi(x)$ and $\beta : U \rightarrow GL(m, \mathbb{R})$ such that $\tilde{g} \circ \psi = \frac{\partial \psi}{\partial x} g \beta$, which implies

$$\ker(\tilde{E} \circ \psi) = \text{span}\{\tilde{g}_1, \dots, \tilde{g}_m\} \circ \psi = \text{span}\left\{ \frac{\partial \psi}{\partial x} g_1, \dots, \frac{\partial \psi}{\partial x} g_m \right\} = \frac{\partial \psi}{\partial x} \ker E$$

and $q = \tilde{q}$ (since $\dim \ker \tilde{E} = \tilde{m} = m = \dim \tilde{E}$). We can deduce from the above equation that there exists $Q_1 : U \rightarrow GL(q, \mathbb{R})$ such that

$$\tilde{E}_1 \circ \psi = Q_1 E_1 \left(\frac{\partial \psi}{\partial x} \right)^{-1}. \tag{33}$$

Subsequently, by $\tilde{f} \circ \psi = \frac{\partial \psi}{\partial x} (f + \gamma h + g\alpha)$ of Definition 3.3, we have

$$(\tilde{E}_1^\dagger \circ \psi)(\tilde{F}_1 \circ \psi) = \frac{\partial \psi}{\partial x} (E_1^\dagger F_1 + \gamma F_2 + g\alpha).$$

Pre-multiply the above equation by $\tilde{E}_1 \circ \psi = Q_1 E_1 \left(\frac{\partial \psi}{\partial x} \right)^{-1}$, to obtain

$$\tilde{F}_1 \circ \psi = Q_1 F_1 + Q_1 E_1 \gamma F_2. \tag{34}$$

Then by $\tilde{h} \circ \psi = \eta h$ of Definition 3.3, we immediately get

$$\tilde{F}_2 \circ \psi = \eta F_2. \tag{35}$$

Now combining (33), (34) and (35), we conclude that Ξ' and $\tilde{\Xi}'$ are ex-equivalent via $\tilde{x} = \psi(x)$ and $Q = \begin{bmatrix} Q_1 & Q_1 E_1 \gamma \\ 0 & \eta \end{bmatrix}$, which implies that $\Xi \overset{ex}{\sim} \tilde{\Xi}$ (since $\Xi \overset{ex}{\sim} \Xi'$ and $\tilde{\Xi} \overset{ex}{\sim} \tilde{\Xi}'$).

Only if. Suppose that locally $\Xi \overset{ex}{\sim} \tilde{\Xi}$ around x_p . It follows that locally $\Xi' \overset{ex}{\sim} \tilde{\Xi}'$ around x_p , which implies that $q = \tilde{q}$. Assume that they are ex-equivalent via $Q : U \rightarrow GL(l, \mathbb{R})$ and $\tilde{x} = \psi(x)$ defined on a neighborhood U of x_p . Let $Q = \begin{bmatrix} Q_1 & Q_2 \\ Q_3 & Q_4 \end{bmatrix}$, where Q_1, Q_2, Q_3 and Q_4 are matrix-valued functions of sizes $q \times q, q \times m, p \times q$ and $p \times p$, respectively. Then by $\begin{bmatrix} Q_1 & Q_2 \\ Q_3 & Q_4 \end{bmatrix} \begin{bmatrix} E_1 \\ 0 \end{bmatrix} = \begin{bmatrix} \tilde{E}_1 \circ \psi \\ 0 \end{bmatrix} \frac{\partial \psi}{\partial x}$, we can deduce that $Q_3 = 0$ and Q_1, Q_4 are invertible matrices. Then we have

$$\begin{bmatrix} Q_1 & Q_2 \\ 0 & Q_4 \end{bmatrix} \begin{bmatrix} E_1 \\ 0 \end{bmatrix} = \begin{bmatrix} \tilde{E}_1 \circ \psi \\ 0 \end{bmatrix} \frac{\partial \psi}{\partial x}, \quad \begin{bmatrix} Q_1 & Q_2 \\ 0 & Q_4 \end{bmatrix} \begin{bmatrix} F_1 \\ F_2 \end{bmatrix} = \begin{bmatrix} \tilde{F}_1 \circ \psi \\ \tilde{F}_2 \circ \psi \end{bmatrix},$$

which implies

$$\tilde{E}_1 \circ \psi = Q_1 E_1 \left(\frac{\partial \psi}{\partial x} \right)^{-1}, \quad \tilde{F}_1 \circ \psi = Q_1 F_1 + Q_2 F_2, \quad \tilde{F}_2 \circ \psi = Q_4 F_2. \tag{36}$$

Thus by $\text{Im } g(x) = \ker E(x) = \ker E_1(x)$ and $\text{Im } \tilde{g}(x) = \ker \tilde{E}(\tilde{x}) = \ker \tilde{E}_1(\tilde{x})$, and using (36), we have

$$\tilde{g} \circ \psi = \frac{\partial \psi}{\partial x} g \beta \tag{37}$$

for some $\beta : U \rightarrow GL(m, \mathbb{R})$. Moreover, there exists $\alpha : U \rightarrow \mathbb{R}^m$ such that

$$\begin{aligned} \tilde{f} \circ \psi &= \tilde{E}_1^\dagger \circ \psi \tilde{F}_1 \circ \psi \stackrel{(36)}{=} \frac{\partial \psi}{\partial x} E_1^\dagger Q_1^{-1} Q_1 F_1 + Q_2 F_2 \\ &= \frac{\partial \psi}{\partial x} E_1^\dagger Q_1^{-1} (Q_1 F_1 + Q_2 F_2 + Q_1 E_1 g \alpha) \\ &= \frac{\partial \psi}{\partial x} (f + E_1^\dagger Q_1^{-1} Q_2 y + g \alpha). \end{aligned} \tag{38}$$

In addition, we have

$$\tilde{h} \circ \psi = \tilde{F}_2 \circ \psi \stackrel{(36)}{=} Q_4 F_2 = Q_4 h. \tag{39}$$

Finally, it can be seen from (37), (38), and (39) that $\Sigma \stackrel{\text{sys}}{\approx} \tilde{\Sigma}$ via $\tilde{x} = \psi(x)$, $\alpha, \beta, \gamma = E_1^\dagger Q_1^{-1} Q_2$ and $\eta = Q_4$. \square

Proof of Proposition 3.8. We first show that the sequence of submanifolds M_k^c of the geometric reduction method of the DAE Ξ and the sequence N_k^c of the zero dynamics algorithm of any control system $\Sigma = (f, g, h) \in \mathbf{Expl}(\Xi)$ locally coincide. Suppose that $\text{rank } E(x) = \text{const.} = q$ in a neighborhood U_1 of x_p . Then there always exists an invertible matrix $Q(x)$ defined on U_1 such that $E_1(x)$ of $Q(x)E(x) = \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix}$ is of full row rank q for all $x \in U_1$, denote $Q(x)F(x) = \begin{bmatrix} F_1(x) \\ F_2(x) \end{bmatrix}$. Recall, see Remark 3.7, that N_k of the zero dynamics algorithm are well-defined for any $\Sigma \in \mathbf{Expl}(\Xi)$ and that N_k are the same for all control systems belonging to $\mathbf{Expl}(\Xi)$. So the choice of an excitation system makes no difference for N_k . We may choose a control system $\Sigma = (f, g, h) \in \mathbf{Expl}(\Xi)$, given by $f(x) = E_1^\dagger(x)F_1(x)$, $\text{Im } g(x) = \ker E(x)$, $h(x) = F_2(x)$. By the definition of M_1 (see (4)) and $N_1 = h^{-1}(0)$, we have

$$\begin{aligned} M_1^c &= M_1 \cap U_1 = \{x \in U_1 : Q(x)F(x) \in \text{Im } Q(x)E(x)\} = \left\{x \in U_1 : \begin{pmatrix} F_1(x) \\ F_2(x) \end{pmatrix} \in \text{Im} \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix}\right\} \\ &= \{x \in U_1 : F_2(x) = 0\} = \{x \in U_1 : h(x) = 0\} = N_1 \cap U_1 = N_1^c. \end{aligned}$$

For $k > 1$, suppose $M_{k-1}^c = N_{k-1}^c$. Then by (4) and (18), we have

$$\begin{aligned} M_k &= \{x \in M_{k-1}^c : Q(x)F(x) \in Q(x)E(x)T_x M_{k-1}^c\} = \left\{x \in M_{k-1}^c : \begin{pmatrix} F_1(x) \\ F_2(x) \end{pmatrix} \in \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix} T_x M_{k-1}^c\right\} \\ &= \{x \in M_{k-1}^c : F_1(x) \in E_1(x)T_x M_{k-1}^c\} \\ &= \{x \in M_{k-1}^c : f(x) + \ker E_1(x) \subseteq T_x M_{k-1}^c + \ker E_1(x)\} \\ &= \{x \in N_{k-1}^c : f(x) \in T_x N_{k-1}^c + \mathcal{G}(x)\} = N_k, \end{aligned}$$

and thus $M_k^c = N_k^c$. If either one among (A1) and (A2) is satisfied, then by $N_k^c = M_k^c$, we can easily deduce the other one and thus (A1) and (A2) are equivalent. Then by Proposition 2.7, $M^* = M_{k^*}^c$ is a locally maximal invariant submanifold and by Proposition 6.1.1 of [27], $N^* = N_{k^*}^c$ is a local maximal output zeroing submanifold. Moreover, we have locally $M^* = N^*$ (since locally $M_k^c = N_k^c$). Now under the assumption that $\dim E(x)T_x M^* = \text{const.}$ for all $x \in M^*$ around x_p , by Theorem 2.20, Ξ is internally regular if and only if $\dim M^* = \dim E(x)T_x M^*$, i.e., $\ker E(x) \cap T_x M^* = 0$, locally $\forall x \in M^*$ around x_p . Thus by $N^* = M^*$ and $\ker E(x) = \mathcal{G}(x)$, we have that Ξ is internally regular (around x_p) if and only if $\mathcal{G}(x_p) \cap T_{x_p} N^* = 0$. \square

Proof of Theorem 3.13. (i) \Rightarrow (ii): Suppose in a neighborhood U of x_p that $\text{rank } E(x) = q$ and $\mathcal{G}(x) = \ker E(x) = \text{span}\{g_1(x), \dots, g_m(x)\}$ is involutive, where g_1, \dots, g_m are independent vector fields on U and $m = n - q$. Then by the involutivity of \mathcal{G} , there exist local coordinates $\tilde{x} = (\tilde{x}_1, \tilde{x}_2) = \psi(x)$, where $\tilde{x}_1 = (\tilde{x}_1^1, \dots, \tilde{x}_1^q)$ and $\tilde{x}_2 = (\tilde{x}_2^1, \dots, \tilde{x}_2^{n-q})$, such that $\text{span}\{d\tilde{x}_1^1, \dots, d\tilde{x}_1^q\} = \text{span}\{d\tilde{x}_1\} = \mathcal{G}^\perp$ (Frobenius theorem [3]), where \mathcal{G}^\perp denotes the co-distribution which annihilates \mathcal{G} . Note that in the \tilde{x} -coordinates, the distribution

$$\ker \tilde{E}(\tilde{x}) = \ker \left(E(x) \left(\frac{\partial \psi(x)}{\partial x} \right)^{-1} \right) = \frac{\partial \psi(x)}{\partial x} \mathcal{G}(x) = \text{span}\{\tilde{g}_1(\tilde{x}), \dots, \tilde{g}_m(\tilde{x})\},$$

where $\tilde{g}_i \circ \psi = \frac{\partial \psi}{\partial x} g_i$, $i = 1, \dots, m$. Now let \tilde{g} be a matrix whose columns consist of \tilde{g}_i , for $i = 1, \dots, m$. It follows that $\text{rank } \tilde{g}(\tilde{x}) = m$ around $\tilde{x}_0 = \psi(x_0)$. By $\text{span}\{d\tilde{x}_1\} = \mathcal{G}^\perp$, we have $\langle d\tilde{x}_1, \tilde{g}_i \rangle = 0$, for $i = 1, \dots, m$. Thus $\tilde{g}(\tilde{x})$ is of the form $\tilde{g}(\tilde{x}) = \begin{bmatrix} 0 \\ \tilde{g}_2(\tilde{x}) \end{bmatrix}$, where $\tilde{g}_2 : \psi(U) \rightarrow \mathbb{R}^{m \times m}$. Since $\text{rank } \tilde{g}(\tilde{x}) = m$, it can be seen that $\tilde{g}_2(\tilde{x})$ is an invertible matrix, which implies by $\text{Im } \tilde{g}(\tilde{x}) = \ker \tilde{E}(\tilde{x})$ that $\tilde{E}(\tilde{x})$ has to be of the form $\tilde{E}(\tilde{x}) = [\tilde{E}_1(\tilde{x}) \ 0]$, where $\tilde{E}_1 : \psi(U) \rightarrow \mathbb{R}^{l \times m}$. Thus in the \tilde{x} -coordinates, $\tilde{\Xi} = (\tilde{E}, \tilde{F})$ admits the following form:

$$[\tilde{E}_1(\tilde{x}) \ 0] \begin{bmatrix} \dot{\tilde{x}}_1 \\ \dot{\tilde{x}}_2 \end{bmatrix} = \tilde{F}(\tilde{x}),$$

where $\tilde{F} \circ \psi = F$. Now by $\text{rank } E(x) = q$, we get $\text{rank } [\tilde{E}_1(\tilde{x}) \ 0] = \text{rank } E(x) = q$ (the coordinate transformation preserves the rank). Thus by Dolezal’s theorem [30], see also [31], there exists a smooth map $Q : \psi(U) \rightarrow GL(l, \mathbb{R})$ such that $Q(\tilde{x})\tilde{E}(\tilde{x}) = Q(\tilde{x})[\tilde{E}_1(\tilde{x}) \ 0] = \begin{bmatrix} \tilde{E}_1^1(\tilde{x}) & 0 \\ 0 & 0 \end{bmatrix}$, where $\tilde{E}_1^1 : \psi(U) \rightarrow \mathbb{R}^{q \times q}$. Since $Q(\tilde{x})$ preserves the rank of $\tilde{E}(\tilde{x})$, we have $\text{rank } \tilde{E}_1^1(\tilde{x}) = q$. Therefore, $\tilde{E}_1^1(\tilde{x})$ is an invertible matrix. Now let $Q'(\tilde{x}) = \begin{bmatrix} (\tilde{E}_1^1(\tilde{x}))^{-1} & 0 \\ 0 & I_m \end{bmatrix} Q(\tilde{x})$ and denote $Q'(\tilde{x})\tilde{F}(\tilde{x}) = \begin{bmatrix} F_1(\tilde{x}) \\ F_2(\tilde{x}) \end{bmatrix}$. It is seen that, via $\tilde{x} = \psi(x)$ and Q' , the system Ξ is locally ex-equivalent to $\tilde{\Xi} = (Q'\tilde{E}, Q'\tilde{F})$, where $Q'\tilde{E} \circ \psi = Q'E \left(\frac{\partial \psi}{\partial x} \right)^{-1} = \begin{bmatrix} I_q & 0 \\ 0 & 0 \end{bmatrix}$. Clearly, $\tilde{\Xi}$ is a semi-explicit DAE.

(ii) \Rightarrow (iii): Suppose that Ξ is locally ex-equivalent to Ξ^{SE} of the form (2) around x_p . Then, any control system $\Sigma \in \mathbf{Expl}(\Xi)$ is locally sys-equivalent to $\Sigma' \in \mathbf{Expl}(\Xi^{SE})$ below (by Theorem 2.20):

$$\Sigma' : \begin{cases} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} F_1(x_1, x_2) \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ I_m \end{bmatrix} v, \\ y = F_2(x_1, x_2). \end{cases}$$

Suppose that $\Sigma \stackrel{\text{sys}}{\sim} \Sigma'$ via $z = (z_1, z_2) = \psi(x)$, α, β and $\gamma = \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix}$, then

$$\Sigma : \begin{cases} \begin{bmatrix} \dot{z}_1 \\ \dot{z}_2 \end{bmatrix} = \frac{\partial \psi(x)}{\partial x} \left(\begin{bmatrix} F_1(x) \\ 0 \end{bmatrix} + \begin{bmatrix} \gamma_1(x) \\ \gamma_2(x) \end{bmatrix} y + \begin{bmatrix} 0 \\ I_m \end{bmatrix} (\alpha(x) + \beta(x)v) \right), \\ \tilde{y} = \eta(x)F_2(x), \end{cases}$$

where $x = \psi^{-1}(z)$. By Definition 3.12, Σ can always be fully reduced to (by a coordinates change and a feedback transformation)

$$\begin{cases} \dot{x}_1 = F_1(x_1, x_2) + \gamma_1(x_1, x_2)F_2(x_1, x_2), \\ y = \eta(x_1, x_2)F_2(x_1, x_2), \end{cases}$$

where x_2 is the new control.

(iii) \Rightarrow (i): Suppose (iii) holds. Then $\mathbf{Expl}(\Xi)$ is not empty implies that $E(x)$ has constant rank around x_p . By Definition 3.12, if a control system $\Sigma \in \mathbf{Expl}(\Xi)$ can be fully reduced, then $\mathcal{G} = \ker E(x) = \text{span} \{g_1, \dots, g_m\}$ is involutive. \square

Appendix C. Proof of Theorem 4.1

Claim. *If assumptions (A1)-(A3) of Theorem 4.1 are satisfied, then the point x_p is a regular point of the zero dynamics algorithm (rank conditions (i), (ii), (iii) of Proposition 6.1.3 of [27] are satisfied) for any control system $\Sigma \in \mathbf{Expl}(\Xi)$. If so, we use Proposition 6.1.5 of [27] with a small modification: there exist local coordinates $(z, z^*) = (z_1, \dots, z_m, z^*)$, where $z_i = (z_i^1, \dots, z_i^{\rho_i})$, such that Σ is the following form*

$$\left\{ \begin{array}{ll} \begin{array}{l} y_1 = z_1^1 \\ \dot{z}_1^1 = z_1^2 + \sigma_1^1 v \\ \dots \\ \dot{z}_1^{\rho_1-1} = z_1^{\rho_1} + \sigma_1^{\rho_1-1} v \\ \dot{z}_1^{\rho_1} = \alpha_1 + \beta_1 v \end{array} & \begin{array}{l} y_i = z_i^1, \quad \text{for } 3 \leq i \leq m \\ \dot{z}_i^1 = z_i^2 + \sum_{s=1}^{i-1} \delta_{i,s}^1 (\alpha_s + \beta_s v) + \sigma_i^1 v \\ \dots \\ \dot{z}_i^{\rho_i-1} = z_i^{\rho_i} + \sum_{s=1}^{i-1} \delta_{i,s}^{\rho_i-1} (\alpha_s + \beta_s v) + \sigma_i^{\rho_i-1} v \\ \dot{z}_i^{\rho_i} = \alpha_i + \beta_i v \end{array} \\ \begin{array}{l} y_2 = z_2^1 \\ \dot{z}_2^1 = z_2^2 + \delta_{2,1}^1 (\alpha_1 + \beta_1 v) + \sigma_2^1 v \\ \dots \\ \dot{z}_2^{\rho_2-1} = z_2^{\rho_2} + \delta_{2,1}^{\rho_2-1} (\alpha_1 + \beta_1 v) + \sigma_2^{\rho_2-1} v \\ \dot{z}_2^{\rho_2} = \alpha_2 + \beta_2 v \end{array} & \begin{array}{l} z^* = f^*(z, z^*) + g^*(z, z^*)v, \end{array} \end{array} \right. \tag{40}$$

where $\delta_{i,s}^j \equiv 0$ for $1 \leq j < \rho_s, 1 \leq s \leq i - 1$.

- (i) Note that in (40), $\rho_1 \leq \rho_2 \leq \dots \leq \rho_m$ and the matrix $\beta = (\beta_1, \dots, \beta_m)$ is invertible at x_p . The functions σ_k satisfy $\sigma^k|_{N_k} = 0$ for $k = 1, \dots, \rho_i - 1$, where

$$N_k = \{(z, z^*) : z_i^j = 0, 1 \leq i \leq m, 1 \leq j \leq k\}.$$

- (ii) There are two differences between system (40) and the zero dynamics form of Proposition 6.1.3 of [27], where the functions $\sigma_1^1, \dots, \sigma_1^{\rho_1-1}$ are not present and all the functions $\delta_{i,s}^j$ can be nonzero. However, in (40), $\sigma_1^1, \dots, \sigma_1^{\rho_1-1}$ vanish on N_1, \dots, N_{ρ_1-1} , respectively, but may not outside, and $\delta_{i,s}^j \equiv 0$ for $1 \leq j < \rho_s, 1 \leq s \leq i - 1$.

Proof of the Claim. We will prove that assumptions (A1), (A2), (A3) of Theorem 4.1 correspond to the rank conditions (i), (ii), (iii) of Proposition 6.1.3 in [27]. By the assumption of Theorem 4.1 that $\text{rank } E(x) = \text{const.}$ around x_p , we have $\mathbf{Expl}(\Xi)$ is not empty. Now, in order to compare the two algorithms (the geometric reduction algorithm in section 2.2 for Ξ and the zero dynamics algorithm in [27] for $\Sigma \in \mathbf{Expl}(\Xi)$), we use the same notations as in the algorithm of section 2.2.

Then for a control system $\Sigma = (f, g, h) \in \mathbf{Expl}(\Xi)$, we have $f(x) = (\tilde{E}_1^1)^\dagger \tilde{F}_1^1(x)$, $\text{Im } g(x) = \ker E(x) = \ker \tilde{E}_1^1(x)$, $h(x) = \tilde{F}_1^2(x)$. The zero dynamics algorithm for Σ can be implemented in the following way:

Step 1: by (A2) of Theorem 4.1, we get $Dh(x) = D\tilde{F}_1^2(x)$ has constant rank $n - n_1$ around x_p (condition (i) of Proposition 6.1.3 in [27]). Thus $h^{-1}(0)$ can be locally expressed as $N_1^c = \{x : H_1(x) = 0\}$, where $H_1 = \psi_1(x) = (\psi_1^1, \dots, \psi_1^{n-n_1})$.

Step k ($k > 1$): By the proof of Proposition 3.8, we have $N_{k-1}^c = M_{k-1}^c$, which is

$$N_{k-1}^c = M_{k-1}^c = \{x : H_{k-1}(x) = 0\},$$

where $H_{k-1} = (\psi_0, \dots, \psi_{k-1})$. By the zero dynamic algorithms, N_k consists of all $x \in N_{k-1}^c$ such that

$$L_f H_{k-1}(x) + L_g H_{k-1}(x)u = 0.$$

Then by assumption (A1) of Theorem 4.1, we can deduce that

$$\dim(\ker E \cap \ker dH_{k-1})(x) = \dim(\text{span}\{g_1, \dots, g_m\} \cap \ker dH_{k-1})(x) = \text{const.}, \quad (41)$$

for all $x \in M_{k-1}^c$ around x_p . Now by $\dim \ker E(x) = \text{const.}$ around x_p (implied by $\text{rank } E(x) = \text{const.}$), we get

$$\dim \text{span}\{g_1, \dots, g_m\}(x) = \text{const.} \quad (42)$$

locally around x_p . By (41) and (42), we get $\text{rank } L_g H_{k-1}(x) = \text{const.}$ for all $x \in M_{k-1}^c$ around x_p (condition (ii) of Proposition 6.1.3 in [27]).

Since $\text{rank } L_g H_{k-1}(x) = \text{const.}$, there exists a basis matrix $R_{k-1}(x)$ of the annihilator of the image of $L_g H_{k-1}(x)$, that is $R_{k-1}(x)L_g H_{k-1}(x) = 0$. Thus N_k^c can be defined by

$$N_k^c = \{x \in U_k : H_{k-1}(x) = 0, R_{k-1}(x)L_f H_{k-1}(x) = 0\}.$$

Notice that by the geometric reduction algorithm, we have

$$M_k^c = \{x \in U_k : H_{k-1}(x) = 0, \tilde{F}_k^2(x) = 0\}.$$

By $N_k^c = M_k^c$ and the fact that the rank of the differential of $(H_{k-1}(x), \tilde{F}_k^2(x))$ is constant around x_p (assumption (A2) of Theorem 4.1), it follows that the rank of the differential of $\begin{bmatrix} H_{k-1}(x) \\ R_{k-1}(x)L_f H_{k-1}(x) \end{bmatrix}$ is constant around x_p (condition (i) of Proposition 6.1.3 in [27]).

Assumption (A3) of Theorem 4.1 that $\dim E(x)T_x M^* = \dim M^*$, locally around x_p , implies

$$\text{span} \{g_1(x_p), \dots, g_m(x_p)\} \cap T_{x_p} N^* = 0.$$

Finally, by $N^* = \{x : H_{k^*}(x) = 0\}$, it follows that the matrix $L_g H_{k^*}(x_p)$ has rank m (condition (iii) of Proposition 6.1.3 in [27]). \square

Proof of Theorem 4.1. Observe that by assumption (A3) and Theorem 2.20(iii), we have that Ξ is internally regular. Then by Claim, we have that x_p is a regular point of the zero dynamics algorithm for any control system $\Sigma \in \text{Expl}(\Xi)$. Thus there exist local coordinates (z, z^*) such that Σ is in the form (40) around x_p . Notice that the matrix $\beta = (\beta_1, \dots, \beta_m)$ is invertible at x_p and the functions $\sigma_i^k|_{N_k^c} = 0$, for $1 \leq i \leq m, 1 \leq k \leq \rho_i - 1$, which implies $\sigma_i^k \in \mathbf{I}^k$, where \mathbf{I}^k is the ideal generated by $z_i^j, 1 \leq i \leq m, 1 \leq j \leq k$ in the ring of smooth functions of z_b^a and z_c^* . Then for system (40), using the feedback transformation $\tilde{v} = \alpha + \beta v$, where $\alpha = (\alpha_1, \dots, \alpha_m)$, we get

$$\left\{ \begin{array}{l} y_i = z_i^1, \quad i = 1, \dots, m, \\ \dot{z}_i^1 = z_i^2 + \sum_{s=1}^{i-1} \delta_{i,s}^1 \tilde{v}_s + a_i^1 + b_i^1 \tilde{v}, \\ \dots \\ \dot{z}_i^{\rho_i-1} = z_i^{\rho_i} + \sum_{s=1}^{i-1} \delta_{i,s}^{\rho_i-1} \tilde{v}_s + a_i^{\rho_i-1} + b_i^{\rho_i-1} \tilde{v}, \\ \dot{z}_i^{\rho_i} = \tilde{v}_i, \\ \dot{z}^* = \tilde{f}^*(z, z^*) + \tilde{G}^*(z, z^*) \tilde{v}, \end{array} \right. \tag{43}$$

where $\tilde{f}^* = f^* - \bar{g}\beta^{-1}\alpha, \tilde{G}^* = g^*\beta^{-1}$, and where $a_i^k = -\sigma_i^k\beta^{-1}\alpha, b_i^k = \sigma_i^k\beta^{-1}$, for $1 \leq i \leq m, 1 \leq k \leq \rho_i - 1$ and by $\sigma_i^k \in \mathbf{I}^k$, we have $a_i^k, b_i^k \in \mathbf{I}^k$.

Recall from (40) that the functions $\delta_{i,s}^j = 0$ for $1 \leq j < \rho_s, 1 \leq s \leq i - 1$. Then if the function $\delta_{i,\bar{s}}^j \neq 0, j = \rho_{\bar{s}} + k$, for a certain $1 \leq \bar{s} \leq i - 1$ and a certain $0 \leq k \leq \rho_i - 1 - \rho_{\bar{s}}$, we show that, via suitable changes of coordinates and output multiplications, the nonzero function $\delta_{i,\bar{s}}^{k+\rho_{\bar{s}}}$ can be eliminated. Namely, define the new coordinates (and keep the remaining coordinates unchanged):

$$\tilde{z}_i^{k+1} = z_i^{k+1} - \delta_{i,\bar{s}}^{\rho_{\bar{s}}+k} z_{\bar{s}}^1, \quad \tilde{z}_i^{k+2} = z_i^{k+2} - \delta_{i,\bar{s}}^{\rho_{\bar{s}}+k} z_{\bar{s}}^2, \quad \dots, \quad \tilde{z}_i^{k+\rho_{\bar{s}}} = z_i^{k+\rho_{\bar{s}}} - \delta_{i,\bar{s}}^{\rho_{\bar{s}}+k} z_{\bar{s}}^{\rho_{\bar{s}}},$$

we have (notice that below $\delta_{\bar{s},s}^1 \equiv 0$ for $1 \leq s \leq \bar{s} - 1$)

$$\begin{aligned} \dot{z}_i^{k+1} &= z_i^{k+2} + \sum_{s=1}^{i-1} \delta_{i,s}^{k+1} \tilde{v}_s + a_i^{k+1} + b_i^{k+1} \tilde{v} - (\delta_{i,\bar{s}}^{\rho_s+k})' z_{\bar{s}}^1 - \delta_{i,s}^{\rho_s+k} (z_{\bar{s}}^2 + a_{\bar{s}}^1 + b_{\bar{s}}^1 \tilde{v} + \sum_{s=1}^{\bar{s}-1} \delta_{\bar{s},s}^1 \tilde{v}_s) \\ &= (z_i^{k+2} - \delta_{i,\bar{s}}^{\rho_s+k} z_{\bar{s}}^2) + (a_i^{k+1} - (\delta_{i,\bar{s}}^{\rho_s+k})' z_{\bar{s}}^1 - \delta_{i,s}^{\rho_s+k} a_{\bar{s}}^1) + (b_i^{k+1} - \delta_{i,s}^{\rho_s+k} b_{\bar{s}}^1) \tilde{v} + \sum_{s=1}^{i-1} \delta_{i,s}^{k+1} \tilde{v}_s \\ &= \tilde{z}_i^{k+2} + \tilde{a}_i^{k+1} + \tilde{b}_i^{k+1} \tilde{v} + \sum_{s=1}^{i-1} \delta_{i,s}^{k+1} \tilde{v}_s, \end{aligned}$$

where $(\delta_{i,\bar{s}}^{\rho_s+k})'$ denotes the derivative of $\delta_{i,\bar{s}}^{\rho_s+k}(x(t))$ with respect to t , and $\tilde{a}_i^{k+1} = a_i^{k+1} - (\delta_{i,\bar{s}}^{\rho_s+k})' z_{\bar{s}}^1 - \delta_{i,s}^{\rho_s+k} a_{\bar{s}}^1$, $\tilde{b}_i^{k+1} = b_i^{k+1} - \delta_{i,s}^{\rho_s+k} b_{\bar{s}}^1$, and it is clear that $\tilde{a}_i^{k+1}, \tilde{b}_i^{k+1} \in \mathbf{I}^{k+1}$. Then via similar calculations, we have

$$\dot{z}_i^{k+j} = \tilde{z}_i^{k+j+1} + \tilde{a}_i^{k+j} + \tilde{b}_i^{k+j} \tilde{v} + \sum_{s=1}^{i-1} \delta_{i,s}^{k+j} \tilde{v}_s, \quad 2 \leq j \leq \rho_{\bar{s}} - 1,$$

for some $\tilde{a}_i^{k+j}, \tilde{b}_i^{k+j} \in \mathbf{I}^{k+j}$. Moreover, we have

$$\begin{aligned} \dot{z}_i^{k+\rho_{\bar{s}}} &= z_i^{k+\rho_{\bar{s}}+1} + \sum_{s=1}^{i-1} \delta_{i,s}^{k+\rho_{\bar{s}}} \tilde{v}_s + a_i^{k+\rho_{\bar{s}}} + b_i^{k+\rho_{\bar{s}}} \tilde{v} - (\delta_{i,\bar{s}}^{\rho_{\bar{s}}+k})' z_{\bar{s}}^{\rho_{\bar{s}}} - \delta_{i,\bar{s}}^{\rho_{\bar{s}}+k} \tilde{v}_{\bar{s}} \\ &= z_i^{k+\rho_{\bar{s}}+1} + (a_i^{k+\rho_{\bar{s}}} - (\delta_{i,\bar{s}}^{\rho_{\bar{s}}+k})' z_{\bar{s}}^{\rho_{\bar{s}}}) + b_i^{k+1} \tilde{v} + \sum_{s=1}^{i-1} \delta_{i,s}^{k+\rho_{\bar{s}}} \tilde{v}_s - \delta_{i,\bar{s}}^{k+\rho_{\bar{s}}} \tilde{v}_{\bar{s}} \\ &= z_i^{k+\rho_{\bar{s}}+1} + \tilde{a}_i^{k+\rho_{\bar{s}}} + \tilde{b}_i^{k+\rho_{\bar{s}}} \tilde{v} + \sum_{s=1}^{\bar{s}-1} \delta_{i,s}^{k+\rho_{\bar{s}}} \tilde{v}_s + \sum_{s=\bar{s}+1}^{i-1} \delta_{i,\bar{s}}^{k+\rho_{\bar{s}}} \tilde{v}_s, \end{aligned}$$

where the functions $\tilde{a}_i^{k+\rho_{\bar{s}}}, \tilde{b}_i^{k+\rho_{\bar{s}}} \in \mathbf{I}^{k+\rho_{\bar{s}}}$. Thus in the above formula, the nonzero function $\delta_{i,\bar{s}}^{k+\rho_{\bar{s}}}$ is eliminated. Note that if $k = 0$, then the change of coordinate $\tilde{z}_i^1 = z_i^1 - \delta_{i,\bar{s}}^{\rho_{\bar{s}}} z_{\bar{s}}^1$ transforms the first equation $y_i = z_i^1$ of (43) into $y_i = \tilde{z}_i^1 + \delta_{i,\bar{s}}^{\rho_{\bar{s}}} z_{\bar{s}}^1$. We define a new output $\tilde{y}_i = y_i - \delta_{i,\bar{s}}^{\rho_{\bar{s}}} z_{\bar{s}}^1 = y_i - \delta_{i,\bar{s}}^{\rho_{\bar{s}}} y_{\bar{s}}$ (which is actually an output multiplication of the form $\tilde{y}_i = \eta_i y$) such that the first equation of (43) becomes $\tilde{y}_i = \tilde{z}_i^1$.

Repeat the above construction to eliminate all nonzero functions $\delta_{i,s}^j$ for $j \geq \rho_s, 1 \leq s \leq i - 1$. Then system (43) becomes the following control system

$$\tilde{\Sigma} : \begin{cases} \tilde{y}_i = \tilde{z}_i^1, & i = 1, \dots, m, \\ \dot{\tilde{z}}_i^1 = \tilde{z}_i^2 + \tilde{a}_i^1 + \tilde{b}_i^1 \tilde{v}, \\ \dots \\ \dot{\tilde{z}}_i^{\rho_i-1} = \tilde{z}_i^{\rho_i} + \tilde{a}_i^{\rho_i-1} + \tilde{b}_i^{\rho_i-1} \tilde{v}, \\ \dot{\tilde{z}}_i^{\rho_i} = \tilde{v}_i, \\ \dot{z}^* = \tilde{f}^*(z, z^*) + \tilde{G}^*(z, z^*) \tilde{v}, \end{cases}$$

where $a_i^k, b_{i,s}^k \in \mathbf{I}^k$ for $1 \leq k \leq \rho_i - 1$. It is clear that $\Sigma \overset{\text{sys}}{\sim} \tilde{\Sigma}$ (we used coordinates changes, feedback transformations and output multiplications to transform Σ into $\tilde{\Sigma}$). Then consider the last row of every subsystem of $\tilde{\Sigma}$, which is $\dot{z}_i^{\rho_i} = \tilde{v}_i$. By deleting this equation in every subsystem and setting $y_i = 0$ for $i = 1, \dots, m$, and replacing the vector \tilde{v} by \dot{z}^ρ , we transform $\tilde{\Sigma}$ into a DAE $\tilde{\Xi}$ below. It is straightforward to see that $\tilde{\Sigma} \in \mathbf{Expl}(\tilde{\Xi})$, where

$$\tilde{\Xi} : \left\{ \begin{aligned} & \begin{bmatrix} 0 & & & & \\ & \ddots & & & \\ & & \ddots & & \\ & & & 1 & 0 \end{bmatrix} \begin{bmatrix} \dot{z}_i^1 \\ \dot{z}_i^2 \\ \vdots \\ \dot{z}_i^{\rho_i} \end{bmatrix} \\ & -\tilde{G}^*(\tilde{z}, z^*) \dot{z}^\rho + \dot{z}^* \end{aligned} \right. = \begin{bmatrix} \dot{z}_i^1 \\ \dot{z}_i^2 \\ \vdots \\ \dot{z}_i^{\rho_i} \end{bmatrix} + \begin{bmatrix} 0 \\ \tilde{a}_i^1 + \tilde{b}_i^1 \dot{z}^\rho \\ \vdots \\ \tilde{a}_i^{\rho_i-1} + \tilde{b}_i^{\rho_i-1} \dot{z}^\rho \end{bmatrix}, \quad i = 1, \dots, m, \\ & = \tilde{f}^*(\tilde{z}, z^*). \end{bmatrix}$$

Finally, by Theorem 3.6 and $\Sigma \overset{\text{sys}}{\sim} \tilde{\Sigma}$, we have that $\Xi \overset{\text{ex}}{\sim} \tilde{\Xi}$ and that $\tilde{\Xi}$ is in the **NWF1**, given by (20). \square

References

- [1] P.J. Rabier, W.C. Rheinboldt, Theoretical and numerical analysis of differential-algebraic equations, in: P.G. Ciarlet, J.L. Lions (Eds.), Handbook of Numerical Analysis, Volume VIII, Elsevier Science, Amsterdam, the Netherlands, 2002, pp. 183–537.
- [2] R. Rianza, Differential-Algebraic Systems. Analytical Aspects and Circuit Applications, World Scientific Publishing, Basel, 2008.
- [3] S. Kobayashi, K. Nomizu, Foundations of Differential Geometry, vol. 1, Interscience, 1963.
- [4] R. Rianza, DAEs in circuit modelling: a survey, in: A. Ilchmann, T. Reis (Eds.), Surveys in Differential-Algebraic Equations I, Differential-Algebraic Equations Forum, Springer-Verlag, Berlin-Heidelberg, 2013, pp. 97–136.
- [5] G. Byrne, P. Ponzi, Differential-algebraic systems, their applications and solutions, Comput. Chem. Eng. 12 (1988) 377–382.
- [6] C.C. Pantelides, D. Gritsis, K.R. Morison, R.W.H. Sargen, The mathematical modeling of transient systems using differential-algebraic equations, Comput. Chem. Eng. 12 (1988) 449–454.
- [7] P.J. Rabier, W.C. Rheinboldt, Nonholonomic Motion of Rigid Mechanical Systems from a DAE Viewpoint, vol. 68, Society for Industrial and Applied Mathematics, 2000.
- [8] P. Betsch, P. Steinmann, A DAE approach to flexible multibody dynamics, Multibody Syst. Dyn. 8 (2002) 365–389.
- [9] W. Blajer, K. Kołodziejczyk, Control of underactuated mechanical systems with servo-constraints, Nonlinear Dyn. 50 (2007) 781–791.
- [10] S. Reich, On a geometrical interpretation of differential-algebraic equations, Circuits Syst. Signal Process. 9 (1990) 367–382.
- [11] S. Reich, On an existence and uniqueness theory for nonlinear differential-algebraic equations, Circuits Syst. Signal Process. 10 (1991) 343–359.
- [12] P.J. Rabier, W.C. Rheinboldt, A geometric treatment of implicit differential-algebraic equations, J. Differ. Equ. 109 (1994) 110–146.
- [13] Y. Chen, S. Trenn, On geometric and differentiation index of nonlinear differential-algebraic equations, IFAC-PapersOnLine 54 (2021) 186–191.
- [14] T. Berger, Controlled invariance for nonlinear differential-algebraic systems, Automatica 64 (2016) 226–233.
- [15] T. Berger, The zero dynamics form for nonlinear differential-algebraic systems, IEEE Trans. Autom. Control 62 (2017) 4131–4137.
- [16] Y. Chen, S. Trenn, W. Respondek, Normal forms and internal regularization of nonlinear differential-algebraic control systems, Int. J. Robust Nonlinear Control 31 (2021) 6562–6584.
- [17] Y. Chen, S. Trenn, An approximation for nonlinear differential-algebraic equations via singular perturbation theory, IFAC-PapersOnLine 54 (2021) 187–192.
- [18] Y. Chen, W. Respondek, Geometric analysis of differential-algebraic equations via linear control theory, SIAM J. Control Optim. 59 (2021) 103–130.

- [19] L. Kronecker, Algebraische Reduction der Schaaren bilinearer Formen, Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften zu Berlin, 1890, pp. 1225–1237.
- [20] F.R. Gantmacher, *The Theory of Matrices (Volume II)*, Chelsea, New York, 1959.
- [21] K. Weierstraß, Zur Theorie der bilinearen und quadratischen Formen, Berl. Monatsb., 1868, pp. 310–338.
- [22] J.J. Loiseau, K. Özçaldıran, M. Malabre, N. Karcanias, Feedback canonical forms of singular systems, *Kybernetika* 27 (1991) 289–305.
- [23] T. Berger, A. Ilchmann, S. Trenn, The quasi-Weierstraß form for regular matrix pencils, *Linear Algebra Appl.* 436 (2012) 4052–4069.
- [24] T. Berger, S. Trenn, The quasi-Kronecker form for matrix pencils, *SIAM J. Matrix Anal. Appl.* 33 (2012) 336–368.
- [25] P. Rouchon, M. Fliess, J. Lévine, Kronecker’s canonical forms for nonlinear implicit differential systems, in: *System Structure and Control*, Pergamon, 1992, pp. 248–251.
- [26] A. Kumar, P. Daoutidis, *Control of Nonlinear Differential Algebraic Equation Systems with Applications to Chemical Processes*, Chapman and Hall/CRC Research Notes in Mathematics, vol. 397, Chapman and Hall, Boca Raton, FL, 1999.
- [27] A. Isidori, *Nonlinear Control Systems*, 3rd ed., Communications and Control Engineering Series, Springer-Verlag, Berlin, 1995.
- [28] H. Nijmeijer, A.J. van der Schaft, *Nonlinear Dynamical Control Systems*, Springer-Verlag, Berlin-Heidelberg-New York, 1990.
- [29] K.-T. Wong, The eigenvalue problem $\lambda T x + Sx$, *J. Differ. Equ.* 16 (1974) 270–280.
- [30] V. Doležal, The existence of a continuous basis of a certain linear subspace of e_r which depends on a parameter, *Čas. Pěst. Mat.* 089 (1964) 466–469.
- [31] K.A. Grasse, A vector-bundle version of a theorem of V. Doležal, *Linear Algebra Appl.* 392 (2004) 45–59.
- [32] T. Berger, T. Reis, Regularization of linear time-invariant differential–algebraic systems, *Syst. Control Lett.* 78 (2015) 40–46.
- [33] R. Marino, W. Respondek, A. van der Schaft, Equivalence of nonlinear systems to input-output prime forms, *SIAM J. Control Optim.* 32 (1994) 387–407.
- [34] A.S. Morse, Structural invariants of linear multivariable systems, *SIAM J. Control* 11 (1973) 446–465.
- [35] Y. Chen, W. Respondek, From Morse triangular form of ODE control systems to feedback canonical form of DAE control systems, *J. Franklin Inst.* 358 (2021) 8556–8592.
- [36] Y. Chen, *Geometric Analysis of Differential-Algebraic Equations and Control Systems: Linear, Nonlinear and Linearizable*, Ph.D. thesis, Normandie Université, INSA de Rouen, 2019.