

University of Groningen

Transporters involved in uptake of di- and tricarboxylates in *Bacillus subtilis*

Krom, Bastiaan P.; Warner, Jessica B.; Konings, Wil N.; Lolkema, Juke S.

Published in:

Antonie Van Leeuwenhoek: International Journal of General and Molecular Microbiology

DOI:

[10.1023/A:1024445131925](https://doi.org/10.1023/A:1024445131925)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2003

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Krom, B. P., Warner, J. B., Konings, W. N., & Lolkema, J. S. (2003). Transporters involved in uptake of di- and tricarboxylates in *Bacillus subtilis*. *Antonie Van Leeuwenhoek: International Journal of General and Molecular Microbiology*, *84*(1), 69-80. <https://doi.org/10.1023/A:1024445131925>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Transporters involved in uptake of di- and tricarboxylates in *Bacillus subtilis*

Bastiaan P. Krom, Jessica B. Warner, Wil N. Konings and Juke S. Lolkema*

Groningen Biomolecular Sciences and Biotechnology Institute, Department of Microbiology, University of Groningen, Kerklaan 30, 9751 NN Haren, The Netherlands; *Author for correspondence (e-mail: j.s.lolkema@biol.rug.nl; phone: (31)-50-3632150; fax: (31)-50-3632154)

Received 10 June 2002; accepted in revised form 21 November 2002

Key words: *Bacillus subtilis*, Di- and tricarboxylates, Regulation of transport, TCA cycle, Transport

Abstract

Di- and tricarboxylates found as intermediates in the tricarboxylic acid cycle can be utilized by many bacteria and serve as carbon and energy source under aerobic and anaerobic conditions. A prerequisite for metabolism is that the carboxylates are transported into the cells across the cytoplasmic membrane. *Bacillus subtilis* is able to metabolize many di- and tricarboxylates and in this overview the available data on all known and putative di- and tricarboxylate transporters in *B. subtilis* is summarized. The *B. subtilis* transporters, that are of the secondary type, are discussed in the context of the protein families to which they belong. Available data on biochemical characterization, regulation of gene expression and the physiological function is summarized. It is concluded that in *B. subtilis* multiple transporters are present for tricarboxylic acid cycle intermediates.

Introduction

Di- and tricarboxylates found as intermediates in the tricarboxylic acid cycle (TCA cycle, citric acid cycle, or Krebs cycle) can be metabolized by many bacteria and serve as carbon and energy source. Under aerobic conditions the substrates are oxidized to CO₂ and H₂O by running a full or partial TCA cycle during which free energy is conserved in the form of reducing equivalents that are used to generate proton motive force (pmf) and ATP in oxidative phosphorylation. Under anaerobic conditions multiple pathways exist. The C₄-dicarboxylic acids fumarate and L-malate, that is dehydrated to fumarate by fumarase, serve as energy source through fumarate respiration which leads to the generation of a pmf, high enough to drive the synthesis of ATP by F₀F₁-ATPases (for review see Jansusch et al. 2002, Kroger et al. 2002). Succinate produced during fumarate respiration is exchanged for external fumarate without the cost of metabolic energy.

The tricarboxylate citrate is used under anaerobic

conditions following different pathways (for enterobacteriae review by Bott 1997). In all cases, citrate is first converted to oxaloacetate by citrate lyase. Bacteria that lack this enzyme, as is the case for *Bacillus subtilis*, do not grow anaerobically on citrate. After the conversion of citrate to oxaloacetate several pathways are possible. In *Escherichia coli*, oxaloacetate is converted to malate and, subsequently, to fumarate by malate dehydrogenase and fumarase, respectively. Fumarate is metabolized further by fumarate respiration. A co-substrate like glucose is required to supply the reducing equivalents. In *Klebsiella pneumoniae*, no co-substrate is needed. Oxaloacetate is converted to pyruvate and carbon dioxide by oxaloacetate decarboxylase. This membrane embedded enzyme functions as a Na⁺ pump, thereby conserving the free energy released in the decarboxylation reaction in the form of a sodium ion gradient (for review see Dimroth et al. 2001). The sodium ion motive force is used to drive the uptake of citrate, or can be converted into a pmf by the action of Na⁺/H⁺ antiporters. Subsequently, pyruvate is converted to

acetyl-CoA, which is converted to acetylphosphate and, finally, to acetate, a process which yields one mol of ATP per mol of citrate (Bott 1997).

Lactic acid bacteria use fermentative pathways to metabolize the di- and tricarboxylates malate and citrate. In *Lactococcus lactis* malate is decarboxylated to yield lactate. The enzyme is cytoplasmic and, in contrast to the membrane bound Na^+ -pump in *K. pneumoniae*, not energy conserving. The uptake of malate is coupled to the exit of its decarboxylation product lactate (precursor/product exchange) which results in the formation of a membrane potential of physiological polarity. Furthermore, during the decarboxylation step a cytoplasmic proton is consumed which results in a pH gradient over the cell membrane. Taken together the pathway, termed malolactic fermentation, results in a pmf that is large enough to drive ATP synthesis via the F_0F_1 -ATPase (Poolman et al. 1991; Bandell et al. 1997). In a similar, but more complex manner, citrate is metabolized in *Leuconostoc mesenteroides* and *L. lactis* by citrolactic fermentation (Marty-Teysset et al. 1996). Similar as in malolactic fermentation, divalent citrate is taken up in exchange for monovalent lactate. The pathway in the cytosol converting citrate into lactate requires three different enzymes and requires co-metabolism of glucose (Marty-Teysset et al. 1995, 1996).

For all metabolic pathways described above it is essential that the di- and tricarboxylates be transported across the cytoplasmic membrane. All known and putative transporters of di- and tricarboxylates are secondary transporters; they derive energy from the chemiosmotic gradient of protons or sodium ions to accumulate the substrate against a concentration gradient. Typical for the anaerobic breakdown pathways is the coupling of the uptake of the growth substrate to the excretion of the end product of the pathway, i.e.

fumarate/succinate exchange, citrate/succinate exchange, citrate/lactate exchange, and malate/lactate exchange.

Di- and tricarboxylate uptake in *B. subtilis* has been studied in the seventies and early eighties of the last century. For C_4 -dicarboxylates two uptake systems were described. One is a L-malate uptake system that is induced by L-malate in the medium. It was reported to be responsible for growth on L-malate as sole carbon and energy source (Willecke and Lange 1974). The second system is less specific and was claimed to transport L-malate, succinate and fumarate. It was found in cells grown on yeast extract (Ghei and Kay 1973), in the presence of citrate (Ghei and Kay 1972), or expressed constitutively (Bisschop et al. 1975). The latter study in membrane vesicles showed that the transporter was of the secondary type. Similarly, it was demonstrated that a secondary transporter catalyzes the uptake of the tricarboxylate citrate in *B. subtilis* (Bergsma and Konings 1983). Remarkably, uptake of citrate was dependent on the availability of divalent metal ions (Warner et al. 2002b). These early reports offer no information on the identity of the transporters that are responsible for the uptake of the solutes. The availability of modern molecular biology techniques has greatly facilitated the identification and characterization of the proteins involved in specific metabolic pathways. Moreover, the complete genome sequence of *B. subtilis* (Kunst et al. 1997) yielded valuable information on all the transport capabilities of this bacterium (Saier et al. 2002).

In this review an overview is given of the di- and tricarboxylate transporters found in *B. subtilis* and the protein families to which these transporters belong (summarized in Table 1). Available data on functional properties, regulation of transcription and physiological function of the individual proteins are discussed.

Table 1. Transporter families and individual transporters of *B. subtilis* 168 discussed in this review.

| Family | TC system ^a | Transporter | Gene | GI # ^b |
|--------|------------------------|-------------|-------------|-------------------|
| MeCit | 2.A.11. CitMHS | CitM | <i>yflN</i> | 16077828; 2443237 |
| | | CitH | <i>yxiQ</i> | 16080957 |
| | | YraO | <i>yraO</i> | 16079739; 2108284 |
| DctA | 2.A.23. DAACS | DctP | <i>ydbH</i> | 16077514 |
| 2HCT | 2.A.24. CCS | MaeN | <i>yufR</i> | 16080210; 7446780 |
| | | CimH | <i>yxkJ</i> | 16080928; 7446781 |
| | | MleN | <i>yqkI</i> | 16079413; 1731094 |
| DASS | 2.A.47. DASS | YflS | <i>yflS</i> | 16077824 |

^a Transport Commission family number and name (Saier 2000)^b NCBI Protein database GI number (www.ncbi.nlm.nih.gov).

MeCit family

The MeCit family currently consists of 12 known proteins that are found exclusively in bacteria. Some are found in typical pathogenic bacteria, like *Staphylococcus aureus* and *Neisseria meningitidis*, others in typical soil bacteria like *Streptomyces coelicolor* or *B. subtilis*. Two members of the family have been characterized in some detail, CitM and CitH of *B. subtilis*. The transporters transport tricarboxylates in complex with divalent cations. Both were shown to be proton-coupled symporters. The hydrophathy profiles of members of the family contain 11 to 12 hydrophobic regions, suggesting the same number of putative transmembrane segments (TMSs). The membrane topology of the family has not been determined experimentally. *B. subtilis* contains in total three proteins that belong to the MeCit family, besides CitM and CitH, the protein coded by open reading frame *yraO*, which has not been characterized.

Characterization

CitM and CitH, designated YfIN and YxiQ in the Subtilist database, respectively, and YraO of *B. subtilis* share 52 to 61% identical residues and, additionally, 18 to 25% similar residues. CitM and CitH have been successfully cloned and functionally expressed in *E. coli* (Boorsma et al. 1996; Krom et al. 2000). However, functional expression of *yraO* in *E. coli* has failed so far (B.P. Krom, unpubl.). Using right-side-out (RSO) membrane vesicles, CitM and CitH were shown to be secondary transporters that are driven by the transmembrane pH gradient, ΔpH , and by the membrane potential, $\Delta\Psi$, inside negative (Boorsma et al. 1996). This observation indicates that at least one net positive charge is transported into the cell.

At first, CitH was identified as a transporter for free citrate (Boorsma et al. 1996). However, a more recent study has shown that CitH is a Ca^{2+} -citrate transporter that also accepts Ba^{2+} and Sr^{2+} complexed to citrate but not free citrate (Krom et al. 2000). The apparent affinities (K_{app}) for the different complexes were very similar, ranging from 33 to 50 μM . In contrast, the maximal rates of transport were at least 3 fold higher for Ca^{2+} -citrate than for Sr^{2+} -citrate or Ba^{2+} -citrate. It was shown that Ca^{2+} is co-transported into the cell.

CitM has been shown to transport Mg^{2+} -citrate, but also Mn^{2+} , Co^{2+} , Ni^{2+} and Zn^{2+} -citrate. Similar as

observed for CitH, the K_{app} 's of the cloned transporter in *E. coli* were very similar for the different complexes ranging from 35 to 63 μM and are in line with the K_{app} of 40 μM measured in *B. subtilis* vesicles for Mg-citrate (Bergsma and Konings 1983). The maximal rate of transport ranged from 214 to 661 pmol/mg protein \cdot min for Mn^{2+} -citrate and Co^{2+} -citrate, respectively (Krom et al. 2000). It has been shown that the divalent cation is transported into the cell, i.e. the complex of metal and citrate is the transported species. Besides complexes of citrate with divalent metal ions, CitM also transports isocitrate in complex with divalent metal ions (Warner et al. 2002b).

Summarizing, both CitM and CitH transport a metal-citrate complex, but with complementary metal specificity. It is possible that the two transporters distinguish metal ion-citrate complexes based on the ionic radii of the metal ions. CitH would transport bulky ions with ionic radii greater than 0.98 Å, while CitM would transport the smaller ions (Krom et al. 2000). Since transport by both proteins was shown to be electrogenic and the Me-citrate complex is monovalent anionic, at least two protons are co-transported with the Me-citrate complex.

Regulation of expression

Expression of the *citH* gene was studied using the LacZ-fusion approach by which the promoter region of *citH* was fused upstream of the gene for β -galactosidase (*lacZ*). Growth of the recombinant strain on plates in the presence of many different di- and tricarboxylates revealed a low level of expression, but no clear induction pattern. Similarly, a low level of expression was observed in exponentially growing cells which correlated with the low citrate transport activity in the presence of Ca^{2+} , Sr^{2+} and Ba^{2+} in the cells (Krom et al. 2000). On the other hand, a DNA array study using 24 two-component regulatory systems present on the genome of *B. subtilis* indicated that the YufLM sensor kinase-response regulator pair would be involved in CitH induction (Kobayashi et al. 2001). Interestingly, many genes induced via YufLM are members of the ComK regulon involved in competence (Kobayashi et al. 2001). Possibly CitH expression is induced during the development of competence.

The regulation of expression of *citM* has been studied extensively. *CitM* expression requires the presence of Mg^{2+} and citrate or isocitrate in the

medium (Warner et al. 2000; Yamamoto et al. 2000). Mn^{2+} -, Ni^{2+} -, Zn^{2+} - and Co^{2+} -citrate were also capable of induction, albeit to lower levels. The induction is mediated by a classical two-component signal transduction system, consisting of a sensor, CitS encoded by *yflR*, and a response regulator, CitT encoded by *yflQ*, both of which are located upstream of *citM* (Figure 1). CitT is the DNA binding transcriptional regulator and has been shown to bind to two regions in the promoter region of *citM* (Yamamoto et al. 2000). The induction by citrate is overruled by the presence of carbon sources that are more easy to metabolize, like glucose, glycerol and inositol (Warner et al. 2000). Also the combination of succinate and glutamate and some component(s) in Luria Bertani broth were shown to repress CitM expression (Warner et al. 2002a, 2003). Repression is mediated by the classical Carbon Catabolite Repression system (CCR) as has been shown for numerous other proteins in *B. subtilis* (Hueck and Hillen 1995). A functional Catabolite Responsive Element (CRE) is located within the promoter region of CitM (Miwa et al. 2000; Warner et al. 2000; Yamamoto et al. 2000).

The third member of the MeCit family in *B. subtilis*, encoded by *yraO* gene, could not be shown to be expressed under any of many conditions tested using the LacZ promoter fusion approach (B.P. Krom, unpubl.). Either this gene is expressed under very special conditions, or the gene is cryptic.

Physiological function

B. subtilis grown in continuous cultures with glucose as sole carbon source results in acid production caused by overflow of the TCA-cycle, and consequently, rapid acidification of the medium which inhibits growth (Goel et al. 1995). Co-metabolism of glucose and citrate resulted in lower acid production and a higher growth yield due to a better coordination between glycolysis and the TCA cycle. It has been

proposed that elevated levels of Ca^{2+} in the cell might result in inhibition of pyruvate kinase activity, thus slowing down glycolysis. A specific Ca^{2+} -citrate transporter was thought to be responsible for this elevated level of Ca^{2+} in the cell. If this were indeed the case, CitH may very well be this Ca^{2+} -citrate uptake system.

CitM is the transporter involved in the growth of *B. subtilis* on citrate (Yamamoto et al. 2000) and isocitrate (Warner et al. 2002b) as sole carbon source. A CitM deletion strain lost the ability to grow on citrate and isocitrate and both carbon sources induced expression of the transporter when present in the medium. Catabolite repression by other medium components exerted on CitM expression further confirm the role of CitM during growth on citrate and isocitrate. The effect of CitM expression in the presence of citrate complexed to toxic heavy metals like Ni^{2+} , Zn^{2+} and Co^{2+} , on the growth of *B. subtilis* dramatically increased the toxic effects of these metals (Krom et al. 2002). The fact that CitM expression is induced best by the Mg^{2+} -citrate complex as compared to the Ni^{2+} , Zn^{2+} and Co^{2+} -citrate complexes, together with the toxic effects in the presence of these heavy metal-citrate complexes, suggest that Mg^{2+} -citrate is the natural substrate for CitM.

Bergsma and Konings studied Me^{2+} -dependent citrate uptake in membrane vesicles prepared from *B. subtilis* W23 cells grown in the presence of citrate (Bergsma and Konings 1983). At that time it was not recognized that citrate transport in *B. subtilis* could be mediated by at least three different transporter proteins and the observed metal dependence is best explained by similar levels of expression of both CitM and CitH. The uptake of citrate was high in the presence of both Mg^{2+} and Ca^{2+} . In *B. subtilis* strain 168, the contribution of the Ca^{2+} -dependent uptake was much smaller (Krom et al. 2000), suggesting a different regulation of expression in the W23 and 168 strains.




Figure 1. Schematic overview of the region on the *B. subtilis* genome that contains the *citM* gene. Arrows indicate confirmed or putative promoter regions, while terminators of transcription are indicated by loop structures. *CitST* code for the sensor kinase and response regulator respectively, *citM* codes for the Mg^{2+} -dependent citrate transporter.

DctA family

In bacteria, four families of secondary transporters specific for C_4 -dicarboxylates can be distinguished; the DctA family, members of which are expressed under aerobic conditions, and the three Dcu families, DcuA, DcuB and DcuC, members of which are expressed under anoxic or anaerobic conditions. No homologues of either one of the Dcu families were identified in the genome of *B. subtilis*, while one homologue of the DctA family was found, and, unfortunately, designated DctP (Asai et al. 2000). The DctA family is a subfamily of the large dicarboxylate/ aminoacid:cation symporter (DAACS) family of transporters found in bacteria, archaea and eukaryotes (Saier 2000; Janausch et al. 2002). The bacterial DctA subfamily homologues share at least 40% identical residues (Slotboom et al. 1999a) and are found in Gram-positive and Gram-negative aerobic or facultative anaerobic bacteria. All members use either H^+ or Na^+ as coupling ion and are around 450 amino acids in length (for a recent review see Janausch et al. 2002). *Rhizobium meliloti* DctA is reported to contain 12 TMS based on a series of LacZ fusions (Jording and Puhler 1993). Members of the DctA subfamily share between 27% and 40% amino acid sequence identity with members of the glutamate transporter subfamily of the DAACS family. The latter proteins contain 8 TMSs and two reentrant loop-like structures that resemble the pore-loop structures found in ion channels (Grunewald et al. 1998; Slotboom et al. 1999b, 2001). Because of significant evolutionary relationship between the two subfamilies, a similar membrane topology most likely applies to members of the DctA subfamily.

Characterization

To date no cloning and functional characterization of DctP of *B. subtilis* has been reported. Growth studies

(see below) suggest that DctP transports fumarate and succinate and, possibly, L-malate. A study by Bisschop *et al.* using membrane vesicles prepared from cells grown under conditions that later were shown to result in expression of DctP revealed apparent affinities for fumarate, succinate and L-malate of 7.5 μM , 4.3 μM , and 13.5 μM , respectively (Bisschop et al. 1975). Competitive inhibition of uptake by the three substrates suggested that they were all transported by one and the same transporter. It cannot be excluded that transporters other than DctP might have been involved in the uptake of L-malate as well. Transport of the C_4 -dicarboxylates in the membrane vesicles was sensitive to uncouplers suggesting pmf driven transport.

Regulation of expression

The *dctP* gene is transcribed both in a single mRNA and polycistronically together with two genes coding for the two-component signal transduction system DctSR (Figure 2) (Asai et al. 2000). The two-component system DctSR (the *ydbFG* genes) is required for induction of DctP together with the product of the *dctB* gene that is located immediately upstream of the *dctSRP* operon. DctB is homologous to the periplasmic binding protein of the binding protein dependent secondary transporter family (or TRAP family (for review see Kelly and Thomas 2001)), but is not needed for transport catalyzed by DctP. It was postulated that DctB might bind the substrate with high affinity and, subsequently, would activate the DctSR two-component system (Rabus et al. 1999; Asai et al. 2000).

Expression of *dctP* can be found in minimal medium in the presence of a small amount of yeast extract. Interestingly induction is prevented by the addition of mM concentrations of L-malate in the growth medium (Asai et al. 2000) while this may be one of the substrates of the transporter. It is remark-




Figure 2. Schematic overview of the region on the *B. subtilis* genome that contains the *dct* operon. Arrows indicate the direction of transcription, terminators of transcription are indicated by loop structures. *DctB* codes for the periplasmic binding protein, *dctSR* for the sensor kinase and response regulator, respectively, and *dctP* codes for the transporter

able to see that the substrates succinate and fumarate do not induce transcription of the transporter above levels found when *B. subtilis* was grown on minimal medium with a small amount of yeast extract. Perhaps the amount of inducer in yeast extract is high enough to induce gene expression.

A functional CRE sequence located in the promoter region of *dctP* has been identified (Miwa et al. 2000) indicating that expression is under control of Carbon Catabolite Repression. In an older study it was demonstrated that malate uptake activity was repressed by the presence of glucose in the medium (Ghei and Kay 1973).

Physiological function

DctP is required for growth of *B. subtilis* on fumarate and succinate, but not on L-malate. A possible explanation for this phenotype might be that multiple systems are capable of L-malate uptake while probably only one system is responsible for the uptake of succinate and fumarate. The auxotrophic phenotypes for succinate and fumarate, *suc*⁻ and *fum*⁻, respectively, were obtained in strains in which the two-component system was deleted (*dctS*⁻, *dctR*⁻) as well as in the strain in which the binding protein was deleted (*dctB*⁻). DctP is likely to be the C₄-dicarboxylate transporter that is expressed in cells grown on yeast extract (Ghei and Kay 1972, 1973; Willecke and Lange 1974; Bisschop et al. 1975). Interestingly, in one of these studies a dicarboxylate binding protein has been identified, purified and characterized (Kay 1981). The binding protein had an apparent molecular mass of 41 kDa and could be extracted from membrane vesicles with detergents indicating the presence of a membrane anchor. The purified binding protein was shown to bind L-malate. The predicted molecular mass of DctB is 40 kDa. It is quite possible that DctB is the C₄-dicarboxylate binding protein described by Kay (Kay 1981).

2HCT family

The 2-hydroxycarboxylate transporter (2HCT) family currently consists of 17 members that are solely found in bacteria. Some members of the 2HCT family have been studied extensively. The family contains Na⁺- and H⁺-symporters (CitS of *K. pneumoniae*, MaeN and CimH of *B. subtilis* and MaeP of *Streptococcus*

bovis (van der Rest et al. 1992; Kawai et al. 1997; Wei et al. 2000)), but also citrate/lactate and malate/lactate exchangers found in lactic acid bacteria (CitP of *Lc. mesenteroides* and MleP of *L. lactis* (Marty-Teyssset et al. 1995; Bandell et al. 1997)). The latter transporters are involved in secondary metabolic energy generation under physiological conditions (citrolactic and malolactic fermentation, respectively (Poolman et al. 1991; Lolkema et al. 1996; Marty-Teyssset et al. 1996)). Recently, a second member of the family was described in *K. pneumoniae*, termed CitW, that catalyzes citrate/acetate exchange (Kästner et al. 2002). Substrate specificity studies revealed that the symporters in the family are very specific, transporting only citrate or L-malate, while the exchangers catalyze transport of a wide range of 2-hydroxycarboxylates (Bandell et al. 1997).

The hydrophathy profile of the amino acid sequence of the members of the family shows 12 hydrophobic regions long enough to traverse the membrane. Studies of CitS of *K. pneumoniae* revealed that 11 of these span the membrane (van Geest et al. 1999; van Geest and Lolkema 2000). The hydrophobic region between TMSs V and VI is located externally. Studies on the exchangers CitP of *L. mesenteroides* and MleP of *L. lactis* revealed that the last cytoplasmic loop and the C-terminal outgoing TMS XI are involved in substrate specificity (Bandell and Lolkema 2000b). A conserved arginine in TMS XI was shown to directly interact with one of the carboxylates of the substrate (Bandell and Lolkema 2000a, Krom et al. 2003). Conserved residues in the cytoplasmic loop preceding TMS XI were shown to be accessible from the outside of the cell. It is proposed that this cytoplasmic loop is part of the structure that forms the translocation pathway through the protein.

Characterization

Two members of the 2HCT family, MaeN and CimH, are found in *B. subtilis* and have been functionally characterized. MaeN, designated YufR in the Subtilist database, shows the highest sequence identity to MaeP of *S. bovis*, a H⁺ dependent L-malate transporter (Kawai et al. 1997). MaeN was cloned and functionally expressed in *E. coli* (Wei et al. 2000) and shown to code for a Na⁺ coupled L-malate transporter. *E. coli* cells deficient in malate transport could be complemented for transport of L-malate by MaeN expression. The cells were unable to grow on minimal medium containing fumarate, succinate or citrate,

indicating that these di- and tricarboxylates are not substrates of MaeN (Wei et al. 2000).

CimH, designated YxkJ in the Subtilist database, has been cloned and functionally expressed in *E. coli* (Krom et al. 2001). *E. coli* cells deficient either in malate or citrate transport could be complemented by expression of CimH, indicating that both citrate and L-malate are substrates of this transporter. CimH is capable of citrate and L-malate symport in membrane vesicles, while L-citramalate binds to the protein but is not transported. Kinetic analysis revealed a high affinity for citrate, but a low maximal rate, while the affinity for L-malate was low, but the maximal rate high. Therefore, CimH is a high affinity, low capacity citrate transporter and a low affinity, high capacity L-malate transporter. Citrate is transported in the free anionic state, in contrast to two other citrate transporters of *B. subtilis* (see MeCit family) that require divalent cations for transport. Like most 2-HCT members, CimH is highly stereoselective, recognizing the S-, but not the R-enantiomers of malate and citramalate. CimH catalyzes electroneutral H⁺-symport; transport is driven by a pH gradient ΔpH and not by the membrane potential $\Delta\Psi$. Since at pH 6, the condition of the experiments, divalent citrate and malate are the most prevalent anionic species, most likely two protons are co-transported.

Regulation of expression

Expression of *maeN* is induced by L-malate in the medium (Asai et al. 2000). The *yufLM* genes that code for a two component sensor system and that are located upstream of the *maeN* gene would be a potential candidate for the sensor system (Figure 3). Based on amino acid sequence identity the sensor, YufL, is clustered in the CitA family of sensors for C₄-dicarboxylates (Janausch et al. 2002) together with the C₄-carboxylate sensors DcuS of *E. coli* (35% identical residues), and citrate sensors CitS (31% identity) and CitA (25% identity) of *B. subtilis* and *K.*

pneumoniae, respectively (Kaspar et al. 1999; Yamamoto et al. 2000). Nevertheless, a DNA array study did not identify *maeN* as a potential target for YufL regulation. In fact, none of 24 two component regulatory systems analyzed could be linked to induction of MaeN expression (Kobayashi et al. 2001). The uptake of malate is repressed by the presence of glucose in the medium (Ghei and Kay 1973). Although not included in an evaluation of 126 putative CRE-sequences in *B. subtilis* (Miwa et al. 2000) a putative CRE-sequence is present in front of *maeN* with only two mismatches (Figure 4) with the consensus CRE-sequence (Weickert and Chambliss 1990).

A *B. subtilis* strain containing a fusion of the promoter region of *cimH* and the gene coding for β -galactosidase (LacZ fusion) did not show induction of expression on solid nor in liquid medium supplemented with a variety of di- or tricarboxylates, including citrate and L-malate. Cells harvested in the exponential growth phase grown on a minimal medium (Krom et al. 2000) showed a low citrate uptake activity in the presence of the chelator EDTA, suggesting a low, constitutive level of expression. The promoter of *cimH* contains a functional CRE sequence suggesting that expression is subject to Carbon Catabolite Repression (Miwa et al. 2000). Interestingly, at the position of the ribosome binding site (RBS) in front of *cimH* of the 2HCT family and *citH* of the MeCit family, a completely conserved stretch of ten nucleotides are present at exactly the same distance from the translation start site (Figure 4). Only two other genes on the *B. subtilis* genome share this sequence at the same position; *dctB*, the C₄-dicarboxylate periplasmic binding protein involved in induction of *dctP* (see DctA subfamily), and *mleA*, encoding the malolactic enzyme found in the *mleAN* operon (see NhaC family below). Possibly, this sequence constitutes a rather long conserved RBS. Alternatively it might represent some sort of regulatory region. The relevance of this observation is not clear.



Figure 3. Schematic overview of the region on the *B. subtilis* genome that contains the *maeN* gene. Arrows indicate the direction of transcription, putative terminators of transcription are indicated by loop structures. *YufLM* encode the sensor kinase and response regulator, respectively, *maeN* encodes the malate transporter.

A

```

CRE      TGWAANCGNTNWCA
          ||| ||| ||| ||| |||
maeN     TGTTAACGCTTTCT

```

B

```

cimH     CTTAACATAGTGAACAAT-TTCACTGCATCAAG--AATAGGGGAGG-TTTTTCAATG
citH     ---AAGGATGGAAGCGCTTACCAAAAAAGCTGTTGAATAGGGGAGG-TCATATCATG
dctB     TTTACGAATGTCATAGCC-CCCATTATCCTTAAC--ATAGGGGAGG-GCTTTTCATG
mleA     ---ACAATACAGTTAAA-GCCGAAAAAAGCTAGGCTAATAGGGGAGGATCACTGAATG
          *                *                *****                ***

```

Figure 4. Panel A. Alignment of consensus CRE sequence upstream of the *amyE* gene and the putative CRE sequence found in the *maeN* promoter region. N and W represent any base and A or T, respectively. Vertical lines indicate identical bases. Panel B. Alignment of the 50 base pairs preceding the ATG start codon (underlined) of four genes involved in carboxylate transport and metabolism in *B. subtilis*. *CimH* codes for a citrate/malate transporter, *citH* for a Ca^{2+} -citrate transporter, *dctB* for a dicarboxylate binding protein, and *mleA* for a malolactic enzyme. The alignment was prepared using the Clustal X program (Jeanmougin et al. 1998). Conserved nucleotides are marked with an asterix.

Physiological function

The 2HCT family contains proteins that function as symporters or precursor/product exchangers. The latter are involved in the generation of metabolic energy in lactic acid bacteria, (Poolman et al. 1991; Lolkema et al. 1996; Marty-Teysset et al. 1996). MaeN and CimH probably function as L-malate and citrate/L-malate symporters, respectively, under physiological conditions (Wei et al. 2000; Krom et al. 2001). The observation that CimH is a high affinity, low capacity citrate transporter and a low affinity, high capacity L-malate transporter could mean that the physiological function of CimH is two-fold. Firstly, CimH might function in the uptake of citrate under conditions where citrate in the medium is present in its uncomplexed state. Secondly, CimH might function as an L-malate transporter at high concentrations of L-malate, while another transporter, possibly DctP, takes over at lower concentrations. Microorganisms are known to use both high and low affinity transporters for the same substrate that are differentially expressed dependent on the external concentration of the substrate (e.g. the iron transporters of *Candida albicans* (Ramanan and Wang 2000)).

NhaC family

Na^+/H^+ antiporters are very abundant in nature and believed to play a role in pH and Na^+ homeostasis (Padan and Schuldiner 1994; Padan et al. 2001). The

proteins are distributed over a number of different gene families, but in *B. subtilis* only two members of the NhaC family are present (Saier et al. 2002). The NhaC family of Na^+/H^+ antiporters is relatively small, with members found in bacteria and in archaea. The proteins are around 460 amino acids long and predicted to contain 12 putative TMSs. The two members in *B. subtilis*, YheL and MleN, are the only members that have been characterized to some extent (Ito et al. 1997; Wei et al. 2000). YheL functions as a genuine Na^+/H^+ antiporter (not further discussed here), and MleN couples Na^+/H^+ exchange to the exchange of malate and lactate (Wei et al. 2000).

Characterization

MleN, designated YqkI in the Subtilist database, was cloned and functionally expressed in *E. coli*. Everted membrane vesicles expressing MleN took up lactate in symport with Na^+ ions, but only in the presence of intravesicular L-malate. Since the exchange between lactate and L-malate was electroneutral, it was concluded that MleN catalyzes the exchange of divalent L-malate and two H^+ for monovalent lactate and one Na^+ ion (Wei et al. 2000).

Regulation of expression

The *mleN* gene is found together with the *mleA* gene in an operon structure (Figure 5). MleA codes for malolactic enzyme that converts malate in a single step into lactate and CO_2 . No information is available



Figure 5. Schematic representation of the DNA region that contains the *mle* operon. The arrow indicates the direction of transcription. A putative terminator of transcription is indicated with a loop structure. *MleN* encodes the transporter, *mleA* encodes malolactic enzyme.

on the regulation of transcription of the operon. Mostly based on the proposed physiological function (see below), it was suggested that *mleN* expression would be induced under low pmf conditions. The *mleA* gene is one of the four genes on the *B. subtilis* genome that is preceded by the conserved putative RBS depicted in Figure 4.

Physiological function

The organization of MleN together with malolactic enzyme MleA in an operon structure suggests a similar pathway for the breakdown of malate as observed in lactic acid bacteria. Malate is taken up, decarboxylated yielding lactate and carbon dioxide, and lactate leaves the cell in exchange for malate. In lactic acid bacteria, malolactic fermentation functions in proton motive force generation (Poolman et al. 1991) or resistance against lactate under acidic conditions (Magni et al. 1999). It was proposed that in *B. subtilis* the pathway might function in Na^+ resistance under conditions of low pmf by the following mechanism. H_2 -malate would enter the cell via malate-lactate exchange catalyzed by MleN and be converted into lactate and CO_2 by MleA. Lactate⁻ would leave the cell in symport with Na^+ and in exchange for H_2 -malate in an electroneutral transport step. Apart from the chemical conversion, the net result would be the exchange of one cytoplasmic Na^+ ion for one H^+ , i.e. the decarboxylation reaction catalyzed by MleA would drive the extrusion of Na^+ catalyzed by MleN. It is proposed that *B. subtilis* uses this pathway as an alternative Na^+/H^+ antiport mechanism in Na^+ resistance under conditions of low pmf (Wei et al. 2000). *B. subtilis* grown on malate under conditions of low pmf requires MleN activity for growth.

Miscellaneous transporters

The genome of *B. subtilis* (Kunst et al. 1997) contains many proteins with unknown functions. Based on homology to proteins with known function, putative transporters of di- and tricarboxylates may be as-

signed. One additional ORF, *yflS*, was identified that could potentially code for a transporter involved in growth of *B. subtilis* on di- and tricarboxylates. Two other ORF's coding for YoaB and CsbX are annotated in the Subtilist database as putative α -ketoglutarate transporters, but are probably not.

On the genome of *B. subtilis*, *yflS* is located close to the gene that codes for the Mg^{2+} -citrate transporter CitM (see Figure 1). The product of the *yflS* gene belongs to the ubiquitous family of divalent anion: Na^+ symporters (DASS (Saier et al. 2002)). All members are around 450 amino acids in size and contain 12 hydrophobic regions indicative of 12 putative TMSs (Pos et al. 1998). Members of the family transport inorganic anions, like sulfate and phosphate, but also organic anions like di- and tricarboxylates and acidic amino acids. The product of *yflS* is homologous to CitT, the citrate-succinate antiporter of *E. coli* that is expressed under anaerobic conditions (Pos et al. 1998). CitT is located in an operon involved in the metabolism of citrate under anaerobic conditions, similar to the operon for anaerobic citrate metabolism in *K. pneumoniae* (Bott 1997). These operons contain the genes coding for citrate lyase that converts citrate into oxaloacetate, an essential step in anaerobic citrate metabolism; these genes are not present on the *B. subtilis* genome. Therefore, *B. subtilis* is unlikely to be capable of anaerobic growth on citrate and YflS to be a citrate-succinate antiporter as is CitT of *E. coli*. YflS has the highest identity with SOT1 found in chloroplasts of *Spinacia oleraceae* (35% identity to CitT, 53% to SOT1). SOT1 has been extensively characterized (Weber et al. 1995) and was shown to import α -ketoglutarate in exchange for stromal malate. Besides malate also succinate, fumarate and α -ketoglutarate could be used as counter ions. Possibly YflS transports one of these substrates.

The two homologous gene products YoaB and CsbX have been annotated in the Subtilist database (<http://genolist.pasteur.fr/SubtiList/>) to transport α -ketoglutarate. A BLAST search (Altschul et al. 1997) reveals the highest homology to the ribitol transporter, RbtT of *E. coli* and *K. pneumoniae* and the arabinitol transporter DalT of *K. pneumoniae* and no significant

homology to known α -ketoglutarate transporters like KgtP of *E. coli*. Since no experimental data are available to support the annotation, we tentatively conclude that both CsbX and YoaB are not involved in transport of di- or tricarboxylates in *B. subtilis*.

Conclusion

Transport of TCA cycle intermediates in *B. subtilis* is catalyzed by both proton and sodium dependent secondary transporters. A variety of protein families are involved and overlapping substrate specificities between several transporters has been found (Table 2). Multiple transport systems could be involved in growth on a certain substrate as carbon and energy source.

At least three and possibly five transporters (Table 2) transport citrate. Because *i.* CitM expression is induced by citrate in the medium, *ii.* induction is repressed via Carbon Catabolite Repression, and *iii.* the CitM deletion strain shows a *cit⁻* phenotype, CitM is the transporter responsible for growth on citrate as sole carbon and energy source under standard conditions. The exact roles of CitH and CimH remain unclear, however both proteins are functional citrate transporters that, under standard growth conditions, are expressed at very low levels. Until the inducing conditions for both transporters are found, their exact role in *B. subtilis* remains unclear. YraO, the third member of the MeCit family in *B. subtilis* is likely to be a citrate transporter as well, but expression of the ORF was not observed, nor has it been possible to heterologously express the gene coding for YraO. YfIS is a homologue of CitT of *E. coli*, a citrate/succinate antiporter that is expressed under anaerobic conditions. Though the *B. subtilis* genome does not contain the known genes coding for citrate lyase, necessary for anaerobic citrate degradation, anaerobic growth of

B. subtilis has recently been reported (for review see Nakano and Zuber (1998)). Possibly, YfIS plays a role under these growth conditions.

Isocitrate is transported by CitM in the MeCit family. A CitM deletion strain of *B. subtilis* could not utilize isocitrate as carbon source (Warner et al. 2002b). Therefore, CitM is the transporter responsible for growth on isocitrate in *B. subtilis*. Isocitrate transport by CitM is probably dependent on the presence of the same divalent metal ions as the transport of citrate (Warner et al. 2002b). Since CitH and YraO share a high sequence identity to CitM it is possible that both proteins are also capable of isocitrate transport. However, unlike CitM, CitH was not induced by isocitrate in the medium (B.P. Krom, unpubl.).

Succinate and fumarate are both transported by DctP. A DctP knock-out strain has a *suc⁻* and a *fum⁻* phenotype indicating that DctP is the transporter involved in growth of *B. subtilis* on succinate and fumarate. Remarkably, neither succinate nor fumarate seems to be involved in the induction of DctP and the inducer that activates the two-component system DctSR remains to be identified.

The involvement of the different transporters in the metabolism of L-malate in *B. subtilis* is not exactly clear. At least four but possibly five different transport proteins are capable of L-malate transport. DctP might be involved in growth on L-malate based on studies in *B. subtilis* membrane vesicles (Bisschop et al. 1975), but since a DctP deletion strain does not show a *mal⁻* phenotype (Asai et al. 2000), DctP is clearly not the only transporter involved in growth on L-malate. Repression of DctP expression by millimolar concentrations of L-malate in the medium suggests that DctP may be involved at low L-malate concentrations. The low K_m for L-malate transport reported by Bisschop et al. (1975) and the dependence of the sensing system that regulates expression of the gene on a binding protein (DctB) would support such a function. In contrast, expression of the Na^+ -malate symporter MaeN in the 2HCT family is induced by L-malate in the medium and, therefore, could take over uptake of L-malate at higher concentration. Uptake of L-malate has been shown to be subject to carbon catabolite repression and the promoter region of both the *dctP* and *maeN* genes contain CRE-sequences. CimH in the 2HCT family is not induced by L-malate in the medium. The kinetic parameters, high affinity constant and maximal rate, suggest that CimH functions at high external L-malate concentrations. MleN also transports L-malate, but does so under conditions of low pmf in an exchange reaction with

Table 2. Transporters involved in transport of TCA cycle intermediates in *B. subtilis* 168.

| Substrate | Confirmed transporter | Putative transporter |
|-------------------------|---------------------------|----------------------|
| citrate | CitM, CitH and CimH | YraO and YfIS |
| cis-aconitate | – | – |
| isocitrate | CitM | CitH and YraO |
| α -ketoglutarate | – | – |
| succinate | DctP | YfIS |
| fumarate | DctP | YfIS |
| malate | MaeN, MleN, DctP and CimH | YfIS |
| oxaloacetate | – | – |

Na-lactate. MleN is partly responsible for growth on L-malate under these specific circumstances, but its main function is thought to be in Na⁺ resistance under conditions of low pmf (Wei et al. 2000). Finally, YfjS is potentially involved in malate transport since it is homologous to SOT1 found in chloroplasts of *Spinacia oleracea*, which transports, amongst others, L-malate.

In our survey of transporters of TCA cycle intermediates in *B. subtilis* we did not identify any putative or known transport systems for oxaloacetate, cis-aconitate and α -ketoglutarate. The overview provided in the paper shows clearly that *B. subtilis* is capable of using a range of di- and tricarboxylates. This can be explained by the fact that *B. subtilis* is a soil bacterium that has to be able to survive under very different growth conditions utilizing every carbon and energy source it may encounter.

Acknowledgements

This research was supported by the Ministry of Economic Affairs, in the framework of the IOP Milieutechnologie/Zware metalen IZW97404.

References

- Altschul S.F., Madden T.L., Schaffer A.A., Zhang J., Zhang Z., Miller W. et al. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* 25: 3389–3402.
- Asai K., Baik S.H., Kasahara Y., Moriya S. and Ogasawara N. 2000. Regulation of the transport system for C₄-dicarboxylic acids in *Bacillus subtilis*. *Microbiology* 146: 263–271.
- Bandell M., Ansanay V., Rachidi N., Dequin S. and Lolkema J.S. 1997. Membrane potential-generating malate (MleP) and citrate (CitP) transporters of lactic acid bacteria are homologous proteins. Substrate specificity of the 2-hydroxycarboxylate transporter family. *J. Biol. Chem.* 272: 18140–18146.
- Bandell M. and Lolkema J.S. 2000a. Arg425 of the citrate transporter CitP is responsible for high affinity binding of di- and tricarboxylates. *J. Biol. Chem.* 275: 39130–39136.
- Bandell M. and Lolkema J.S. 2000b. The conserved C-terminus of the citrate (CitP) and malate (MleP) transporters of lactic acid bacteria is involved in substrate recognition. *Biochemistry* 39: 13059–13067.
- Bergsma J. and Konings W.N. 1983. The properties of citrate transport in membrane vesicles from *Bacillus subtilis*. *Eur. J. Biochem.* 134: 151–156.
- Bisschop A., Doddema H. and Konings W.N. 1975. Dicarboxylic acid transport in membrane vesicles from *Bacillus subtilis*. *J. Bacteriol.* 124: 613–622.
- Boorsma A., van der Rest M.E., Lolkema J.S. and Konings W.N. 1996. Secondary transporters for citrate and the Mg²⁺-citrate complex in *Bacillus subtilis* are homologous proteins. *J. Bacteriol.* 178: 6216–6222.
- Bott M. 1997. Anaerobic citrate metabolism and its regulation in enterobacteria. *Arch. Microbiol.* 167: 78–88.
- Dimroth P., Jockel P. and Schmid M. 2001. Coupling mechanism of the oxaloacetate decarboxylase Na⁺ pump. *Biochim. Biophys. Acta* 1505: 1–14.
- Ghei O.K. and Kay W.W. 1972. A dicarboxylic acid transporter system in *Bacillus subtilis*. *FEBS Letters* 20: 137–140.
- Ghei O.K. and Kay W.W. 1973. Properties of an inducible C₄-dicarboxylic acid transport system in *Bacillus subtilis*. *J. Bacteriol.* 114: 65–79.
- Goel A., Lee J., Domach M.M. and Ataai M.M. 1995. Suppressed acid formation by cofeeding of glucose and citrate in *Bacillus* cultures: emergence of pyruvate kinase as a potential metabolic engineering site. *Biotechnol. Prog.* 11: 380–385.
- Grunewald M., Bendahan A. and Kanner B.I. 1998. Biotinylation of single cysteine mutants of the glutamate transporter GLT-1 from rat brain reveals its unusual topology. *Neuron* 21: 623–632.
- Hueck C.J. and Hillen W. 1995. Catabolite repression in *Bacillus subtilis*: a global regulatory mechanism for the gram-positive bacteria? *Mol. Microbiol.* 15: 395–401.
- Ito M., Guffanti A.A., Zemsy J., Ivey D.M. and Krulwich T.A. 1997. Role of the *nhaC*-encoded Na⁺/H⁺ antiporter of alkaliphilic *Bacillus firmus* OF4. *J. Bacteriol.* 179: 3851–3857.
- Janausch I.G., Zientz E., Tran Q.H., Kroger A. and Uden G. 2002. C₄-dicarboxylate carriers and sensors in bacteria. *Biochim. Biophys. Acta* 1553: 39–56.
- Jeanmougin F., Thompson J.D., Gouy M., Higgins D.G. and Gibson T.J. 1998. Multiple sequence alignment with Clustal X. *Trends Biochem. Sci.* 23: 403–405.
- Jording D. and Puhler A. 1993. The membrane topology of the *Rhizobium meliloti* C₄-dicarboxylate permease (DctA) as derived from protein fusions with *Escherichia coli* K12 alkaline phosphatase (PhoA) and beta-galactosidase (LacZ). *Mol. Gen. Genet.* 241: 106–114.
- Kaspar S., Perozzo R., Reinelt S., Meyer M., Pfister K., Scapozza L. et al. 1999. The periplasmic domain of the histidine autokinase CitA functions as a highly specific citrate receptor. *Mol. Microbiol.* 33: 858–872.
- Kästner C.N., Schneider K., Dimroth P. and Pos K.M. 2002. Characterization of the citrate/acetate antiporter CitW of *Klebsiella pneumoniae*. *Arch. Microbiol.* 177: 500–506.
- Kawai S., Suzuki H., Yamamoto K. and Kumagai H. 1997. Characterization of the L-malate permease gene (*maeP*) of *Streptococcus bovis* ATCC 15352. *J. Bacteriol.* 179: 4056–4060.
- Kay W.W. 1981. Purification by affinity chromatography of a membrane dicarboxylate binding protein from *Bacillus subtilis*. *Can. J. Microbiol.* 27: 795–800.
- Kelly D.J. and Thomas G.H. 2001. The tripartite ATP-independent periplasmic (TRAP) transporters of bacteria and archaea. *FEMS Microbiol. Rev.* 25: 205–424.
- Kobayashi K., Ogura M., Yamaguchi H., Yoshida K., Ogasawara N., Tanaka T. et al. 2001. Comprehensive DNA microarray analysis of *Bacillus subtilis* two-component regulatory systems. *J. Bacteriol.* 183: 7365–7370.
- Kroger A., Biel S., Simon J., Gross R., Uden G. and Lancaster C.R. 2002. Fumarate respiration of *Wolinella succinogenes*: enzymology, energetics and coupling mechanism. *Biochim. Biophys. Acta* 1553: 23–38.
- Krom B.P., Warner J.B., Konings W.N. and Lolkema J.S. 2000.

- Complementary metal ion specificity of the metal-citrate transporters CitM and CitH of *Bacillus subtilis*. *J. Bacteriol.* 182: 6374–6381.
- Krom B.P., Aardema R. and Lolkema J.S. 2001. *Bacillus subtilis* YxkJ is a secondary transporter of the 2-hydroxycarboxylate transporter family that transports L-malate and citrate. *J. Bacteriol.* 183: 5862–5869.
- Krom B.P., Huttinga H., Warner J.B. and Lolkema J.S. 2002. Impact of the Mg^{2+} -citrate transporter CitM on heavy metal toxicity in *Bacillus subtilis*. *Arch. Microbiol.* 178: 370–375.
- Krom B.P. and Lolkema J.S. 2003. Conserved residues R420 and Q428 in a cytoplasmic loop of the citrate/malate transporter CimH of *Bacillus subtilis* are accessible from the external face of the membrane. *Biochemistry* 42: 467–474.
- Kunst F., Ogasawara N., Moszer I., Albertini A.M., Alloni G., Azevedo V. et al. 1997. The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*. *Nature* 390: 249–256.
- Lolkema J.S., Poolman B. and Konings W.N. 1996. Secondary transporters and metabolic energy generation in bacteria. In: Konings W.N., Kaback H.R. and Lolkema J.S. (eds), *Transport processes in eukaryotic and prokaryotic organisms*. Elsevier, Amsterdam, pp. 229–260.
- Magni C., de Mendoza D., Konings W.N. and Lolkema J.S. 1999. Mechanism of citrate metabolism in *Lactococcus lactis*: resistance against lactate toxicity at low pH. *J. Bacteriol.* 181: 1451–1457.
- Marty-Teyssset C., Lolkema J.S., Schmitt P., Divies C. and Konings W.N. 1995. Membrane potential-generating transport of citrate and malate catalyzed by CitP of *Leuconostoc mesenteroides*. *J. Biol. Chem.* 270: 25370–25376.
- Marty-Teyssset C., Posthuma C., Lolkema J.S., Schmitt P., Divies C. and Konings W.N. 1996. Proton motive force generation by citrolactic fermentation in *Leuconostoc mesenteroides*. *J. Bacteriol.* 178: 2178–2185.
- Miwa Y., Nakata A., Ogiwara A., Yamamoto M. and Fujita Y. 2000. Evaluation and characterization of catabolite-responsive elements (*cre*) of *Bacillus subtilis*. *Nucleic Acids Res.* 28: 1206–1210.
- Nakano M.M. and Zuber P. 1998. Anaerobic growth of a “strict aerobe” (*Bacillus subtilis*). *Annu. Rev. Microbiol.* 52: 165–190.
- Oehr P. and Willecke K. 1974. Citrate- Mg^{2+} transport in *Bacillus subtilis*. Studies with 2-fluoro-L-erythro-citrate as a substrate. *J. Biol. Chem.* 249: 2037–2042.
- Padan E. and Schuldiner S. 1994. Molecular physiology of the Na^+/H^+ antiporter in *Escherichia coli*. *J. Exp. Biol.* 196: 443–456.
- Padan E., Venturi M., Gerchman Y. and Dover N. 2001. Na^+/H^+ antiporters. *Biochim. Biophys. Acta.* 1505: 144–157.
- Poolman B., Molenaar D., Smid E.J., Ubbink T., Abee T., Renault P.P. et al. 1991. Malolactic fermentation: electrogenic malate uptake and malate/lactate antiport generate metabolic energy. *J. Bacteriol.* 173: 6030–6037.
- Pos K.M., Dimroth P. and Bott M. 1998. The *Escherichia coli* citrate carrier CitT: a member of a novel eubacterial transporter family related to the 2-oxoglutarate/malate translocator from spinach chloroplasts. *J. Bacteriol.* 180: 4160–4165.
- Rabus R., Jack D.L., Kelly D.J. and Saier M.H. Jr. 1999. TRAP transporters: an ancient family of extracytoplasmic solute-receptor-dependent secondary active transporters. *Microbiology* 145: 3431–3445.
- Ramanan N. and Wang Y. 2000. A high-affinity iron permease essential for *Candida albicans* virulence. *Science* 288: 1062–1064.
- Saier M.H. Jr. 2000. A functional-phylogenetic classification system for transmembrane solute transporters. *Microbiol. Mol. Biol. Rev.* 64: 354–411.
- Saier M.H. Jr., Goldman S.R., Maile R.R., Moreno M.S., Weyler W., Yang N. et al. 2002. Transport capabilities encoded within the *Bacillus subtilis* genome. *J. Mol. Microbiol. Biotechnol.* 4: 37–67.
- Slotboom D.J., Konings W.N. and Lolkema J.S. 1999a. Structural features of the glutamate transporter family. *Microbiol. Mol. Biol. Rev.* 63: 293–307.
- Slotboom D.J., Sobczak I., Konings W.N. and Lolkema J.S. 1999b. A conserved serine-rich stretch in the glutamate transporter family forms a substrate-sensitive reentrant loop. *Proc. Natl. Acad. Sci. U.S.A.* 96: 14282–14287.
- Slotboom D.J., Konings W.N. and Lolkema J.S. 2001. The structure of glutamate transporters shows channel-like features. *FEBS Lett.* 492: 183–186.
- van der Rest M.E., Siewe R.M., Abee T., Schwarz E., Oesterheld D. and Konings W.N. 1992. Nucleotide sequence and functional properties of a sodium-dependent citrate transport system from *Klebsiella pneumoniae*. *J. Biol. Chem.* 267: 8971–8976.
- van Geest M. and Lolkema J.S. 2000. Membrane topology of the Na^+ /citrate transporter CitS of *Klebsiella pneumoniae* by insertion mutagenesis. *Biochim. Biophys. Acta* 1466: 328–338.
- van Geest M., Nilsson I., von Heijne G. and Lolkema J.S. 1999. Insertion of a bacterial secondary transport protein in the endoplasmic reticulum membrane. *J. Biol. Chem.* 274: 2816–2823.
- Warner J.B., Krom B.P., Magni C., Konings W.N. and Lolkema J.S. 2000. Catabolite repression and induction of the Mg^{2+} -citrate transporter CitM of *Bacillus subtilis*. *J. Bacteriol.* 182: 6099–6105.
- Warner J.B. and Lolkema J.S. 2002a. LacZ-promoter fusions: the effect of growth. *Microbiology* 148: 1241–1243.
- Warner J.B. and Lolkema J.S. 2002b. Growth of *Bacillus subtilis* on citrate and isocitrate is supported by the Mg^{2+} -citrate transporter CitM. *Microbiology* 148: 3413–3421.
- Warner J.B. and Lolkema J.S. 2003. CcpA independent regulation of expression of the Mg^{2+} -citrate transporter *citM* gene by arginine metabolism in *Bacillus subtilis*. *J. Bacteriol.* 185: 854–859.
- Weber A., Menzlaff E., Arbingner B., Gutensohn M., Eckerskorn C. and Flugge U.I. 1995. The 2-oxoglutarate/malate translocator of chloroplast envelope membranes: molecular cloning of a transporter containing a 12-helix motif and expression of the functional protein in yeast cells. *Biochemistry* 34: 2621–2627.
- Wei Y., Guffanti A.A., Ito M. and Krulwich T.A. 2000. *Bacillus subtilis* YqkI is a novel malic/ Na^+ -lactate antiporter that enhances growth on malate at low protonmotive force. *J. Biol. Chem.* 275: 30287–30292.
- Weickert M.J. and Chambliss G.H. 1990. Site-directed mutagenesis of a catabolite repression operator sequence in *Bacillus subtilis*. *Proc. Natl. Acad. Sci. U.S.A.* 87: 6238–6242.
- Willecke K. and Lange R. 1974. C_4 -dicarboxylate transport in *Bacillus subtilis* studied with 3-fluoro-L-erythro-malate as a substrate. *J. Bacteriol.* 117: 373–378.
- Yamamoto H., Murata M. and Sekiguchi J. 2000. The CitST two-component system regulates the expression of the Mg -citrate transporter in *Bacillus subtilis*. *Mol. Microbiol.* 37: 898–912.