

University of Groningen

Gene expression in Hodgkin lymphoma. The role of miR-155 in normal and malignant B cells

Kluiver, Joost Laurens

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kluiver, J. L. (2006). *Gene expression in Hodgkin lymphoma. The role of miR-155 in normal and malignant B cells*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Papers

- (1) Maggio EM, van den Berg A, Visser L, Diepstra A, Kluiver J, Emmens R, et al. Common and differential chemokine expression patterns in rs cells of NLP, EBV positive and negative classical Hodgkin lymphomas. *Int J Cancer* 2002 Jun 10;99(5):665-72.
- (2) Maggio E, van den Berg A, Diepstra A, Kluiver J, Visser L, Poppema S. Chemokines, cytokines and their receptors in Hodgkin's lymphoma cell lines and tissues. *Ann Oncol* 2002;13 Suppl 1:52-6.
- (3) van den Berg A, Kroesen BJ, Kooistra K, de Jong D, Briggs J, Blokzijl T, et al. High expression of B-cell receptor inducible gene BIC in all subtypes of Hodgkin lymphoma. *Genes Chromosomes Cancer* 2003 May;37(1):20-8.
- (4) Kluiver J, Poppema S, de Jong D, Blokzijl T, Harms G, Jacobs S, et al. BIC and miR-155 are highly expressed in Hodgkin, primary mediastinal and diffuse large B cell lymphomas. *J Pathol* 2005 Oct;207(2):243-9.
- (5) Rust R, Harms G, Blokzijl T, Boot M, Diepstra A, Kluiver J, et al. High expression of Mcl-1 in ALK positive and negative anaplastic large cell lymphoma. *J Clin Pathol* 2005 May;58(5):520-4.
- (6) Poppema S, Kluiver JL, Atayar C, van den Berg A, Rosenwald A, Hummel M, et al. Report: workshop on mediastinal grey zone lymphoma. *Eur J Haematol Suppl* 2005 Jul;(66):45-52.
- (7) Kluiver J, Haralambieva E, de Jong D, Blokzijl T, Jacobs S, Kroesen BJ, et al. Lack of BIC and microRNA miR-155 expression in primary cases of Burkitt lymphoma. *Genes Chromosomes Cancer* 2006 Feb;45(2):147-53.
- (8) Atayar C, Kok K, Kluiver J, Bosga A, van den Berg E, van der Vlies P, et al. BCL6 ABR break and homozygous deletion of 17q24 in the nodular lymphocyte predominance type Hodgkin lymphoma derived cell line DEV. *Hum Pathol* 2006;in press.
- (9) Rust R, Kluiver J, Visser L, Harms G, Blokzijl T, Kamps WA, et al. Gene expression study in dendritic/Langerhans cells and Langerhans cell histiocytosis cases. *J Pathol* 2006;in press.
- (10) Kluiver J, Kok K, de Jong D, Blokzijl T, Harms G, van der Vlies P, et al. Global correlation of genome and transcriptome changes in classical hodgkin lymphoma. Submitted 2006.
- (11) Kluiver J, Kroesen BJ, Poppema S, van den Berg A. The role of microRNAs in normal hematopoiesis and hematopoietic malignancies. Submitted 2006.
- (12) Kluiver J, van den Berg A, de Jong D, Blokzijl T, Harms G, Bouwman E, et al. Regulation of primary microRNA BIC transcription and processing in Burkitt lymphoma. Submitted 2006.

Abstracts

- (1) Kluiver J, Schwering I, de Jong D, Küppers R, Poppema S, van den Berg A. Serial analysis of gene expression in Hodgkin lymphoma. *Eur J Haematol Suppl*, 6th International Symposium on Hodgkin Lymphoma, Cologne, Germany, 2004 Sep 18;73:B16.
- (2) Kluiver J, van den Berg A, Blokzijl T, Harms G, de Jong D, Atayar C, et al. TARC and BIC are expressed in a subset of primary mediastinal B-cell lymphomas. *Eur J Haematol Suppl*, 6th International Symposium on Hodgkin Lymphoma, Cologne, Germany, 2004 Sep 18;73:B15.
- (3) van den Berg A, Kluiver J, de Jong D, Koekoek T, Blokzijl T, Poppema S, et al. High expression of microRNA BIC / miR-155 in all subtypes of Hodgkin lymphoma. *Blood*, 46th annual symposium of the American Society of Hematology, San Diego, USA, 2004 Nov 16;104: 429.
- (4) Kluiver J, Schwering I, de Jong D, Küppers R, Poppema S, van den Berg A. Serial analysis of gene expression revealed consistent downregulation of more than 100 genes in Hodgkin lymphoma. *Blood*, 46th annual symposium of the American Society of Hematology, San Diego, USA, 2004 Nov 16;104: 2488.
- (5) Kluiver J, Poppema S, de Jong D, Blokzijl T, Harms G, Jacobs S, et al. BIC and miR-155 are highly expressed in Hodgkin, primary mediastinal and diffuse large B-cell lymphomas. *Blood*, 47th annual symposium of the American Society of Hematology, San Diego, USA, 2005 Nov 16;106: 970.
- (6) Kluiver J, Haralambieva E, de Jong D, Blokzijl T, Jacobs S, Kroesen BJ, et al. Lack of BIC and microRNA miR-155 expression in primary cases of Burkitt lymphoma. *Blood*, 47th annual symposium of the American Society of Hematology, San Diego, USA, 2005 Nov 16;106: 1922.

