

University of Groningen

## Implantology and the severely resorbed edentulous mandible

Stellingsma, C.; Vissink, A.; Meijer, H.J.A.; Raghoobar, G.M.

*Published in:*  
Critical reviews in oral biology & medicine

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2004

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Stellingsma, C., Vissink, A., Meijer, H. J. A., & Raghoobar, G. M. (2004). Implantology and the severely resorbed edentulous mandible. *Critical reviews in oral biology & medicine*, 15(4), 240-248.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# IMPLANTOLOGY AND THE SEVERELY RESORBED EDENTULOUS MANDIBLE

C. Stellingsma\*

A. Vissink

H.J.A. Meijer<sup>1</sup>

C. Kuiper

G.M. Raghoobar

Department of Oral and Maxillofacial Surgery and Maxillofacial Prosthetics, University Hospital, PO Box 30.001, 9700 RB Groningen, the Netherlands; and <sup>1</sup>Department of Oral Function and Prosthetic Dentistry, Faculty of Medical Sciences, University of Groningen; \*corresponding author, c.stellingsma@kchir.azg.nl

**ABSTRACT:** Patients with a severely resorbed edentulous mandible often suffer from problems with the lower denture. These problems include: insufficient retention of the lower denture, intolerance to loading by the mucosa, pain, difficulties with eating and speech, loss of soft-tissue support, and altered facial appearance. These problems are a challenge for the prosthodontist and surgeon. Dental implants have been shown to provide a reliable basis for fixed and removable prostheses. This has resulted in a drastic change in the treatment concepts for management of the severely resorbed edentulous mandible. Reconstructive, pre-prosthetic surgery has changed from surgery aimed to provide a sufficient osseous and mucosal support for a conventional denture into surgery aimed to provide a sufficient bone volume enabling implants to be placed at the most optimal positions from a prosthetic point of view. The aim of this paper is to review critically the literature on procedures related to the severely resorbed edentulous mandible and dental implant treatment. The study includes the transmandibular implant, (short) endosseous implants, and reconstructive procedures such as distraction osteogenesis, augmentation of the mandibular ridge with autogenous bone, and bone substitutes followed by the placement of implants. The number of patients participating in a study, the follow-up period, the design of the study, the degree of mandibular resorption, and the survival rate of the dental implants all are considered evaluation parameters. Although numerous studies have described the outcome results of dental implants in the edentulous mandible, there have been few prospective studies designed as randomized clinical trials that compare different treatment modalities to restore the severely resorbed mandible. Therefore, it is not yet possible to select an evidence-based treatment modality. Future research has to be focused on long-term, detailed follow-up clinical trials before scientifically based decisions in treating these patients can be made. This will contribute to a higher level of care in this field.

**Key words.** Dental implant, mandible, edentulous patient, resorption, denture problems.

## Introduction

Severe atrophy of the inferior alveolar process and underlying basal bone often results in problems with a lower denture. These problems include insufficient retention of the lower denture, intolerance to loading by the mucosa, pain, difficulties with eating and speech, loss of soft-tissue support, and altered facial appearance. These problems are a challenge for the prosthodontist and surgeon.

In the decades preceding the broad clinical use of dental implants, numerous surgical techniques were developed to improve the starting point for successful prosthetic rehabilitation of the edentulous patient (Jennings, 1989). The main techniques were sulcoplasties (Hillerup, 1979, 1994; Davis and Davis, 1995) and grafting procedures (Härle, 1975; Curtis and Ware, 1977; De Koomen *et al.*, 1979; Lekkas and Wes, 1981; Peterson, 1983; Kent and Jarcho, 1995). Although these techniques provided an enlarged denture-bearing area, thereby contributing to improvement of the retention and stability of the lower denture, most of these techniques only temporarily improved the retention and stability of the lower denture. In addition, a considerable rate of morbidity had to be dealt with (Stoelinga *et al.*, 1986).

Since dental implants have been shown to provide a reliable basis for fixed and removable prostheses, reconstructive pre-prosthetic surgery has changed from surgery aimed to provide a sufficient osseous and mucosal support for a conventional denture into surgery aimed to provide a sufficient bone volume to enable implants to be placed at the most optimal

positions from a prosthetic point of view. This treatment is generally accepted for the moderate to severely resorbed edentulous mandible. However, the use of implants in the extremely resorbed mandible, and the selection of a reconstructive surgical procedure to facilitate reliable placement of implants in such a resorbed mandible are still subjects of discussion in the literature. Therefore, the aim of this paper was to review critically the literature on procedures related to implant treatment of the edentulous mandible, with special emphasis on the extremely resorbed edentulous mandible. In the present discussion, an extremely resorbed edentulous mandible is defined as a mandibular height in the symphyseal area of 12 mm or less as measured on a standardized lateral cephalogram.

## Definition of Dental Implants

Dental implants, as discussed in this review, are prosthetic devices of alloplastic material implanted into the oral tissues beneath the mucosal and/or periosteal layer, and on/or within the bone, to provide retention and support for a fixed or a removable prosthesis. Although dental implants may be classified by their silhouette or geometric form (*i.e.*, fin, screw, cylinder, basket, root-form), in this review dental implants will be discussed according to their anchorage component (the dental implant body) as this relates to the bone that provides support and stability (Van Blarcom, 1999). The three basic types that will be discussed in this review are eposteal dental implants, transosteal dental implants, and endosteal dental implants.

## Search of the Literature

This paper provides a comprehensive review of human studies published in international peer-reviewed literature up to May, 2003, regarding procedures related to implant treatment of the severely resorbed edentulous mandible. A MEDLINE search was completed along with a manual search to locate relevant literature. Publications presented in abstract form were ignored. The following MESH terms for the search in MEDLINE were used: dental implants, edentulous mandible, augmentation, distraction, atrophied, and transmandibular. The number of patients participating in a study, follow-up period, the design of the study, degree of mandibular resorption, and the survival rate of the implants were all considered evaluation parameters.

## Success vs. Survival

In evaluation studies, the mere presence of dental implants in the oral cavity can be defined as 'survival' of the implant. When certain other criteria—such as radiographic aspects, clinical mobility, and functional aspects—are taken into consideration, the extent of 'success' can be defined. These 'success' criteria are not uniform, and several criteria have been proposed (Albrektsson *et al.*, 1986; van Steenberghe, 1997). Because the majority of evaluation studies concerning the use of dental implants in the severely resorbed edentulous mandible do not use the same 'success' criteria, and other studies merely report 'survival' percentages, comparison of studies is hindered. For that reason, study 'survival' percentages will be given in the present discussion, unless stated otherwise.

## Epoosteal Dental Implants

Epoosteal implants are dental implants that receive their primary bone support by leaning on the residual bone of the mandible. The subperiosteal dental implant, also known as the subperiosteal frame, is the epoosteal implant system most used in this category (Van Blarcom, 1999). Other, rarely used, systems are the intramucosal inserts and ramus frames (Kerley *et al.*, 1981; Worthington and Rubenstein, 1998). These earlier forms of implant rehabilitation could be successful, but long-term studies are lacking. For that reason it can be assumed that this therapy is either not successful in the long-term, is used by few clinicians, or is surpassed by new materials and new techniques.

The subperiosteal frame was introduced by Dahl in 1943. The technique was refined by Goldberg (Goldberg and Gershkoff, 1949), and later on by Linkow (Linkow *et al.*, 1998). Patients had to undergo two surgical interventions. During the first operation, the surgeon uncovered the bony edentulous alveolar process and the surrounding basal mandibular bone by raising a mucoperiosteal flap. Subsequently, an impression was made of the denture-bearing area. A custom-made frame, made of a cobalt-chromium alloy, was placed subperiosteally during the second operation. Fixed or removable prostheses could be connected to several transmucosal posts.

The concept of a subperiosteal frame was innovative. Non-biological materials were inserted into human tissues with open communication with the oral environment, thereby creating transmucosal posts. Fixed or removable prostheses could be anchored to these posts. Although retrospective studies reported ten-year survival rates of between 60% and 75% (Young *et al.*, 1983), various structural problems were clinically experienced, including epithelial ingrowth, dehiscence of the implant, infection, and paresthesia of the mental nerve (Garefis, 1978; Bodine

*et al.*, 1996). These major drawbacks resulted in removal of the implant in more than 60% of the patients examined during a 20-year follow-up study (Yanase *et al.*, 1994), although cases have been reported with long-term (over 25 years) success (Kurtzman and Schwartz, 1995; Morrow *et al.*, 2000).

Specific information about the clinical performance of this system in relation to the level of resorption of the inferior alveolar process or comparison with other implant systems is, to the best of our knowledge, not available in the literature. Today, the technique of applying a subperiosteal frame has been practically abandoned, because successful rehabilitation in a high proportion of the cases over a long-term period is apparently inferior to that achieved with other systems, such as transosseous and endosseous systems (Adell *et al.*, 1981). Moreover, the morbidity of the transosseous and endosseous systems is less than that of the subperiosteal frame.

## Transosteal Dental Implants

Transosteal or transosseous dental implants are implants composed of a metal plate and transosteal pins or posts. The metal plate is held with retentive pins or screws fixed to the inferior border of the mandible. This metal plate supports the transosteal pins/posts that penetrate the full thickness of the mandible and project into the mouth in the inter-foraminal area (Van Blarcom, 1999). The transosseous dental implants used in humans are the 'staple bone implant' system and the 'transmandibular implant' system (TMI).

### STAPLE BONE IMPLANT SYSTEM

The staple bone implant system was developed as an alternative to subperiosteal frames, because of the major complications that were encountered in the clinical application of subperiosteal frames (Small, 1975, 1980). The main objectives in designing the staple bone implant system were to reduce forces on the implant and to make thin transmucosal perforations. To prevent overloading of this implant system, a tissue-borne overdenture has to be made with stress-breaking attachments to stabilize the denture. The staple bone implant consists of a baseplate with two or four (parallel) transosseous pins and from two to five retentive pins (or screws) to stabilize the baseplate to the inferior border. The implant is made of a titanium alloy to allow for osseointegration (Small *et al.*, 1995).

The mandibular staple bone implant has been evaluated in several retrospective studies that have reported survival rates of between 86% and 100% (Small and Misiak, 1986; Small, 1993; Meijer *et al.*, 1998). The most common complications are gingival hyperplasia, crestal bone loss, and infections around the transmucosal part of the transosseous pins. Serious, but rarely observed, complications are fracture or mobility of a transosseous pin, and fracture of the mandible. The transosseous pins, rigidly connected to the baseplate, are always parallel to each other. This makes prosthetics in general easier, but limits this procedure's application in compromised situations, because individual angulation of the pins is not possible. In contrast, individual placement of transosteal posts is possible in the TMI system, making placement of the TMI in compromised situations possible. Another disadvantage of the staple bone implant system is that it is difficult to remove, because of the bony integration of the mushroom-shaped retentive pins. Although no specific study has been performed concerning the survival of the mandibular staple bone implant in the extremely resorbed mandible, Meijer *et al.* (1998) reported a tendency of

higher failure rates in patients with a mandibular height less than 12 mm. A similar tendency to increased failure rate in a severely resorbed mandible has been reported for the transmandibular implant system (Versteegh *et al.*, 1995).

### TRANSMANDIBULAR IMPLANT SYSTEM

The transmandibular implant system (TMI) was especially developed for the extremely atrophied mandible (Bosker, 1986). Although in both the original and subsequent reports on TMI research the term 'extremely atrophied mandible' was never defined, the majority of the patients included in these studies had an anterior mandibular bone height of less than 12 mm (Bosker and Van Dijk, 1989; Maxson *et al.*, 1989; Bosker *et al.*, 1991b). The TMI consists of a baseplate, five cortical screws, and four transosseous posts. In contrast to most other implant systems made of a titanium alloy, all TMI components are made of Implanor® (Cendres et Métaux, Bien-Brille, Switzerland), a gold alloy containing 70% gold, 5% platinum, 12.8% silver, and 12.2% copper. It is claimed that this bioinert material osseointegrates into human bone, although histologic studies are, to our knowledge, limited to animal studies (Arvier *et al.*, 1989).

Like the staple bone implant system, the TMI is inserted by an extra-oral approach while the patient is under general anesthesia. The baseplate is fixed to the inferior border of the mandible with the cortical screws. The transosseous posts, connected to the baseplate, perforate the mandible and the oral mucosa and are post-operatively connected to each other with a bar equipped with two distal cantilevers (Powers *et al.*, 1994). Three months after placement, an implant-supported overdenture is usually constructed.

The TMI has been frequently evaluated (Bosker and Van Dijk, 1989; Maxson *et al.*, 1989; Bosker *et al.*, 1991b; Versteegh *et al.*, 1995; Kwakman *et al.*, 1996; Meijer *et al.*, 2001; Verhoeven *et al.*, 2001; Paton *et al.*, 2002). High survival rates (95%-100%) have been reported (Bosker and Van Dijk, 1989; Maxson *et al.*, 1989; Powers *et al.*, 1989; Bosker *et al.*, 1991b), but other studies report lower (56%-75%) survival rates (Versteegh *et al.*, 1995; Kwakman *et al.*, 1996; Meijer *et al.*, 2001; Verhoeven *et al.*, 2001; Paton *et al.*, 2002). These differences can be ascribed to the different definitions used for complications and failures in the

various studies and to different protocols for placement of the implant and construction of the overdenture, and could be related to the experiences of the surgical and prosthetic teams (Van Pelt, 1997; Powers, 2001; Paton *et al.*, 2002). An overview of studies that have evaluated the transmandibular implant system is given in Table 1. Complications include infections, loss of osseointegration of the transosseous posts, fracture of posts, hyperplasia of the oral mucosa, peri-implant bone loss, disturbances of the mental nerve, and prosthetic complications such as bar and clip corrections.

Several studies have concluded that the TMI is especially suitable for the extremely resorbed mandible, although these studies do not provide information regarding mandibular height (Bosker and Van Dijk, 1989; Betts *et al.*, 1995). In a comparative retrospective study, however, an opposite conclusion was reached, namely, that the failure rate was higher in extremely resorbed mandibles (bone height less than 12 mm) (Versteegh *et al.*, 1995). A similar tendency was observed for the staple bone implant (Meijer *et al.*, 1998).

To date, two prospective studies have compared the TMI with endosseous implants (Geertman *et al.*, 1996; Stellingsma *et al.*, 2004a,b). With regard to the severely resorbed mandible, the one-year results showed no significant differences between the two systems (Geertman *et al.*, 1996), but thereafter, significantly more complications were reported with the TMI system. These complications included loss of osseointegration, infection, and non-fitting superstructures. After six years, a survival rate of 97% was reported for the endosseous implants *vs.* a survival rate of 72% for the TMI group (Meijer *et al.*, 2001). Also, with regard to the extremely resorbed mandible, short endosseous implants perform significantly better than the transmandibular implant (Stellingsma *et al.*, 2004a,b).

When the prescribed prosthodontic protocol was strictly adhered to, controlled bone growth in the mandible has been reported to occur (Powers *et al.*, 1994; Bosker and Powers, 1995). The explanation given for this phenomenon is that the bending forces in the area distal to the lateral implants that develop during functional loading of the implant might serve as a mechanical stimulus for bone (re)modeling processes. There are reports of bone growth in the areas adjacent to the implant of up to 9 mm

(Bosker *et al.*, 1991a). This so-called 'rejuvenation' has been claimed to occur in large groups of patients (Bosker *et al.*, 1991a; Betts *et al.*, 1993; Powers *et al.*, 1994). However, the evaluation instrument that was used, *i.e.*, panoramic radiographs, is questionable; standardization of the recording technique was not done, and this makes comparison of subsequent radiographs hazardous due to distortion and magnification errors (Batenburg *et al.*, 1997). Moreover, statistical analysis of the data is lacking, and consequently the

**TABLE 1**  
**Overview of the Literature Concerning the Evaluation of the Transmandibular Implant System**

| Author                            | # of Patients | Follow-up Period (yrs) | Retro/Prospective Study | Mandibular Height (mm) | Type of (evaluative) Radiograph | Survival Rate (%)     |
|-----------------------------------|---------------|------------------------|-------------------------|------------------------|---------------------------------|-----------------------|
| Bosker and van Dijk, 1989         | 368           | 0.5-12                 | retrospective           | 4-20                   | panoramic                       | 97 <sup>b</sup>       |
| Maxson <i>et al.</i> , 1989       | 190           | 0.25-5                 | retrospective           | 4-18                   | panoramic                       | 95 <sup>b</sup>       |
| Powers <i>et al.</i> , 1989       | 13            | 2                      | prospective             | — <sup>a</sup>         | panoramic                       | 100 <sup>b</sup>      |
| Bosker <i>et al.</i> , 1991b      | 1356          | 0.5-13                 | retrospective           | 4-24                   | panoramic                       | 96 <sup>b</sup>       |
| Versteegh <i>et al.</i> , 1995    | 37            | 2.3-6.5                | retrospective           | 7.5-16.0               | panoramic                       | 74.8                  |
| Meijer <i>et al.</i> , 2001       | 30            | 6                      | prospective             | 8-15                   | panoramic                       | 72.0                  |
| Verhoeven <i>et al.</i> , 2001    | 70            | 3-13                   | retrospective           | ≤ 14                   | panoramic                       | 84.8-100 <sup>b</sup> |
| Paton <i>et al.</i> , 2002        | 58            | 5-15                   | retrospective           | 5-16                   | panoramic                       | 56                    |
| Stellingsma <i>et al.</i> , 2003b | 20            | 2                      | prospective             | 6-12                   | oblique lateral                 | 93.8                  |

<sup>a</sup> Not specifically stated.

<sup>b</sup> In these studies, replacement of an individual implant post due to, *e.g.*, fracture or loss of osseointegration is considered to be a reversible complication that does not reduce the survival rate, and was therefore not scored as a failure. In the other studies, all reporting lower survival rates, such a complication was scored as a failure.

interpretation of the results is subject to considerable bias. Bone growth has also been reported in other studies, although not to as great an extent as in earlier studies (Kwakman *et al.*, 1997; Verhoeven *et al.*, 2001). Again, this differing result may be due to the different protocols used (Powers, 2001).

In addition to providing retention and stability to the lower denture and therefore rehabilitating masticatory function, it is possible (Bosker and Wardle, 1999) to reconstruct the function and appearance of the lower face following the insertion of the transmandibular implant (Powers and Bosker, 1996). By using an extra-oral approach, the surgeon relocated the position of several facial muscles and, additionally, removed redundant skin and fat. The results were evaluated in 146 patients: The reported satisfaction was claimed to be high, although this subjective increase in satisfaction was not objectively assessed by means of, *e.g.*, a validated questionnaire, and no pre-treatment data were recorded. Thus, there is a high risk of an interpretation bias.

### **Endosteal Dental Implants**

Prior to the evolution of transosseous implant systems, which were exclusively used in the edentulous mandible, there was the development of endosteal implant systems capable of replacing one or more teeth in the partial or complete edentulous mandible or maxilla. An endosteal or endosseous dental implant is a dental implant placed into the alveolar and/or basal bone of the mandible or maxilla and transecting only one cortical plate. With regard to the edentulous mandible, the apical part of the endosseous implant occasionally extends into the caudal cortical plate. An implant placed this way is still considered 'endosseous'. In contrast to transosseous systems, where the retentive components form a unit, these endosseous implants can be regarded as solitary components, so that adjustments or replacement of individual implants is possible. This is a major advantage compared with the transosseous systems where, in cases of complications, a complete revision of the entire system is needed.

The endosseous dental implant is composed of an anchorage component, termed the 'endosseous dental implant body', which ideally is within the bone, and a retentive component, termed the 'endosseous dental implant abutment'. Descriptions of the dental implant body that use silhouette or geometric forms—such as cylinder, conical, screw, or blade—may be used as adjectives to enhance the understanding of the geometry of endosseous dental implants (Van Blarcom, 1999).

The Swedish research group led by Brånemark and the Swiss research group led by Schroeder were the first to study the direct contact between bone and titanium endosseous implants (Brånemark *et al.*, 1969; Schroeder *et al.*, 1976). This direct bone-to-implant contact, a phenomenon called 'osseointegration' (Brånemark *et al.*, 1977), led to the development of various endosseous implant systems that could be used in a clinical setting. It became clear that both the geometric design and the surface conditions of the implant, and a meticulous surgical technique combined with an optimal condition of the implant site were prerequisites for successful osseointegration. Geometric designs were, among other reasons, developed to make possible selection with respect to location and/or application (Buser *et al.*, 1994; Mericske-Stern *et al.*, 2000). This way, one can choose an implant design to reach the optimal conditions for both functional and esthetic rehabilitation of the patient. From studies that have focused on surface properties of endosseous implants, it can be concluded that these properties not only play an important role in qualitative and quantitative

aspects of the bone-implant interface, but are also decisive in the time needed to reach a certain level of osseointegration (Buser *et al.*, 1998; Cochran *et al.*, 2002).

The clinical use for endosseous implants in prosthetic dentistry is obvious. The first clinical results published by Swedish research groups (Brånemark *et al.*, 1977) showed favorable survival rates. After confirmation of these results by Zarb and his co-workers, the use of titanium endosseous implants was widely accepted (Zarb and Symington, 1983). Today, endosseous titanium implants are utilized in both partially and completely edentulous patients.

Application of endosseous implants in the edentulous mandible has changed the treatment concepts enormously; with the use of these implants, it is possible to provide retention for fixed and removable prostheses. This kind of treatment improves oral function and has considerable patient satisfaction (Boerrigter *et al.*, 1995; Bakke *et al.*, 2002).

### **FIXED BRIDGES AND REMOVABLE OVERDENTURES ON ENDOSSEOUS IMPLANTS**

The concept of installing five or six endosseous implants in the interforaminal region, followed by the construction of a fixed bridge, was developed by the Brånemark group and has been evaluated in several studies (Adell *et al.*, 1981; Albrektsson *et al.*, 1986; Naert *et al.*, 1992; Quirynen *et al.*, 1992; Lindquist *et al.*, 1996). Provided that a strict protocol is followed, it is a reliable treatment option, and the survival rates of the endosseous implants are high (between 90 and 98%).

It is not always possible or advisable to install five or six endosseous implants in the edentulous mandible. Therefore, the treatment concept of a removable overdenture anchored to two to four endosseous implants was introduced. The superstructure connecting the implants with the overdenture can be divided into ball attachments, clip-bar attachments, magnet attachments, and a milled bar with precision attachments (Davis and Davis, 1995). Although the differences in functional aspects are minimal, patients prefer implant-supported prostheses (Tang *et al.*, 1997; Van Kampen *et al.*, 2002).

The treatment concept of the mandibular overdenture retained by endosseous implants has been evaluated in several studies (Wismeijer *et al.*, 1995; Batenburg *et al.*, 1998a,b; Sadowsky, 2001). Studies that have focused on clinical behavior and radiological aspects confirm that this kind of treatment is very predictable in showing high survival rates (> 90%) of the implants, along with healthy peri-implant tissues, on condition that a high level of oral hygiene is maintained (Geertman, 1995; Boerrigter *et al.*, 1997; Roynesdal *et al.*, 1998). With respect to patient satisfaction and psychosocial functioning, it is clear that patients regard implant-supported mandibular dentures as very beneficial (Kent, 1992; Bouma *et al.*, 1997; Locker, 1998; Raghoebar *et al.*, 2000b; Stellingsma *et al.*, 2003). These results are comparable with those for implant-supported bridges (Adell *et al.*, 1990; Johns *et al.*, 1992; Hemmings *et al.*, 1994). The choice between a fixed bridge and a removable overdenture in the edentulous mandible is dependent on several factors. Not only are anatomic factors such as the interforaminal space and inter-maxillary relations important, but oral hygiene and speech-related factors play a role as well. Finally, patient-related factors such as costs and the preference for a fixed or removable prosthesis must also be considered.

### **SHORT ENDOSSEOUS IMPLANTS**

The placement of short endosseous implants is another option to

treat the extremely resorbed mandible. In the case of severe ridge atrophy and short implants (< 12 mm), the ratio between implant length and the distance to the occlusal plane is compromised, resulting in unfavorable biomechanics. Since the latter could jeopardize long-term osseointegration, this mode of treatment is not widely used (Brånemark *et al.*, 1985; Worthington, 1992). There are some reports, however, concerning the use of short endosseous implants in the extremely resorbed mandible. The scale of these studies is limited, and comparison with other modalities is limited to the studies by Geertman *et al.* (1996) and Stellingsma *et al.* (2003a,b). Nevertheless, it is an attractive treatment option because of the relatively simple surgical procedure and limited morbidity (Triplett *et al.*, 1991; Keller, 1995; Geertman *et al.*, 1996; Bruggenkate *et al.*, 1998; Friberg *et al.*, 2000; Stellingsma *et al.*, 2000, 2004a,b; Deporter *et al.*, 2002). Survival rates vary from 88 to 100%. Recently, Deporter *et al.* (2002) reported excellent ten-year outcomes from the use of short endosseous implants to support mandibular overdentures: a ten-year implant survival of 92.7% and an average annual bone loss limited to 0.03 mm since the first year of implant placement. Although more randomized clinical trials are needed, survival rates thus far are comparable with those of implants in less severely resorbed edentulous mandibles (Batenburg *et al.*, 1998b). An overview of the literature concerning the use of short endosseous implants in the edentulous mandible is given in Table 2. In a randomized clinical trial that compared three treatment modalities (transmandibular implant, augmentation of the mandible with an autologous bone graft followed by placement of four endosseous implants, and the placement of four short endosseous implants) for the extremely resorbed edentulous mandible, it was concluded that treatment with short endosseous implants is the treatment of choice due to the minimal complications, the high survival rate, the stable bone-implant interface, and the fact that patients can be treated in an outpatient clinic setting (Stellingsma *et al.*, 2004a,b). The major complication of treatment with short endosseous implants is the risk of a (partial) fracture of the mandible (Mason *et al.*, 1990; Triplett *et al.*, 1991). Even though the incidence of this complication was shown to be rare (Raghoebar *et al.*, 2000c), treatment is often difficult if a fracture does occur (Tolman and Keller, 1991).

### Grafting Procedures

In the case of severe atrophy of the edentulous mandible, it is possible to augment the mandible prior to the placement of endosseous implants. Various techniques and materials have

been developed to increase mandibular height. Onlay techniques as well as interposition of the graft in the inter-foraminal area are used. Autogenous materials, such as bone and cartilage, and allogenic materials, such as hydroxyapatite or bone substitutes, as well as combinations of these materials, are used for ridge augmentation (Stoelinga *et al.*, 1986; Vanassche *et al.*, 1988; Tolman, 1995). Depending on the clinical conditions, endosseous implants can be inserted at the same treatment session or after the graft has been incorporated for 3-4 months.

The advantage of a one-stage procedure is that the graft and the implant can be placed at the same time, thereby eliminating a second operation. An important disadvantage is that the positioning and angulation of the implants are more complicated, thereby making this one-stage procedure undesirable from a prosthetic point of view (Bell *et al.*, 2002).

Another drawback of the one-step reconstruction with onlay bone grafts and endosseous implants (Keller and Tolman, 1992; Vermeeren *et al.*, 1996; Verhoeven *et al.*, 1997) is the unpredictable resorption of the grafted bone around the implants (Vermeeren *et al.*, 1996). Resorption of the graft is less extensive than in the onlay technique, when one interposes a bone graft in the inter-foraminal area, in combination with the placement of endosseous implants in a one-stage (Lew *et al.*, 1991; Keller and Tolman, 1992) or a two-stage procedure (Satow *et al.*, 1997; Stellingsma *et al.*, 1998; Bell *et al.*, 2002). In particular, when the two-stage procedure is used, the bone-implant interface can, for the most part, be preserved (Gratz *et al.*, 1994; Stellingsma *et al.*, 1998). Although most studies report the use of an intra-oral grafting approach for the edentulous mandible, a submental extra-oral approach to prevent oral contamination of the graft has also been used (Lew *et al.*, 1991; Bell *et al.*, 2002).

With both onlay or interposed bone grafts, hydroxyapatite can be used as a filler, alongside autologous bone, to achieve the desired volume and contour for the augmented mandible (Haers *et al.*, 1991; McGrath *et al.*, 1996). Complications that are experienced with the use of hydroxyapatite are migration, displacement, and dehiscence of hydroxyapatite particles that can cause mucosal erosions (Kent *et al.*, 1986). Stabilization of endosseous implants with hydroxyapatite as a primary retentive material does not seem adequate; it is just not a substitute for viable bone (Kent and Jarcho, 1995). However, to restore the extremely resorbed mandible, hydroxyapatite can be used posteriorly in the lateral parts, with dental implants placed in the inter-foraminal area (Lew *et al.*, 1991).

The most significant complications that occur following

**TABLE 2**  
**Overview of the Literature Concerning the Use of Short Endosseous Implants in the Edentulous Mandible**

| Author                            | # of Patients/<br># of Implants | Follow-up<br>Period (yrs) | Retro/Prospective<br>Study | Mandibular<br>Height (mm) | Type of (evaluative)<br>Radiograph | Survival<br>Rate (%) | Type of<br>Prosthesis |
|-----------------------------------|---------------------------------|---------------------------|----------------------------|---------------------------|------------------------------------|----------------------|-----------------------|
| Triplett <i>et al.</i> , 1991     | 28/130                          | 1-5                       | retrospective              | < 10                      | — <sup>a</sup>                     | 94                   | fixed, removable      |
| Keller, 1995                      | 52/260                          | 1-10                      | prospective                | < 10                      | panoramic, lateral                 | 93                   | fixed, removable      |
| Kwakman <i>et al.</i> , 1996      | 29/ 58                          | 5                         | prospective                | 8-15                      | panoramic                          | 100                  | removable             |
| Bruggenkate <i>et al.</i> , 1998  | 126/253                         | 1-7                       | retrospective              | — <sup>a</sup>            | panoramic, periapical              | 94                   | fixed, removable      |
| Stellingsma <i>et al.</i> , 2000  | 17/ 68                          | 5                         | retrospective              | < 12                      | oblique lateral                    | 88                   | removable             |
| Friberg <i>et al.</i> , 2000      | 49/260                          | 1-14                      | prospective                | — <sup>a</sup>            | panoramic, Scanora®                | 92.3                 | fixed/removable       |
| Stellingsma <i>et al.</i> , 2003b | 20/ 80                          | 2                         | prospective                | 6-12                      | oblique lateral                    | 100                  | removable             |

<sup>a</sup> Not specifically stated.

grafting procedures in the mandible are sensory disturbances of the mental nerve, wound dehiscence, infections of the grafted area, and, with autogenous bone grafts, donor area morbidity (Tolman, 1995). An overview of the literature concerning augmentation of the edentulous mandible in combination with endosseous implants and an implant-retained mandibular overdenture is given in Table 3. As discussed previously, the placement of short endosseous implants gives results at least as predictable as those achieved with the insertion of longer endosseous implants after a grafting procedure or the use of a transmandibular implant system (Stellingsma *et al.*, 2004a,b). Future randomized clinical trials are needed to prove the hypothesis that short endosseous implants are the treatment of choice to rehabilitate patients with an extremely resorbed edentulous mandible.

### Distraction Osteogenesis

Besides grafting techniques, distraction osteogenesis can be performed to improve the starting point for the placement of implants in the inter-foraminal area of the severely resorbed edentulous mandible (Chin and Toth, 1996; Hidding *et al.*, 1999). Distraction osteogenesis is a technique of gradual bone-lengthening, allowing natural healing mechanisms to generate new bone. When applied to the reconstruction of a severely resorbed edentulous mandible, an osteotomy in the interforaminal area of the mandible is made, after which the distraction device is placed. Five to seven days after surgery, active distraction is started at a rate of 0.5 to 1 mm *per* day. Between four and eight weeks after the last day of active distraction, mineralization of the newly formed bone matrix in the distraction area has progressed sufficiently to allow for the placement of endosseous implants with sufficient primary stability. During the next three months, the implants are left unloaded to allow for further mineralization and remodeling of the distracted area (Raghoobar *et al.*, 2000a, 2002). In comparison with grafting procedures, the advantages of distraction osteogenesis are the absence of donor site morbidity, the presence of vital bone in the distraction area, and the gain of soft tissues. Possible complications of the distraction technique for the edentulous (severely resorbed) mandible are fracture of the mandible, infection, and necrosis of the superior fragment, but such complications are rarely reported in the literature.

The results of distraction osteogenesis have been evaluated in several studies (Urbani *et al.*, 1999; McAllister, 2001; Raghoobar *et al.*, 2002). The short-term clinical, radiographic, and histomorphologic results are very promising. There is some evidence for the assumption that, in the near future, distraction osteogenesis can develop into a reliable tool for augmentation of the anterior segment of a severely resorbed edentulous mandible. However, because long-term results of this technique are still unavailable, some caution still needs to be exercised in the recommendation of this mode of treatment in general practice. Comparison of the distraction method with other techniques—like the placement of a transmandibular implant, augmentation of the edentulous mandible in combination with implants, or the placement of short endosseous implants—will facilitate an assessment of the efficacy of distraction osteogenesis, in combination with endosseous implants in the treatment of the severely resorbed mandible. Such studies have not yet been published.

### Conclusions

Dental implantology has evolved from an experimental to a mature evidence-based discipline. It is currently a valuable treatment modality in the prosthetic treatment of edentulous patients. Numerous techniques have been developed for the use of dental implants for retention and stabilization of fixed or removable mandibular dentures. Today, the options for the restoration of the extremely resorbed mandible with implants can be categorized as follows:

- (1) the use of (short) endosseous implants in combination with either a fixed or removable prosthesis;
- (2) augmentation of the mandible by means of distraction techniques or grafting procedures, followed by the placement of endosseous implants in combination with either a fixed or removable prosthesis; and
- (3) the installation of a transosseous implant system in combination with a removable prosthesis.

Although numerous studies have been published about the outcomes of dental implants in the edentulous mandible, there are still certain questions that have to be answered. For example, comparative studies, designed as prospective clinical trials to evaluate various treatment modalities for the extremely resorbed mandible with dental implants, are still scarce. In fact, they are

**TABLE 3**

#### Overview of the Literature Concerning Augmentation of the Edentulous Mandible in Combination with Endosseous Implants and an Implant-retained Mandibular Overdenture

| Author                            | # of Patients/<br># of Implants | Follow-up<br>Period (yrs) | Retro/Prospective<br>Study | Mandibular<br>Height (mm) | Immediate/Delayed<br>Implant Placement | Type of (evaluative)<br>Radiograph | Survival<br>Rate (%) | Type of<br>Graft                          |
|-----------------------------------|---------------------------------|---------------------------|----------------------------|---------------------------|--|------------------------------------|----------------------|---|
| Lew <i>et al.</i> , 1991          | 10/43                           | 1-3                       | prospective                | 4-8                       | immediate                              | panoramic                          | 93                   | onlay <sup>a</sup> , bone/HA <sup>b</sup> |
| Keller and Tolman, 1992           | 7/32                            | 1-4                       | retrospective              | < 7                       | immediate                              | panoramic                          | 94                   | onlay, bone                               |
| Gratz <i>et al.</i> , 1994        | 23/78                           | < 5                       | prospective                | 15-23                     | immediate                              | lateral                            | 98                   | interposition, bone                       |
| McGrath <i>et al.</i> , 1996      | 18/36                           | 1-3                       | retrospective              | 6-12                      | immediate                              | panoramic                          | 92                   | onlay, bone/HA                            |
| Vermeeren <i>et al.</i> , 1996    | 31/78                           | 5                         | prospective                | < 8                       | immediate                              | panoramic                          | 90                   | onlay, bone/HA                            |
| Verhoeven <i>et al.</i> , 1997    | 13/26                           | 2-4                       | prospective                | 7-12                      | immediate                              | oblique lateral                    | 100                  | onlay, bone                               |
| Satow <i>et al.</i> , 1997        | 32/73                           | 1-7                       | prospective                | < 12                      | delayed                                | lateral, panoramic                 | 95                   | interposition, bone/HA                    |
| Stellingsma <i>et al.</i> , 1998  | 10/40                           | 2-5                       | retrospective              | 9-14                      | delayed                                | panoramic                          | 100                  | interposition, bone                       |
| Bell <i>et al.</i> , 2002         | 14/60                           | 2                         | retrospective              | 6-10                      | delayed                                | panoramic                          | 100                  | onlay <sup>a</sup> , bone                 |
| Stellingsma <i>et al.</i> , 2003b | 20/80                           | 2                         | prospective                | 6-12                      | delayed                                | oblique lateral                    | 88                   | interposition, bone                       |

<sup>a</sup> Onlay graft *via* extra-oral submental approach.

<sup>b</sup> HA = Hydroxyapatite.

limited to the studies by Geertman *et al.* (1996) and Stellingsma *et al.* (2004a,b). Future research concerning implant treatment of the extremely resorbed edentulous mandible should be focused not only on long-term, detailed follow-up clinical trials in which clinical and radiographic aspects are analyzed, but also on the evaluation of the restoration of function and other patient-based parameters. Only by taking all of the factors into account can one arrive at evidence-based decisions in treating these patients, thereby contributing to a higher level of care in this field.

## REFERENCES

- Adell R, Lekholm U, Rockler B, Brånemark PI (1981). A 15-year study of osseointegrated implants in the treatment of the edentulous jaw. *Int J Oral Surg* 10:387-416.
- Adell R, Eriksson B, Lekholm U, Brånemark PI, Jemt T (1990). Long-term follow-up study of osseointegrated implants in the treatment of totally edentulous jaws. *Int J Oral Maxillofac Implants* 5:347-359.
- Albrektsson T, Zarb G, Worthington P, Eriksson AR (1986). The long-term efficacy of currently used dental implants: a review and proposed criteria of success. *Int J Oral Maxillofac Implants* 1:11-25.
- Arvier J, Scott J, Goss A, Wilson D, Tideman H (1989). Biological and clinical evaluation of the transmandibular implant. *Austr Dent J* 34:254-259.
- Bakke M, Holm B, Gotfredsen K (2002). Masticatory function and patient satisfaction with implant-supported mandibular overdentures: a prospective 5-year study. *Int J Prosthodont* 15:575-581.
- Batenburg RH, Stellingsma K, Raghoobar GM, Vissink A (1997). Bone height measurements on panoramic radiographs: the effect of shape and position of edentulous mandibles. *Oral Surg Oral Med Oral Pathol Radiol Endod* 84:430-435.
- Batenburg RHK, Meijer HJA, Raghoobar GM, van Oort RP, Boering G (1998a). Mandibular overdentures supported by two Brånemark, IMZ or ITI Bonefit implants. A prospective comparative study. One year results. *Clin Oral Implant Res* 9:374-383.
- Batenburg RHK, Meijer HJA, Raghoobar GM, Vissink A, van Oort RP, Boering G (1998b). Review of the literature on mandibular overdentures supported by endosseous implants. *Int J Oral Maxillofac Implants* 13:539-545.
- Bell RB, Blakey GH, White RP, Hillebrand DG, Molina A (2002). Staged reconstruction of the severely atrophic mandible with autogenous bone graft and endosteal implants. *J Oral Maxillofac Surg* 60:1135-1141.
- Betts NJ, Barber HD, Powers MP, Wu L, Hennig T, Fonseca RJ (1993). Osseous changes following placement of the transmandibular implant system in edentulous mandibles. *Implant Dent* 2:11-17.
- Betts NJ, Powers MP, Barber HD (1995). Reconstruction of the severely atrophic edentulous mandible with the transmandibular implant system. *J Oral Maxillofac Surg* 53:295-304.
- Bodine RL, Yanase RT, Bodine A (1996). Forty years of experience with subperiosteal implant dentures in 41 edentulous patients. *J Prosthet Dent* 75:33-44.
- Boerrigter EM, Geertman ME, van Oort RP, Bouma J, Raghoobar GM, Van Waas MA, *et al.* (1995). Patient satisfaction with implant-retained mandibular overdentures. A comparison with new complete dentures not retained by implants: a multicentre randomized clinical trial. *Br J Oral Maxillofac Surg* 33:282-288.
- Boerrigter EM, van Oort RP, Raghoobar GM, Stegenga B, Schoen PJ, Boering G (1997). A controlled clinical trial of implant-retained mandibular overdentures: clinical aspects. *J Oral Rehabil* 24:182-190.
- Bosker H (1986). The transmandibular implant (dissertation). Utrecht, The Netherlands: University of Utrecht.
- Bosker H, Powers MP (1995). The transmandibular reconstruction system. In: Reconstructive preprosthetic oral and maxillofacial surgery. Fonseca R, Davis WH, editors. Philadelphia: W.B. Saunders Company, pp. 565-668.
- Bosker H, Van Dijk L (1989). The transmandibular implant: a 12-year follow-up study. *J Oral Maxillofac Surg* 47:442-450.
- Bosker H, Wardle LM (1999). Muscular reconstruction to improve the deterioration of facial appearance and speech caused by mandibular atrophy: technique and case reports. *Br J Oral Maxillofac Surg* 37:277-284.
- Bosker H, Jordan RD, Powers MP, Van Pelt AWJ (1991a). Bone induction and bone loss by use of the TMI. *Oral Surg Diagn* 2:18-26.
- Bosker H, Jordan RD, Sindet-Petersen S, Koole R (1991b). The transmandibular implant: a 13-year survey of its use. *J Oral Maxillofac Surg* 49:482-492.
- Bouma J, Boerrigter EM, van Sonderen E, Boering G (1997). Psychosocial effects of implant-retained overdentures. *Int J Oral Maxillofac Implants* 12:515-522.
- Brånemark PI, Adell R, Breine U, Hansson BO, Lindstrom J, Ohlsson A (1969). Intra-osseous anchorage of dental prostheses. I. Experimental studies. *Scand J Plast Reconstr Surg* 3:81-100.
- Brånemark PI, Hansson BO, Adell R, Breine U, Lindstrom J, Hallen O, *et al.* (1977). Osseointegrated implants in the treatment of the edentulous jaw. Experience from a 10-year period. *Scand J Plast Reconstr Surg* 16:1-132.
- Brånemark PI, Zarb G, Albrektsson T (1985). Tissue-integrated prostheses. Osseointegration in clinical dentistry. Chicago: Quintessence Publishing.
- Bruggenkate CM ten, Asikainen P, Foitzik C, Krekeler G, Sutter F (1998). Short (6-mm) nonsubmerged dental implants: results of a multicenter clinical trial of 1 to 7 years. *Int J Oral Maxillofac Implants* 13:791-798.
- Buser D, Weber HP, Brägger U, Balsiger C (1994). Tissue integration of one-stage implants: three-year results of a prospective longitudinal study with hollow cylinder and hollow screw implants. *Quintessence Int* 25:679-686.
- Buser D, Nydegger T, Hirt HP, Cochran DL, Nolte LP (1998). Removal torque values of titanium implants in the maxilla of miniature pigs. *Int J Oral Maxillofac Implants* 13:611-619.
- Chin M, Toth BA (1996). Distraction osteogenesis in maxillofacial surgery using internal devices: review of five cases. *J Oral Maxillofac Surg* 54:45-53.
- Cochran DL, Buser D, ten Bruggenkate CM, Weingart D, Taylor TM, Bernard JP, *et al.* (2002). The use of reduced healing times on ITI implants with a sandblasted and acid-etched (SLA) surface: early results from clinical trials on ITI SLA implants. *Clin Oral Implants Res* 13:144-153.
- Curtis TA, Ware WH (1977). Autogenous bone graft procedures for atrophic edentulous mandibles. *J Prosthet Dent* 38:366-379.
- Dahl GSA (1943). Om möjligheten för implantation i kaken av metallskelet som das eller retention för fasta eller avtagbara protoster [The possibilities to implant metal skeletons in the jaws for retention of fixed or removable prostheses]. *Odontol Tidskr* 52:440-446.
- Davis WH, Davis CL (1995). Soft-tissue procedures. In: Reconstructive preprosthetic oral and maxillofacial surgery. Fonseca R, Davis WH, editors. Philadelphia: W.B. Saunders Company, pp. 743-820.
- De Koomen HA, Stoeltinga PJ, Tideman H, Huybers TJ (1979). Interposed bone-graft augmentation of the atrophic mandible (a progress report). *J Maxillofac Surg* 7:129-135.
- Deporter D, Watson P, Pharoah M, Todescan R, Tomlinson G (2002). Ten-year results of a prospective study using porous-surfaced dental implants and a mandibular overdenture. *Clin Implant Dent Relat Res* 4:183-189.
- Friberg B, Grondahl K, Lekholm U, Brånemark PI (2000). Long-



- term follow-up of severely atrophic edentulous mandibles reconstructed with short Brånemark implants. *Clin Implant Dent Relat Res* 2:184-189.
- Garefis PN (1978). Complete mandibular subperiosteal implants for edentulous mandibles. *J Prosthet Dent* 39:670-677.
- Geertman ME (1995). Implant-retained mandibular overdentures; clinical evaluation, satisfaction and mastication (dissertation). Nijmegen, The Netherlands: University of Nijmegen.
- Geertman ME, Boerrigter EM, Van Waas MAJ, van Oort RP (1996). Clinical aspects of a multicenter clinical trial of implant-retained mandibular overdentures in patients with severely resorbed mandibles. *J Prosthet Dent* 75:194-204.
- Goldberg NI, Gershkoff A (1949). The implant lower denture. *Dental Digest* 55:490-494.
- Gratz KW, Sailer HF, Oechslin CK (1994). Results after interforaminal mandibular sandwich procedure in combination with titanium screw implants. *Oral Maxillofac Surg North Am* 6:689-698.
- Haers PEJ, van Straaten W, Stoeltinga PJW, De Koomen HA, Blijdorp PA (1991). Reconstruction of the severely resorbed mandible prior to vestibuloplasty or placement of endosseous implants. A 2 to 5 year follow-up. *Int J Oral Maxillofac Surg* 20:149-154.
- Härle F (1975). Visor osteotomy to increase the absolute height of the atrophied mandible. A preliminary report. *J Maxillofac Surg* 3:257-260.
- Hemmings KW, Schmitt A, Zarb GA (1994). Complications and maintenance requirements for fixed prostheses and overdentures in the edentulous mandible: a 5-year report. *Int J Oral Maxillofac Implants* 9:191-196.
- Hidding J, Lazar F, Zoller JE (1999). Erste Ergebnisse bei der vertikalen Distraktionsosteogenese des atrophischen Alveolarkamms [First results of vertical distraction osteogenesis of the atrophied mandible]. *Mund Kiefer Gesichtschir* 3(Suppl 1):79-83.
- Hillerup S (1979). Preprosthetic vestibular sulcus extension by the operation of Edlan and Mejchar. A 2-year follow-up study-I. *Int J Oral Surg* 8:333-339.
- Hillerup S (1994). Preprosthetic surgery in the elderly. *J Prosthet Dent* 72:551-558.
- Jennings DE (1989). Treatment of the mandibular compromised ridge: a literature review. *J Prosthet Dent* 61:575-579.
- Johns RB, Jemt T, Heath MR, Hutton JE (1992). A multicenter study of overdentures supported by Brånemark implants. *Int J Oral Maxillofac Implants* 7:513-522.
- Keller EE (1995). Reconstruction of the severely atrophic edentulous mandible with endosseous implants: a 10-year longitudinal study. *J Oral Maxillofac Surg* 53:305-320.
- Keller EE, Tolman DE (1992). Mandibular ridge augmentation with simultaneous onlay iliac bone graft and endosseous implants: a preliminary report. *Int J Oral Maxillofac Implants* 7:176-184.
- Kent G (1992). Effects of osseointegrated implants on psychological and social well-being: a literature review. *J Prosthet Dent* 68:515-518.
- Kent JN, Jarcho M (1995). Ridge augmentation procedures with hydroxyapatite. In: Reconstructive preprosthetic oral and maxillofacial surgery. Fonseca R, Davis WH, editors. Philadelphia: W.B. Saunders Company, pp. 853-933.
- Kent JN, Finger IM, Quinn JH, Guerra LR (1986). Hydroxylapatite alveolar ridge reconstruction: clinical experiences, complications, and technical modifications. *J Oral Maxillofac Surg* 44:37-49.
- Kerley TR, Phillips RM, Mulherin DR, English J, Schow CE Jr (1981). The ramus frame implant. *J Oral Surg* 39:415-420.
- Kurtzman GM, Schwartz K (1995). The subperiosteal implant as a viable long-term treatment modality in the severely atrophied mandible: a patient's 40-year case history. *J Oral Implantol* 21:35-39.
- Kwakman JM, Voorsmit RACA, Van Waas MAJ, Freihofer HPM, Geertman ME (1996). Transmandibular implant versus intramobile cylinder implants: a randomized, prospective clinical trial. *Int J Oral Maxillofac Surg* 25:433-438.
- Kwakman JM, Van Waas MAJ, Hagens M, Voorsmit RACA (1997). Bone level changes in patients with transmandibular implants. *J Oral Maxillofac Surg* 55:15-18.
- Lekkas K, Wes BJ (1981). Absolute augmentation of the extremely atrophic mandible (a modified technique). *J Maxillofac Surg* 9:103-107.
- Lew D, Hinkle RM, Unhold GP, Shroyer JV, Stutes RD (1991). Reconstruction of the severely atrophic edentulous mandible by means of autogenous bone grafts and simultaneous placement of osseointegrated implants. *J Oral Maxillofac Surg* 49:228-233.
- Lindquist LW, Carlsson GE, Jemt T (1996). A prospective 15-year follow-up study of mandibular fixed prostheses supported by osseointegrated implants. Clinical results and marginal bone loss. *Clin Oral Implants Res* 7:329-336.
- Linkow LI, Wagner JR, Chanavaz M (1998). Tripodal mandibular subperiosteal implant: basic sciences, operational procedures, and clinical data. *J Oral Implantol* 24:16-36.
- Locker D (1998). Patient-based assessments of the outcomes of implant therapy: a review of the literature. *Int J Prosthodont* 11:453-461.
- Mason ME, Triplett RG, Van Sickels JE, Parel SM (1990). Mandibular fractures through endosseous cylinder implants: report of cases and review. *J Oral Maxillofac Surg* 48:311-317.
- Maxson BB, Sindet-Petersen S, Tideman H, Fonseca R, Zijlstra G (1989). Multicenter follow-up study of the transmandibular implant. *J Oral Maxillofac Surg* 47:785-789.
- McAllister BS (2001). Histologic and radiographic evidence of vertical ridge augmentation utilizing distraction osteogenesis: 10 consecutively placed distractors. *J Periodontol* 72:1767-1779.
- McGrath CJR, Schepers SHW, Blijdorp PA, Hoppenreijts TJM, Erbe M (1996). Simultaneous placement of endosteal implants and mandibular onlay grafting for treatment of the atrophic mandible. *Int J Oral Maxillofac Surg* 25:184-188.
- Meijer HJ, van Oort RP, Raghoobar GM, Schoen PJ (1998). The mandibular staple bone plate: a long-term retrospective evaluation. *J Oral Maxillofac Surg* 56:141-145.
- Meijer HJ, Geertman ME, Raghoobar GM, Kwakman JM (2001). Implant-retained mandibular overdentures: 6-year results of a multicenter clinical trial on 3 different implant systems. *J Oral Maxillofac Surg* 59:1260-1268.
- Mericske-Stern RD, Taylor TD, Belser U (2000). Management of the edentulous patient. *Clin Oral Implants Res* 11(Suppl 1):108-125.
- Morrow LA, Smith PW, McCord JF (2000). Case report: restorative maintenance of prostheses stabilised by non-endosseous implants. *Eur J Prosthodont Rest Dent* 8:53-56.
- Naert I, Quirynen M, van Steenberghe D, Darius P (1992). A study of 589 consecutive implants supporting complete fixed prostheses. Part II: Prosthetic aspects. *J Prosthet Dent* 68:949-956.
- Paton G, Fuss J, Goss AN (2002). The transmandibular implant: a 5- and 15-year single-center study. *J Oral Maxillofac Surg* 60:851-857.
- Peterson LJ (1983). Augmentation of the mandibular residual ridge by a modified visor osteotomy. *J Oral Maxillofac Surg* 41:332-338.
- Powers MP (2001). Discussion; implant-retained mandibular overdentures: 6-year results of a multicenter clinical trial on 3 different implant systems. *J Oral Maxillofac Surg* 59:1269.
- Powers MP, Bosker H (1996). Functional and cosmetic reconstruction of the facial lower third associated with placement of the transmandibular implant system. *J Oral Maxillofac Surg* 54:934-942.
- Powers MP, Maxson B, Scott RF, Fonseca R (1989). The transmandibular implant: a 2-year prospective study. *J Oral Maxillofac Surg* 47:679-683.
- Powers MP, Bosker H, van Pelt H, Dunbar N (1994). The transmandibular implant: from progressive bone loss to controlled

- bone growth. *J Oral Maxillofac Surg* 52:904-910.
- Quirynen M, Naert I, van Steenberghe D, Nys L (1992). A study of 589 consecutive implants supporting complete fixed prostheses. Part I: Periodontal aspects. *J Prosthet Dent* 68:655-663.
- Raghoobar GM, Heydenrijk K, Vissink A (2000a). Vertical distraction of the severely resorbed mandible. The Groningen distraction device. *Int J Oral Maxillofac Surg* 29:416-420.
- Raghoobar GM, Meijer HJA, Stegenga B, Van't Hof M, van Oort RP, Vissink A (2000b). Effectiveness of three treatment modalities for the edentulous mandible. A five-year randomized clinical trial. *Clin Oral Implant Res* 11:195-201.
- Raghoobar GM, Stellingsma K, Batenburg RH, Vissink A (2000c). Etiology and management of mandibular fractures associated with endosteal implants in the atrophic mandible. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 89:553-559.
- Raghoobar GM, Liem RSB, Vissink A (2002). Vertical distraction of the severely resorbed edentulous mandible. A clinical, histological and electron microscopic study of 10 treated cases. *Clin Oral Implant Res* 13:558-565.
- Roynesdal AK, Ambjornsen E, Stovne S, Haanaes HR (1998). A comparative clinical study of three different endosseous implants in edentulous mandibles. *Int J Oral Maxillofac Implants* 13:500-505.
- Sadowsky SJ (2001). Mandibular implant-retained overdentures: a literature review. *J Prosthet Dent* 86:468-473.
- Satow S, Slagter AP, Stoelinga PJ, Habets LL (1997). Interposed bone grafts to accommodate endosteal implants for retaining mandibular overdentures. A 1-7 year follow-up study. *Int J Oral Maxillofac Surg* 26:358-364.
- Schroeder A, Pohler O, Sutter F (1976). Gewebsreaktion auf ein Titan-Hohlzylinderimplantat mit Titan-Spritzschichtoberfläche [Tissue reaction to titanium plasma sprayed hollow cylinder implants]. *Schweiz Monat Zahn* 86:713-727.
- Small IA (1975). Metal implants and the mandibular staple bone plate. *J Oral Surg* 33:571-585.
- Small IA (1980). The mandibular staple bone plate for the atrophic mandible. *Dent Clin North Am* 24:565-570.
- Small IA (1993). The fixed mandibular implant: a 6-year review. *J Oral Maxillofac Surg* 51:1206-1210.
- Small IA, Misiek D (1986). A sixteen-year evaluation of the mandibular staple bone plate. *J Oral Maxillofac Surg* 44:60-66.
- Small IA, Helfrick JF, Stines AV (1995). The fixed mandibular implant. In: Reconstructive preprosthetic oral and maxillofacial surgery. Fonseca R, Davis WH, editors. Philadelphia: W.B. Saunders Company, pp. 518-564.
- Stellingsma C, Raghoobar GM, Meijer HJ, Batenburg RH (1998). Reconstruction of the extremely resorbed mandible with interposed bone grafts and placement of endosseous implants. A preliminary report on outcome of treatment and patients' satisfaction. *Br J Oral Maxillofac Surg* 36:290-295.
- Stellingsma C, Meijer HJ, Raghoobar GM (2000). Use of short endosseous implants and an overdenture in the extremely resorbed mandible: a five-year retrospective study. *J Oral Maxillofac Surg* 58:382-387.
- Stellingsma K, Bouma J, Stegenga B, Meijer HJ, Raghoobar GM (2003). Satisfaction and psychosocial aspects of patients with an extremely resorbed mandible treated with implant-retained overdentures. *Clin Oral Implants Res* 14:166-172.
- Stellingsma C, Raghoobar GM, Meijer HJA, Stegenga B, de Bont LGM (2004a). The extremely resorbed mandible: two-year results of a comparative, prospective study of three treatment modalities. Part I: clinical results. *Int J Oral Maxillofac Implants* (accepted).
- Stellingsma C, Meijer HJA, Raghoobar GM, Stegenga B, Kuiper C (2004b). The extremely resorbed mandible: two-year results of a comparative, prospective study of three treatment modalities. Part II: radiographic results. *Int J Oral Maxillofac Implants* (accepted).
- Stoelinga PJ, Blijdorp PA, Ross RR, De Koomen HA, Huybers TJ (1986). Augmentation of the atrophic mandible with interposed bone grafts and particulate hydroxylapatite. *J Oral Maxillofac Surg* 44:353-360.
- Tang L, Lund JP, Tache R, Clokie CM, Feine JS (1997). A within-subject comparison of mandibular long-bar and hybrid implant-supported prostheses: psychometric evaluation and patient preference. *J Dent Res* 76:1675-1683.
- Tolman DE (1995). Reconstructive procedures with endosseous implants in grafted bone: a review of the literature. *Int J Oral Maxillofac Implants* 10:275-294.
- Tolman DE, Keller EE (1991). Management of mandibular fractures in patients with endosseous implants. *Int J Oral Maxillofac Implants* 6:427-436.
- Triplett RG, Mason ME, Alfonso WF, McAnear JT (1991). Endosseous cylinder implants in severely atrophic mandibles. *Int J Oral Maxillofac Implants* 6:264-269.
- Urbani G, Lombardo G, Santi E, Consolo U (1999). Distraction osteogenesis to achieve mandibular vertical bone regeneration: a case report. *Int J Periodont Rest Dent* 19:321-331.
- Van Blarcom CW (1999). The glossary of prosthodontic terms. 7th ed. *J Prosthet Dent* 81:39-110.
- Van Kampen FM, Van Der Bilt BA, Cune MS, Bosman F (2002). The influence of various attachment types in mandibular implant-retained overdentures on maximum bite force and EMG. *J Dent Res* 81:170-173.
- Van Pelt AWJ (1997). Has the TMI prosthetic protocol been changed? (letter to the editor). *Int J Oral Maxillofac Surg* 26:394.
- van Steenberghe D (1997). Outcomes and their measurement in clinical trials of endosseous oral implants. *Ann Periodontol* 2:291-298.
- Vanassche BJ, Stoelinga PJ, De Koomen HA, Blijdorp PA, Schoenaers JH (1988). Reconstruction of the severely resorbed mandible with interposed bone grafts and hydroxylapatite. A 2-3 year follow-up. *Int J Oral Maxillofac Surg* 17:157-160.
- Verhoeven JW, Cune MS, Terlou M, Zoon MA, de Putter C (1997). The combined use of endosteal implants and iliac crest onlay grafts in the severely atrophic mandible: a longitudinal study. *Int J Oral Maxillofac Surg* 26:351-357.
- Verhoeven JW, Cune MS, Van Kampen FM, Koole R (2001). The use of the transmandibular implant system in extreme atrophy of the mandible; a retrospective study of the results in two different hospital situations. *J Oral Rehabil* 28:497-506.
- Vermeeren JJJF, Wismeijer D, Van Waas MAJ (1996). One-step reconstruction of the severely resorbed mandible with onlay bone grafts and endosteal implants. *Int J Oral Maxillofac Surg* 25:112-115.
- Versteegh PA, Beek GJv, Slagter AP, Ottervanger JP (1995). Clinical evaluation of mandibular overdentures supported by multiple-bar fabrication: a follow-up study of two implant systems. *Int J Oral Maxillofac Implants* 10:595-603.
- Wismeijer D, van Waas MAJ, Vermeeren JJJF (1995). Overdentures supported by ITI implants: a 6.5-year evaluation of patient satisfaction and prosthetic aftercare. *Int J Oral Maxillofac Implants* 10:744-749.
- Worthington P (1992). Clinical aspects of severe mandibular atrophy. In: Advanced osseointegration surgery. Worthington P, Brånemark PI, editors. Chicago: Quintessence Books, pp. 119-122.
- Worthington P, Rubenstein JE (1998). Problems associated with the atrophic mandible. *Dent Clin North Am* 42:129-160.
- Yanase RT, Bodine RL, Tom JF, White SN (1994). The mandibular subperiosteal implant denture: a prospective survival study. *J Prosthet Dent* 71:369-374.
- Young LJ, Michel JD, Moore DJ (1983). A twenty-year evaluation of subperiosteal implants. *J Prosthet Dent* 49:690-694.
- Zarb GA, Symington JM (1983). Osseointegrated dental implants: preliminary report on a replication study. *J Prosthet Dent* 50:271-276.