

University of Groningen

## Self-affine case for the roughness effect on the frictional force in boundary lubrication

Palasantzas, George

*Published in:*  
Physical Review E

*DOI:*  
[10.1103/PhysRevE.50.4256](https://doi.org/10.1103/PhysRevE.50.4256)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1994

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Palasantzas, G. (1994). Self-affine case for the roughness effect on the frictional force in boundary lubrication. *Physical Review E*, 50(5). <https://doi.org/10.1103/PhysRevE.50.4256>

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## Self-affine case for the roughness effect on the frictional force in boundary lubrication

George Palasantzas

Physics Department, Northeastern University, Boston, Massachusetts 02115

(Received 5 July 1994)

We comment on the analytic expressions obtained by Daikhin and Urbakh [Phys. Rev. E **49**, 1424 (1994)] for the case where the wall roughness is described by self-affine structure for the frictional forces in confining geometry systems.

PACS number(s): 68.45.-v, 68.15.+e

Very recently Daikhin and Urbakh [1] have shown that the presence of roughness essentially *increases* the frictional force in confining systems, and also can lead to the time dependence of the friction. We investigate here the static part of the frictional force between two walls separated by a thin liquid film in the case where the wall roughness is characterized by a self-affine structure. Furthermore, we shall try to derive the corresponding analytical expressions. Daikhin and Urbakh [1] considered the case of walls bounded by rough surfaces with a Gaussian height-height correlation  $C(R) = \sigma^2 e^{-R^2/\xi^2}$ , which in the case of weak roughness  $\sigma \ll \xi$  and wall separation  $d \ll \xi$  yielded analytical expressions. The parameters  $\sigma$  and  $\xi$  are respectively the normal rms roughness and the in-plane correlation length.

However, a wide variety of surfaces and interfaces occurring in nature are well represented by a kind of roughness associated with self-affine fractal scaling [2]. If a surface exhibits self-affine roughness [3], the height difference correlation function  $g(R)$  will scale as  $g(R) \propto R^{2H}$  if  $R \ll \xi$ , and  $g(R) \approx 2\sigma^2$  if  $R \gg \xi$ . The function  $g(R)$  is related to the height-height correlation  $C(R)$  by  $g(R) = 2\sigma^2 - 2C(R)$ . The exponent  $0 < H < 1$  is referred to as the "roughness" exponent [4,5]. Small values of  $H \sim 0$  represent extremely jagged or irregular surfaces, while large values of  $H \sim 1$  represent smooth hill-valley structures. In fact, the Gaussian correlation can be considered as a limiting case of the fractal correlation  $C(R) \sim e^{-(R/\xi)^{2H}}$  for  $H=1$ . In our calculations for self-affine fractals, we shall consider the  $k$ -correlation model that in  $k$ -space has the form

$$C(\mathbf{k}) = (2\pi)\sigma^2\xi^2 / (1 + ak^2\xi^2)^{1+H}$$

with  $a = (1/2H)[1 - (1 + ak_c^2\xi^2)^{-H}]$  for  $H > 0$ , and  $a = 1/2\ln(1 + ak_c^2\xi^2)$  for  $H = 0$ , as well as  $k_c = \pi/a_0$ , with  $a_0$  the atomic spacing [5].

Let us denote the velocity of motion of the upper wall by  $V_0$  and the liquid viscosity by  $n$ . The static part of the frictional force is given by [1]

$$F_{st} = (nV_0/d) \left[ 1 + (2\pi)^{-2} d^{-2} \sum_{i=1}^2 \int C_i(\mathbf{k}) a(kd, \phi) d^2\mathbf{k} \right], \quad (1)$$

with  $\int_0^{2\pi} a(kd, \phi) d\phi \approx \frac{5}{2}(2\pi) - 7(kd)^2/10$  for  $d \ll \xi_i$ . The

indices  $i=1,2$  refer to the upper and lower walls, and  $nV_0/d$  represents the frictional force for flat walls. The fulfillment of the condition  $d \ll \xi$  is not sufficient for the case of power law roughness to ensure experimentally realizable values for  $d \sim 5$  nm in order that  $F_{st} \geq nV_0/d$ . In fact, depending on the value of  $H$  the perturbation expansion of  $a(kd, \phi)$  has to continue to higher order terms. For self-affine fractals  $F_{st}$  is given by

$$F_{st} = \frac{nV_0}{d} \left( 1 + \frac{1}{d^2} F_1 - F_2 \right), \quad (2)$$

$$F_1 = \frac{5}{2} \sum_{i=1}^2 \sigma_i^2, \quad (3)$$

$$F_2 = \frac{7}{10} \sum_{i=1}^2 \frac{\sigma_i^2}{2a_i\xi_i^2} \times \left\{ \frac{1}{a_i(1-H_i)} [(1 + a_i k_{ci}^2 \xi_i^2)^{1-H_i} - 1] - 2 \right\},$$

where its applicability up to second order in  $a(kd, \phi)$  is limited to wall separations  $d \ll d_c = (F_1/F_2)^{1/2} (F_1 \gg d^2 F_2)$ . In order to gain a feeling of the effect, of  $H$ , as well as the applicability of the previous results to nanoscale roughness, we present some estimations of  $d_c$ . In the following, we shall consider  $H_{1,2} = H$  for illustration purposes. For  $\xi_{1,2} = 1000$  nm and  $\sigma_{1,2} = 100$  nm,  $d_c \geq 36$  nm for  $H \geq 0.6$ . For  $\xi_{1,2} = 100$  nm and  $\sigma_{1,2} = 10$  nm,  $d_c \geq 9$  nm for  $H \geq 0.6$ . Therefore, the previous estimations show that for  $H < 0.5$  higher order terms [or the exact expressions for  $a(kd, \phi)$ ] are required in order for experimentally reasonable values of  $d \ll d_c$  to be feasible ( $d \sim 5$  nm [1,6] for wall roughness on the nanoscale range).

The dependence of the frictional force on the particular choice of the correlation function is given by its Fourier  $C_i(k)$  in Eq. (1). (For various forms of the correlation function see Refs. [5] and [7].) From Eq. (3), the term  $F_2$  includes, besides the ratio  $\sigma_i^2/\xi_i^2$ , the roughness exponent  $H$ . If we consider  $H = 0.5$ , for  $a_0 \ll \xi$  the analytic  $k$ -correlation model used in this study yields the simple correlation function  $C(R) = \sigma^2 e^{-R/\xi}$  [5]. For  $\sigma_{1,2} = 10$  nm,  $\xi_{1,2} = 200$  nm, and  $d = 5$  nm, we obtain, for  $H = 0.5$   $F_1/d^2 = 20.0$  and  $F_2 = 7.32$ , as well as for

$H=0.8$   $F_1/d^2=20.0$  and  $F_2=0.4$ . Therefore, for a fixed ratio  $\sigma/\xi \ll 1$  the larger the  $H$  the smaller is  $F_2$ , and as a result the dominant contribution to the frictional force comes from  $\sigma$ . However, for small values of  $H$  the term  $F_2$  increases, resulting in higher sensitivity of the frictional force to the roughness exponent  $H$ .

In conclusion, our simple calculations show that in order to account for nanoscale roughness in confining geometry systems, the effect of possible surface fractality

has to be taken into account in the perturbation expansions. Furthermore, we gave the expressions for this effect in terms of the theory developed by Daikhin and Urbakh [1].

It is pleasure to acknowledge partial support from the NSF Grant No. DMR-9204022, and the hospitality of the VSM-Lab at Katholieke Universiteit Leuven (Belgium).

- 
- [1] L. Daikhin and M. Urbakh, *Phys. Rev. E* **49**, 1424 (19994), see also references therein for a review.  
[2] For a review, see P. Meakin, *Phys. Rep.* **235**, 189 (1993).  
[3] F. Family and T. Vicsek, *Dynamics of Fractal Surfaces* (World Scientific, Singapore, 1991).  
[4] G. Palasantzas, *Phys. Rev. E* **49**, 1740 (1994); J. Krim and J. O. Indekeu, *ibid.* **48**, 1576 (1993).

- [5] G. Palasantzas, *Phys. Rev. B* **48**, 14472 (1993); **49**, 5785 (1994).  
[6] J. N. Israelachvili, *Intermolecular Forces and Surface Forces*, 2nd ed. (Academic, London, 1991).  
[7] G. Palasantzas and J. Krim, *Phys. Rev. B* **48**, 22873 (1994).