

University of Groningen

Lanthanide mediated activation of C-H and C-X bonds

Deelman, Berth Jan

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1994

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Deelman, B. J. (1994). *Lanthanide mediated activation of C-H and C-X bonds*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.


Chapter 6

Reaction of $(Cp^*_2LnH)_2$ ($Ln = Y, La$) and $Cp^*_2Y(2-C_6H_4CH_2NMe_2)$ with Esters and Amides. Molecular Structure of $[Cp^*_2Y)_2(\mu-OCMe=CHC(OEt)O)]_2$

Introduction

The recent interest in the use of organolanthanides for organic synthesis¹ has led to the discovery of some useful C-C bond formation reactions mediated by these complexes. For instance aldol coupling of enolizable ketones could be induced by $Cp^*_2LnCH(SiMe_3)_2$ as has been found by Heeres.² Samarium based C-C bond forming reactions with aldehydes, ketones and acid chlorides have been reported by Kagan *et al.* and have a broad scope.³ Recently, interesting work on stereoselective aldol condensations using lanthanide alkoxides has been published.⁴



In this study we focus on the reactions of $(Cp^*_2LnH)_2$ ($Ln = Y$ (1a), La (1b)), and the intramolecularly stabilized aryl $Cp^*_2Y(2-C_6H_4CH_2NMe_2)$ (2)⁵ with esters and amides since condensation of these compounds by yttrium or lanthanide complexes is to our knowledge unprecedented. The intramolecularly stabilized aryl complex was chosen because of the lower steric congestion at the yttrium center compared to that in $Cp^*_2YCH(SiMe_3)_2$.⁶ Esters are especially interesting because of the possible addition-elimination sequence depicted in eq 1, which would lead to the alkylation of esters to produce ketones. Another possibility would be enolization of the ester leading to Claisen type condensation with a second equivalent of ester. In addition, we are interested in the effect of the intramolecularly coordinated amine function of 2 on its reactivity.



This work has been performed in collaboration with F. Wierda. The X-ray structure determination was performed by A. Meetsma.

Results and Discussion

Activation of Esters and Amides by $(\text{Cp}^*_2\text{LnH})_2$ ($\text{Ln} = \text{Y}$ (1a) and La (1b)). Both hydrides 1a and 1b reacted with ethyl acetate to form rapidly the alkoxide complexes $\text{Cp}^*_2\text{LnOEt}$ ($\text{Ln} = \text{Y}$ (3a) and La (3b)) through a reductive ester cleavage (eq 2). Even at -80°C no intermediates could be detected ($^1\text{H-NMR}$). With ethyl benzoate an analogous ester cleavage was observed resulting in a 1 : 1 mixture of alkoxides 3 and 4 (eq 3). No metallation of the phenyl ring takes place, which is remarkable when compared with other functionalized arenes PhX ($\text{X} = \text{OMe}$, SMe , NMe_2 , CH_2NMe_2 , PMe_2 , $\text{PPh}_2=\text{CH}_2$) which are metallated very easily in the ortho position by the same system.⁷ It is clear that activation of the carbonyl group by nucleophilic attack of the hydride ligand is the dominant reaction taking place here.




The first step in the ester reduction is most likely insertion of the carbonyl group in the Ln-H bond as has been observed for 1b with di-*t*-butyl ketone to form $\text{Cp}^*_2\text{LaOCH}(\text{tBu})_2$ ^{2b} (Scheme I). Elimination of the ethoxy group and formation of 3a and aldehyde HC(O)R seems to be a plausible second step. The aldehyde formed can then be attacked by another molecule of 1 to form $\text{Cp}^*_2\text{LnOCH}_2\text{R}$.


To test whether activation of the carbonyl group is still the main reaction with α , β -unsaturated esters, 1a and 1b were allowed to react with ethyl acrylate. This resulted in fast polymerization of the acrylate whereas the bulk of 1a or 1b remained intact. This indicates that only minor amounts of 1a and 1b are involved in the catalytic polymerization which can be explained by slow initiation. The poly(ethyl acrylate) formed is apparently completely atactic as follows from the $^1\text{H-NMR}$ spectrum, which shows two types of methylene and methine protons.⁸ However, it is clear that alkoxide formation, which would lead to complete deactivation of the catalyst, is not taking place here. From this it can be concluded that insertion of C-C double bonds into Ln-H can in principle compete successfully with alkoxide elimination. Since polymerization of methyl metacrylate (MMA) has

been investigated extensively⁹ we did not look further into this aspect of organo-lanthanide chemistry.


Scheme I



The $-\text{NR}_2$ groups of organic amides are poor leaving groups compared to alkoxy groups of esters. We therefore anticipated that reaction with *N,N*-dimethylacetamide would lead to the carbonyl insertion products of **1a** and **1b**. However the reaction was not clean owing to competitive α -H abstraction and aldol condensation of acetamide (eq 4). This resulted in a mixture of carbonyl insertion product **5** (40 %) and condensation product **6** (60 %). No attempts were made to isolate the aldol coupled product since the reaction of **2** with *N,N*-dimethylacetamide produced **6** in much better yield (*vide infra*).



Activation of Esters and Amides by $\text{Cp}^*_2\text{Y}(2\text{-C}_6\text{H}_4\text{CH}_2\text{NMe}_2)$ (2). In contrast to the hydrides 1a and 1b which react by nucleophilic attack on the carbonyl function of esters, the aryl 2 behaves like a base and abstracts an $\alpha\text{-H}$ of ethyl acetate which subsequently leads to $[\text{Cp}^*_2\text{Y}(\mu\text{-OCMe}=\text{CHC}(\text{OEt})\text{O})]_2$ (7) as the isolated product (eq 5). Formation of this compound can be explained as follows (Scheme II). The enolate $\text{Cp}^*_2\text{YOC}(\text{OEt})=\text{CH}_2$, which is formed by $\alpha\text{-H}$ abstraction from ethyl acetate, enters a condensation with another molecule of ethyl acetate similar to the mechanism proposed for aldol condensations with $\text{Cp}^*_2\text{LnCH}(\text{SiMe}_3)_2$.^{2b} The condensation product then eliminates Cp^*_2YOEt producing a $\beta\text{-keto}$ ester and turning the overall process into a Claisen condensation. The final step is enolization of the $\beta\text{-keto}$ ester by 2 to form 7.




Compound 7 could be isolated in good yield and was fully characterized. Also the elimination product Cp^*_2YOEt could be detected in the mother liquor as the ethyl acetate adduct after isolation of 7. The IR spectrum of 7 exhibits a band at 1568 cm^{-1} indicative of a severely reduced carbonyl bond order. Compared to $\beta\text{-keto}$ esters for which $\nu_{\text{C}=\text{O}}$ occurs at 1650 cm^{-1} ,¹⁰ this value is significantly shifted to lower frequency. This carbonyl shift can be explained by coordination to the Lewis acidic yttrium center which causes reduction of the carbonyl double bond character. Also the delocalized bonding within the $\text{OCMe}=\text{CHC}(\text{OEt})\text{O}$ causes significant reduction of the $\text{C}=\text{O}$ bond order (*vide infra*). The low carbonyl stretching frequency compares well with those observed for tris acetylacetonate complexes $\text{Ln}(\text{acac})_3(\text{H}_2\text{O})_n$ (1600 cm^{-1}).¹¹

The molecular structure of 7 was determined by X-ray diffraction and the resulting geometry is depicted in Figure 1. For selected bond distances and angles see Table I. The molecule is a dimer consisting of two equivalent $\text{Cp}^*_2\text{Y}(\mu\text{-OCMe}=\text{CHC}(\text{OEt})\text{O})$ units. The arrangement of the Cp^* ligands on yttrium with a $\text{Cp}^*\text{-Y-Cp}^*$ angle of 138.6° is normal for bent ytrocene compounds.¹² The Y1-O3a

is significantly longer than the terminal Y-OR distances in the alkoxides $\text{Cp}^*\text{Y}(\text{OC}_6\text{H}_3^t\text{Bu}_2)_2$ (2.096(4) and 2.059(3))¹³ and $[\text{Cp}^*\text{Y}(\mu\text{-O}^t\text{Bu})(\text{O}^t\text{Bu})_2]$ (1.995(10) and 2.018(9)).¹⁴ However, the almost linear angle Y1-O3a-C23, which is due to π overlap of the oxygen lone pairs with metal orbitals, was also observed for one of the aryloxy ligands in $\text{Cp}^*\text{Y}(\text{OC}_6\text{H}_3^t\text{Bu}_2)_2$. In fact the Y1-O3a distance is intermediate between the Y-O distances in these alkoxides and that of the enolate $[(\text{C}_5\text{H}_4\text{Me})_2\text{Y}(\mu\text{-OCH}_2=\text{CH}_2)]_2$ (2.275(3) and 2.290(2) Å).¹⁵

Scheme II






Figure 1. *PLUTO* drawing of $[\text{Cp}^*_2\text{Y}(\mu\text{-OCMe}=\text{CHC}(\text{OEt})\text{O})]_2$ (7).

The Y1-O1 distance is significantly longer than Y1-O3a whereas the O1-C21 distance is slightly shorter than O3a-C23. This can be interpreted as a carbonyl function interacting datively with yttrium. The fact that Y1, O1, C21, C22, C23, and O3 lie almost within the same plane, as follows from the torsion angles O3a-Y1-O1-C21, C22-C21-O1-Y1, O1-C21-C22-C23 and C21-C22-C23-O3a, shows that there is significant π overlap between these atoms resulting in delocalized bonding. This is supported by the distances O1-C21, C21-C22, C22-C23, O3-C23 which all indicate significant double bond character (bonding distances for pure double and single bonds: C-C = 1.54 Å, C=C = 1.34 Å, C-O = 1.43 Å, C=O = 1.20 Å).¹⁶ The delocalized bonding within the $\mu\text{-OCMe}=\text{CHC}(\text{OEt})\text{O}$ ligands could also explain the elongation of the Y-O3a distance relative to the Y-O distances of $\text{Cp}^*\text{Y}(\text{OC}_6\text{H}_3^t\text{Bu}_2)_2$ and $[\text{Cp}^*\text{Y}(\mu\text{-O}^t\text{Bu})(\text{O}^t\text{Bu})]_2$.


By using a more sterically hindered ester, ethyl-2-methylpropanoate, we tried to isolate the anticipated enolate intermediate. This ester has still one acidic $\alpha\text{-H}$ proton but due to the methyl groups in the α -position, nucleophilic attack of the enolate on a second equivalent of ethyl 2-methylpropanoate would be sterically highly unfavorable. However, we did not find the expected enolate complex $\text{Cp}^*_2\text{Y}(\text{O}-\text{C}(\text{OEt})=\text{CMe}_2)$. Instead the product obtained was $\text{Cp}^*_2\text{Y}(\text{OC}(2-$

$C_6H_4CH_2NMe_2=CMe_2(Me_2CHCOEt)$ (**8**) (eq 6), which could be fully characterized by standard techniques.

Table I. Selected Distances (Å) and Angles (°) for $[Cp^*_2Y(\mu-O CMe=CHC(OEt)O)]_2$ (7**).^a**


Y1-O1	2.292(2)	O3a-C23	1.281(4)
Y1-O3a	2.179(2)	C21-C22	1.406(4)
O1-C21	1.260(3)	C22-C23	1.385(3)
O2-C21	1.341(4)	C23-C24	1.513(4)
O2-C25	1.439(4)	C25-C26	1.509(5)
O1-Y1-O3a	103.30(7)	O2-C21-C22	117.9(2)
Y1-O1-C21	137.8(2)	C21-C22-C23	129.3(3)
C21-O2-C25	118.2(2)	O3a-C23-C22	119.8(3)
C23-O3a-Y1a	174.59(18)	O3a-C23-C24	116.8(2)
O1-C21-O2	119.3(3)	C22-C23-C24	123.4(3)
O1-C21-C22	122.8(3)	O2-C25-C26	106.0(3)
Ct1-Y1-Ct2	138.6 ^b		
O3a-Y1-O1-C21	6.7(3)	O1-C21-C22-C23	176.7(3)
C22-C21-O1-Y1	-3.6(5)	C21-C22-C23-O3a	177.9(3)

^a label "a" indicates symmetry operation: -x, -y, 2-z. ^b Ct1 = C1-C5, Ct2 = C11-C15




The presence of the benzyldimethylamine function in **8** suggests that the first step is insertion of the carbonyl function of the ester in the Y-aryl bond of **2** (Scheme III). Next the ethoxy group can be eliminated to form Cp^*_2YOEt and $\text{Me}_2\text{CHC}(\text{O})\text{C}_6\text{H}_4\text{CH}_2\text{NMe}_2$. This is supported by the presence of Cp^*_2YOEt in the reaction mixture ($^1\text{H-NMR}$). The ketone produced can then be converted into the enolate by another equivalent of **2** and the final product is formed by complexation of one molecule of ester. This reaction shows that by using a sterically more hindered ester, we have indeed prevented aldol condensation but apparently also the enolization of the ester to $\text{Cp}^*_2\text{YOC}(\text{OEt})=\text{CMe}_2$ is more difficult. Instead, insertion of the carbonyl group of the ester into the Y-aryl bond of **2** is taking place. Although the C-C coupling reaction is now blocked, the OEt elimination, stimulated by the electrophilicity of yttrium, is still possible yielding the observed product **8**.

Scheme III



With *N,N*-dimethylacetamide regular aldol condensation was observed to form $\text{Cp}^*_2\text{Y}[\text{OC}(\text{Me})(\text{NMe}_2)\text{CH}_2\text{C}(\text{O})\text{NMe}_2]$ (**6**) (eq 7). In contrast to reaction with **1a** the carbonyl addition product was not formed which indicates that **2** is less nucleophilic and behaves like a base. As stated before, the $-\text{NMe}_2$ function is a poor leaving group which prevents NMe_2 elimination and formation of the β -keto amide. Apparently enolization with another equivalent of **2** is more difficult since formation of the enolate is not taking place here.



Concluding Remarks

In this study we have seen that organo-yttrium complexes can function as versatile synthetic tools in organic chemistry. The C-C bond forming aldol and Claisen condensations could offer opportunities for the development of catalysts for these reactions. However, a lot of work needs to be done to reach this goal. In particular the stereochemistry with α -substituted carbonyls needs to be controlled to make these reactions useful in organic synthesis. Therefore, the development of complexes which induce diastereoselective and enantioselective C-C forming condensations remains to be worked out. In this study we have shown that organo-yttrium compounds could serve as promising candidates for transformations of this type.

Experimental Section

General Considerations. General procedures, techniques and instrumentation were described in chapters 2 and 4. Compounds **1a**,⁶ **1b**,¹⁷ $\text{Li}(2\text{-C}_6\text{H}_4\text{CH}_2\text{NMe}_2)^{\text{sa}}$ and $(\text{Cp}^*_2\text{YCl})_2$ ¹⁸ were prepared as described. Reagents ethyl acetate, ethyl benzoate, ethyl acrylate, *N,N*-dimethylacetamide and ethyl 2-methyl propanoate

were distilled and stored over molecular sieves (4 Å). Solvents were distilled from Na/K alloy and degassed prior to use.

Reactions of $(Cp^*_2LnH)_2$ ($Ln = Y(1a), La(1b)$) with Ethyl Acetate. Ethyl acetate (1.4 μ L, 0.015 mmol) was added to a solution of 11 mg (0.015 mmol) **1a** in 0.5 mL of cyclohexane- d_{12} . 1H -NMR spectroscopy after 5 min at room temperature showed the quantitative formation of **3a**. For 1H -NMR data of **3a** see chapter 5. A similar procedure was used for reaction of **1b** with 0.5 equivalent of ethyl acetate per **La**. Quantitative formation of **3b** was observed within several minutes at room temperature. 1H -NMR (300 MHz, benzene- d_6): δ 3.87 (broad s, $lw_{1/2} = 20$ Hz, 2H, $LaOCH_2$), 2.07 (s, 30H, C_5Me_5), 1.24 (t, $^3J_{HH} = 7.12$ Hz, 3H, OCH_2CH_3).

Reactions of $(Cp^*_2LnH)_2$ ($Ln = Y(1a), La(1b)$) with Ethyl Benzoate. Similar to the reaction of **1a** with ethyl acetate, 1 equivalent of ethyl benzoate per **Y** was added to a solution of **1a** in cyclohexane- d_{12} . 1H -NMR showed the quantitative formation a 1 : 1 : 1 mixture of **3a**, **4a**, and ethyl benzoate. 1H -NMR for **4a** (200 MHz, cyclohexane- d_{12}): δ 7.60 (d, $^3J_{HH} = 7.0$ Hz, 2H, ortho H), 7.34 (t, $^3J_{HH} = 6.8$ Hz, 2H, meta H), 5.36 (s, 2H, OCH_2) 2.07 (s, 30H, C_5Me_5), para H not found due to overlap with signals of unreacted ester. In a similar procedure for **1b**, using 1 equivalent of ethyl benzoate per **La**, quantitative formation of 1 : 1 : 1 molar mixture of **3b**, **4b** and ethyl benzoate was observed within 5 min at room temperature. 1H -NMR for **4b** (200 MHz, benzene- d_6): δ 7.55 (d, $^3J_{HH} = 6.8$ Hz, 2H, ortho H), 7.16 (t, $^3J_{HH} = 6.8$ Hz, 2H, meta H), 5.32 (s, 2H, OCH_2), 2.12 (s, 30 H, C_5Me_5), para H not found due to overlap with signals of unreacted ester.

Reaction of **1a with Ethyl Acrylate.** To a solution of 0.025 g (0.035 mmol) of **1a** in 20 mL of cyclohexane was added 1.40 mmol of ethyl acrylate. The reaction mixture was stirred for 2 h at room temperature after which the reaction was quenched with 1 mL of methanol. After the mixture was filtrated and volatiles were removed, a sticky substance remained. 1H -NMR (200 MHz, chloroform- d_1): δ 4.11 (q, $^3J_{HH} = 6.8$ Hz, OCH_2CH_3), 2.30 (broad s, $CH-CH_2-CH$), 1.95 (broad s, $CH_2-CH-CH_2$), 1.65 (broad s, $CH-CH_2-CH$), 1.52 (broad s, CH_2-CH), 1.25 (t, $^3J_{HH} = 6.8$ Hz, OCH_2CH_3), relative integrated intensities 4 : 2 : 1 : 2 : 1 : 6 respectively.

Reaction of 1a with *N,N*-Dimethylacetamide. To a stirred solution of 0.20 g (0.28 mmol) of 1a in 30 mL of pentane was added 104 μ L (1.12 mmol) *N,N*-dimethylacetamide. Gas evolution was observed. After 5 min at room temperature stirring was stopped and the mixture was left to crystallize at room temperature. Yield: 0.236 g. ^1H - and ^{13}C -NMR showed that this material consisted of 5 (0.23 mmol) and 6 (0.34 mmol). ^1H -NMR for 5 (200 MHz, benzene- d_6): 5.01 (q, $^3J_{\text{HH}} = 5.6$ Hz, 1H, OCH(Me), 2.35 (s, 6H, NMe $_2$), 2.14 (s, 30H, C $_5$ Me $_5$), 1.26 (d, $^3J_{\text{HH}} = 5.6$ Hz, 3H, OCHMe). ^{13}C -NMR for 5 (75.4 MHz, benzene- d_6): 114.97 (s, C $_5$ Me $_5$), 87.61 (dd, $^1J_{\text{CH}} = 144$ Hz, $^2J_{\text{CY}} = 5$ Hz, YOCH), 39.17 (t, $^1J_{\text{CH}} = 134$ Hz, NMe $_2$), 18.08 (q, $^1J_{\text{CH}} = 125$ Hz, OCHMe), 11.68 (q, $^1J_{\text{CH}} = 125$ Hz, C $_5$ Me $_5$). For NMR data of 6, see synthesis of this compound from ethyl acetate and 2.

Cp* $_2$ Y(2-C $_6$ H $_4$ CH $_2$ NMe $_2$) (2). A suspension of 15.9 g (20 mmol) of (Cp* $_2$ YCl) $_2$ and 5.68 g (40 mmol) of Li-2-C $_6$ H $_4$ CH $_2$ NMe $_2$ in 50 mL of toluene was stirred at room temperature for 15 h. Volatiles were removed in vacuum and the remaining solid was stripped 3 times with pentane. Extraction with pentane and crystallization at -80 $^\circ\text{C}$ afforded 9.82 g (20 mmol, 50 %) of 2 as yellow crystals. NMR data were identical to those reported before.⁷

[Cp* $_2$ Y(μ -OCMe=CHC(OEt)O)] $_2$ (7). To a stirred solution of 1.13 g (2.29 mmol) of 2 in 20 mL of benzene was added 0.45 mL (4.6 mmol) of ethyl acetate. The mixture was heated to 50 $^\circ\text{C}$ for 3 h. Cooling to room temperature gave white crystals which were washed with benzene. Yield: 0.44 g (0.45 mmol, 39 %). IR (cm $^{-1}$): 1730 (m), 1568 (s), 1537 (m), 1396 (sh), 1323 (m), 1271 (s), 1067 (m), 1020 (m). ^1H -NMR (300 MHz, benzene- d_6): δ 5.37 (s, 1H, C=CH), 4.16 (q, $^3J_{\text{HH}} = 7.0$ Hz, 2H, OCH $_2$), 2.46 (s, 3H, C(O)CH $_3$), 2.01 (s, 30H, C $_5$ Me $_5$), 1.10 (t, $^3J_{\text{HH}} = 7.3$ Hz, 3H, OCH $_2$ CH $_3$). ^{13}C -NMR (75.4 MHz, benzene- d_6): δ 190.08 (d, $^2J_{\text{CY}} = 5$ Hz, Y-O-C), 174.01 (d, $^2J_{\text{CY}} = 2$ Hz, Y-O=C), 116.42 (s, C $_5$ Me $_5$), 91.26 (d, $^1J_{\text{CH}} = 154$ Hz, C=CH), 61.14 (t, $^1J_{\text{CH}} = 145$ Hz, OCH $_2$), 28.23 (q, $^1J_{\text{CH}} = 127$, C(O)Me), 14.40 (q, $^1J_{\text{CH}} = 127$ Hz, OCH $_2$ CH $_3$), 11.06 (q, $^1J_{\text{CH}} = 126$ Hz, C $_5$ Me $_5$). Anal. Calcd for C $_{26}$ H $_{39}$ O $_3$ Y: C, 63.93; H, 8.05; Y, 18.20. Found: C, 63.80; H, 8.29; Y, 18.43. The mother liquor was evaporated to dryness and the residue was identified as Cp* $_2$ YOEt(MeCOOEt) and traces of 7 by ^1H -NMR. ^1H -NMR (300 MHz, benzene- d_6): δ 4.19 (q, $^3J_{\text{HH}} = 6.8$ Hz, 2H, YOCH $_2$), 3.97 (q, $^3J_{\text{HH}} = 7.3$ Hz, C(O)OCH $_2$), 2.04 (s, 30H, C $_5$ Me $_5$), 1.97 (s, 3H,

C(O)Me), 1.32 (t, $^3J_{\text{HH}} = 6.8$ Hz, 3H, YOCH_2CH_3), 0.88 (t, $^3J_{\text{HH}} = 7.3$ Hz, 3H, OCH_2CH_3). Crystal data for 7: $[\text{C}_{26}\text{H}_{39}\text{O}_3\text{Y}]_2$, $M = 977.00$, triclinic with $a = 10.129(1)$ Å, $b = 10.650(1)$ Å, $c = 12.093(1)$ Å, $\alpha = 77.642(6)^\circ$, $\beta = 79.402(7)^\circ$, $\gamma = 86.408(9)^\circ$, $V = 1252.2(2)$ Å³, space group $P\bar{1}$, $Z = 1$, $\mu(\text{Mo K}\alpha) = 23.6$ cm⁻¹. Anisotropic least-squares refinement based on 4871 reflections converged to $R_F = 0.034$ and $wR = 0.041$ for 428 refined parameters.

Cp*₂Y(OC(2-C₆H₄CH₂NMe₂)=CMe₂)(Me₂CHCOOEt) (8). To a stirred solution of 0.42 g (0.85 mmol) of 2 in 10 mL of toluene was added 250 μL (1.87 mmol) of ethyl 2-methylpropanoate. The reaction mixture was heated at 100 °C for 3 d. Volatiles were evaporated in vacuum and the residue was washed with pentane. Crystallization at -80 °C afforded 0.13 g (0.19 mmol, 22 %) of white crystals. IR (cm⁻¹): 3056 (w), 2780 (m), 2723 (w), 1663 (s), 1642 (s), 1591 (w), 1568 (w), 1536 (w), 1404 (m), 1308 (s), 1292 (s), 1265 (w), 1213 (m), 1196 (m), 1175 (s), 1094 (s), 1068 (w), 1028 (s), 949 (w), 895 (w), 858 (m), 777 (s), 638 (m), 567 (w), 532 (w), 422 (w). ¹H-NMR (300 MHz, benzene-*d*₆): δ 7.83 (d, $^3J_{\text{HH}} = 7.6$ Hz, 1H, aryl H), 7.46 (d, $^3J_{\text{HH}} = 6.8$ Hz, 1H, aryl H), 7.10 (m, 2H, aryl H), 3.79 (m, 4H, overlapping OCH₂ and NCH₂ signals), 2.29 (s, 6H, NMe₂), 2.03 (s, 30H, C₅Me₅), 1.94 (s, 3H, Me), 1.54 (s, 3H, Me), 0.71 (m, 9H, overlapping signals of OCH₂CH₃ and CHMe₂), CHMe₂ signal not found, presumably due to overlap with the C₅Me₅ signal. ¹³C-NMR (75.4 MHz, benzene-*d*₆): δ 184.76 (s, C=O), 151.40 (Y-OC), 143.82 (s, aryl C), 137.95 (s, aryl C), 130.2 (d, $^1J_{\text{CH}} = 149$ Hz, aryl CH), 127.75 (aryl C, overlapping with solvent signal), 126.17 (d, $^1J_{\text{CH}} = 147$ Hz, aryl CH), 116.28 (s, C₅Me₅), 97.25 (s, =CMe₂), 62.95 (t, $^1J_{\text{CH}} = 148$ Hz, OCH₂), 61.44 (t, $^1J_{\text{CH}} = 132$ Hz, NCH₂), 46.05 (q, $^1J_{\text{CH}} = 132$ Hz, NMe₂), 35.31 (d, $^1J_{\text{CH}} = 136$ Hz, CHMe₂), 20.17 (q, $^1J_{\text{CH}} = 124$ Hz, Me), 19.86 (q, $^1J_{\text{CH}} = 124$ Hz, Me), 19.42 (q, $^1J_{\text{CH}} = 128$ Hz, CHMe₂), 13.62 (q, $^1J_{\text{CH}} = 128$ Hz, Me), 11.68 (q, $^1J_{\text{CH}} = 125$ Hz, C₅Me₅), one aryl CH not found due to overlap with solvent signal. Anal. Calcd for C₃₉H₆₀O₃NY: C, 68.91; H, 8.90; Y, 13.08. Found: C, 68.11; H, 8.85; Y, 13.74.

Cp*₂Y(OC(Me)(NMe₂)CH₂C(O)NMe₂) (6). To a stirred solution of 1.33 g (2.69 mmol) of 2 in 10 mL of benzene was added 0.50 mL (5.2 mmol) of *N,N*-dimethylacetamide. The reaction mixture was stirred for 15 h at 50 °C and volatiles were removed in vacuum. The residue was washed with pentane and crystallization from toluene at -80 °C yielded 0.59 g (1.13 mmol, 42 %) of white crystals. IR (cm

¹): 1606 (s), 1587 (s), 1410 (m), 1325 (m), 1264 (w), 1224 (m), 1215 (m), 1018 (s), 985 (m), 594 (m). ¹H-NMR (200 MHz, benzene-*d*₆): δ 3.21 (s, 2H, C(O)CH₂), 2.77 (s, 6H, NMe₂), 2.45 (s, 3H, Me), 2.17 (s, 30H, C₅Me₅), 1.82 (s, 3H, NMe), 1.80 (s, 3H, NMe). ¹³C-NMR (75.4 MHz, benzene-*d*₁₂): δ 167.50 (s, C=O), 115.96 (s, C₅Me₅), 60.99 (t, ¹J_{CH} = 155 Hz, C(O)CH₂), 40.44 (q, ¹J_{CH} = 134 Hz, NMe₂), 37.46 (q, ¹J_{CH} = 139 Hz, NMe), 35.98 (q, ¹J_{CH} = 140 Hz, NMe), 21.90 (q, ¹J_{CH} = 130 Hz, Me), 11.53 (q, ¹J_{CH} = 125 Hz, C₅Me₅), Y-O-C carbon not found. Anal. Calcd for C₂₈H₄₇O₂N₂Y: C, 63.14; H, 8.89; N, 5.26; Y, 16.69. Found: C, 63.47; H, 8.99; N, 5.44; Y, 17.13.

References and Notes

- ¹ See for instance: (a) Molander, G.A.; *Chem. Rev.* **1992**, *92*, 29. (b) Kagan, H.B.; Sasaki, M.; Collin, J. J. *Pure Appl. Chem.* **1988**, *60*, 1725. (c) Molander G.A. In *The Chemistry of the Metal-Carbon Bond*, Patai, S.; Hartley, F.R., Eds., John Wiley and Sons: Chichester: England, 1989.
- ² (a) Heeres, H.J. *Thesis*, University of Groningen: Groningen, The Netherlands, 1990. (b) Heeres, H.J.; Maters, M.; Teuben, J.H.; Helgesson, G.; Jagner, S. *Organometallics* **1992**, *11*, 350.
- ³ Bied, C.; Collin, J.; Kagan, H.B. *Tetrahedron* **1992**, *48*, 3877.
- ⁴ Makioka, Y.; Nakagawa, I.; Taniguchi, Y.; Takaki, K.; Fujiwara, Y. *J. Org. Chem.* **1993**, *58*, 4771 and references cited herein.
- ⁵ (a) Manzer, L.E.; *J. Am. Chem. Soc.* **1978**, *100*, 8068. (b) Rausch, M.D.; Foust, D.F.; Rogers, R.D.; Atwood, J.L. *J. Organomet. Chem.* **1984**, *265*, 241.
- ⁶ Den Haan, K.H.; Wielstra, Y.; Teuben, J.H. *Organometallics* **1987**, *6*, 2053.
- ⁷ Booiij, M.; Deelman, B.-J.; Duchateau, R.; Postma, D.S.; Meetsma, A.; Teuben, J.H. *Organometallics* **1993**, *12*, 3531.
- ⁸ Wayland, B.B.; Poszmik, G. *Organometallics* **1992**, *11*, 3534.
- ⁹ Yasuda, H.; Yamamoto, H.; Yokota, K.; Miyake, S.; Nakamura, A. *J. Am. Chem. Soc.* **1992**, *114*, 4908.
- ¹⁰ See for instance: Hesse, M.; Meier, H.; Zeeh, B. *Spektroskopische Methoden in der organischen Chemie*, 3 Aufl., Georg Thieme Verlag: Stuttgart, Germany, 1987.
- ¹¹ Birnbaum, E.R.; Forsberg, J.H.; Marcus, Y. In *Gmelin Handbook of Inorganic Chemistry*, Moeller, T.; Schleitner-Rust, E., Eds., Springer Verlag: Berlin, 1981, System nr. 39, part D3, pp 85-93.

- 12 Evans, W.J. *J. Organomet. Chem.* **1990**, *433*, 79.
- 13 Schaverien, C.J.; Frijns, J.H.G.; Heeres, H.J.; Van den Hende, J.R.; Teuben, J.H.; Spek, A.L. *J. Chem. Soc., Chem. Commun.* **1991**, 642.
- 14 Evans, W.J.; Boyle, T.J.; Ziller, J.W. *Organometallics* **1993**, *12*, 3998.
- 15 Evans, J. W.; Dominguez, R.; Hanusa, T. P. *Organometallics* **1986**, *5*, 1291.
- 16 See for instance: Huheey, J.E. *Inorganic Chemistry*, Harper and Row: New York, 1983, 3rd ed, appendix E.
- 17 Jeske, G.; Lauke, H.; Mauermann, H.; Swepston, P.N.; Schumann, H.; Marks, T.J. *J. Am. Chem. Soc.* **1985**, *107*, 8091.
- 18 Evans, W.J.; Peterson, T.T.; Rausch, M.D.; Hunter, W.E.; Zhang, H.; Atwood, J.L. *Organometallics* **1985**, *4*, 554.