

University of Groningen

## On the interrelationships between morphology and movement in the tail of cichlid fish *Tilapia nilotica* (L.).

Videler, Johannes Jozef

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1975

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Videler, J. J. (1975). *On the interrelationships between morphology and movement in the tail of cichlid fish *Tilapia nilotica* (L.)*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

ON THE INTERRELATIONSHIPS BETWEEN  
MORPHOLOGY AND MOVEMENT IN THE TAIL OF  
THE CICHLID FISH *TILAPIA NILOTICA* (L.)

by

J. J. VIDELER

(Zoological Laboratory, University of Groningen, The Netherlands)

SUMMARY

1. An account is given of the morphological structures in the tail region of the "typical fish" *Tilapia nilotica* (fam. Cichlidae), emphasizing the mechanical properties of the caudal skeleton, the vertical septum, the joints between the fin rays and the body, the muscles and tendons and the skin fascia. Two large cartilaginous plates, situated in the vertical septum, are described for the first time.
2. Detailed descriptions of the transformations of the caudal peduncle and fin during short series of lateral undulations of the body and fin are presented.
3. Mutual comparisons of the morphological descriptions with the descriptions of the movements show how structures in the tail are suited to the transfer of propulsive forces, generated by the oscillating tail fin, from the fin to the body. The way structures transfer oscillating movements from the body to the fin and the way muscle activity and several properties of the joints are able to affect the final form of the fin stroke are discussed.
4. A brief account is given of electromyographic data collected and of some problems concerning the use of electromyography in the tail region.
5. It is suggested that the skin fascia collaborates with the myotomes and myosepts to bend the body and the fin.

CONTENTS

I. Introduction . . . . .	144
II. Methods . . . . .	145
III. Terminology and definitions . . . . .	146
1. Names of the skeleton and parts of the fin . . . . .	146
2. Names of soft parts like muscles, cartilage and connective tissue . . . . .	147
3. Names of planes and movements . . . . .	150
IV. Morphology . . . . .	152
1. Morphological technics . . . . .	152
2. The skeleton, fin rays and raylets . . . . .	152
3. The vertical septum and the connection with the caudal fin . . . . .	155
4. The muscles and tendons . . . . .	161
4.1. Marginal muscles . . . . .	161
4.2. Profundal muscles . . . . .	163
4.3. Superficial muscles and horizontal septum . . . . .	165
5. The skin . . . . .	167
V. Movements of the tail . . . . .	168
1. Technical approach . . . . .	169
2. Description of movements in the frontal planes . . . . .	172