

University of Groningen

The application of alkyllithium and alkylmagnesium reagents in cyclo(thia)phosphazene chemistry

Winter, Herman

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1986

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Winter, H. (1986). *The application of alkyllithium and alkylmagnesium reagents in cyclo(thia)phosphazene chemistry*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

$(\text{NPI}^i\text{Pr}(\text{NSOPh})_2$

ic, Pna2₁

075

$2\theta < 50^\circ$

arization effects and
decay; absorption using
min. and max. applied
factor (on F) 0.91 and
tively.

terson and Fourier
finement on F by block-
ast-squares techniques
ropic thermal parame-
e non-hydrogen atoms.
ms were introduced at
1.0 Å from their cen-
assuming sp² or sp³
on and refined in the
with an overall iso-
perature factor.

Summary

This thesis describes a study of the reactivity of cyclo(thia)phosphazenes towards alkyllithium and alkyl Grignard reagents. Attention is focused on three types of starting materials, viz. $(\text{NPF}_2)_n(\text{NSOX})_{3-n}$, $(\text{NPCl}_2)_n(\text{NSOX})_{3-n}$ ($n=1,2,3$; $X=\text{F}, \text{Cl}, \text{Ph}$) and $(\text{NPCl}_2)_4$.

Chapter I provides information about the characteristics of cyclo(thia)phosphazenes; the literature on reactions of $(\text{NPX}_2)_n$ ($n=3,4$; $X=\text{F}, \text{Cl}$) with the above-mentioned organometallic compounds is reviewed.

During attempts to synthesize fluorocyclothia-phosphazenes in the presence of KF/18-crown-6 ether in acetonitrile hydrolysis phenomena were observed. Chapter II describes reactions of $(\text{NPCl}_2)_n(\text{NSOX})_{3-n}$ ($n=1,2,3$; $X=\text{Cl}, \text{Ph}$) with H₂O in the presence of KCl and 18-crown-6 affording potassium crown ether salts of hydroxy derivatives of the ring compounds. Both mono- and disubstituted compounds could be isolated. It appears that in the case of $(\text{NPCl}_2)_3$ a non-geminal substitution pattern occurs, whereas $(\text{NPCl}_2)_2\text{NSOCl}$ reacts according to a geminal pathway. The latter observation has been confirmed by crystal structure determination of the compound $(\text{KC}_{12}\text{H}_{24}\text{O}_6)^+(\text{NHPU}_2\text{NPCl}_2\text{NSOCl})^-$.

Reactions of $\text{NPCl}_2(\text{NSOX})_2$ and $(\text{NPCl}_2)_2\text{NSOX}$ ($X=\text{F}, \text{Cl}, \text{Ph}$) with KSO₂F (Chapter III) lead to a variety of fluorocyclothiaphosphazenes, the ³¹P and ¹⁹F NMR spectra of which are analyzed by means of spectrum simulation. Trans-NPF₂(NSOPh)₂ reacts with ^tBuLi via nucleophilic substitution to give (1 α ,3 β ,5 α)-NPF^tBu(NSOPh)₂. No reactivity is observed towards ^tBuMgCl.

Chapter IV and V provide a detailed discussion

of the reactions of $(\text{NPCl}_2)_3$, trans and cis- $\text{NPCl}_2(\text{NSOPh})_2$ and $(\text{NPCl}_2)_2\text{NSOPh}$ with RLi ($\text{R}=\text{Me}, ^t\text{Bu}$) and 2-propanol. The reaction products, e.g.

$\text{NP}(\text{H})\text{O}^i\text{Pr}(\text{NPCl}_2)_n(\text{NSOPh})_{2-n}$ ($n=0,1,2$) are supposed to arise from a metal-halogen exchange process. Compound $(1\alpha,3\beta,5\alpha)\text{-NP}(\text{H})\text{O}^i\text{Pr}(\text{NSOPh})_2$ was characterized by means of an X-ray structure determination. The tetramer $(\text{NPCl}_2)_4$ also reacts with MeLi according to a metal-halogen exchange process, leading to unprecedented hydridocyclophosphazenes and bi(cyclophosphazenes); the X-ray structure determination of the bi(cyclophosphazene) $(\text{N}_4\text{P}_4\text{Cl}_6\text{Me})_2$ is discussed (Chapter VI).

Chapter VII reports the reactivity of trans- $\text{NPCl}_2(\text{NSOPh})_2$ towards $(\text{Bu}_3\text{PCuI})_4/\text{RMgCl}$ ($\text{R}=\text{Pr}, ^i\text{Pr}$) or RMgCl ($\text{R}=\text{Me}, ^t\text{Bu}$); again a metal-halogen exchange process is observed. The X-ray structure determinations of $(1\alpha,3\beta,5\alpha)\text{-NP}^n\text{Pr}^i\text{Pr}(\text{NSOPh})_2$ and $(1\alpha,3\beta,5\alpha)\text{-NPI}^i\text{Pr}(\text{NSOPh})_2$, in combination with that of cis- $\text{NPCl}_2(\text{NSOPh})_2$, provide information about the effects of the substituents on the ring skeleton. The difference in electronegativity between the sulfur and phosphorus centres within a PNS segment can be correlated with the difference between the NP and NS bond lengths.

The reactivity of the hydridocyclophosphazenes $\text{NP}(\text{H})\text{R}(\text{NP}(\text{OCH}_2\text{CF}_3)_2)_2$ ($\text{R}=\text{O}^i\text{Pr}, \text{Me}$) is discussed in Chapter VIII. Reactions of these compounds with p-nitrobenzaldehyde in the presence of small amounts of Et_3N or quinine afford $\text{NP}(\text{CHOHC}_6\text{H}_4\text{NO}_2)\text{R}(\text{NP}(\text{OCH}_2\text{CF}_3)_2)_2$. Asymmetric catalysis in the case of quinine is demonstrated.