

University of Groningen

## Cholinergic neurotransmission in human and guinea pig airways Function and dysfunction of muscarinic receptor subtypes

Berge, Ronald Egbert Joseph ten

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

1994

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Berge, R. E. J. T. (1994). *Cholinergic neurotransmission in human and guinea pig airways Function and dysfunction of muscarinic receptor subtypes*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## SUMMARY

The diameter of the airways, and as a result airway resistance, are controlled by the contractile tone of smooth muscle layers within the airway wall. In asthma, there is shortness of breath (bronchial obstruction), wheezing and coughing, as a result of increased contractile tone of the airway smooth muscle. In addition, other factors such as oedema of the airway wall or increased mucus production by airway submucosal glands contribute to the obstruction. In allergic asthma, inhalation of a critical amount of allergen provokes an early allergic response, which in most humans consists of severe shortness of breath during 1 to 3 hours. In guinea pigs, the experimental animal used in the studies described in this thesis, this phase lasts for 2 to 5 hours (**Chapters 7 and 8**). After the early phase, there is an interval of several hours without obstruction, followed at approximately 8 hours after antigen challenge by the late obstructive reaction that lasts for 10 to 15 hours. Both after the early and after the late phase there is increased susceptibility of the airways to non-specific non-allergic, physical, chemical and pharmacological stimuli; this bronchial hyperreactivity (BHR) is one of the major characteristics of asthma. Both the airway obstruction and BHR are partly caused by increased contractility of the smooth muscle tissue.

The most important contractile nervous system in the airways is the parasympathetic or cholinergic (named after its neurotransmitter acetylcholine) system. A further contractile nerve system is the so called excitatory non-adrenergic non-cholinergic (e-NANC) system; it releases contractile neuropeptides called tachykinins like substance P and neurokinin A. Relaxation of airway smooth muscle (and thus dilation of the airways) can be produced by circulating adrenaline or by the neurotransmitter of adrenergic nerves (sympathetic nervous system), noradrenaline. In connection to airway smooth muscle contraction, mediators of the inhibitory NANC system can also be released, leading to decrease of contractile tone (**Chapter 1**). The vagal nerve pathways of the cholinergic system consist of preganglionic nerves, that run from the central nervous system down to the ganglion, and short postganglionic nerves that project at only a few nanometers distance from the smooth muscle cells. Muscarinic receptors have been found to be involved in vagal neurotransmission in ganglia (postjunctional  $M_1$  subtype receptors), as well as in negative feedback of neurotransmitter release (by prejunctional  $M_2$  subtype receptors on postganglionic nerve endings) and in the contraction of smooth muscle (via postganglionic postjunctional  $M_3$  subtype receptors). In **Chapters 3 to 8** of this thesis, the role of these muscarinic receptor subtypes in the airways of guinea pig and man, and the putative alterations in their function under conditions of allergic bronchoconstriction and bronchial hyperreactivity, have been investigated.

In **Chapter 3**, the function of the prejunctional  $M_2$  receptors in isolated guinea pig tracheal ring preparations was examined. To this end, the rings were mounted for isotonic recording, and twitch contractions were elicited by electrical field stimulation (EFS) (30Hz, 0.5ms, 8V over the electrodes, 5s per min). It was found that low concentrations of  $M_2$ -selective muscarinic receptor antagonists (gallamine, methoctramine, AF-DX 116, AQ-RA 741) increased the twitch contractions - as a result of blockade of the negative feedback loop - whereas higher concentrations decreased twitch contractions through blockade of postjunctional  $M_3$  receptors. It was observed that the extent of the increase in twitch height induced by these  $M_2$ -selective antagonists was related to their selectivity for  $M_2$  over  $M_3$  receptors, and that the concentration dependence of this effect was related to their  $M_2$  receptor affinity. From the additional examination of non- and  $M_3$ -selective antagonists (which only produced twitch decrease) it was inferred that the inhibition of the twitch contractions at higher concentrations was strongly correlated to the  $M_3$  receptor affinity of all types of antagonist. The findings in this chapter show that putative dysfunctions of  $M_2$  and  $M_3$  receptors in sensitised and/or challenged animals can be studied using this experimental set-up.

The functional presence of ganglionic  $M_1$  receptors that had been suggested to be involved in facilitatory neurotransmission in human lungs had not been satisfactorily demonstrated in the guinea pig at the start of our experiments (**Chapter 4**); in fact such receptors were postulated in sympathetic and parasympathetic ganglia, but also denied in the latter. In our experiments, we used isolated preparations of the main bronchus which were carefully dissected with the supplying vagus nerve left intact. Twitch contractions were alternately elicited by (1) EFS, which releases acetylcholine from the vagus nerve endings without the intervention of the ganglia, and (2) electrical stimulation of the vagus nerve (VS) which includes neurotransmission through the ganglia. The inhibitory effect of the  $M_1$  receptor selective muscarinic antagonist pirenzepine on VS- and EFS-induced twitch contractions was established under various conditions. Using identical stimulation conditions for VS and EFS (30Hz, 0.5ms, 8V over the electrodes, 5s per min), VS-induced contractions were smaller than EFS-induced contractions, probably due to ganglionic filtering, and pirenzepine was 2.3-fold selective in antagonising VS-induced contractions, which small selectivity does not readily indicate the involvement of  $M_1$  receptors in vagal bronchoconstriction. This was confirmed by experiments in which stimulation frequency for EFS was decreased to obtain twitch contractions that matched those elicited by VS; under these conditions pirenzepine showed no preferential inhibition of VS-induced twitches at all. However, high affinity inhibition of VS-induced twitches was achieved when stimulation was prolonged (for both VS and EFS) to reach plateau contractions, in the presence of  $\beta$ -adrenoceptor and  $M_2$  receptor blockade to exclude any feedback regulation, and when the major component of ganglionic transmission, i.e. nicotinic receptors, were partially blocked using hexamethonium. It thus appears that muscarinic  $M_1$  receptors are indeed

operational in  
and are invo  
indicated to e  
of importance  
not.

In **Ch**  
of endogenou  
endings in iso  
the function o  
endogenous a  
indicate that  
differences in  
been reported  
of buffer, an  
chromatograp  
relatively hig  
neostigmine),  
antagonist at  
agonist metha  
conditions sy  
receptors are  
activated. Un  
absence of n  
indeed inhibi  
pilocarpine,  
These result  
through prej  
neurotransmi

In **Ch**  
autoreceptors  
from thoraco  
strips from t  
of bronchi a  
method desc  
(gallamine, A  
in guinea pig  
in all prepar  
terminal bro  
smaller bron  
bronchial pre  
atropine in a

in isolated guinea  
were mounted for  
l field stimulation  
as found that low  
nists (gallamine,  
contractions - as a  
ner concentrations  
receptors. It was  
these M<sub>2</sub>-selective  
tors, and that the  
otor affinity. From  
ich only produced  
tractions at higher  
y of all types of  
ons of M<sub>2</sub> and M<sub>3</sub>  
g this experimental

been suggested to  
been satisfactorily  
**Chapter 4**); in fact  
ganglia, but also  
tions of the main  
nerve left intact.  
ases acetylcholine  
, and (2) electrical  
ssion through the  
carinic antagonist  
hed under various  
30Hz, 0.5ms, 8V  
maller than EFS-  
nzepine was 2.3-  
electivity does not  
constriction. This  
FS was decreased  
r these conditions  
at all. However,  
stimulation was  
he presence of β-  
ilation, and when  
rs, were partially  
eceptors are indeed

operational in parasympathetic ganglia of the guinea pig bronchus, as in human lungs, and are involved in the amplification of postganglionic impulses. Future studies are indicated to establish whether alterations in the involvement of these M<sub>1</sub> receptors are of importance in bronchial hyperreactivity, be it in response to allergic reactions or not.

In **Chapter 5**, experiments were aimed at measuring directly the concentrations of endogenous (i.e. not radiolabeled) acetylcholine that are released from vagus nerve endings in isolated guinea pig trachea, in order to obtain *direct* information concerning the function of inhibitory M<sub>2</sub> autoreceptors on these nerve endings; the measurement of endogenous acetylcholine may have relevant advantages, since there is evidence to indicate that radiolabeling does not affect all pools of acetylcholine equally and differences in modulation of radiolabeled and endogenous acetylcholine release have been reported. To achieve this, tracheas were electrically stimulated in a small volume of buffer, and samples were immediately analysed using high performance liquid chromatography with electrochemical detection (HPLC/ECD). It was found that under relatively high stimulation intensity (30Hz, 0.5ms, 8V, for 5 min in the presence of neostigmine), resulting in high levels of acetylcholine release, the muscarinic receptor antagonist atropine more than doubled this release, whereas the muscarinic receptor agonist methacholine did not show any effect. These results suggested that under these conditions synaptic concentrations of acetylcholine are that high that prejunctional M<sub>2</sub> receptors are maximally activated; consequently, they can be blocked but not further activated. Under mild stimulation conditions (16Hz, 0.1ms, 8V, for 5 min in the absence of neostigmine), resulting in 20-fold lower acetylcholine levels, methacholine indeed inhibited this release; this was similarly true for the partial muscarinic agonist pilocarpine, which under high stimulation behaved as an antagonist (i.e. like atropine). These results showed that the negative feedback control of acetylcholine release through prejunctional M<sub>2</sub> autoreceptors is directly dependent on the synaptic levels of neurotransmitter, and thus on nerve activity.

In **Chapter 6**, the presence and function of prejunctional inhibitory muscarinic autoreceptors in human airway preparations (derived from human lung tissue obtained from thoracotomies and transplant donors) was investigated. Isolated smooth muscle strips from trachea and main bronchus and ring preparations from various generations of bronchi and bronchioli (3.5-0.7 mm I.D.) were subjected to EFS, and using the method described in Chapter 3, it was studied whether M<sub>2</sub>-selective antagonists (gallamine, AQ-RA 741) would potentiate twitch contractions as previously observed in guinea pig trachea. Remarkably, such increase in twitch contractions was observed in all preparations from central airways (trachea, main bronchus), and in half of the terminal bronchi (1-2 mm I.D.); no prejunctional M<sub>2</sub> receptor function was found in smaller bronchi (<1 mm I.D.). In addition, the direct release of acetylcholine from bronchial preparations (1.5-3.5 mm I.D.) was significantly increased by the addition of atropine in all tests performed. In conclusion, prejunctional inhibitory M<sub>2</sub> autoreceptors

are functionally present in human airways, but it remains to be established why there is (individual) variability in peripheral airways.

In **Chapter 7** it was investigated to what extent prejunctional  $M_2$  receptor function in guinea pig trachea is changed after the event of an allergic asthmatic reaction. In these experiments we made use of an animal model of allergic asthma developed in our laboratory, in which lung function is being measured on-line in conscious, unrestrained guinea pigs that have been sensitised to and subsequently challenged by inhalation with the allergen ovalbumin; lung function is monitored as pleural pressure using a small fluid-filled balloon-cannula implanted by microsurgery. As in human asthmatics, these animals develop early and late phase obstructive reactions in response to antigen; in between (5-8 hours after antigen challenge) lung function is normal. Immediately after the early and after the late reaction (i.e. 6 and 24 hours after challenge, respectively), bronchial responsiveness to histamine was measured. Prejunctional muscarinic  $M_2$  receptor function was investigated using the method described in chapter 3, in isolated tracheal preparations obtained from animals that had been terminated just after the early or late response. It was observed that after the early allergic reaction, when BHR was at its highest level,  $M_2$  receptors on cholinergic nerve terminals were largely dysfunctional, because the potentiation of the twitch responses by all  $M_2$ -selective antagonists (gallamine, methoctramine, AQ-RA 741, AF-DX 116) was significantly and markedly decreased compared to controls. Moreover, the degree of dysfunction was correlated to the magnitude of the early response. After the late reaction, dysfunction was already restored to a large extent, but BHR was still present, although smaller than after the early response. These results strongly suggest that BHR to histamine after the early allergic response is partly due to the fact that prejunctional inhibitory  $M_2$  autoreceptors are severely dysfunctional, since it has been shown that histamine activates sensory nerve endings in the airways leading to vagal reflex activity and, consequently, to acetylcholine release from vagal nerve endings. Dysfunctional  $M_2$  receptors would lead to increased acetylcholine release and thus to increased bronchoconstriction. Recently, it was indeed shown by Santing that inhaled ipratropium bromide strongly decreased BHR to histamine in our guinea pigs. Finally, it should be mentioned that postsynaptic  $M_2$  receptors in the hearts of the animals were not affected by the allergic responses, indicating that inhalation of antigen only induces these effects locally.

In **Chapter 8**,  $M_2$  receptor function after the allergic response was investigated again, but now *in vivo*, in conscious, unrestrained animals, in order to establish the extent and time course of the putative dysfunction in the intact respiratory tract, and under physiological conditions. Guinea pigs were subjected to microsurgery to implant an intrapleural balloon-cannula (as in Chapter 7) as well as a bipolar stimulation electrode around the right vagus nerve; in contrast to other studies using anaesthetized animals in which the vagus nerve was not sectioned, native nervous activity was preserved. In this way, electrical stimulation of the nerve with stepwise-increasing

frequencies and  
monitored as i  
autoreceptors w  
inhaled in re  
bronchoconstric  
prejunctional re  
frequency-respo  
a result of blo  
experiments we  
observed that g  
still blocked co  
were rather sim  
chapter, BHR to  
and smaller (bu  
those obtained i  
airways develop  
early allergic re  
strongly contrib  
sensory nerve en

The mech  
allergic response  
an inflammatory  
has been shown  
basic protein, v  
reversible (allo  
receptors were f  
as in the *in vivo*

Finally, i  
NANC neurotra  
of prejunctional  
identity of these  
(30Hz, 0.5ms,  
contraction and  
release of the ne  
adrenoceptor ag  
studied both on  
NANC contracti  
receptor antagon  
contractions ind  
was found that c  
adrenergic agon

established why there is  
functional  $M_2$  receptor  
of an allergic asthmatic  
model of allergic asthma  
being measured on-line in  
and subsequently  
function is monitored as  
performed by microsurgery.  
late phase obstructive  
(antigen challenge) lung  
reaction (i.e. 6 and 24  
hours) to histamine was  
investigated using the  
obtained from animals  
was observed that after  
level,  $M_2$  receptors on  
the potentiation of the  
methoctramine, AQ-RA  
compared to controls.  
magnitude of the early  
response to a large extent,  
response. These results  
response is partly due to  
ly dysfunctional, since  
in the airways leading  
arise from vagal nerve  
acetylcholine release and  
shown by Santing that  
in our guinea pigs.  
in the hearts of the  
that inhalation of  
response was investigated  
in order to establish the  
of the respiratory tract, and  
microsurgery to implant  
bipolar stimulation  
using anaesthetized  
nervous activity was  
stepwise-increasing

frequencies induced increasing levels of bronchoconstriction in conscious animals, monitored as increase in respiration amplitude. The function of prejunctional  $M_2$  autoreceptors was studied using the  $M_2$ -selective antagonist gallamine which, when inhaled in relatively low concentrations, increased nerve stimulation-induced bronchoconstriction at each frequency, as a result of the blockade of these inhibitory prejunctional receptors; with higher concentrations of gallamine, the left-shift of the frequency-response curve was revealed and the maximum obstruction was depressed as a result of blockade of postjunctional (smooth muscle)  $M_3$  receptors. When these experiments were repeated (in the same animal) after the early allergic response, it was observed that gallamine no longer potentiated nerve-induced bronchoconstriction, but still blocked constriction in the highest dose. After the late reaction, gallamine effects were rather similar to control, i.e. before antigen challenge. As in the previous chapter, BHR to histamine was measured and found to be rather high after the early and smaller (but still significant) after the late response. These results fully confirmed those obtained in isolated tracheal preparations *in vitro*, in showing that guinea pig airways develop a severe dysfunction of prejunctional  $M_2$  receptors already during the early allergic response after antigen challenge. The resulting vagal hyperactivity may strongly contribute to the bronchial hyperreactivity to inhaled substances stimulating sensory nerve endings.

The mechanism underlying the almost complete  $M_2$  receptor dysfunction after an allergic response has been proposed to be related to the eosinophilic granulocyte, being an inflammatory cell that infiltrates the airways already during the early response. It has been shown that these cells produce and release a cytotoxic protein called major basic protein, which appears to block muscarinic  $M_2$  (but not  $M_3$ ) receptors in a reversible (allosteric) fashion. In agreement with this, postjunctional smooth muscle  $M_3$  receptors were found to be unaffected by the allergic responses, in the *in vitro* as well as in the *in vivo* studies.

Finally, in **Chapter 9** it was investigated to what extent cholinergic and e-NANC neurotransmission in guinea pig main bronchus are under the inhibitory control of prejunctional  $\beta$ -adrenoceptors; this has been postulated by others but the subtype identity of these  $\beta$ -adrenoceptors remained controversial. EFS was performed for 20s (30Hz, 0.5ms, 8V over the electrodes) to produce both a rapid cholinergic twitch contraction and a slow, 15-min e-NANC wave-like contraction, which results from the release of the neuropeptide transmitter neurokinin A. The effects of  $\beta_2$ - and  $\beta_3$ -selective adrenoceptor agonists (fenoterol and salbutamol, and BRL 37344, respectively) were studied both on the height of the cholinergic twitch and on the magnitude of the e-NANC contraction, in the absence and as well as in the presence of the  $\beta_2$ -selective receptor antagonist ICI 118,551; these inhibitory effects were compared to those on contractions induced by exogenous methacholine and neurokinin A, respectively. It was found that e-NANC contractions were inhibited by lower concentrations of the  $\beta$ -adrenergic agonists compared to neurokinin A-induced contractions, and that this

response only involved  $\beta_2$  receptors; such preferential inhibition of EFS-induced contractions was not observed for the cholinergic pathway. It was concluded that the release of e-NANC neurotransmitter(s) in guinea pig main bronchus is under the inhibitory control of prejunctional  $\beta_2$ -adrenoceptors, but that there is no such control on cholinergic nerve endings.

In conclusion, the studies described in this thesis have demonstrated the functional presence of  $M_1$  receptors in parasympathetic ganglia of isolated guinea pig airways, and of postganglionic, prejunctional  $M_2$  autoreceptors in guinea pig and human airways, both by indirect (contraction) and direct (endogenous acetylcholine release) measurement. Most importantly, it was shown both *in vitro* and *in vivo* that the prejunctional  $M_2$  receptors become dysfunctional after the early allergic response in the guinea pig isolated trachea and in the respiratory tract of the intact (freely moving) animal, but recover during the late phase, which indicates that  $M_2$  receptor dysfunction may contribute to bronchial hyperresponsiveness to sensory stimuli after the early but not after the late response following a single allergen challenge. It remains to be established whether dysfunction (i.e. increased reactivity) of ganglionic  $M_1$  receptors contributes to BHR as well. Similarly, further investigations are necessary to establish the precise relationship between  $M_2$  receptor dysfunction in human asthmatics (which has been observed by studying  $SO_2$ -induced bronchoconstriction) and the observations made in this thesis.

The new methods developed in this study may be of great value for future investigations on prejunctional control mechanisms of acetylcholine release in isolated human and guinea pig airways and the regulation of vagally induced bronchoconstriction in conscious, unrestrained guinea pigs, both under physiological and pathophysiological conditions. Especially investigations on the direct and indirect role of the e-NANC and i-NANC systems are challenging.

Mens  
produceren  
gassen tuss  
luchtwege  
(longblaasje  
bronchiën r  
van belang  
mond en al  
eventuele so  
luchtwege  
wand van d

Astm  
door korta  
astmatici,  
belemmerd,  
spierweefse  
luchtwege  
zich vaak  
allergeen e  
duurt bij m  
(hoofdstuk  
openbaart  
allergeen e  
vroeg als  
aspecifieke  
deze bronc  
obstructie e  
spierlaag.

Het  
of parasym  
afgifte var  
overdracht)  
zogenaamd  
deze zenuv  
weefsel (ve  
circulerend  
afkomstige  
van het inl  
verlagen (t