

University of Groningen

## Boundary value problems in diffraction theory and lifting surface theory

Boersma, Johannes

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*

2008

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Boersma, J. (2008). *Boundary value problems in diffraction theory and lifting surface theory*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

## § 1. Introduction

The present paper deals with a number of boundary value problems in diffraction theory and wing lifting surface theory with special reference to circular disks and apertures and to slits.

The problems of the diffraction of an incident wave (scalar or electromagnetic) by a circular disk, circular aperture, strip or slit have received considerable attention during the last decades. For a survey of the various methods of solution the reader is referred to the review articles of Bouwkamp [8] and of Hönl, Maue and Westpfahl [19]. In this paper we are especially interested in low-frequency expansions for the field quantities, valid when  $ka$  or  $kb$  is small, where  $k$  is the wave number,  $a$  is the radius of the disk or aperture,  $2b$  is the width of the strip or slit. These expansions will be obtained from Fredholm integral equations of the second kind, the kernels of these integral equations being small when  $ka$  or  $kb$  is small.

The reduction of the scalar diffraction problem for a circular disk or circular aperture to Fredholm integral equations has been performed by several authors. We mention Magnus [34], Jones [23], Bazer and Brown [2], Bazer and Hochstadt [3], Heins and MacCamy [17], Noble [41], Collins [9e, f] and Williams [51].

Bazer and Brown [2], Bazer and Hochstadt [3], Collins [9e, f] start from certain integral representations for the transmitted or scattered wave. These integral representations which contain an unknown function, are designed to satisfy all conditions of the problem but one. The latter condition leads to a Fredholm integral equation of the second kind for the unknown function. A related integral equation with the same kernel was derived before by Jones [23], using a very complicated method. The above-mentioned integral representations may be considered as generalizations of similar representations exploited to solve some problems in potential theory, compare Love [31], Green and Zerna [14], Collins [9]. In an appendix to [3b] Bazer and Hochstadt show their integral representations to be related to a representation for