

University of Groningen

The Smallest Resonance Energy Transfer Acceptor for Tryptophan

Broos, Jaap; Pas, Hendri H.; Robillard, George T.

Published in:
Biophysical Journal

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2002

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Broos, J., Pas, H. H., & Robillard, G. T. (2002). The Smallest Resonance Energy Transfer Acceptor for Tryptophan. *Biophysical Journal*.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

THE SMALLEST RESONANCE ENERGY TRANSFER ACCEPTOR FOR TRYPTOPHAN

Jaap Broos, Erwin P.P. Vos, Hendri H. Pas, George T. Robillard; University of Groningen, Nijenborgh 4, Groningen, 9747 AG Netherlands

Diazirine compounds have found widespread use as photoactivatable reagents. In this work we report on the extension of the utility of diazirine ligands in protein science. The diazirine moiety, which absorb between 330-380 nm, is used as an acceptor for tryptophan (Trp) and Trp analogs, like 5-fluoroTrp, in resonance energy transfer (RET) experiments. Distances up to 18 X can be determined with high accuracy. The principle is demonstrated for a diazirine derivative of mannitol, 2-azi-2-deoxy-D-arabino-hexitol (azi-mannitol) and a dozen single-Trp-containing mutants of the membrane bound mannitol transporter, EII^{mtl}, from *E. coli*. The diazirine moiety is photostable under the experimental conditions. The ambiguity about the exact position of the chromophoric group relative to the ligand due to the use of a flexible linker is absent in azi-mannitol, since the chromophoric group is an integral part of the substrate. The small size of the diazirine group allows its isosteric introduction in biological ligands without a large effect on the biological activity. In conclusion, the versatility of Trp as an intrinsic spectroscopic probe of proteins and the small size of the diazirine group make this a very attractive donor-acceptor couple for accurate RET distance information in protein chemistry.