

University of Groningen

A comprehensive estimate of recent carbon sinks in China using both top-down and bottom-up approaches

Jiang, Fei; Chen, Jing M; Zhou, Lingxi; Ju, Weimin; Zhang, Huifang; Machida, Toshinobu; Ciais, Philippe; Peters, Wouter; Wang, Hengmao; Chen, Baozhang

Published in:
Scientific Reports

DOI:
[10.1038/srep22130](https://doi.org/10.1038/srep22130)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Jiang, F., Chen, J. M., Zhou, L., Ju, W., Zhang, H., Machida, T., Ciais, P., Peters, W., Wang, H., Chen, B., Liu, L., Zhang, C., Matsueda, H., & Sawa, Y. (2016). A comprehensive estimate of recent carbon sinks in China using both top-down and bottom-up approaches. *Scientific Reports*, 6, [22130].
<https://doi.org/10.1038/srep22130>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

SCIENTIFIC REPORTS



OPEN

A comprehensive estimate of recent carbon sinks in China using both top-down and bottom-up approaches

Received: 26 August 2015
Accepted: 01 February 2016
Published: 29 February 2016

Fei Jiang¹, Jing M. Chen^{1,2}, Lingxi Zhou³, Weimin Ju¹, Huifang Zhang⁴, Toshinobu Machida⁵, Philippe Ciais⁶, Wouter Peters^{7,8}, Hengmao Wang¹, Baozhang Chen⁴, Lixin Liu³, Chunhua Zhang^{1,10}, Hidekazu Matsueda⁹ & Yousuke Sawa⁹

Atmospheric inversions use measurements of atmospheric CO₂ gradients to constrain regional surface fluxes. Current inversions indicate a net terrestrial CO₂ sink in China between 0.16 and 0.35 PgC/yr. The uncertainty of these estimates is as large as the mean because the atmospheric network historically contained only one high altitude station in China. Here, we revisit the calculation of the terrestrial CO₂ flux in China, excluding emissions from fossil fuel burning and cement production, by using two inversions with three new CO₂ monitoring stations in China as well as aircraft observations over Asia. We estimate a net terrestrial CO₂ uptake of 0.39–0.51 PgC/yr with a mean of 0.45 PgC/yr in 2006–2009. After considering the lateral transport of carbon in air and water and international trade, the annual mean carbon sink is adjusted to 0.35 PgC/yr. To evaluate this top-down estimate, we constructed an independent bottom-up estimate based on ecosystem data, and giving a net land sink of 0.33 PgC/yr. This demonstrates closure between the top-down and bottom-up estimates. Both top-down and bottom-up estimates give a higher carbon sink than previous estimates made for the 1980s and 1990s, suggesting a trend towards increased uptake by land ecosystems in China.

The carbon balance of China is characterized by the World's highest emissions of CO₂ from fossil fuel use and substantial carbon sequestration in intensively managed ecosystems. The large land area of China (6.4% of the global land mass), coupled to its rapid economic development, its large food production, and its recent large-scale afforestation practices puts its carbon cycle in the center of current global carbon cycle research. Top-down atmospheric inversions^{1,2} have used globally distributed stations measuring atmospheric CO₂ mole fraction observations to provide estimates of surface-atmosphere CO₂ fluxes over large (>10⁶ km²) spatial areas. One limitation of inversions is the insufficient density of atmospheric stations over continental regions. In this study, we derive new top-down calculations of China's CO₂ budget, by combining new atmospheric CO₂ observations within and around China, with two independent atmospheric inversion systems. We additionally conduct a synthesis of the bottom-up carbon budget of China's terrestrial ecosystems to gauge the convergence between these independent streams of information.

¹Jiangsu Provincial Key Laboratory of Geographic Information Science and Technology, International Institute for Earth System Science, Nanjing University, Nanjing 210046, China. ²Department of Geography, University of Toronto, Ontario M5S3G3, Canada. ³Chinese Academy of Meteorological Sciences, China Meteorological Administration, Beijing 100081, China. ⁴State Key Laboratory of Resources and Environment Information System, Institute of Geographic Sciences and Natural Resources Research, Chinese Academy of Sciences, Beijing 100101, China. ⁵National Institute for Environmental Studies, 305-8506 Tsukuba, Japan. ⁶Laboratoire des Sciences du Climat et de l'Environnement, CEA CNRS UVSQ, 91191 Gif sur Yvette, France. ⁷Department of Meteorology and Air Quality, Wageningen University and Research Center, 6708 PB Wageningen, The Netherlands. ⁸University of Groningen, Centre for Isotope Research, 9747 AG Groningen, The Netherlands. ⁹Geochemical Research Department, Meteorological Research Institute, Tsukuba 305-0052, Japan. ¹⁰School of Geography and Planning, Ludong University, Yantai 264025, China. Correspondence and requests for materials should be addressed to J.M.C. (email: jing.chen@utoronto.ca)




Figure 1. Inverted carbon sinks in China during 2006–2009 from two inversion systems. Bayesian Inversion (BI) and Carbon Tracker-China (CTC). Values have been adjusted with the national CO₂ emissions from fossil fuel burning, cement manufacture, and gas flaring of 1.90 PgC yr⁻¹ during 2006–2009 reported by the Carbon Dioxide Information Analysis Center¹⁵. Blue: constrained only with global CO₂ datasets; orange: constrained with additional China Meteorological Administration (CMA)'s measurements (3 sites); and green: constrained with additional CMA and CONTRAIL aircraft CO₂ measurements.

Results

Top-down estimate. Atmospheric inversions³, quantify net CO₂ fluxes at the surface of the Earth, based on transport models and atmospheric CO₂ observations. In this process, a higher density of observations allows more detailed estimates of fluxes. In China, only one high altitude monitoring station (Mt Waliguan) in the western part of China⁴ has been available to constrain the published estimates up until 2006. Since then, the Chinese Meteorological Administration (CMA) installed three additional surface GHG monitoring stations⁵. In addition, CO₂ measurements on board of passenger aircraft, with vertical profiles at selected airport locations and horizontal transects at the cruising altitude of aircraft, have been acquired over Asia and Europe⁶ since July 2005 by the Comprehensive Observation Network for Trace gases by AirLiner project (CONTRAIL). These CO₂ observations form the basis for the revised top-down estimate of the CO₂ budget of China.

We use two well-established inversion systems, a nested Bayesian inversion (BI) system⁷ and the CarbonTracker-China (CTC) system⁸ to estimate CO₂ fluxes in China during the 2000s. Details of both systems are provided in the Supplementary Material. Using the inversion systems with only the Mt Waliguan CO₂ record as constraint, we estimate over China a net sink of atmospheric CO₂ during 2006–2009, excluding CO₂ emissions from fossil fuels and cement. The CO₂ sink estimates are of 0.29 ± 0.21 and 0.20 ± 0.36 PgC yr⁻¹ (1-sigma posterior Gaussian uncertainties), respectively in each inversion. These mean values are close to previous inversion estimates, in the range of 0.16–0.35 PgC yr⁻¹ during 1996–2009^{9–12}. When the three new CMA stations are assimilated into the BI and CTC systems, the inverted terrestrial CO₂ sink in China increases to 0.43 ± 0.19 and 0.29 ± 0.35 PgC yr⁻¹, respectively. When both CMA and CONTRAIL data are assimilated, the sink further increases to 0.51 ± 0.18 and 0.39 ± 0.33 PgC yr⁻¹, respectively (Fig. 1). The two inversion systems thus consistently show that when new CO₂ measurements within or around China are included, the inverted CO₂ sink in China gets larger and its uncertainty is reduced. With the new CO₂ data added, the mean inverted CO₂ sink in China is 0.45 ± 0.25 PgC yr⁻¹, which is a higher than previous inversions. In inversions, the inferred sinks depend on the value being assumed for CO₂ emissions from fossil fuel burning and cement production (FFCO₂). There is a rather large uncertainty of FFCO₂ in China, as evidenced by differences between published estimates^{13,14}. In this study, we used as a reference FFCO₂ from the Carbon Dioxide Information Analysis Center (CDIAC)¹⁵, of 1.90 PgC yr⁻¹ during 2006–2009 (Table 1). Using the value of FFCO₂ recently produced by Liu *et al.*¹⁴, based on a downward revision of the carbon content of coal burned in China, which is 9% lower than CDIAC, would lead to a mean inverted CO₂ sink in China of 0.28 ± 0.25 PgC yr⁻¹.

Bottom-up estimate. The top-down estimates of the carbon sources and sinks excluding FFCO₂ should equal the change in carbon stocks in the various reservoirs involved in carbon exchange. Various methods, often referred to as bottom-up, have been developed to estimate these carbon stock changes. In order to evaluate specifically the new top-down estimate of China's terrestrial ecosystems, we updated bottom-up carbon exchange estimate to cover the period of 2006–2009.

We reconstructed carbon stock changes of vegetation and soil in China during 2000s (Table 2). For vegetation carbon stock change, forest is the most important biome. Based on the 6th (1999–2003) and 7th (2004–2008) national forest inventories, Zhang *et al.*¹⁶, Guo *et al.*¹⁷ and Pan *et al.*¹⁸ estimated that forest biomass carbon stock accumulated at a rate of 0.174, 0.104, and 0.115 PgC yr⁻¹ during the 2000s, respectively. We use the mean and standard deviation of these three estimates, which is 0.13 ± 0.038 PgC yr⁻¹. This is larger than the value of 0.075 ± 0.035 PgC yr⁻¹ reported by Piao *et al.*⁹ during the 1980s and 1990s, suggesting that forest biomass carbon gains in 2000s significantly increased from 1980s and 1990s. In addition, short rotation forests and bamboo

	BI	CTC
Prior bio flux	-0.10 ± 0.26	-0.092 ± 0.49
Fire emission	0.010	0.022
Fossil fuel emission	1.94	2.01
Optimized bio flux (Case_1) ¹	-0.34 ± 0.21	-0.33 ± 0.36
Optimized bio flux (Case_2) ¹	-0.48 ± 0.19	-0.42 ± 0.35
Optimized bio flux (Case_3) ¹	-0.56 ± 0.18	-0.51 ± 0.33
CDIAC	1.90	1.90
Adjusted bio flux (Case_1) ²	-0.29 ± 0.21	-0.20 ± 0.36
Adjusted bio flux (Case_2) ²	-0.44 ± 0.19	-0.29 ± 0.35
Adjusted bio flux (Case_3) ²	-0.51 ± 0.18	-0.39 ± 0.33

Table 1. Prior, optimized, and adjusted carbon flux from the inversion systems in China (PgC yr⁻¹) for the period 2006–2009 (positive values represent carbon source, negative values represent carbon sink). Case_1: inversion result constrained with global CO₂ datasets only; Case_2: result of additional constraint with China Meteorological Administration (CMA)'s measurements (3 sites); Case_3: result of further constraint with CONTRAIL aircraft CO₂ measurements. ¹inverted using inversion systems, and exclude fossil fuel and biomass burning CO₂ emissions. ²further adjusted with the national CO₂ emission reported in CDIAC, only exclude fossil fuel CO₂ emissions (Adjusted bio flux = Fossil fuel emission + Fire emission + Optimized bio flux – CDIAC).

Category		Method	Area (1.0e6 ha)	Carbon balance (PgC yr ⁻¹)	Period	Ref.
Vegetation	Forest stands	Inventory	149	0.174	1999–2008	16
		Inventory	156	0.115	2000–2007	18
		Inventory	149	0.104	1999–2008	17
	Forest ave.		151	0.13 ± 0.04		
	Economic forests	Inventory	21	0.00	1999–2008	16
		Inventory	21	0.00	1999–2008	17
	Economic Forest ave.		21	0.00		
	Bamboo	Inventory	5.1	0.013	1999–2008	16
		Inventory	5.1	0.005	1999–2008	17
	Bamboo ave.	Inventory	5.1	0.009 ± 0.006		
	Woodlands	Inventory	5.4	-0.002 ± 0.001	1999–2008	16
	Shrub	Inventory	49.5	0.019 ± 0.013	1999–2008	16
	Tree on non-forest lands			-0.001 ± 0.001	1999–2008	16
Grass	Inventory	331	0.007 ± 0.003	1980s,1990s	9	
Subtotal			0.17 ± 0.060			
Soil	Forest	InTEC model	155	0.068 ± 0.034	1999–2008	This study
		Inventory	156	0.060 ± 0.030	2000–2007	18
	Forest ave.		155	0.064 ± 0.030		
	Shrub	Statistic model	215	0.039 ± 0.009	1980s,1990s	9
		Process model	141	0.012 ± 0.005	1981–2000	20
	Shrub ave.			0.026 ± 0.019		
	Crop	Aggregate	130	0.021 ± 0.004	1980s,1990s	21
Grass	Aggregate	331	0.005 ± 0.002	1980s,1990s	21	
Subtotal			0.12 ± 0.060			
Total			0.29 ± 0.12			

Table 2. Carbon accumulated in China's terrestrial ecosystems during 2000s.

plantations were estimated to have accumulated 0.009 ± 0.006 PgC yr⁻¹ in total^{16,17}, and woodlands, shrub, tree on non-forest lands were estimated to have a sink of 0.016 ± 0.011 PgC yr⁻¹ in total¹⁶. Due to lack of more recent research results for grasslands, we use the same estimate than Piao *et al.*⁹ of 0.007 ± 0.003 PgC yr⁻¹. In total, vegetation biomass in China accumulated 0.17 ± 0.060 PgC yr⁻¹ of carbon during 2000s.

For soil carbon stock (SOC) change, we first estimate the rate of change of forest soil carbon stock, include dead wood, litter and soil carbon, to be 0.068 ± 0.034 PgC yr⁻¹ during 2000s using the InTEC model¹⁹. This estimate is consistent with the value of 0.060 ± 0.030 PgC yr⁻¹ for 2000–2007 estimated using ratios of soil carbon to vegetation biomass by Pan *et al.*¹⁸, but much higher than the value of 0.004 ± 0.015 PgC yr⁻¹ in the statistical

models of Piao *et al.*⁹ for 1996–2005. Since Piao's statistical models were only able to explain 23–29% of the observed forest lands soil carbon variations, and their result is one order of magnitude lower than those of this study and Pan *et al.*¹⁸, we do not adopt Piao's result in this study. We use the midpoint of the InTEC model and Pan *et al.*¹⁸ of $0.064 \pm 0.030 \text{ PgC yr}^{-1}$ as the rate of SOC accumulation in Chinese forests. For shrub lands, Piao *et al.*⁹ estimated the changes of SOC, using a statistical model, to be $0.039 \pm 0.009 \text{ PgC yr}^{-1}$ during 1982–1999, and using a process model, Tian *et al.*²⁰ reported that shrub SOC accumulated an average $0.012 \pm 0.005 \text{ PgC yr}^{-1}$ from 1981 to 2001. We use the average of these two studies of $0.026 \pm 0.019 \text{ PgC yr}^{-1}$ for this biome. For cropland and grassland SOC, we directly use the estimates of $0.021 \pm 0.004 \text{ PgC yr}^{-1}$ and $0.005 \pm 0.002 \text{ PgC yr}^{-1}$, respectively, reported by Huang *et al.*²¹. In Total, this gives a bottom-up SOC accumulation rate of $0.12 \pm 0.060 \text{ PgC yr}^{-1}$ during the 2000s. Therefore, the bottom-up estimate is a net carbon accumulation in land ecosystems of $0.29 \pm 0.12 \text{ PgC yr}^{-1}$ in China.

Consistent top-down and bottom-up estimates. The bottom-up estimate of carbon stock change in vegetation and soil is still much lower than the inversion results. That is because inland waters, ocean and wood products are also reservoirs for terrestrial carbon and the inverted CO_2 sink is also influenced by CO_2 from the oxidization of net imported products and reduced carbon compounds (RCC) emitted from fossil fuels and ecosystems. We then try to reconcile top-down and bottom-up results as follows:

$$\begin{aligned} \text{Land sink}_{\text{top-down}} &= \text{inverted } \text{CO}_2 \text{ sink} \\ &\quad - \text{fossil fuel RCC transferred to global atmosphere} \\ &\quad - \text{fossil fuel RCC deposited to land surface} \\ &\quad - \text{biogenic RCC transferred to global atmosphere} + \text{net import} \end{aligned} \quad (1)$$

$$\begin{aligned} \text{Land sink}_{\text{bottom-up}} &= \text{carbon stock change} + \text{accumulation in products} \\ &\quad + \text{burial in aquatic sediments} + \text{delivery to ocean} \end{aligned} \quad (2)$$

$$\begin{aligned} \text{NEP} &= \text{land sink} + \text{biomass burning} + \text{CO}_2 \text{ outgassing} + \text{biogenic RCC emission} \\ &\quad - \text{biogenic RCC deposition} \end{aligned} \quad (3)$$

where biogenic RCC includes RCC from biomass burning and biogenic volatile organic carbon emissions. Net ecosystem productivity (NEP) is the difference between photosynthesis and respiration in terrestrial ecosystems, which is also estimated from the top-down and bottom-up results to provide a full picture of the carbon cycle in China. All items in the above equations are positive. Equations (2) and (3) are straightforward to understand, and details of equation (1) are given in the Supplementary Material.

Fossil fuel emission inventories, i.e., CDIAC, are based on CO_2 emission factors that include direct emissions of CO_2 from fossil fuels and emissions of RCC, e.g., carbon monoxide (CO), methane (CH_4) and non-methane volatile organic carbons (NMVOCs) that are later oxidized into CO_2 ²². When the total fossil fuel emission is treated as all CO_2 emission, as done in most inversion studies including ours, the contribution of the emission to the regional near surface CO_2 concentration is overestimated. That is because, after emission to the atmosphere, NMVOCs is first oxidized to CO, which is subsequently oxidized to CO_2 . The NMVOCs oxidation process is typically fast (hours), while the CO oxidation process is rather slow (1–2 months). CH_4 is also oxidized to CO_2 at a very slow rate. Generally, these oxidation processes will occur during the air mass transport, and therefore non- CO_2 carbon species emitted from one region (e.g., China) will transform into CO_2 globally rather than locally. Hence, the treatment of non- CO_2 carbon species as direct CO_2 emission in inversions will tend to overestimate the contribution of fossil fuel emission to CO_2 concentration over China in inversions, causing overestimation of the inverted carbon sink, i.e. too high sinks needed to offset CO_2 gradients due to too high emissions²³. During 2006–2009, China emitted RCC at a rate of $0.102 \pm 0.007 \text{ PgC yr}^{-1}$, including $0.072 \pm 0.005 \text{ PgC yr}^{-1}$ of CO, $0.019 \pm 0.001 \text{ PgC yr}^{-1}$ of NMVOCs and $0.011 \pm 0.001 \text{ PgC yr}^{-1}$ of CH_4 on average, roughly 14% of these emissions were converted to CO_2 in the boundary layer over China, 12% were deposited to the land surface, and the remaining 74% were transported to the global atmosphere²⁴. Therefore, the “fossil fuel RCC transferred to global atmosphere” term in Eq. 1 is $0.076 \pm 0.0050 \text{ PgC yr}^{-1}$, and the “fossil fuel RCC deposited to land” term in Eq. 1 is $0.012 \pm 0.0010 \text{ PgC yr}^{-1}$.

The Global Fire Emission Database (GFED) biomass burning emission dataset, explicitly separates CO_2 , CO, CH_4 and NMVOC emissions. Based on GFED v3.1²⁵, emissions of CO_2 and RCC from biomass burning are 0.016 and $0.0020 \text{ PgC yr}^{-1}$, respectively over China. In addition, land ecosystems also directly release biogenic RCC, including NMVOC and CH_4 . Their emissions over China are estimated to be 0.021 ± 0.010 and $0.027 \pm 0.013 \text{ PgC yr}^{-1}$, respectively. The “biogenic RCC emission” term in Eq. 3 is $0.050 \pm 0.024 \text{ PgC yr}^{-1}$, and the “biogenic RCC deposition” term in Eq. 3 is $0.0060 \pm 0.0020 \text{ PgC yr}^{-1}$, taken as 12% of the sum. The “biogenic RCC transferred to global atmosphere” term in Eq. 1 is $0.037 \pm 0.018 \text{ PgC yr}^{-1}$, which is 74% of the biogenic RCC emission. It is a negative adjustment to the top-down land sink estimate because 74% of the biogenic RCC (carbon source) is lost to the global atmosphere and not captured by the inversion²⁴.

The net imports of forest and crop products from outside China, which are decomposed in China and become additional sources of carbon to the atmosphere are included in the top-down sink estimates, but should be subtracted from it to be compared with the bottom-up ecosystem carbon stock change estimate, which does not count wood and crop products stocks. Moreover, carbon accumulated in forest products is a net accumulation of carbon that should be included in the bottom-up estimate²⁶. Based on the Food and Agriculture Organization




Figure 2. Terrestrial carbon balance in China. Atmospheric signal A: inversion result (exclude CO₂ emissions from fossil fuels) constrained with global CO₂ datasets only; A+: result of additional constraint with China Meteorological Administration (CMA)’s measurements (3 sites); A++: result of further constraint with CONTRAIL aircraft CO₂ measurements; A1: result after considering the fossil fuels RCC emission and transformation; A2: result after correcting for the biogenic RCC lateral transport; A3: result after correcting for the net import through international trade. Land signal: carbon accumulated in China’s ecosystems; L1, result after considering the accumulation in harvested wood products; L2: result after considering the carbon burial and transport to ocean; A4, L3: result after considering the CO₂ outgassing from inland waters; A5, L4: result after correcting for the biomass burning CO₂ emission; A6, L5: result after correcting for the net RCC emission from ecosystems. The lower dotted blue curve indicates adjustments made to atmospheric inversion results without the added data. From land sink to NEP estimates, there are three major adjustments.

of the United Nations (FAO) statistical databases²⁷, we estimate that 0.012 PgC of wood and 0.019 PgC of food were imported into China every year during 2000s. The net imported food is assumed to be fully consumed and oxidized to CO₂ in the same year, while only a portion of the net imported wood products is fully oxidized. The remainder goes into long-term products and is slowly oxidized over time. Using the method of Winjum *et al.*²⁸, we calculate that during 2006–2009, the CO₂ emission due to net wood import is 0.006 PgC yr⁻¹, and the net accumulation of carbon in wood products made by local harvests in China during 2000s is 0.005 PgC yr⁻¹. Therefore, the “net import” term in Eq. 1 is 0.025 PgC yr⁻¹, and the “accumulation in products” term in Eq. 2 is 0.0050 PgC yr⁻¹.

Inland aquatic systems are now considered as a significant component of land-atmosphere CO₂ fluxes^{29,30,31}. Globally, about 2.1 PgC yr⁻¹ of carbon are transported from terrestrial landscape to inland waters, in which 1.7 PgC yr⁻¹ is from soil erosion and 0.4 PgC yr⁻¹ is from rock weathering³². In this study, we estimate that inland waters of China annually receive 0.12 ± 0.06 PgC yr⁻¹ of carbon from land, in which 0.105 ± 0.052 PgC yr⁻¹ is from eroded soils and 0.015 ± 0.008 PgC yr⁻¹ is from rock weathering (half of the exported dissolved inorganic carbon (DIC)²⁴). During the inland water carbon transport processes, about 0.020 ± 0.010 PgC yr⁻¹ of carbon is buried in aquatic sediments, and 0.062 ± 0.030 PgC yr⁻¹ is returned to the atmosphere, and the remainder of 0.038 ± 0.019 PgC yr⁻¹ is delivered to the coastal ocean, including 0.008 ± 0.004 PgC yr⁻¹ of total organic carbon (TOC) and 0.030 ± 0.015 PgC yr⁻¹ of DIC. The amounts of carbon that are transported to ocean, buried in the sediment and outgassed to the atmosphere in China are estimated to be all lower than those in Europe, but the relative fractions among burial, outgassing, and transport to the ocean are close to those in Europe²⁴. Therefore, the “burial in aquatic sediments” term in Eq. 2 is 0.020 ± 0.010 PgC yr⁻¹, the “delivery to ocean” term in Eq. 2 is 0.023 ± 0.010 PgC yr⁻¹ (i.e., TOC + DIC/2), and the “CO₂ outgassing” term in Eq. 3 is 0.062 ± 0.030 PgC yr⁻¹.

By including the various fluxes outlined above (see details in the Supplementary Material), the land sink estimates by the top-down and bottom-up methods are adjusted to 0.35 ± 0.23 (mean range 0.29–0.41) PgC yr⁻¹ and 0.33 ± 0.14 PgC yr⁻¹, respectively, and the corresponding NEP estimates are 0.47 ± 0.28 PgC yr⁻¹ and 0.45 ± 0.19 PgC yr⁻¹ (Fig. 2).

Discussion

The top-down and bottom-up estimates are consistent within their respective uncertainties. A full picture of the carbon cycle of China is shown in Fig. 3. However, considerable sources of systematic uncertainties still exist in these estimates, and the real uncertainties of both top-down and bottom-up estimates are likely higher than those calculated above.

The top-down results for south and southwest China are very uncertain (Supplementary Fig. S3, Fig. S4), although the results for eastern and northern China from different inversion systems are consistent within their uncertainties. Generally, significant and spatially explicit constraints on fluxes can be obtained in locations near and immediately upwind of surface measurements³³. In south and southwest China there are no local surface CO₂

Administration (CAMS/CMA)⁵, and aircraft CO₂ measurements from Nov 2005 to Dec 2009 over Eurasian by the Comprehensive Observation Network for Trace gases by AirLiner (CONTRAIL) project⁶ are used in both systems. The three CAMS/CMA sites are all regional background stations, which are located in Northeast China (LFS), North China (SDZ), and East China (LAN), and with altitudes of 330, 293 and 139 m, respectively. The air intake height is 10 m above ground level for all three sites. The measurements in these stations are sampled and analyzed using the recommended methods of WMO/GAW, and the accuracy is comparable with that of NOAA/ESRL⁵.

Simulation for the soil carbon fluxes over forest land. The Integrated Terrestrial Ecosystem C-budget (InTEC) model¹⁹, which is a regional C-budget model, is used to simulate the soil carbon fluxes over forest land. It combines the CENTURY model for soil C and nutrient dynamics and Farquhar's leaf biochemical model for canopy-level annual photosynthesis implemented using a temporal and spatial scaling scheme. In this study, the InTEC model is run from 1901 to 2012. The simulation region covers the whole China, with a horizontal resolution of 1 km × 1 km. LAI, NPP, forest cover and stand age data in 2005; climate data during 1901–2012, nitrogen deposition data during 1901–2010, soil data, and CO₂ data during 1901–2012 were used to driving the InTEC model. No forest management was considered. Forest disturbance was considered according to the stand age. The simulation results for 2006 to 2009 are used in this study.

RCC emissions in China. The Asian anthropogenic emission inventory for 2006 for the NASA INTEX-B Mission, and the Multi-resolution Emission Inventory for China (MEIC) for 2008 and 2010 are used to calculate the fossil fuel RCC emissions in China. The Global Fire Emission Data (GFED) V3.1 is used to calculate the biomass burning RCC emissions. The biogenic NMVOCs emissions are adopted from literature review. The CH₄ emissions is from a top-down estimate by Klinger *et al.*³⁶.

Carbon transport by rivers. The carbon delivered to the ocean through rivers include dissolve organic carbon (DOC), dissolve inorganic carbon (DIC) and particulate organic carbon (POC). Nine Chinese exorheic rivers are considered, including the Yangtze River, Yellow River, Pearl River, Huai River, Hai River, Liao River, Songhua River, Qiantang River and Min River. For the Yangtze River, Yellow River and Pearl River, we use the observations by respectively Wu *et al.*³⁷, Ran *et al.*³⁸, and Zhang *et al.*³⁹. For the Hai and Liao rivers, we use the observations by Xia and Zhang⁴⁰, and for the other 4 rivers there are no observations available to date. We use several simple methods to estimate the transport: for DOC, we use the mean concentration of Yellow and Yangtze River, for DIC, we use the mean DIC/DOC ratio observed in the five rivers, and for POC we use an empirical formula⁴¹.

The CO₂ outgassing from inland waters in China is calculated based on limited observations of CO₂ outgassing rates in the past decade and the water surface area is reported by the National Bureau of Statistics in China. For rivers and steams, the observations in Pearl River, Yangtze River and Yellow River are used^{42–44}; for reservoirs, the observations in five reservoirs in Yangtze River are used⁴⁵; and for natural lakes, the average of global natural lakes⁴⁶ is adopted directly.

The carbon burial in lakes and reservoirs are estimated using the data reported by Gui *et al.*⁴⁷ and Dong *et al.*⁴⁸, which both covered lakes in the middle and lower reaches of the Yangtze River Basin. The mean rate of these two studies is about two times the global mean rate. For the reservoirs, due to lack of observations, we assume that the carbon burial rate in Chinese reservoirs is also about two times of the global mean rate.

Trade of food and wood. The import and export data of food and wood products from the FAO statistical databases²⁷ are used. The food products include cereals, roots, sugar, soybeans and pulses, oil crops, vegetables, fruits, coffee and teas. The wood products include sawn wood, wood-based panels, paper and paperboard, recovered paper, other industry roundwood, and wood fuel and charcoal. Every year, the food products and the wood products of wood fuel and charcoal are assumed to be totally consumed and oxidized to CO₂, while the other wood products are partially oxidized and partially go into uses or long-term storage, the CO₂ release by these products are calculated using the method of Winjum *et al.*²⁸.

Carbon accumulated in wood products. This carbon is calculated using the local production data reported in the FAO statistical databases and the method of Winjum *et al.*²⁸.

References

- Peters, W. *et al.* An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker. *PNAS* **104**, 18925–18930 (2007).
- Peylin, P. *et al.* Global atmospheric carbon budget: results from an ensemble of atmospheric CO₂ inversions. *Biogeosciences* **10**, 6699–6720 (2013).
- Gurney, K. R. *et al.* Towards robust regional estimates of CO₂ sources and sinks using atmospheric transport models. *Nature* **415**, 626–630 (2002).
- GLOBALVIEW-CO₂. Cooperative Atmospheric Data Integration Project—Carbon Dioxide. NOAA ESRL, Boulder, Colorado. Data product accessed at <http://dx.doi.org/10.3334/OBSPACK/1002> (2013).
- Liu, L. *et al.* The characteristics of atmospheric CO₂ concentration variation of the four national background stations in China. *Sci. China Ser. D* **52**, 1857–1863 (2009).
- Machida, T. *et al.* Worldwide measurements of atmospheric CO₂ and other trace gas species using commercial airlines. *J Atmos Oceanic Technol* **25**, 1744–1754 (2008).
- Jiang, F. *et al.* Nested Atmospheric Inversion for the Terrestrial Carbon Sources and Sinks in China. *Biogeosciences* **10**, 5311–5324 (2013).
- Zhang, H. F. *et al.* Net terrestrial CO₂ exchange over China during 2001–2010 estimated with an ensemble data assimilation system for atmospheric CO₂. *J Geophys Res* **119**, 3500–3515 (2014).
- Piao, S. L. *et al.* The carbon balance of terrestrial ecosystems in China. *Nature* **458**, 1009–1013 (2009).

10. Rayner, P. J. *et al.* Two decades of terrestrial carbon fluxes from a carbon cycle data assimilation system (CCDAS). *Global Biogeochem Cycles* **19**, GB2026 (2005).
11. Peters, W. *et al.* Seven years of recent European net terrestrial carbon dioxide exchange constrained by atmospheric observations. *Global Change Biol* **16**, 1317–1337 (2010).
12. Zhang, S. P. *et al.* Global carbon assimilation system using a local ensemble Kalman filter with multiple ecosystem models. *J. Geophys. Res. Biogeosci* **119**, 2171–2187 (2014).
13. Guan, D., Liu, Z., Geng, Y., Lindner, S. & Hubacek, K. The gigatonne gap in China's carbon dioxide inventories. *Nature Clim Change* **2**, 672–675 (2012).
14. Liu, Z. *et al.* Reduced carbon emission estimates from fossil fuel combustion and cement production in China. *Nature* **524**, 335–338 (2015).
15. Boden, T. A., Marland, G. & Andres, R. J. Global, Regional, and National Fossil-Fuel CO₂ Emissions. doi: 10.3334/CDIAC/00001_V2010 (2014).
16. Zhang, C. H. *et al.* China's forest biomass carbon sink based on seven inventories from 1973 to 2008. *Climatic Change* **118**, 933–948 (2013).
17. Guo, Z. D. *et al.* Spatio-temporal changes in biomass carbon sinks in China's forests during 1977–2008. *Sci China Life Sci* **56**, 661–671 (2013).
18. Pan, Y. *et al.* A large and persistent carbon sink in the world's forests. *Science* **333**, 988–993 (2011).
19. Chen, W., Chen, J. & Cihlar, J. An integrated terrestrial ecosystem C-budget model based on changes in disturbance, climate, and atmospheric chemistry. *Ecological Modelling* **135**, 55–79 (2000).
20. Tian, H. *et al.* China's terrestrial carbon balance: Contributions from multiple global change factors. *Global Biogeochem Cycles* **25**, GB1007 (2011).
21. Huang, Y., Sun, W. J., Zhang, W. & Yu, Y. Q. Changes in soil organic carbon of terrestrial ecosystems in China: A mini-review. *Sci China Life Sci* **53**, 766–775 (2010).
22. Nassar, R. *et al.* Modeling global atmospheric CO₂ with improved emission inventories and CO₂ production from the oxidation of other carbon species. *Geosci Model Dev* **3**, 689–716 (2010).
23. Suntharalingam, P. *et al.* Influence of reduced carbon emissions and oxidation on the distribution of atmospheric CO₂: Implications for inversion analyses. *Global Biogeochemical Cycles* **19**, GB4003 (2005).
24. Ciais, P. *et al.* The impact of lateral carbon fluxes on the European carbon balance. *Biogeosciences* **5**, 1259–1271 (2008).
25. Randerson, J. T., van der Werf, G. R., Giglio, L., Collatz, G. J. & Kasibhatla, P. S. Global Fire Emissions Database, Version 3 (GFEDv3.1). Data set. Available on-line <<http://daac.ornl.gov/>> from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA. doi: 10.3334/ORNLDAAAC/1191 (2013) (Date of access: 11/08/2014).
26. Pacala, S. W. *et al.* Consistent land- and atmosphere-based US carbon sink estimates. *Science* **292**, 2316–2320 (2001).
27. Food and Agriculture Organization of the United States, FAO: Statistics. <<http://faostat3.fao.org>>, (2013) (Date of access: 25/08/2014).
28. Winjum, J. K., Brown, S. & Schlamadinger, B. Forest harvests and wood products: sources and sinks of atmospheric carbon dioxide. *Forest Science* **44**, 272–284 (1998).
29. Cole, J. J. *et al.* Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget. *Ecosystems* **10**, 171–184 (2007).
30. Regnier, P. *et al.* Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nature Geosci.* **6**, 597–607 (2013).
31. Raymond, P. A. *et al.* Global carbon dioxide emissions from inland waters. *Nature* **503**, 355–359 (2013).
32. Ciais, P. *et al.* Carbon and Other Biogeochemical Cycles. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA (2013).
33. Wells, K. C. *et al.* Simulation of atmospheric N₂O with GEOS-Chem and its adjoint: evaluation of observational constraints. *Geosci. Model Dev.* **8**, 3179–3198 (2015).
34. Zhang, H. *et al.* Inclusion of soil carbon lateral movement alters terrestrial carbon budget in China. *Scientific Reports* **4**, 7247, doi: 10.1038/srep07247 (2014).
35. Le Quéré, C. *et al.* Global carbon budget 2013. *Earth Syst Sci Data* **6**, 235–263 (2014).
36. Kirschke, S. *et al.* Three decades of global methane sources and sinks. *Nature Geosci* **6**, 813–823 (2013).
37. Wu, Y. *et al.* Sources and distribution of carbon within the Yangtze River system. *Estuar Coast Shelf Sci* **71**, 13–25 (2007).
38. Ran, L. S. *et al.* Spatial and seasonal variability of organic carbon transport in the Yellow River, China. *Journal of Hydrology* **498**, 76–88 (2013).
39. Zhang, L. K., Qin, X. Q., Yang, H., Huang, Q. B. & Liu, P. Y. Transported Fluxes of the Riverine Carbon and Seasonal Variation in Pearl River Basin. *Environmental Science* **34**, 3025–3034 (2013).
40. Xia, B. & Zhang, L. J. Carbon distribution and fluxes of 16 rivers discharging into the Bohai Sea in summer. *Acta Oceanologica Sinica* **30**, 43–54 (2011).
41. Ludwig, W., Probst, J. L. & Kempe, S. Predicting the oceanic input of organic carbon by continental erosion. *Global Biogeochemical Cycles* **10**, 23–41 (1996).
42. Wang, F. S., Wang, Y. C., Zhang, J., Xu, H. & Wei, X. G. Human impact on the historical change of CO₂ degassing flux in River Changjiang. *Geochemical Transactions* **8**, 7 (2007).
43. Li, S. *et al.* Daily CO₂ partial pressure and CO₂ outgassing in the upper Yangtze River basin: A case study of the Longchuan River, China. *Journal of Hydrology* **466–467**, 141–150 (2012).
44. Wang, F. S. *et al.* Carbon dioxide emission from surface water in cascade reservoirs-river system on the Maotiao River, southwest of China. *Atmospheric Environment* **45**, 3827–3834 (2011).
45. Yang, L. *et al.* Spatial and seasonal variability of CO₂ flux at the air-water interface of the Three Gorges Reservoir. *Journal of Environmental Sciences* **25**, 2229–2238 (2013).
46. St Louis, V. L., Kelly, C. A., Duchemin, E., Rudd, J. W. M. & Rosenberg, D. M. Reservoir surfaces as sources of greenhouse gases to the atmosphere: a global estimate. *Bioscience* **50**, 766–775 (2000).
47. Gui, Z. F., Xue, B., Yao, S. C., Wei, W. J. & Yi, S. Organic carbon burial in lake sediments in the middle and lower reaches of the Yangtze River Basin, China. *Hydrobiologia* **710**, 143–156 (2013).
48. Dong, X., Anderson, N. J., Yang, X., Chen, X. & Shen, J. Carbon burial by shallow lakes on the Yangtze floodplain and its relevance to regional carbon sequestration. *Global Change Biology* **18**, 2205–2217 (2012).

Acknowledgements

We kindly acknowledge all atmospheric data providers to the ObsPack version 1.0.2, including the NOAA Cooperative Air Sampling network and those that contribute their data to WDCGG. This work was supported by the National Key Basic Research Development Program of China (Grant No: 2010CB950704, 2010CB950601 and 2010CB833503), National Natural Science Foundation of China (Grant No: 41201194), and the Priority Academic Development Program of Jiangsu Higher Education Institutions. W. Peters was supported by an NWO

VIDI grant (864.08.012) and the Chinese-Dutch collaboration was funded by the China Exchange Program project (12CDP006).

Author Contributions

J.M.C. and F.J. designed research. L.Z. and L.L. provided additional CO₂ data in China. T.M., H.M. and Y.S. provided CONTRAIL CO₂ data. H.Z., W.P. and B.C. run CTC model, C.Z. and W.J. run INTEC model. F.J., J.M.C., P.C., W.P., W.J. and H.W. wrote the paper. All authors reviewed the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Jiang, F. *et al.* A comprehensive estimate of recent carbon sinks in China using both top-down and bottom-up approaches. *Sci. Rep.* **6**, 22130; doi: 10.1038/srep22130 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>